



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

August 28, 2024

Michael Barber
Regulatory Affairs Specialist
SBM Life Science
11000 Weston Parkway, Suite 150
Cary, NC 27513

Subject: Label Amendment - Registration Review Mitigation for Thiencarbazone-methyl & Metsulfuron-methyl
Product Name: Lawn 3FL2 Herbicide GR
EPA Registration Number: 92564-57
Application Dates: October 26, 2020; January 22, 2021; & January 5, 2022
Decision Numbers: 567275 & 569864

Dear Michael Barber:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Thiencarbazone-methyl & Metsulfuron-methyl Interim Decisions, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must

submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr by phone at (202) 566-0636, or via email at carr.caleb@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington", with a stylized flourish at the end.

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label


(Front Panel)

Lawn 3FL2 Herbicide GR

[ABN: Southern Season Long Weed Control for Lawns Granules]

[ABN: Southern Season Long Weed Control for Lawns Ready-to-Spread Granules]

■ Editorial Note - Marketing Claims Positioned Here



ACTIVE INGREDIENTS:

Metsulfuron-methyl -----	0.014%
Thiencarbazone-methyl -----	0.015%
Indaziflam -----	0.024%
Dicamba acid -----	0.102%
OTHER INGREDIENTS: -----	99.845%
TOTAL :	100.000%

[FERTILIZER ANALYSIS X-X-X]

Net Contents [8oz to 50 lb]

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements

Editorial Note - [Bracketed text] is optional language. Text in italics is editorial comment.


(Back Panel)

OPEN Resealable Label for Directions & Precautions **PRESS TO RESEAL**

[The days of multiple applications to kill weeds are over!] Just one application of Lawn 3FL2 Herbicide GR's [patented] [exclusive] [proprietary] (4-in-1) (3-in-1) (2-in-1) formula* kills [over 100] lawn weeds [plus] prevents new lawn weeds for up to 6 months [and feeds and strengthens your lawn!] [It delivers Season Long Weed Control that's simple and easy to use] [and it will not harm your lawn when used as directed]. [No need [to re-apply every couple of weeks] [for frequent re-application] [PLUS, when applied in Early Spring, prevents Crabgrass [Goosegrass] and when applied late summer, controls Annual Bluegrass]]. Ideal for large areas - treats 5,000 sq feet (15 lb size) [or X,XXX sq ft (for XX lb size)]. [Easy-to-Spread Granules]. [Rainproof in [1][4](24)[X] hour]. Controlling [controls] [listed] weeds in your lawn for up to 6 months [plus feeding and strengthening the lawn] [has never been easier]!

[*4-in-1 refers to 4 active ingredients; 3-in-1 refers to 3 herbicide modes of action; and 2-in-1 refers to pre-emergent and post-emergent weed control in one product.]



QUICK FACTS	
Editorial Note – Marketing Claims Positioned Here [see Appendix 2]	
KILLS & PREVENTS For up to 6 Months [PLUS, FEEDS]	Kills [over 100] broadleaf weeds [such as Chickweed, Dandelion, Clover & Plantain] Prevents grassy & broadleaf weeds [such as Crabgrass, Annual Bluegrass, Goosegrass, White Clover, Dandelion, Plantain, and Henbit] <i>[substitute other listed weeds as appropriate]</i> [Feeds and strengthens] the lawn. Contains slow release fertilizer for long lasting feeding.]
WHEN TO [APPLY] [USE]	[Apply] Anytime [in] [early] spring [or fall]
WHERE TO USE	ON ESTABLISHED LAWNS of the turf types listed below: Bermudagrass (except Ormond variety), Centipedegrass, St. Augustinegrass, and Zoysia DO NOT use on Ormond variety of Bermudagrass, Bluegrass, Fescue, Ryegrass, Bentgrass, Carpetgrass, Kikuyugrass, Seashore Paspalum, or Dichondra lawns.
COVERAGE	This bag treats 5,000 sq ft [15 lb size] [or X,XXX sq ft [for XX lb size]]
	For questions or comments, call toll-free 1-877-229-3924 or visit us on-line at www.bioadvanced.com

ACTIVE INGREDIENTS:

Metsulfuron-methyl -----	0.014%
Thiencarbazone-methyl -----	0.015%
Indaziflam -----	0.024%
Dicamba acid -----	0.102%
OTHER INGREDIENTS -----	99.845%
TOTAL:	100.000%

[FERTILIZER ANALYSIS X-X-X]

[Total Nitrogen (N).....X %]

[Available Phosphate (P₂O₅).....X %]

[Soluble Potash (K₂O)X %]

[Derived from commercially available fertilizer] *[Additional technical fertilizer information]*

EPA Reg. No. 92564-57

EPA Est. No.

DIRECTIONS FOR USE	
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.	
BEFORE YOU USE	<ul style="list-style-type: none"> • Use this product only on established, listed southern lawns - new lawns must be at least [3] [or X] months old if sodded and [16] [or X] months old if seeded or sprigged. • Do not apply this product to stressed, weakened or thin lawns, such as lawns that have drought conditions, soil compaction, sandy soils, excessive shade, or that have a history of dead spots early in the season, disease, or insect damage. Do not apply this product during spring 'green-up' while the grass is resuming normal growth after dormancy. This product may delay turf recovery or green-up. • Do not overlap treated areas or make repeat applications within 6 months as over application could damage the lawn. • Avoid application to non-lawn areas such as flower beds, ornamental beds, and gardens. Avoid application of this product near the roots of newly planted ornamentals. • Apply when the air is calm to prevent application to non-target sites. • Allow the lawn to recover between verticutting and using this product, or vice versa. This normally requires about 1 month
HOW TO USE	<ul style="list-style-type: none"> • Use rotary or drop type spreader to apply granules evenly over the lawn. Apply up to a maximum of 3 lbs <i>[or lower rate as specified to match label rate. See Spreader Settings table]</i> of this product per 1,000 sq ft. Do not apply by hand or with hand-held belly grinder applicators. • To be sure you cover lawn areas uniformly, first treat the border of the lawn then fill in the center. Make parallel passes, walking at a steady pace to ensure even distribution of the granules. Do not overlap treated areas. • After applying, activate the residual barrier by irrigating with 1/8" to 1/4" of water, unless rain is expected within 2 days. • Sweep or blow any accidentally applied material off driveways, sidewalks etc. and place back into spreader hopper or bag. <p>Do not make more than two applications per site in any 12 month period. Wait a minimum of 6 months between applications</p>

<p>[FOR BEST RESULTS]</p>	<p>[WHEN TO APPLY]</p> <ul style="list-style-type: none"> • [For best results in controlling existing broadleaf weeds, apply when <ul style="list-style-type: none"> - weeds are small and actively growing - temperatures are between 65°F - 85°F.] • [May be applied to wet or dry lawn. For best results, apply to dew-wetted lawn or water lawn lightly before application to allow product to adhere to weeds.] • [To minimize the potential for winter kill, do not apply this or other fertilizer-containing products to your lawn late in the season - use October 1 as a general guideline or consult your local extension office for the exact date specific to your area and turf type.] • [This product also prevents crabgrass & goosegrass <i>[other listed weeds as appropriate]</i> when applied early spring before the fourth mowing or before the temperatures are regularly in the 80's. This product will not kill existing crabgrass or goosegrass.] • [This product also prevents annual bluegrass (<i>Poa annua</i>) when applied August - November. Earlier applications provide better results. This product will not kill existing annual bluegrass.] • [Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product.] • [Lawns may be seeded [12] [or X] months and sodded [6] [orX] month after applying this product.] <p>[MOWING]</p> <ul style="list-style-type: none"> • [Tall grass should be mowed 1 to 2 days before applying.] • [Do not mow for 1 to 2 days after applying.] • [Clippings from the first three mowing's should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.] <p>[WATERING]</p> <ul style="list-style-type: none"> • [If lawn is dry, water 1 to 2 days before using this product] • [Do not water [for at least [X] [hours] [days] after application] [or] [if rain is expected within [X] (hours) (days)]
<p>HOW IT WORKS</p>	<p>This product contains a proven blend of weed killers to kill existing weeds plus a residual herbicide to keep [them] [weeds] from coming back for up to 6 months. Existing weeds cease growing within hours after application and are completely dead in 1 - 4 weeks [and new weeds will not grow for up to 6 months]. [And, it feeds and strengthens your lawn for quick green-up and long-lasting feeding].</p>

KILLS & PREVENTS

[Weeds presented here to be chosen from Appendix 1.]

[and many other weeds]

SPREADER SETTING

If your spreader is not listed, please call our Consumer Service Hotline toll-free (877) 229-3724 for help.

SPREADER	SETTING
	To deliver 2 lbs per 1,000 sq ft
Scott's SpeedvGreen	4.5
<ul style="list-style-type: none">• Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions.• Tip : 1000 sft ft = 20ft x 50ft• Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Be sure the spreader works properly and spreads granules evenly.	



Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Human and Domestic Animals

- Harmful if absorbed through skin.
- Avoid contact with skin, eyes, or clothing.
- Remove and wash contaminated clothing before reuse.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and waterproof gloves.

After using this product:

- Rinse gloves before removing and wash hands.
- If product gets on skin, wash thoroughly with soap & water.
- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet

+ FIRST AID

If on Skin or Clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 - 20 minutes.
- Call a poison control center or doctor for further treatment advice.

If Swallowed

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.


NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. **You may call toll-free 1-877-229-3763 for medical emergency information.** Active Ingredients: 0.014% Metsulfuron-methyl, (CAS # 74223-64-6), 0.015% Thien carbazon-methyl (CAS # 317815-83-1), 0.024% Indaziflam (CAS # 950782-86-2), and 0.102% Dicamba (CAS # 1918-00-9).

ENVIRONMENTAL HAZARDS

- This product is toxic to fish, aquatic invertebrates and plants. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean water mark. Do not contaminate water when disposing of rinsate or washwater. This product may impact water through spray drift or runoff. Follow directions for use to avoid spray drift and runoff. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams and springs will reduce the potential of this product entering water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.
- This pesticide may impact water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application
- **NON-TARGET ORGANISM ADVISORY:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.
- **SURFACE WATER ADVISORY:** This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of metsulfuron methyl and thien carbazon-methyl from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.
- **GROUNDWATER ADVISORY:** This product is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

- **WINDBLOWN SOIL PARTICLES:** Lawn 3FL2 Herbicide GR has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying Lawn 3FL2 Herbicide GR if prevailing local conditions may be expected to result in off-site movement.

{Note to PM: Product name Lawn 3FL2 Herbicide GR will be replaced with [this product] or ABN on the marketplace label }

STORAGE AND DISPOSAL	
Do not contaminate water, food or feed by storage and disposal	
Pesticide Storage	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
Pesticide Disposal and Container Handling 	<p>If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.</p> <p>If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.</p>

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

***Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.

PRODUCED FOR

**SBM Life Science Corp.
PO Box 525,
Morrisville, NC 27560**

Lawn 3FL2 Herbicide GR (MASTER) Approved 12/5/2023

Appendix 1

KILLS			
Ageratum, Tropic ² Amaranth, Palmer Aster, Heath ³ Aster, Slender Bahigrass ³ Barley, Mouse ² Barnyardgrass ¹ Beggarweed, Creeping Betony, Florida Bittercress ² Bittercress, Hairy Bluegrass, Annual (<i>Poa annua</i>) ¹ Brassbuttons, Southern ⁴ Brome, Foxtail ² BuckWheat, Wild Burclover (California Burclover) ¹ Burnweed, American (Fireweed) Burweed, Lawn (Spurweed) ¹ Carolina False Dandelion Carpetgrass Carpetweed (Indian Chickweed) Carrot, Wild ⁴ Chamberbitter Chamomile, False ⁵ Cheatgrass ² Chickweed, Common ¹ Chickweed, Mouseear ¹ Clover, Hop Clover, Rabbitfoot Clover, White ¹ Clover, White Sweet Crabgrass, Henry ² Crabgrass, Large (Hairy Crabgrass) ² Crabgrass, Smooth ² Cudweed, Linear-leaf (Purple Cudweed) ² Daisy, Oxeye Dallisgrass ³ Dandelion, Common ¹ Dandelion, Cat's Ear ¹ Dandelion, Virginia Dwarf Dichondra, Carolina (Dichondra) ³ Dock, Curly ¹	Dogfennel Dollarweed (Pennywort) Eciipta ² Evening Primrose, Common ² Evening Primrose, Cutlear Facelis (Trampweed) Fescue, Red Fescue, Tall Fieldbane Flatsedge, Rice ² Fleabane Fleabane, Hairy ² Foxtail, Giant ¹ Foxtail, Green ¹ Foxtail, Yellow ¹ Garlic, Wild (Field Garlic) Geranium, Carolina (Wild Geranium) ¹ Goosegrass (Silver Crabgrass) ² Groundsel, Common ² Guineagrass ² Hawksbeard, Asiatic Henbit ¹ Horseweed (Marestail) ¹ Horseweed, Sprawling Ivy, Ground (Creeping Charlie) Johnsongrass Khakiweed ³ Knawel Knotweed, Prostrate ¹ Kochia ² Kyllinga, Green ¹ Lady's Mantle Lambsquarters, Common ³ Lettuce, Wild (Tall Lettuce) Lcivegrass, Gophertail Lovegrass, Tufted ² Madder, Field Mallow, Little ²	Medic, Black ⁵ Millet, Browntop Millet, Common (Proso Millet) Morningglory, Entireleaf Morningglory, Ivyleaf Morningglory, Pitted Mustard, Black ⁵ Mustard, Tansy Mustard, White Mustard, Wild ¹ Nightshade, American Black ² Nightshade, Black Nightshade, Eastern Black Nightshade, Hairy ¹ Nutsedge, Annual ² Oat, Wild Onion, Wild Oxalis ⁵ Panicum, Fall Panicum, Texas Pansy, Field (Johnny Jump-up) ³ Parsley, Wild Parsley-piert Parthenium ² Paspalum, Thin (Bull Paspalum) ³ Pepperweed, Field Phyllanthus, Longstalked ² Pigweed, Prostrate ² Pigweed, Redroot ¹ Plantain, Blackseed Plantain, Bracted Plantain, Broadleaf (Common Plantain) Plantain, Buckhorn (Narrowleaf Plantain) ¹ Plantain, Paleseed ¹	Pokeberry Poorjoe ³ Puncturevine ² Purslane, Common ³ Purslane, Horse Pusley, Florida ² Quackgrass Ragweed, Common ⁴ Ragweed, Giant Rescuegrass ³ Rocket, Londons Rocket, Yellow (Winter Cress) Ryegrass, Italian ² Ryegrass, Perennial ² Sandbur, Field Sesbania, Hamps Shattercane Shepherd's Purse ¹ Sicklepod Sida, Prickly (Teaweed) ⁶ Signalgrass, Broadleaf Smartweed, Pennsylvania Sorrel, Heartwing Sorrel, Red Sowthistle, Annual ² Sowthistle, Spiny Speedwell, Corn ¹ Speedwell, Creeping Spurge, Prostrate ² Spurge, Spotted ² Stinkgrass Sunflower, Common ⁴ Swinecress ¹ Switchgrass Tasselflower ² Thistle, Canada ¹ Thistle, Russian Toadflax, Canada Velvetleaf ⁴ Venus' Looking-glass Vetch, Common Vetch, Purple ⁵ Violet, Field (Wild Pansy) · Waterhemp, Common Willowherb, Panicked ² Woodsorrel, Yellow ⁶

- 1 Kills existing weeds and prevents new weed seedlings from emerging
- 2 Prevents new weed seedlings from emerging
- 3 Suppresses existing weeds and prevents new weed seedlings from emerging
- 4 Kills existing weeds and suppresses new weed seedlings from emerging s Suppresses new weed seedlings from emerging
- 6 Suppresses existing weeds and suppresses new weed seedlings from emerging

Appendix 2

Optional Advertising Claims:

- New [Formulation]!
- [Brand] New [Formula]!
- Breakthrough Formulation
- Breakthrough Convenience
- [Exclusive] [Proprietary] [4-in-1] [2-in-1] Formula*
- Proprietary Formula that Kills Broadleaf Weeds
- Systemic [Herbicide] [Activity]
- Kills to the root.
 - Kills [Current] Weeds to the Root
 - Kills [Controls] even the tough weeds...to the root!
- Kills [##+] Weeds [to the root]
- Kills [Controls] [##] broadleaf weeds [to the root]
- Kills Dandelion, Clover, plus [##] Other listed Weeds
- Kills Over [##] Lawn Weeds in One Easy Step
- Kills Lawn Weeds in One Step
- Kills Broadleaf weeds
 - Kills Both Broadleaf and Grassy Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- [Effectively] Kills Both Broadleaf and Weed [Crab] Grass[es]
- Kills the weeds not the lawn* (when used as directed)
- Won't Harm your Lawn* (when used as directed)
- Won't Harm Lawns* (when used as directed)
- [Exclusive] [Unique] [Proprietary] 2-in-1 Product that Kills [Lawn] Weeds and [Stops] [Prevents] them [from] coming back
- [Exclusive] [Unique] [Proprietary] 3-in-1 Product: Kills Weeds, Prevents New Weeds [up to 6 Months] & Prevents Crabgrass
 - Season Long Weed Control (up to 6 months) (with 1 Application)
 - Season Long (up to 6 months) [Lawn] Weed & [Crab] Grass Control
 - Season Long (up to 6 months) [Lawn] Weed Control [Killer]
 - Season Long Crabgrass Control (up to 6 months)
 - 1 Application Provides Season Long (up to 6 Months) (Lawn) [Weed] Control
 - One application provides up to 6 months [weed] [crabgrass] control
 - [Up to] 6 Month Weed Control [with one application]
 - Kills Weeds [& Prevents New Weeds] [from Coming Back] (up to 6 months)
 - Kills [Current] [Existing] Weeds Up to 6 Months
 - Kills [& Prevents] Weeds Up to 6 Months
 - Keeps weeds out [of your lawn] up to 6 months
 - Prevents [New] Weeds up to 6 Months [with one application]
- Prevents [listed] Grasses Up to 6 Months
- Protects your lawn from weeds up to 6 months
- Tough weed protection [for up to 6 months]
- Weeds don't return [for up to 6 months]
- Weed Free Lawn [up to 6 Months]
- Extended [Weed] Control [Killer]
- Lawn Weed Control is Easier than Ever Before
- Simple, [Convenient], [Easy] to use
- PLUS Prevents Crabgrass!
 - Prevents Crabgrass All Season (up to 6 months)
 - Prevents Crabgrass (up to 6 months)
 - Crabgrass Free Lawn (Up to 6 Months)
 - Prevents Crabgrass, [Goosegrass] [and Annual Bluegrass (Poa annua)] (up to 6 months)
 - (Now) [Plus] Prevents [New] Weeds [Plus] [Crabgrass] [Grassy Weeds] [Poa annua] [Annual Bluegrass]

- [up to 6 months]
- Prevents Annual Bluegrass [*Poa annua*]
- Kills Dollarweed [and other common southern lawn weeds]
- Starts Working Immediately
- Visible Results in 24 Hours
- Rainproof in X Hour(s)
- All Weather Formula (rainproof in 1 hour)
- Guaranteed* not to harm lawns* (when used as directed)
- 100% Satisfaction Guaranteed* (or your money back)
- Easy-to-Spread Granules
- [15 lbs][30 lbs] [XX lbs] treats [up to] [5,000] [10,000] [X,XXX] sq ft
- Treats 5,000 [10,000] [X,XXX] sq ft
- XX% More [FREE] versus XX lb size
- Apply anytime spring or fall
- * (When used as directed. [DO NOT use on] [Ormond variety of Bermudagrass], [Bluegrass], [Fescue], [Bentgrass], [Carpetgrass], [Kikuyugrass], [Seashore Paspalum], [Ryegrass], or [Dichondra] lawns
- (*4-in-1 refers to 4 active ingredients; 3-in-1 refers to 3 herbicide modes of action; and 2-in-1 refers to pre-emergent and post-emergent weed control in one product.)
- Weatherproof Protection

Fertilizer Claims

- Builds thick, green turf from the roots up
- [Feeds and] Strengthens your lawn
- Builds strong roots
- Builds a thick lush green lawn .
- Even feeding for a thicker, greener lawn
- Slow Release Fertilizer/Feeding
- Long Lasting Feeding