



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

92647-25

Date of Issuance:

8/14/18

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Tigris Clop + Flumet

Name and Address of Registrant (include ZIP Code):

Tigris, LLC
204 Village Market Place #173
Morrisville, NC 27560

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Kathryn V. Montague, Product Manager 23
Herbicide Branch, Registration Division (7505P)

Date:

8/14/18

2. You are required to comply with the data requirements described in the DCI Order identified below:

- a. Clopyralid GDCI-117403-1454

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 92647-25.”
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 3/25/2018

If you have any questions, please contact Emily Schmid at 703-347-0189 or by email at schmid.emily@epa.gov.

Enclosure

ACCEPTED

08/14/2018

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 92647-25

[Note to reviewer: [Text] in brackets denotes optional or explanatory language
[Note to reviewer: {Text} in braces denotes where in the final label text will appear
{BOOKLET FRONT PANEL LANGUAGE}]

FLUMETSULAM	GROUP	2	HERBICIDE
CLOPYRALID	GROUP	4	HERBICIDE

Tigris Clop + Flumet [™]

[A selective herbicide for the control of broadleaf weeds in field corn.]

Active Ingredient: (% by weight)

flumetsulam: N-(2,6-difluorophenyl)-5- methyl-1,2,4-triazolo-[1,5a]-pyrimidine-2-sulfonamide 18.5%

clopyralid potassium salt: 3,6-dichloro-2-pyridinecarboxylic acid, potassium salt 60.0%

Other Ingredients:21.5%

Total 100.0%

Acid Equivalent: clopyralid: 3,6-dichloro-2-pyridinecarboxylic acid - 50%

Contains 0.185 lbs. of flumetsulam and 0.5 pound clopyralid acid equivalent per pound of product.

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

See inside label booklet for First Aid, Precautionary Statements and Directions for Use.

EPA Reg. No.: 92647-XX

EPA Est. No.:

Net Weight:

Manufactured For:

Tigris, LLC

1204 Village Market Place

#173

Morrisville, NC 27560

{LANGUAGE INSIDE BOOKLET}

FIRST AID	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

**For Chemical Emergency
Spill, Leak, Fire, Exposure, or Accident
Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

**Precautionary Statements
Hazards to Humans and Domestic Animals
WARNING**

Causes Substantial But Temporary Eye Injury. Harmful If Swallowed, Inhaled, Or Absorbed Through The Skin. May Cause Skin Sensitization Reactions In Certain Individuals.

Do not get in eyes or on clothing. Avoid breathing vapors or spray mist and contact with skin.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

The active ingredients in this product are known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this product where soils are permeable, particularly where the water table is shallow, may result in leaching to groundwater.

Caution should be exercised when handling this product at mixing and loading sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

NOT FOR SALE, USE OR DISTRIBUTION IN NASSAU AND SUFFOLK COUNTIES IN NEW YORK STATE.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Resistance Management

Tigris Clop + Flumet contains two active ingredients, clopyralid and flumetsulam. Clopyralid is classified in the pyridine carboxylic acid chemical class as a Group 4 herbicide, synthetic auxin. Flumetsulam is classified in the triazolopyrimidine chemical class as a Group 2 herbicide, Acetolactate Synthase (ALS) or Acetohydroxy Acid Synthase (AHAS) inhibitor. Any weed population may contain plants naturally resistant to Tigris Clop + Flumet and other Group 4 or 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed. Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of actions for each target weed. If levels of control provided by applications of this product is reduced and cannot be accounted for by factors such as misapplication, abnormal levels of target species or extremes of weather, it may be the case that target species have developed a strain resistant to applications of Tigris Clop + Flumet.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the
- dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

To minimize the occurrence of resistant biotypes, observe the following general weed management practices:

- Scout application site before and after herbicide applications.
- Start with a clean application site, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small.
- Add other herbicides (e.g. a selective and/or a residual herbicide) and cultural practices (e.g. tillage or crop rotation) where appropriate.
- Utilize the specified label rate for the most difficult to control weed in your field. Avoid tank mixtures with other herbicides that reduce this product's efficacy (through antagonism), or tank mixture directions that

encourage application rates of this product below the label directions.

- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from field to field to minimize the spread of weed seed or plant parts.
- Report any incidence of repeated non-performance of this product on a particular weed to local extension specialists, certified crop advisors, or your Tigris, LLC representative.

Product Information

Tigris Clop + Flumet is a selective herbicide for broadleaf weed control in field corn, including high oil and waxy varieties and field corn grown for ensilage. Tigris Clop + Flumet may be applied as a preplant surface, preplant incorporated, preemergence, or postemergence treatment. Soil surface treatments may be applied with water, liquid fertilizer, or impregnated on dry fertilizer. Apply postemergence treatments with water. Absorption of Tigris Clop + Flumet occurs from both shoot and root uptake. Susceptible weeds exposed to Tigris Clop + Flumet stop growing and either die or remain non-competitive with the crop. Tigris Clop + Flumet provides residual control of weeds that may emerge after application. Adequate soil moisture is necessary for optimal activation because uptake and translocation of Tigris Clop + Flumet involves uptake by emerging shoots and/or roots.

Use directions in Tigris, LLC supplemental labeling may supersede directions or limitations in this labeling.

Use Precautions

Handling Precautions if Product is Packaged in Water Soluble Packets: Do not remove water soluble packet from overpack except for immediate use. Do not allow water soluble packet to come into contact with water prior to use. Do not handle water soluble packet with wet hands or wet gloves. Carefully reseal package containing unopened water soluble packets and protect package from moisture.

Application Precautions

- Uneven application or uneven incorporation of Tigris Clop + Flumet can result in erratic weed control or crop injury. Over application may result in crop injury or rotational crop damage from soil residue.

Adverse Weather Conditions Precautions

- **Soil Application Only:** Extended cold, wet conditions (soil temperatures below 50°F and excessive rainfall with wet soil conditions), following soil application of Tigris Clop + Flumet to field corn, which persist during germination and/or early crop development may result in crop injury. Injury symptoms, which include yellowing of leaves and/or crop stunting, are usually temporary and affected corn plants usually recover without affecting yield.
- When applications are made under adverse (dry or cold) conditions or when large weeds or less susceptible species are treated, only weed suppression may be observed. Weed suppression is a visual reduction in weed competition (reduced population, size, and/or vigor) as compared to an untreated area. Degree of control can be increased by applying Tigris Clop + Flumet under favorable growing conditions (i.e., adequate moisture and temperature), and by using a higher rate in the specified rate range.
- Dry weather following preplant surface or preemergence applications of Tigris Clop + Flumet may reduce effectiveness. If sufficient activating rainfall or overhead irrigation does not occur within 7 to 10 days of application, rotary hoe, harrow, or shallowly cultivate to incorporate the herbicide lightly into the soil. Use a preplant incorporated application when a period of dry weather is predicted after application.
- Avoid application when air temperature is near freezing or when freezing conditions are expected for several days following application.
- Postemergence application of Tigris Clop + Flumet to corn that is stressed or damaged by conditions such as cold weather, hot weather (>90°F), hail, drought, water saturated soil, disease, or insects may cause crop injury.

Use Restrictions

- This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.
- Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad.

Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

- Do not apply this product through any type of irrigation system.
- Do not use flood irrigation to apply or incorporate this product.
- Product must be used in a manner that will prevent back-siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

Tank Mixing

Tigris Clop + Flumet may be tank mixed or followed by other overlay or postemergence treatments registered for use on corn to broaden the spectrum of weeds controlled. This product may be applied in tank mix combination with labeled rates of other products provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing is not prohibited by the label of the tank mix product; and (3) the tank mix combination is compatible as determined by a "jar test" described in the "Tank Mix Compatibility Testing" section below. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank Mixing Restrictions:

- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.
- Do not tank mix Tigris Clop + Flumet with Basagran herbicide, Laddock herbicide, or Lightning herbicide as severe crop injury may occur. (See instructions for Postemergence Treatments, Tank Mixing.)
- Do not exceed recommended application rates. Do not tank mix with another pesticide product that contains the same active ingredient as this product unless the label of either tank mix partner specifies the maximum dosages that may be used.
- For products packaged in water soluble packaging, do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment has been adequately cleaned. (See instructions for Sprayer Clean-Out.)

Tank Mix Compatibility Testing: A jar test is recommended prior to tank mixing to ensure compatibility of Tigris

Clop + Flumet and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination must not be used.

For Soil Applications of Tigris Clop + Flumet (Not Applicable to Postemergence Use)

Precautions:

- Use of Tigris Clop + Flumet in soil-applied treatments on soils with less than 1.5% organic matter (O.M.) may result in crop injury. Apply as a soil-treatment to fields which have less than 1.5% O.M. only if the risk of crop injury is acceptable.
- If any herbicide with ALS (acetolactate synthase) inhibition mode of action such as Pursuit, Preview, Canopy, Classic, Scepter, or Squadron herbicide, etc., was applied the previous year, apply Tigris Clop + Flumet to corn only if the rotational restrictions applicable to corn for the preceding product has been met.
- Corn growing in calcareous soils or soils with historically high salt content (soil test results for salinity indicating electrical conductivity greater than 1.0 mmho/cm) may exhibit chlorosis and/or stunting resulting from reduced availability of iron, zinc or other micro nutrients essential for normal crop vigor and growth. The presence of soil-active herbicides, such as Tigris Clop + Flumet may cause additional stress under these conditions resulting in increased leaf chlorosis and/or crop stunting. This added stress may retard crop recovery, especially under conditions of limited rainfall. In fields, which contain calcareous or high salt content soils, growers should plant "IR" or IMR" designated varieties, commonly referred to as "imidazolinone resistant" corn hybrids. On these type soils, the likelihood of crop injury can also be reduced by using the lower end of the recommended rate range for the soil type and/or by applying Tigris Clop + Flumet 10-14 days prior to planting.

Restrictions:

- Corn Planting Depth: Minimum planting depth must be at least 1 1/2 inches.
- Do not soil apply to peat or muck soils as reduced weed control will result. (May be used postemergence.)
- Do not apply to areas where the soil pH is greater than 7.8 as this may result in increased crop injury.
- Do not apply to a soil containing greater than 5% organic matter if the soil pH is below 5.9 as reduced weed control will result.

Soil Insecticide Advisories for Soil Applications of Tigris Clop + Flumet:

When Tigris Clop + Flumet is used for soil applied broadleaf weed control in corn:

- Soil applied organophosphate insecticides should be applied in a T-band or a band to avoid potential crop injury.
- Soil insecticides from other classes of chemistry may be applied in-furrow, T-banded, or banded.
- Terbufos (Counter insecticide products) or phorate (Thimet insecticide products) should not be used.

Soil Insecticide Advisories for Postemergence Applications of Tigris Clop + Flumet:

- Do not apply Tigris Clop + Flumet postemergence if corn was previously treated with Counter insecticide or Thimet insecticide as severe crop injury may result.
- Postemergence applications of Tigris Clop + Flumet to corn previously treated with T-band, band, or in-furrow applications of other organophosphate insecticides such as Lorsban® insecticide, Aztec, Fortress, or Dyfonate insecticides may cause temporary crop injury.

Foliar Insecticide Advisories for Postemergence Applications of Tigris Clop + Flumet

- Do not tank mix Tigris Clop + Flumet with foliar postemergence organophosphate insecticides as severe crop injury may result. To avoid crop injury, apply the foliar organophosphate insecticide treatment at least 10 days before or 10 days after the application of Tigris Clop + Flumet.
- Tigris Clop + Flumet may be tank mixed with non-organophosphate foliar insecticides provided they are

labeled for use with postemergence corn herbicides.

Use with other Products

- Corn previously treated with Tigris Clop + Flumet that is stressed or damaged by conditions such as cold weather, hail, drought, water saturated soil, disease, or insects should not be treated with Accent, Beacon, Permit, Exceed, or Basis herbicides, or other herbicides with ALS inhibition mode of action as this may cause further crop injury.
- Do not foliar apply Tigris Clop + Flumet to corn that exhibits herbicide injury from previous applications made to the current or preceding crop.

Use with Genetically Modified Corn Varieties

- If an "IR" or "IMR" designated hybrid (commonly referred to as "imidazolinone resistant") is planted, any organophosphate insecticide, including Counter or Thimet, can be applied according to label directions without increasing the likelihood of injury to corn from Tigris Clop + Flumet. The adverse interaction between Counter or Thimet insecticide and Tigris Clop + Flumet **does not** occur in corn hybrids identified as "IR" or "IMR". This adverse interaction **does** occur in imidazolinone tolerant "IT", "PT" hybrids which are considered as "standard" hybrids regarding this effect. "IR" or "IMR" hybrids may also be planted to reduce injury to corn from preemergence treatments of Tigris Clop + Flumet on soils with less than 1.5% organic matter or pH greater than 7.8.

Maximum Application Rate

- Do not exceed a total application rate of 6.0 oz per acre of Tigris Clop + Flumet (0.2 lb a.i. of clopyralid and 0.07 lb a.i. of flumetsulam) in a single crop year.
- Multiple applications of Tigris Clop + Flumet within a growing season can be made as a soil application followed by a postemergence application, or as multiple postemergence applications. Do not exceed the cumulative rate of 0.07 lb per acre active ingredient of flumetsulam per single crop year if a postemergence application of Tigris Clop + Flumet is made following a soil application of a flumetsulam-containing herbicide or with a postemergence herbicide containing flumetsulam (See table below to calculate cumulative flumetsulam amount per season.).
- Do not exceed a cumulative amount of 0.25 lb a.i. per acre of clopyralid per single crop year from all sources. (See table below to calculate cumulative clopyralid amount per season.)

Examples: 4.0 ounces of Tigris Clop + Flumet contains 0.047 lb flumetsulam and 0.125 lb clopyralid.
2.9 ounces of Accent Gold herbicide contains 0.035 lb flumetsulam and 0.094 lb clopyralid.

Herbicide	Unit of Measure	Flumetsulam (lb a.i./ unit of measure)	Clopyralid (Acid) (lb a.i./unit of measure)
Hornet	1 ounce	0.0145	0.039
Tigris Clop + Flumet	1 ounce	0.0116	0.031
Python™ WDG	1 ounce	0.05	---
Accent Gold	1 ounce	0.012	0.032
Stinger®	1 fluid ounce	---	0.023

The maximum active ingredient allowed per season in all states:

Flumetsulam = 0.07 lb/acre

Clopyralid = 0.25 lb/acre

NOT FOR SALE, USE OR DISTRIBUTION IN NASSAU AND SUFFOLK COUNTIES IN NEW YORK STATE.

Other Precautions

- **Hybrid Seed Production:** Corn inbred lines grown for hybrid seed production may be injured by Tigris Clop + Flumet. Inbred lines should be thoroughly tested for crop tolerance before treating large acreage. While growers are not prohibited from using Tigris Clop + Flumet on seed corn, **Tigris, LLC will not accept responsibility for any crop injury arising from the use of Tigris Clop + Flumet on field corn grown for seed.**

Other Restrictions

- **Crop Residues from Treated Areas:** Crop residues from treated areas cannot be used for composting or mulching on ground where susceptible crops may be grown the following season. To promote herbicide decomposition, plant material must be evenly incorporated or burned. Adequate moisture is also required to promote breakdown of plant residues, which contain clopyralid.
- **Do not move treated soil.** Avoid situations where soil particles may blow into areas where susceptible crops are grown. The hazard of movement of this product on dust is reduced if treated fields are irrigated or if rain occurs shortly after application.
- **Do not apply under conditions that favor runoff or wind erosion of soil containing Tigris Clop + Flumet to nontarget areas. To prevent off-site movement due to runoff or wind erosion:**
 - Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface must first be settled by rainfall or irrigation.
 - Do not apply to impervious substrates such as paved or highly compacted surfaces or frozen or snow covered ground.
 - Do not apply to soils when saturated with water.
 - Do not use tailwater from the first flood or furrow irrigation of treated fields to treat nontarget crops unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.
- **Do not apply when weather conditions favor drift to nontarget sites.** Spray drift of Tigris Clop + Flumet to emerged soybeans or soil to which soybeans will be planted during the same growing season may cause soybean injury.
- **Read and follow these Advisories to minimize drift to nontarget areas.**
 - Minimize drift by using sufficient spray volume to ensure adequate coverage with large-droplet size sprays.
 - Use low pressure application equipment capable of producing a large-droplet spray. Do not use nozzles that produce a fine-droplet spray. Droplet size has been shown to be the single most important factor affecting drift from ground applications.
 - While increasing droplet size does reduce the potential for spray drift, larger droplets do not eliminate drift if environmental or application conditions are inappropriate for application.
 - Use larger capacity nozzles to increase flow rate rather than increasing spray pressure.
 - Keep height of ground-driven spray booms as low as possible above the target to minimize exposure

to evaporation and wind while still providing good coverage. Applications made late in the growing season with excessive boom heights drastically increase the potential for spray drift.

- Do not apply when wind is gusting or wind speed exceeds 15 mph as uneven spray coverage and drift may result. Avoid application to border rows adjacent to susceptible crops such as soybeans, field peas, or sunflowers under windy conditions unless one of the following drift management steps is taken:

(1) application is made only when the wind direction is such that the susceptible crop is up-wind from the treatment area (wind blowing from the susceptible crop toward the treated crop); or

(2) the applicator leaves an adequate buffer zone between the treated crop and the susceptible crop and coarse or low drift nozzle configurations are used.

A drift control or deposition agent may be used with this product to aid in reducing spray drift due to wind when making applications adjacent to susceptible crops, but may not be effective after prolonged pumping of the spray mix.

- On calm days with little or no wind, check for temperature inversions before making herbicide applications. Temperature inversions occur under calm conditions with little or no wind and air temperature increases with increasing height above the ground. Inversion conditions may be indicated by a layer of fog or mist near the ground and, under clear conditions, may be detected by use of a smoke column. A temperature inversion is indicated when smoke does not rise in a column, but layers at some level above the ground. Do not apply herbicides if temperature inversion conditions exist in the treatment area.

- **Do not apply Tigris Clop + Flumet to sweet corn or popcorn.**
- **Preharvest interval (PHI):** An interval of at least 85 days is required between application of Tigris Clop + Flumet and field corn harvested for grain. If field corn is grown for forage or ensilage, application must occur before corn reaches 20 inches in height or V6 growth stage (whichever occurs first) and an interval of at least 45 days is required between application and harvest.
- **Do not aerially apply Tigris Clop + Flumet unless permitted by EPA approved supplemental labeling.**
- **Avoid all direct or indirect contact with nontarget plants.** Do not apply near desirable vegetation. Allow adequate distance between target area and desirable plants under conditions of application to minimize potential exposure.

Sprayer Cleanup

To avoid injury to or exposure of nontarget crops, thoroughly clean and drain spray equipment used to apply Tigris Clop + Flumet after use. Cleaning should occur as soon as possible after application of Tigris Clop + Flumet. Spray equipment should be cleaned after use with Tigris Clop + Flumet by the following procedure:

1. Drain any remaining Tigris Clop + Flumet from the spray tank and dispose of according to label disposal instructions.
2. Hose down the interior surfaces of the tank. Flush tank, hoses, boom, and nozzles with clean water for 10 minutes. Fill the tank with water and recirculate for 15 minutes. Spray part of the mixture through the hoses, boom, and nozzles and drain the tank. All rinse water must be disposed of in compliance with local, state, and federal guidelines.
3. Fill the tank with water and recirculate for 15 minutes. For optimum cleaning, a tank cleaner such as liquid ammonia (1 gallon per 100 gallons of water) or other commercial tank cleaner is recommended in the second rinse if the spray equipment will be used on crops other than field corn. Spray part of the mixture through the hoses, boom, and nozzles and drain the tank. All rinse water must be disposed of in compliance with local, state, and federal guidelines.
4. Remove the nozzles and screens and clean separately.
5. If the spray equipment will be used on crops other than field corn, repeat steps 1 and 2 again and thoroughly wash the spray mixture from the outside of spray tank and the boom.

Rotational Crop Restrictions

When tank mixing with companion herbicides, follow the most restrictive crop rotation guidelines on the label of each product used.

The following rotational crops may be planted after the indicated interval following application of rates up to 6.0 ounces per acre of Tigris Clop + Flumet:

Numbers within parentheses (-) in table refer to Specific Rotational Crop Requirements below.

Rotational Crop	Interval (Months)
barley, oats, rye, wheat	4
alfalfa (1), dry beans (1), forage grasses (2), lima beans (1), popcorn, rice, seeding of cover crops (3), soybean (1)	10.5
grain, sorghum	12
peas (1, 4), snap beans (1, 4)	18
cotton, peanuts, potatoes, sunflower, sweet corn (5), tobacco	18
sugar beets, canola and all other crops (6)	26

Specific Rotational Crop Requirements:

1. When annual rainfall and/or irrigation is less than 15 inches on soils with less than 2% organic matter, alfalfa, dry beans, lima beans, peas, snap beans, and soybeans must not be planted until 18 months after treatment.
2. Excludes forage grasses grown for commercial seed production.
3. The following cover crops may be planted for establishment of Federal Conservation Reserve Programs and Agricultural Reserve Programs no sooner than 10.5 months following application of Tigris Clop + Flumet at rates up to 4.0 oz per acre: legumes including alfalfa, clovers, crownvetch, birdsfoot trefoil, and lespedeza; and grasses, including big bluestem, little bluestem, switchgrass, Russian wildrye, green needle, smooth bromegrass, Garrison creeping foxtail, canary grass, orchardgrass, intermediate wheatgrass, tall wheatgrass, crested wheatgrass, western wheatgrass and indian grass. Some stand reduction or temporary stunting of legume seedlings is possible. However, Tigris, LLC will not accept responsibility for any crop injury or stand failure in crops established under Federal Conservation Reserve Programs and Agricultural Reserve Programs following use in corn and the subsequent 10.5 month rotational crop restriction. Additionally, Tigris, LLC will not accept responsibility for any crop injury or stand failure of native grasses as a result of inadequate seedbed preparation, erratic germination, lack of seedling vigor, or plant stress from unfavorable environmental conditions.
4. An 18-month crop rotation is required following application of Tigris Clop + Flumet at rates greater than 4 oz per acre. Peas and snap beans may be planted 10.5 months following application of Tigris Clop + Flumet at rates up to 4 oz per acre.
5. Certain sweet corn varieties may be planted 10.5 months following soil or postemergence application of up to 4.0 oz per acre of Tigris Clop + Flumet. This interval applies only to the following varieties of sweet corn grown for processing: Bonus, Challenger, Chase, Cornucopia, Crisp'N Sweet 710, Crisp'N Sweet 710A, DMC 20-04, DMC 20-35, Eliminator, Empire, Excalibur, Excellency, GH 0937, GH 2547, GH 2628, GH 2690, GG 5, GG 8, GG 22, GG 23, GG 40, GG 43, GG 46, GG 55, GG 246, GG 255, GG 256, GG 539, HM 701, 781 Ultra, Lumina, Reward, Sheba, Spirit, Sprint, Viking, and Zenith. The rotational interval is 18 months for other sweet corn varieties not listed here, except as provided in updated listings of sweet corn varieties tolerant to this product.
6. Rotation to sugar beets, canola, and all other crops requires a 26-month rotation interval and a successful

field bioassay.

Field Bioassay Instructions: In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample field conditions such as soil texture, soil pH, drainage, and any other variable that could affect the seed bed of the new crop. Field bioassay at any time between harvest of the treated crop and the planting of the rotational crop. Observe the test crop for herbicidal activity, such as poor stand (effect on seed germination) chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the test rotational crop; plant only a labeled crop or crop listed in the table above for which the rotational interval has clearly been met.

Mixing and Application

Spray Volume

Apply Tigris Clop + Flumet in sufficient spray volume to provide uniform coverage using properly calibrated ground equipment. Apply in a total spray volume of 10 to 60 gallons per acre using low pressure (20-40 lb/sq in). Maintain sufficient agitation during mixing and spraying to ensure a uniform spray mixture. More thorough coverage is possible when making soil applications to minimum or no-till corn by using a total spray volume of 20 or more gallons per acre.

Tigris Clop + Flumet	Acres per Package	
	Acres Per 6 lb Plastic Jug †	Acres per 6 oz Packet (Contains four 1.5 oz units) ^{††}
2.0	48	3.0
3.0	32	2.0
4.0	24	1.5
5.0	19.2	1.2

† If the number of acres to be treated results in the use of a partial container, use the measuring device provided with the container to measure out product according to the scale indicated on the measuring device.

^{††}To calculate the number of 6 oz water soluble packets for your spray mix:

1. Determine the number of acres you wish to spray in the desired application.
2. Divide the number of acres by the acres per packet that falls within the rate range you have chosen. See the above table for broadcast application rates and corresponding acres per packet.
3. The result is the number of packets you are required to add to the spray mix.

If the resulting number of packets is not a whole packet:

1. Do not open the water soluble packets.
2. Round up or down to the nearest whole number of packets and check to make sure that the resulting number of acres per packet falls within the desired rate range for the application.

Sample Calculations:

1. Planned application = 3.0 oz per acre (The acres per packet = 2.0).
2. Assuming 17 acres is to be treated, 17 acres divided by 2.0 acres per packet = 8.5 packets (Round up to 9 packets).

Band Application

Calculate the amount of herbicide needed for band treatment by the formula:

$$\begin{array}{r} \text{Band width in inches} \\ \text{-----} \\ \text{Row width in inches} \end{array} \times \text{Broadcast rate} = \text{Amount needed}$$

per acre per acre of field

Mixing Directions

Tigris Clop + Flumet is a water dispersible granule formulation. Thorough mixing is required.

1. Fill the tank with 1/2 of the total amount of water or liquid fertilizer required for the load.
2. Start agitation system.
3. Add the required amount of Tigris Clop + Flumet directly into the spray tank while agitating. If product is packaged in water soluble packets, open the overpack and add the required number of water soluble packets directly to the spray tank while agitating. (For use of water soluble packaging in liquid fertilizer solutions, see the "Application in Liquid Fertilizer" section of this label for special pre-mixing instructions.) Do not open water soluble packets. Water soluble packets will float on the surface until the water soluble film dissolves and releases the product. Handling packets with hands should be minimized.
4. Continue agitation and complete filling the tank while product disperses in the spray tank solution.

Before spraying make sure Tigris Clop + Flumet is thoroughly mixed in the solution. If product is in water soluble packets, make sure packets have completely disintegrated and product is thoroughly mixed with water. Depending on the water temperature and the degree of agitation, the packet and Tigris Clop + Flumet should be completely dispersed within 5 minutes from the time they were added to the water.

To insure a uniform spray mixture continuous agitation is required during mixing and spraying. Apply within 24 hours after mixing. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying.

Tigris Clop + Flumet in Tank Mix

Tigris Clop + Flumet may be applied in tank mix combination with labeled rates of other products provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing with Tigris Clop + Flumet is not prohibited by the label of the tank mix product. See "Tank Mixing" in "Product Use Precautions" section.

Vigorous, continuous agitation during mixing, filling, and throughout application is required for all tank mixes. Sparger pipe agitators generally provide the most effective agitation in spray tanks. To prevent foaming in the spray tank, avoid stirring or splashing air into the spray mixture.

Mixing Order for Tank Mixes: Fill the spray tank to 1/4 to 1/3 of the total spray volume required with water or liquid fertilizer solution. Start agitation. Add different formulation types in the order indicated below, allowing time for complete mixing and dispersion after addition of each product. Allow extra mixing and dispersion time for dry flowable products.

Add different formulation types in the following order: Tigris Clop + Flumet (slurried if mixing water soluble packets with liquid fertilizer) and other dry flowables; wettable powders; aqueous suspensions, and flowables. Maintain agitation and fill spray tank to 3/4 of total spray volume. Then add emulsifiable concentrates and any solutions.

Note: Spray adjuvants (non-ionic surfactants, crop oil concentrates, methylated seed oil, urea ammonium nitrate,

and ammonium sulfate) required for postemergence foliar applications should be added to the spray tank last.

Finish filling the spray tank. Maintain continuous agitation during mixing, final filling, and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled material may be more difficult to resuspend than when originally mixed.

Line screens in the spray tank should be no finer than 50 mesh (100 mesh is finer than 50 mesh).

Application with Liquid Fertilizer

Bottled Tigris Clop + Flumet is the preferred product form for use in liquid fertilizer. Tigris Clop + Flumet in water soluble packets can be used, but **must be premixed or slurried with water prior to use in liquid fertilizer**. It is important that **all water soluble packet material be totally dissolved before transferring into liquid fertilizer**. Any packet material undissolved in the premix will remain undissolved in the liquid fertilizer solution and could potentially lead to the clogging of screens and nozzles. For best results, use a minimum of 2 pints of water for every 6 oz of Tigris Clop + Flumet water soluble packets. Add the Tigris Clop + Flumet packets while mixing and **continue mixing until all packet material is dissolved** and granules are dispersed. The time needed to completely dissolve the packet material will depend upon water temperature and efficiency of mixing. Increasing the amount of premix water will decrease required mixing time. It is also recommended that premix and rinsate from the premix container be added to the spray tank through a 20-35 mesh screen.

When necessary, a compatibility agent can be used to ensure that Tigris Clop + Flumet mixes properly. The use of appropriate compatibility agents is especially important when tank mixing Tigris Clop + Flumet and other dry flowables, wettable powders, flowables, liquids, aqueous suspensions, or solutions with emulsifiable concentrates in liquid fertilizers. If the emulsifiable concentrate formulation rises to the surface of the fertilizer as an oil ("oils out"), the oil may combine with the wettable powder, flowable, or suspension to form oily curds (viscous phase) which are difficult to disperse. A jar test, utilizing relative proportions of the tank mix ingredients is recommended prior to mixing with liquid fertilizers.

Note: Do not use liquid fertilizer as the carrier when Tigris Clop + Flumet is applied postemergence to corn.

Application with Dry Bulk Fertilizer

Dry bulk fertilizer may be impregnated or coated with Tigris Clop + Flumet. Application of dry bulk fertilizer impregnated with Tigris Clop + Flumet provides weed control equal to the same rates of Tigris Clop + Flumet applied in liquid carriers. Follow label recommendations for Tigris Clop + Flumet regarding rates per acre, crops, special instructions, cautions, and special precautions. Apply 200 to 700 pounds of the fertilizer/herbicide mixture per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential to prevent possible crop injury. Non-uniform application may also result in unsatisfactory weed control. In areas where conventional tillage is practiced, a shallow incorporation of the mixture into the soil may improve weed control.

Most dry fertilizers can be used for herbicide impregnation with Tigris Clop + Flumet. When coated ammonium nitrate and/or limestone are used alone, do not impregnate with Tigris Clop + Flumet; these materials will not absorb the herbicide. Fertilizer blends containing coated ammonium nitrate and/or limestone as a part of the fertilizer mixture can be impregnated.

Compliance with all federal and state regulations relating to blending pesticide mixtures with dry bulk fertilizer, registration, labeling, and application are the responsibility of the individual and/or company offering the fertilizer and chemical mixture for sale.

Impregnation: Tigris Clop + Flumet must be pre-mixed or slurried with water prior to impregnation of dry bulk

fertilizer. For best results, use a minimum of 2 pints of water per 6.0 oz water soluble packet. To make the water slurry, add the required rate of Tigris Clop + Flumet (see formula below) to enough water to give a total volume of at least 6 pints of solution per ton of fertilizer. Make sure the Tigris Clop + Flumet is thoroughly dispersed in the water before spraying onto the fertilizer. Spray nozzles should be placed to provide uniform spray coverage onto the fertilizer. Care should be taken to aim the spray directly onto the fertilizer and avoid spraying the walls of the blender. Use any closed drum, belt, ribbon, or other commonly used dry bulk fertilizer blender.

Calculate amounts of Tigris Clop + Flumet by the following formula:

$$\frac{2,000}{\text{Pounds/acre of fertilizer}} \times \text{Pounds/acre of Hornet} = \text{Pounds of product per ton of fertilizer}$$

Note: Thoroughly clean dry fertilizer blending equipment prior to use with other herbicides. It is important to clean the blender, herbicide spray tank, and spraying apparatus thoroughly. Rinse the sides of the blender and the herbicide tank with water. Then, impregnate the rinsate onto a load of dry fertilizer intended for an approved crop. Use a maximum rate of 1 gallon of rinsate per ton of fertilizer. Follow with 1 to 2 loads of unimpregnated fertilizer in the blender before switching herbicides. The fertilizer application equipment must be empty, clean, and dry before applying any material to crops other than corn.

CROP-SPECIFIC USE DIRECTIONS

FIELD CORN

Soil Applied Treatments

Broadcast Application Rates (Preplant Surface Applied, Preplant Incorporated, Postplant Preemergence, and Spike Stage Treatments)

Soil Texture	Tigris Clop + Flumet (oz/acre)	
	<3.0% Organic matter	>3.0% Organic matter
Coarse	4.0	4.0 - 5.0
Medium or Fine	4.0 - 5.0	5.0

Note: Use the high end of the rate range on soils with greater than 3% organic matter and/or when applications are made 14 to 30 days before planting.

Broadleaf Weeds Controlled by Tigris Clop + Flumet when Soil Applied

Tigris Clop + Flumet will control "triazine tolerant" biotypes of these weeds, commonly know as "triazine resistant".

Note: Numbers within parentheses (-) in weeds list refer to "Use Information for Specific Weeds" below.

amaranth, Palmer	jimsonweed	pigweed, redroot	sida, prickly
anoda, spurred	kochia(1, 5)	pigweed, smooth	smartweed, Pennsylvania
beggarweed, Florida	ladysthumb	poinsettia, wild	spurge, nodding
buckwheat, wild	lambquarters, common	puncturevine	spurge, prostrate
carpetweed	mallow, Venice	purslane, common	spurge, spotted
chickweed, common	morningglory, entireleaf (1)	ragweed, common	sunflower, common
cocklebur, common	morningglory, ivyleaf (1)	purslane, common	thistle, Canada (3)
clover, red	morningglory, tall (1)	ragweed, giant (1)	velvetleaf
henbit	mustard, wild	shepherd's purse	waterhemp species (4, 5)
horseweed (marestail)	nightshade species (2)	sicklepod	wormwood, biennial

Use Information for Specific Weeds:

1. Partially controlled.
2. Control of moderate to heavy infestations of nightshade will be improved with a tank mixture of the appropriate labeled rate of an atrazine pre-mix product or a surface applied acetanilide product such as Dual® II herbicide, Dual II Magnum® herbicide, Surpass® herbicide or Topnotch® herbicide.
3. Burndown control of Canada thistle in minimum and no-till corn only.
4. To aid in control of waterhemp, apply Tigris Clop + Flumet in tank mix combination with the appropriate labeled rate of a surface applied acetanilide product such as Surpass or TopNotch herbicide.
5. Tigris Clop + Flumet will not control ALS resistant or tolerant biotypes of kochia.

Tigris Clop + Flumet may be soil applied as a preplant surface, preplant incorporated, or preemergence treatment. Apply alone or in tank mix combination with a grass control product such as Surpass or Topnotch herbicide.

Tank Mixing Limitations: Tigris Clop + Flumet may be applied in tank mix combination with other products provided (1) the timing and method of application is the same as recommended for Tigris Clop + Flumet; and (2) tank mixing with Tigris Clop + Flumet is not prohibited by the label of the tank mix product. When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels.

Soil Application Directions

Applications may be made from 30 days prior to planting through V6 stage or 20" tall corn, whichever occurs first.

1. **Preplant Incorporated Application:** For best results, apply and incorporate Tigris Clop + Flumet from 0 to 30 days before planting. Preplant incorporated treatments may be applied in water or liquid fertilizer. Uniformly incorporate the herbicide treatment into the top 2 to 3 inches of the final seedbed.
2. **Preplant Surface Applied:** For best results in minimum-tillage or no tillage systems, Tigris Clop + Flumet alone and with certain tank mixtures may be applied up to 30 days before planting. If weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide such as Gramoxone Extra herbicide, Touchdown herbicide, glyphosate (Glyphomax® Plus herbicide or Roundup UltraMAX herbicide). When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels. To the extent possible do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

Canada Thistle Control in Minimum and No-Till Corn: Tigris Clop + Flumet may be applied as a burndown treatment for control of emerged Canada thistle in minimum and no-till corn. The application will result in reduced late season competition. Delay the application until most of the thistle has emerged and averages 4 to 8 inches in height. For applications to Canada thistle, always include crop oil concentrate (See "Adjuvant Systems" in "Postemergence Treatments" section). Tank mix Tigris Clop + Flumet with glyphosate (Glyphomax Plus or Roundup UltraMAX), or sulfosate (Touchdown) and non-ionic surfactant for burndown control of existing grass and annual broadleaf weeds. When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels. Do not apply in tank mixture with Gramoxone Extra as this will result in reduced control of Canada thistle. Do not cultivate for at least 14 days after application to allow for thorough translocation of the herbicide treatment.

Note: Tigris Clop + Flumet will not control Canada thistle that has not emerged at the time of application in minimum or conventional tillage systems.

3. **Burndown Application:** When used as a burndown application, Tigris Clop + Flumet will provide foliar control of broadleaf weeds listed in the "Postemergence Treatments" section of this label and residual control of weeds listed under soil application. Foliar burndown applications should always include crop oil concentrate (see "Adjuvant Systems" in "Postemergence Treatments" section). To broaden the spectrum of weeds controlled, Tigris Clop + Flumet may be tank mixed with other herbicides such as glyphosate (Glyphomax Plus or Roundup UltraMAX), sulfosate (Touchdown), paraquat (Gramoxone Extra), or 2,4-D herbicide, etc. (See tank mixing instructions.)
4. **Preemergence Application:** Apply at the time of planting or after planting, but prior to crop or weed emergence. Adequate soil moisture following application is required for optimum herbicidal activity. For surface applications, rainfall, or overhead sprinkler irrigation is necessary to move Tigris Clop + Flumet into the weed germination zone. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture, and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is generally adequate. If adequate soil moisture is not received within 7 to 10 days after a surface applied treatment, a shallow cultivation is recommended to control established weeds and move the herbicide into the weed germination zone. When adequate soil moisture is received following dry conditions, performance may vary with weed species and the depth of the weed root system in the soil.
5. **Spike Stage Application:** Apply from corn emergence (ground cracking stage) until corn is 2 inches in height and before the first leaf is unfurled. Adequate soil moisture is required for optimum herbicidal activity. For those weeds that have not emerged at the time of application, rainfall or overhead sprinkler irrigation is necessary to move Tigris Clop + Flumet into the weed germination zone. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture, and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is generally adequate. If adequate soil moisture is not received within 7 to 10 days after a surface applied treatment, a shallow cultivation is recommended to control established weeds and mix the herbicide into the weed germination zone. When adequate soil moisture is received following dry conditions, performance may vary with weed species and rooting depth of target weeds.

Tank Mixing

(Preplant Surface Applied, Preplant Incorporated, and Postplant Preemergence Treatments)

Note: When tank mixing with a companion herbicide, read and follow each manufacturer's label for weeds controlled, applicable use directions, precautions, and limitations.

1. **Reduced Rates of Tigris Clop + Flumet Plus Atrazine-Containing Pre-Mix Products**

Reduced rates of Tigris Clop + Flumet can be tank mixed with labeled rates of atrazine-containing pre-mix herbicide products such as Surpass 100 herbicide, FullTime® herbicide, Keystone® herbicide or Keystone LA herbicide for improved control of certain broadleaf weeds not consistently controlled by atrazine pre-mix products. Tigris Clop + Flumet may be applied in tank mix combination with other products provided (1) the timing and method of application is the same as recommended for Tigris Clop + Flumet; and (2) tank mixing with Tigris Clop + Flumet is not prohibited by the label of the tank mix product. When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels. Reduced rates of Tigris Clop + Flumet tank mixed with labeled rates of these atrazine pre-mix products will provide consistent preemergence control of velvetleaf, lambsquarters, pigweed species, waterhemp, and triazine "resistant" varieties (triazine tolerant biotypes) of these species. These tank mixtures will also provide improved control of large-seeded broadleaf weeds such as cocklebur, common ragweed, giant ragweed, common sunflower, and jimsonweed.

On soils with less than 3% organic matter, tank mix Tigris Clop + Flumet at 3.0 oz/A with the recommended label rate of the atrazine pre-mix product. On soils with greater than 3% organic matter, tank mix Tigris Clop + Flumet at 4.0 oz/A with the recommended label rate of the atrazine pre-mix product.

Soil Organic Matter	Tigris Clop + Flumet WDG (oz/acre)	Acres per 6 lb Plastic Jug	Acres per 6 oz Packet (Contains .four 1.5 oz units)
<3%	3.0	32	2
>3%	4.0	24	1.5

2. Tigris Clop + Flumet plus Glyphosate (Glyphomax Plus or Roundup UltraMAX), Gramoxone Extra, or Touchdown for Minimum-tillage or No-tillage Systems

In minimum-tillage or no-tillage situations where corn is planted directly into a cover crop, stale seedbed, or previous crop residues, herbicides such as glyphosate (Glyphomax Plus or Roundup UltraMAX), paraquat (Gramoxone Extra), or sulfosate (Touchdown) may be tank mixed with Tigris Clop + Flumet. Apply in 10 to 60 gallons of water or fluid fertilizer per acre with ground equipment. The higher end of the carrier rate will provide better coverage under high residue situations. When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels.

Application Timing: Apply before, during (behind the planter), or after planting, but before the crop emerges.

Glyphomax Plus or Roundup UltraMAX: See the product label for Glyphomax Plus, Roundup UltraMAX, (or other labeled glyphosate) herbicide label for weeds controlled, recommended rates for specific weeds, and application instructions.

Gramoxone Extra: See the label for Gramoxone Extra for weeds controlled, recommended rates for specific weeds, and application instructions. Do not apply combinations containing Gramoxone Extra in suspension type fertilizers as the activity of the active ingredient paraquat will be reduced.

3. Hornet Plus 2,4-D for Minimum-tillage or No-tillage Systems

Where heavy crop residues exist, add 1.0 to 2.0 pints per acre of an appropriately labeled 3.8 - 4.0 lb a.e. per gallon 2,4-D amine or ester to the spray tank and apply in a volume of carrier capable of providing sufficient coverage of the crop residue. A carrier volume of 20 gallons per acre is recommended in heavy crop residue situations.

As carriers, nitrogen solutions and complete liquid fertilizers applied before corn emergence will enhance burndown of existing weeds and, therefore, are recommended instead of water. Add a crop oil concentrate or non-ionic surfactant at 1.0 to 2.0 quarts per 100 gallons diluted spray or another appropriate surfactant at its recommended rate. Apply before weeds reach 6 inches in height. This tank mixture will not control emerged grasses.

Tigris Clop + Flumet Soil-Applied Followed by Postemergence Treatments:

Broadleaf weeds not controlled by Tigris Clop + Flumet may be controlled with a postemergence herbicide product such as Tigris Clop + Flumet, Banvel herbicide, Clarity herbicide, 2,4-D herbicide, Marksman herbicide, Buctril herbicide, or Beacon herbicide. Read and follow each manufacturer's label for weeds controlled, applicable use directions, precautions, and limitations before use.

Postemergence Treatments

Apply Tigris Clop + Flumet as a postemergence spray at a rate of 2.0 - 5.0 ounces per acre. Use higher rates for control of heavy weed infestations, larger weeds, or when a longer period of residual control is desired. When applied postemergence, Tigris Clop + Flumet must be used with one of the adjuvant systems described below.

Postemergence Application Rates:

Acres Per Package Type	Application Rate (oz/acre) †			
	2.0 oz/acre	3.0 oz/acre	4.0 oz/acre	5.0 oz/acre
Acres per 6 oz Packet (Contains four 1.5 oz units)	3	2	1.5	1.2
Acres per 6 lb jug	48	32	24	19.2

† Refer to Mixing Directions section to determine the number of water soluble packets and total spray volume required for treated acreage.

Application Timing

Apply to actively growing weeds as a broadcast, or band treatment from the time of corn emergence (spike stage) until corn reaches 20 inches in height or the V6 stage whichever occurs first. For optimal control, apply before broadleaf weeds exceed the maximum height listed. Weeds that exceed the maximum height listed may be suppressed and recover after 2 to 3 weeks.

Directed Postemergence Application: Tigris Clop + Flumet may be applied as a directed postemergence application to corn that is 20 to 36 inches in height or has more than 6 leaf collars. Use only drop nozzles and avoid spraying the corn plant by directing the spray as low as possible while allowing for optimal coverage of weeds. Use the highest labeled rates for weeds greater than the maximum size listed on this label. Control of weeds larger than the maximum height listed may vary due to weeds species, stage of growth, and growing conditions. Results may range from complete control to suppression.

- Do not spray into the whorl of corn plants.
- Do not apply to corn more than 36 inches tall.

Note: If field corn is grown for forage or ensilage, application must occur before corn reaches 20 inches in height or V6 growth stage (whichever occurs first) and an interval of at least 45 days is required between application and harvest.

Factors Affecting Weed Control: Apply to actively growing weeds. Extreme growing conditions such as drought, or near freezing temperatures before, at, or following application may result in reduced weed control. Degree of control will depend on coverage of treated weeds and weed susceptibility as well as growing conditions at the time of treatment.

Environmental Conditions and Herbicidal Activity of Tigris Clop + Flumet: Factors in effective weed control with Tigris Clop + Flumet include application rate, weed size, daytime temperature, soil moisture prior to and following application, and use of adjuvants. Best weed control results are obtained when Tigris Clop + Flumet is applied to small, actively growing weeds, when daytime temperatures are warm (70°F or more), and soil moisture is adequate to support active weed growth prior to and following application. If weeds are under drought stress, consider delaying application until more favorable conditions resume. Application when weeds are moisture stressed or taller than the recommended height for control may result in only partial control.

- Tigris Clop + Flumet is rainfast in 2 hours.
- Applications made immediately prior to, during, or immediately following periods of large day/night

temperature fluctuations or where daytime temperatures do not exceed 60°F may decrease weed control.

- Poor weed control may result from applications made to plants under stress from:
 - abnormally hot or cold weather
 - environmental conditions such as drought, water-saturated soils, hail damage, or frost
 - prior herbicide applications

Use of Adjuvants: All postemergence applications of Tigris Clop + Flumet must include 1) a non-ionic surfactant at 0.25% volume/volume (1 qt/100 gal) or 2) crop oil concentrate or methylated seed oil at 1% volume/volume (1 gal/100 gal). Use a good quality surfactant with at least 80% active ingredient (of which at least 50% is actual non-ionic surfactant). Under extremely dry growing conditions, the use of an agriculturally approved sprayable liquid fertilizer or ammonium sulfate, in combination with the non-ionic surfactant or crop oil concentrate or methylated seed oil may enhance control. Use 28%, 30%, or 32% urea ammonium nitrate at 2.5% volume/volume (2.5 gal/100 gal) or 2 to 4 lb of sprayable grade ammonium sulfate per acre.

Note: Do not use liquid fertilizer solutions or suspensions as the total carrier because excessive crop injury may occur. Use only EPA approved surfactants for use on food crops.

Cultivation: For best results, do not cultivate within 10 days before or after application.

Tank Mixing: Tigris Clop + Flumet may be applied in tank mix combination with other products provided (1) the timing and method of application is the same as recommended for Tigris Clop + Flumet; and (2) tank mixing with Tigris Clop + Flumet is not prohibited by the label of the tank mix product; and (3) the tank mix combination is compatible as determined by a "jar test" described in the "Tank Mix Compatibility Testing" section. When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels. For control of grass weeds, Tigris Clop + Flumet may be tank mixed with a postemergence grass herbicide such as Accent or Basis Gold, Option or Steadfast. For an expanded spectrum of broadleaf weed control, Tigris Clop + Flumet may be tank mixed with other postemergence broadleaf herbicides such as atrazine, Banvel, Buctril, Callisto, Clarity, Distinct, or 2,4-D. Tigris Clop + Flumet may also be tank mixed with Glyphomax Plus, Roundup UltraMAX, or other labeled glyphosate formulations for application to Roundup Ready field corn.

Do not post apply Tigris Clop + Flumet in tank mix combination with Basagran, Laddock or Lightning herbicides as severe crop injury may result.

Weeds Controlled And Application Rates for Postemergence Application

(Use higher rates for control of larger weeds and for control of heavy weed infestations.)

Tigris Clop + Flumet will control triazine tolerant biotypes of these weeds, commonly know as "triazine resistant".

Note: Numbers in parentheses (-) within table refer to Specific Use Directions below.

Annual Weed Control			
Application to "Spike" Corn (1)	Postemergence Application After "Spike" Stage of Growth		
4.0 to 5.0 oz/acre	2.0 oz/acre (weeds 1 - 3 in. tall)	3.0 oz/acre (weeds 1 - 6 in. tall)	4.0 oz/acre (weeds 1 - 8 in. tall)
anoda, spurred beggarweed, Florida buckwheat, wild carpetweed chickweed, common cocklebur, common henbit horseweed (marestail) mallow, venice mustard, wild poinsettia, wild puncturevine purslane, common shepherd's purse sida, prickly spurge, nodding spurge, prostrate spurge, spotted sunflower, common velvetleaf	anoda, spurred beggarweed, Florida chickweed, common cocklebur, common henbit horseweed (marestail) mallow, venice mustard, wild poinsettia, wild puncturevine purslane, common shepherd's purse sida, prickly spurge, nodding spurge, prostrate spurge, spotted sunflower, common velvetleaf	anoda, spurred beggarweed, Florida chickweed, common cocklebur, common henbit horseweed (marestail) jimsonweed ladysthumb mallow, venice mustard, wild poinsettia, wild puncturevine purslane, common ragweed, common ragweed, giant shepherd's purse sida, prickly smartweed, Pennsylvania spurge, nodding spurge, prostrate spurge, spotted sunflower, common velvetleaf	anoda, spurred beggarweed, Florida chickweed, common cocklebur, common henbit horseweed (marestail) jimsonweed ladysthumb lettuce, prickly mallow, Venice mustard, wild poinsettia, wild puncturevine purslane, common ragweed, common ragweed, giant shepherd's purse sida, prickly smartweed, Pennsylvania spurge, nodding spurge, prostrate spurge, spotted sunflower, common velvetleaf
Partial Control	Partial Control	Partial Control (weeds <2 in. tall)	Partial Control (weeds <4 in. tall)
morningglory, entireleaf morningglory, ivyleaf morningglory, tall ragweed, giant	jimsonweed ladysthumb ragweed, common ragweed, giant smartweed, Pennsylvania	buckwheat, wild kochia (2) lambquarters, common lettuce, prickly morningglory, entireleaf morningglory, ivyleaf morningglory, tall nightshade species pigweed, redroot pigweed, smooth sicklepod thistle, Russian waterhemp species (2)	buckwheat, wild kochia (2) lambquarters, common morningglory, entireleaf morningglory, ivyleaf morningglory, tall nightshade species pigweed, redroot pigweed, smooth sicklepod thistle, Russian waterhemp species (2)
Biennial and Perennial Weed Control			
Apply 3.0 to 5.0 oz/acre to weeds 3 - 9 inches tall (3,4)			
alfalfa, volunteer artichoke, Jerusalem burdock, common	clover, red clover, sweet dandelion	dock, curly sorrel, red	thistle, Canada (5) wormwood, biennial

Specific Use Directions:

1. **Spike corn:** Apply 5 oz per acre for greater residual control on soils with greater than 3% organic matter. Apply 5 oz per acre to increase the degree of partial control for morningglory species and giant ragweed.
2. Tigris Clop + Flumet will not control ALS resistant or tolerant biotypes.
3. **Biennial and Perennial weeds:** A rate of 4.0 - 5.0 oz per acre will generally provide season-long control. A rate of 3.0 oz per acre will provide control of top growth only. Some regrowth may occur by the end of the season.
4. **Biennial and Perennial weeds:** Do not tank mix with contact herbicides (such as atrazine, metribuzin, or bromoxynil) as reduced weed control will result.
5. **Canada thistle:** For Canada thistle control the following season, expressed as stand reduction, apply 5.0 oz per acre of Tigris Clop + Flumet in tank mix combination with 4.0 oz per acre of Stinger herbicide †

† **Note:** Maximum Use Rate for the active ingredient clopyralid is 0.25 lb per acre. One ounce of Tigris Clop + Flumet contains 0.031 lb of clopyralid. One fluid ounce of Stinger contains 0.023 lb of clopyralid.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 122°F for extended periods of time. If container is damaged or spill occurs, use product immediately or contain with absorbent materials and dispose as waste. Water Soluble Packaging: Packets may become brittle when stored below 32°F. Handle carefully when frozen to avoid breakage or allow package to warm above 32°F before handling.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of onsite according to label use directions or at an approved waste disposal facility.

Nonrefillable rigid containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Nonrefillable nonrigid containers:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refillable rigid containers larger than 5 gal:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container.

Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable rigid containers larger than 5 gal:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows:

Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water.

Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of TIGRIS, LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, TIGRIS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither TIGRIS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

[**Tigris Clop + Flumet**] is a trademark of Tigris, LLC

®Dual MAGNUM and Dual II MAGNUM are registered trademarks of Syngenta

{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

Tigris Clop + FlumetTM

Active Ingredient: (% by weight)
 flumetsulam: N-(2,6-difluorophenyl)-5- methyl-1,2,4-triazolo-
 [1,5a]-pyrimidine-2-sulfonamide 18.5%
 clopyralid potassium salt: 3,6-dichloro-2-
 pyridinecarboxylic acid, potassium salt 60.0%
Other Ingredients 21.5 %
Total 100.0%

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency

Spill, Leak, Fire, Exposure, or Accident Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes Substantial But Temporary Eye Injury. Harmful If Swallowed, Inhaled, Or Absorbed Through The Skin. May Cause Skin Sensitization Reactions In Certain Individuals
 Do not get in eyes or on clothing. Avoid breathing vapors or spray mist and contact with skin.

ENVIRONMENTAL HAZARDS: Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

The active ingredients in this product are known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of

this product where soils are permeable, particularly where the water table is shallow, may result in leaching to groundwater.
 Caution should be exercised when handling this product at mixing and loading sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 122°F for extended periods of time. If container is damaged or spill occurs, use product immediately or contain with absorbent materials and dispose as waste. Water Soluble Packaging: Packets may become brittle when stored below 32°F. Handle carefully when frozen to avoid breakage or allow package to warm above 32°F before handling.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of onsite according to label use directions or at an approved waste disposal facility.

Nonrefillable rigid containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Nonrefillable nonrigid containers:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refillable rigid containers larger than 5 gal:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable rigid containers larger than 5 gal:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows:
 Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

See inside label booklet for additional Precautionary Statements and Directions for Use.

Manufactured for:
 Tigris, LLC
 1204 Village Market Place
 #173
 Morrisville, NC 27560

EPA Reg. No. 92647-XX
EPA Est. No. _____
NET WEIGHT: _____