



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

July 1, 2024

Ailis Gregory, Agent for  
Suntton International Inc.  
c/o Pyxis Regulatory Consulting, Inc.  
4110 136th Street CT NW  
Gig Harbor, Washington 98332

Subject: Pesticide Registration Improvement Act (PRIA) Labeling Amendment – to add use directions for stone fruit to the foliar application table of the label.

Product Name: Sunergist® EBR

EPA Registration Number: 92983-2

EPA Receipt Date: 08/22/23

Action Case Number: 00488896

Dear Ms. Gregory:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

The supplemental labeling contains some new and/or revised uses and/or directions that may be additional to the uses and/or directions found on the label on or attached to the container, but the supplemental labeling does not by itself constitute the complete set of use directions. The complete set of use directions is set forth on the container label as combined with the supplemental labeling.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. **You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling.** In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 § CFR 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S.

Environmental Protection Agency or the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section

12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Cheryl Greene or at [green.cheryl@epa.gov](mailto:green.cheryl@epa.gov) or Gina Burnett at [burnett.gina@epa.gov](mailto:burnett.gina@epa.gov)

Sincerely,

A handwritten signature in black ink, appearing to read 'Gina Burnett', is centered below the 'Sincerely,' text.

Gina Burnett, Senior Regulatory Advisor  
Biochemical Pesticides Branch  
Biopesticides and Pollution Prevention Division (BPPD)  
Office of Pesticide Programs

Enclosure: Stamped Label

**ACCEPTED**

07/01/2024

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No.

92983-2

**Sunergist<sup>®</sup> EBR**

Plant growth regulator

For Agricultural and Non-agricultural use on listed crops

**ACTIVE INGREDIENT**

24-Epibrassinolide..... By wt. 0.01%

**OTHER INGREDIENTS** ..... 99.99%

**Total:** ..... 100.00%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**First Aid**

**If on skin:**

- Take of contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes
- Call a poison control center or doctor for treatment advice.

**If inhaled**

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

**Hot Line Number**

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For general information on product use, etc., call the National Pesticides Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12:00 PM Pacific Standard Time. For emergencies, call the poison control center at 1-800-222-1222.

EPA Reg. No. 92983-2

EPA Est. No. XXXXX-XXX-X

(Batch)(Lot) No.

Manufactured for:

Suntton International Inc.

9127 Highway 431 South, Suite C

Owens Cross Roads, AL 35765

Net Contents:

**PRECAUTIONARY STATEMENT**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Harmful if inhaled. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

SUNERGIST EBR (EPA Reg. No. 92983-2)

Master Label – Version (2) – [EPA Approval Date]

Page **1** of **12**

## PERSONAL PROTECTIVE EQUIPMENT (PPE)

### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves

### USER SAFETY RECOMMENDATION

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all directions on the EPA approved label and any supplemental labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handler may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment, restricted-entry interval, and notification to workers.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls
- Shoes plus socks
- Waterproof gloves

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries, or greenhouse.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

### GENERAL INFORMATION

**SUNERGIST® EBR** is a special formulated product that contains 24-Epibrassinolide plant growth regulator (PGR) found in nature that stimulates seeding and plant growth and improves root development. **SUNERGIST® EBR** improves the physiological and biochemical processes involved in crop growth and development.

- Synergistically acts with other endogenous PGRs
- Promotes cell division and cell elongation
- Promotes seed germination & seedling growth.
- Enhances the plant tolerance to adverse environmental stress (saline, drought, low & elevated temperature)
- Enhances the plant tolerance to adverse biotic stress (Fungi and virus diseases)
- Improvement of photosynthesis and translocation of assimilates
- Better pollination and fruit set and yield increase
- Supports even and early bud break on listed crops
- Enhances quality of produce

### AGRICULTURAL USES

#### SPRAY VOLUME (IN GALLONS) PER ACRE

Crop	Ground		Aerial
	Dilute	Concentrate	
Vegetables, Field Crops	15	15	5
Small fruits, vines, Miscellaneous	100	25	15
Pome Fruits, Stone Fruits Tree/nut crops /nuts & Crops	200	50	20
Citrus	300	100	20

Spray volume applied per acre will differ depending upon the stage of plant growth, equipment used, and the specific crops. Apply sufficient water volume to ensure thorough coverage on entire plants.

Do not apply this product through any type of irrigation system.

**Compatibility:** **SUNERGIST® EBR** compatibility with other agricultural products requires testing for crop safety and performance prior to large-scale use. Products mixed with **SUNERGIST® EBR** must **not** be in alkaline (pH >7) solution. **SUNERGIST® EBR** can be mixed with any product having a pH less than 7. If rain occurs within 2 hours after initial application, a repeat application may be necessary.

**Spray Adjuvant:** Use commercially registered **Non-Ionic** surfactant and penetrant adjuvant at a label rate approved for use on respective crops for all applications of **SUNERGIST® EBR**.

## FOLIAR APPLICATION INSTRUCTIONS

Crop	Amount of product in fl. oz./10 gals water	Application timing	Remarks
<b>Brassica Head and Stem Vegetables (Crop Group 5-16)</b> , including Broccoli, Brussels Sprouts, Cauliflower, Cabbage, Cabbage (Chinese, napa) and Cultivars, varieties, and hybrids of these commodities	0.6 – 1.2	1 <sup>st</sup> . Spray at 3-5 leaf stage or immediately after transplanting / transplant water mixing.  2 <sup>nd</sup> . Spray at early head forming /rosette stage	
<b>Oilseeds (Crop Group 20)</b> , including Borage, Calendula, Canola, Castor oil plant, Chinese tallowtree, Cottonseed, Crambe, Cuphea, Echium, Euphorbia, Evening primrose, Flax seed, Gold of pleasure, Hare's ear mustard, Jojoba, Lesquerella, Lunaria, Meadowfoam, Milkweed, Mustard seed, Niger seed, Oil radish, Poppy seed, Rapeseed, Rose hip, Safflower, Sesame, Stokes aster, Sunflower, Sweet rocket, Tallowwood, Tea oil plant, Vernonia, Cultivars and varieties and/or hybrids of these	0.6 – 1.2	1 <sup>st</sup> . Seedling stage  2 <sup>nd</sup> 50% blooming stage	
<b>Cereal Grains (Crop Group 15)</b> , including  Corn (Sweet, field and popcorn)  Wheat, Sorghum, Barley, Buckwheat, Millet (pearl and proso), Oats, Rye, Teosinte, and Triticale  Rice, Wild rice	0.6 – 1.2  0.6 – 1.2  0.6 – 1.2	1 <sup>st</sup> . Spray at V2-V3 2 <sup>nd</sup> . V10-11 stage-Trumpt stage  1 <sup>st</sup> . Tiller stage 2 <sup>nd</sup> . Flag stage 3 <sup>rd</sup> . Anthesis for tolerance on heat, cold and low light stresses  1 <sup>st</sup> . Tiller stage 2 <sup>nd</sup> . Flag leaf 3 <sup>rd</sup> . Anthesis	Reduce bare tips on corn  Application on anthesis enhances yield.  Helping even tiller and heading  Increase grain sets  To alleviate the heat stress and enhance pollination and set.

Crop	Amount of product in fl. oz./10 gals water	Application timing	Remarks
<b>Cotton</b>	0.6 – 1.2	1 <sup>st</sup> . At 4-6 true leaf stage 2 <sup>nd</sup> . 80% blooming 3 <sup>rd</sup> . 3-4 days after petal fall to prevent boll shedding	For cold tolerance on needed bases
<b>Cucurbit Vegetables (Crop Group 9)</b> , including Chayote, Chinese waxgourd, Citron melon, Gherkin, Gourd (edible), <i>Momordica</i> spp., Muskmelon, Cantaloupe, Pumpkin, Summer Squash, Winter Squash, Watermelon and Honeydew	0.6 – 1.2	1 <sup>st</sup> . At initial blooming 2 <sup>nd</sup> -4 <sup>th</sup> . 3-4 times at 7 days interval	
<b>Cucumber (greenhouse)</b>	0.6 – 1.2	1 <sup>st</sup> . After transplanting 2 <sup>nd</sup> . Every 7 – 10 days thereafter with watering	
<b>Fruiting vegetables (Crop Group 8-10)</b> , including African Eggplant, Bush tomato, Cocona, Currant tomato, Eggplants, Garden huckleberry, Goji berry, Groundcherry, Martynia, Naranjilla, Okra, Pea eggplant, Pepino, Peppers (bell, nonbell), Roselle, Scarlet eggplant, Sunberry, Tomatillo, Tomato, Tree tomato, and cultivars, varieties and/or hybrids of these	0.6 – 1.2	1 <sup>st</sup> . 5-7 days after first flower 2 <sup>nd</sup> . 10-14 days after first 3 <sup>rd</sup> . 10-14 days after 2 <sup>nd</sup> application	Spray 2 – 3 days prior to anticipated heat stress
<b>Leafy Vegetables (Crop Group 4-16)</b> , including Amaranth (Chinese), Amaranth (leafy), Arugula, Aster (Indian), Blackjack, Broccoli (Chinese), Broccoli raab, Cabbage (Abyssinian, Chinese, bok choy; and seakale), Cat's whiskers, Cham-chwi, Cham-na-mul, Chervil, Chipilin, Crysanthemum, Cilantro, Collards, Corn salad, Cosmos, Cress (garden and upland), Dandelion, Gang-gwi, Dillweed, Dock, Dol-nam-mul, Ebolo, Endive, Escarole,	0.6 – 1.2	1 <sup>st</sup> . Early seedling stage 2 <sup>nd</sup> . 10-14 day after 1 <sup>st</sup> spray 3 <sup>rd</sup> . 10-14 days after 2 <sup>nd</sup> spray	

Crop	Amount of product in fl. oz./10 gals water	Application timing	Remarks
<p>Fameflower, Feather cockscomb, Head lettuce, Good King Henry, Hanover salad, Huauzontle, Leaf lettuce, Jute, Kale, Lettuce (bitter), Maca, Mizuna, Mustard greens, Orach, Parsley, Plantain (buckthorn), Primrose (English), Purslane (garden and winter), Radicchio, Radish (leaves), Rape greens, Rocket (wild), Shepherd's purse, Spinach, Spinach (Malabar, New Zealand and tanier), Swiss chard, Turnip greens, Violet (Chinese, leaves), Watercress, and Cultivars, varieties, and hybrids of these</p> <p><b>Celery</b></p>			
<p><b>Legume Vegetables (Crop Group 6)</b> (Succulent or dry), including Bean (<i>Lupinus</i> spp.), Bean (<i>Phaseolus</i> spp.), Bean (<i>Vigna</i> spp.), Broad bean, Chickpea, Guar, Jackbean, Lablab bean, Lentil, Peas (<i>Pisum</i> spp.), Pigeon pea, Soybean, Soybean (immature seed), Sword bean</p>	0.6 – 1.2	<p>1<sup>st</sup>. Early seedling stage</p> <p>2<sup>nd</sup>. Full bloom -early pad stage</p> <p>3<sup>rd</sup>. 10-14 days after 2<sup>nd</sup> spray</p>	
<b>Peanut</b>	0.6 – 1.2	<p>1<sup>st</sup>. At initial blooming</p> <p>2<sup>nd</sup>. At early R2</p> <p>3<sup>rd</sup>. At early R3</p>	<p>Tank mix with fungicide</p> <p>Tank mix with fungicide</p>
<p><b>Tuber Vegetables (Crop Group 1)</b>, including Arracacha, Arrowroot, Artichoke (Chinese, Jerusalem), Beet (garden, sugar), Burdock (edible), Canna (edible), Carrot, Cassava (bitter, sweet), Celeriac (celery root), Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen, Ginger, Ginseng, Horseradish,</p>	0.6 – 1.2	<p>1<sup>st</sup>. 6-8 leaf stage</p> <p>2<sup>nd</sup>. 12-16 leaf stage for small tuber varieties</p> <p>2<sup>nd</sup>. Tuber formation stage for large tuber varieties</p>	



Crop	Amount of product in fl. oz./10 gals water	Application timing	Remarks
Leren, Parsley (turnip-rooted), Parsnip, Potato, Radish, Radish (oriental), Rutabaga, Salsify (oyster plant, black, Spanish), Skirret, Sweet potato, Tanier, Turmeric, Turnip, Yam bean (jicama, manioc pea) and Yam (true)			
<b>Citrus fruits (Crop Group 10-10)</b> including Australian desert lime, Australian finger lime, Australian round lime, Brown River finger lime, Calamondin, Citron, Citrus hybrids, Grapefruit, Japanese summer grapefruit, Kumquat, Lemon, Lime, Mediterranean mandarin, Mount White lime, New Guinea wild lime, Orange (sour, sweet), Pummelo, Russel River lime, Satsuma mandarin, Sweet lime, Tachibana orange, Tahiti lime, Tangelo, Tangerine (Mandarin), Tangor, Trifoliate orange, Uniq fruit, and Cultivars, varieties and/or hybrids of these	6.0 – 12.0 fl. oz. in 100 gals of water	1 <sup>st</sup> . 5-10 days after first flower 2 <sup>nd</sup> . Apply when fruit size is 5 mm	To increase doubles Additional sprays may be needed.
<b>Berries and small fruit (Crop Group 13-07)</b> Including:  Amur river grape, Aronia berry, Bayberry, Bearberry, Bilberry, Blackberry, Blueberry(highbush, lowbush), Buffalo currant, Buffaloberry, Che, Chilean guava, Chokecherry, Cloudberry, Cranberry, Currant (black, red), Elderberry, European barberry, Gooseberry, Highbush cranberry, Honeysuckle (edible), Jostaberry, Juneberry, Kiwifruit (fuzzy, hardy), Lingonberry, Maypop, Mountain pepper berries, Mulberry, Muntries, Native currant, Partridgeberry,	0.6 – 1.2	1 <sup>st</sup> . Bud break spray 2 <sup>nd</sup> . 50-80% flowering stage 3 <sup>rd</sup> . Small fruit stage	Additional spray 2-3 days before forecasted heat stress

Crop	Amount of product in fl. oz./10 gals water	Application timing	Remarks
Phalsa, Pincherry, Raspberry (black, red), Riberry, Salal, Schisandra berry, Sea buckthorn, Serviceberry, Strawberry, Wild raspberry, and Cultivars, varieties and/or hybrids of these  Wine Grape  Table Grape	0.6 – 1.2  0.6 – 1.2	1 <sup>st</sup> . 50-80 % flowering stage 2 <sup>nd</sup> Early small fruit stage (2-5 mm)  1 <sup>st</sup> . At bud break 2 <sup>nd</sup> . At small fruit sizing 3 <sup>rd</sup> . At berry softening	Tank mix with GA3 Enhance ripening
<b>Tree nuts (Crop Group 14-12),</b> Including  African nut-tree, Almond, Beechnut, Brazil nut, Brazilian pine, Bunya, Bur oak, Butternut, Cashew, Cajou nut, Candlenut, Cashew, Chestnut, Chinquapin, Coconut, Coquito nut, Dika nut, Ginkgo, Guiana chestnut, Hazlenut (Filbert), Heartnut, Hickory nut, Japanese horse-chestnut, Macadamia nut, Mongongo nut, Monkey-pot, Monkey puzzle nut, Okari nut, Pachira nut, Peach palm nut, Pecan, Pequi, Pili nut, Pine nut, Pistachio, Sapucaia nut, Tropical almond, Walnut (black and English), Yellowhorn, Cultivars, varieties and/or hybrids of these	6.0 – 12.0/100 gals of water	1 <sup>st</sup> Application 50-80% blooming 2 <sup>nd</sup> . Small nut stage	Improve pollination & net set and quality
<b>Bulb Vegetables (Crop Group 3-07),</b> including Chive (fresh leaves), Chive (Chinese, fresh leaves), Daylily (bulb), Elegans hosta, Fritillaria (bulb), Fritillaria (leaves), Garlic (bulb), Garlic (great headed, bulb), Garlic (Serpent, bulb), Kurrat, Lady's leek, Leek <i>Allium porrum</i> L., Leek (wild), Lily	0.6 – 1.2	1 <sup>st</sup> . early seedling stage 2 <sup>nd</sup> . Bulb forming stage	



## DRENCH APPLICATION OF SUNERGIST® EBR ON CROPS

Sunergist EBR may be applied through transplanting machine at time of transplanting and through irrigation systems after transplanting.

Crop	Amount of SUNERGIST® EBR (fl. oz.)	Application Timing
<b>Brassica Head and Stem Vegetables (Crop Group 5-16)</b> including Broccoli, Brussels sprouts, Cabbage, Cabbage (Chinese, napa), Cauliflower, and Cultivars, varieties, and hybrids of these <b>Brassica Leafy Greens Subgroup (Crop Subgroup 4-16B)</b> including Arugula, Broccoli (Chinese) Broccoli raab, Cabbage (abyssinian), Cabbage (Chinese, bok choy), Cabbage (seakale), Collards, Cress (garden and upland), hanover salad, kale, maca (leaves) mizuna, mustard greens, radish (leaves), rape greens, rocket (wild), shepherd's purse, turnip greens, watercress, cultivars, varieties, and hybrids of these commodities.	0.6 fl. oz. per 10 gallons water	1 <sup>st</sup> treatment at time of transplant. 2 <sup>nd</sup> treatment 10-14 days after first application.  Spray immediately after transplanting.

## IN-FURROW APPLICATION OF SUNERGIST® EBR ON CROPS

Crop	Amount of SUNERGIST® EBR (fl. oz.)	Application Timing
Field Corn Grain Sorghum Wheat, Barley, Oats, Rye Other in-furrow at planting crops	1.2 -1.8 fl. oz. per acre Or 0.6 fl. oz. per 10 gallons water	Apply at planting in the seed furrow or 2 inches beside and 2 inches below seed or with a strip till machine 3 inches below the seed.  Can be applied with or without starter fertilizer.

## SEED TREATMENT

**SUNERGIST® EBR** is not intended for commercial seed treatment application. Commercial Application: The application of Seed Treatments to seed in commercial facilities for commercial sale or gain (as opposed to application on-farm and in planters).

Apply **SUNERGIST® EBR** as a seed treatment at the rate of 1.0-1.5 fl. oz per 100 lbs of seed. Sunergist® EBR may be applied as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment or grower's hopper box. This slurry (totaling 6 fluid ounces) may be composed of water or in combination with EPA registered broad spectrum seed treatment fungicide/Insecticide or plant nutritional products.

All seed treated with this product must be colored with an EPA-approved dye or colorant that imparts an unnatural color to the seed to help prevent the inadvertent use of treated seeds for food, feed or oil purposes.

## TURF

**For Sod Grass:** Apply **SUNERGIST® EBR** by commercial ground sprayer using 20-40 gallons of water per acre. Use 1.2 fl. oz. per 10 gallons of water at a 1:2,000 dilution rate.

**For Turfgrass:** For use on St. Augustine, Bermuda, Centipede, Bluegrass, Rye, and Fescue: Use 0.12-0.60 fl. oz./1-5 gallons of water per 1,000 sq. ft. For Dichondra: Use 0.6-1.2 fl. oz./1-5 gallons of water per 1,000 sq. ft.

Make application at 2-week intervals for a total of 5 applications during the growing season.

## ORNAMENTALS

For greenhouse and nursery ornamentals such as foliage plants, bedding and flowering plants, woody ornamentals, garden grown fruit trees, and garden grown food crops, follow the use directions in the table below:

For Plants Less Than 2 Years Old	For Plants Greater Than 2 Years Old
Dilute 0.6 fl. oz. of <b>SUNERGIST® EBR</b> in 10 gallons of water (1:2000 dilution rate). Apply by using a commercial greenhouse/nursery hand sprayer. Repeat application at 10- to 14-day intervals when required. Apply 1-2 weeks prior to sale. Uniform and thorough coverage is necessary.	Dilute 1.2 fl. oz. of <b>SUNERGIST® EBR</b> in 10 gallons of water (1:1,000 dilution rate) Apply by using a commercial greenhouse/nursery hand sprayer. Repeat application at 10- to 14-day intervals when required. Apply 1-2 weeks prior to sale. Uniform and thorough coverage is necessary.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a cool, dry locked place out of the reach of children and out of direct sunlight.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. Triple rinse (or equivalent). Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## LIMITED WARRANTY AND DISCLAIMER

NOTICE: **Suntton International Inc.** warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use. Buyer assumes all risks of use and handling which is a variance in any way with the directions herein. **Suntton International Inc.** makes no other express or implied warranty of fitness or merchantability. In no case shall **Suntton International Inc.** or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SUNTTON INTERNATIONAL INC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SUNTTON INTERNATIONAL INC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

**Suntton International Inc.**, and Seller offer this product and the Buyer and user accept it, subject to the foregoing Limited Warranty and Disclaimer which may be varied only by agreement in writing signed by a duly authorized representative of Suntton International, Inc.

# Sunergist<sup>®</sup> EBR

Plant growth regulator

EPA Reg. No. 92983-2

## Supplemental Label for Cucurbit Vegetables

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### FOLIAR APPLICATION INSTRUCTIONS

Crop	Amount of product in fl. oz./10-gals water	Application timing	Remarks
<b>Cucurbit Vegetables (Crop Group 9)</b> including Chayote, Chinese waxgourd, Citron melon, Gherkin, Gourd (edible), Momordica spp., Muskmelon, Cantaloupe , Pumpkin, Summer Squash, Winter Squash, Watermelon and Honeydew	0.6 – 1.2	1 <sup>st</sup> . At initial blooming 2 <sup>nd</sup> – 4 <sup>th</sup> . At 7 – 14 days interval	
<b>Cucumber (greenhouse)</b>	0.6 – 1.2	1 <sup>st</sup> . After transplanting 2 <sup>nd</sup> . Every 7 – 10 days thereafter with watering	

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in possession of the user at the time of application.

Read the label affixed to the container for Sunergist<sup>®</sup> EBR before applying.

Use of Sunergist<sup>®</sup> EBR according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Sunergist<sup>®</sup> EBR.

# ***Sunergist***<sup>®</sup> EBR

Plant growth regulator

EPA Reg. No. 92983-2

## **Supplemental Label for Peanut**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### **FOLIAR APPLICATION INSTRUCTIONS**

<b>Crop</b>	<b>Amount of product in fl. oz./10-gals water</b>	<b>Application timing</b>	<b>Remarks</b>
<b>Peanut</b>	0.6 – 1.2	1 <sup>st</sup> . At initial blooming 2 <sup>nd</sup> . At early R2 3 <sup>rd</sup> . At early R3	Tank-mix with fungicide Tank-mix with fungicide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in possession of the user at the time of application.

Read the label affixed to the container for Sunergist<sup>®</sup> EBR before applying.

Use of Sunergist<sup>®</sup> EBR according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Sunergist<sup>®</sup> EBR.



# ***Sunergist***<sup>®</sup> EBR

Plant growth regulator

EPA Reg. No. 92983-2

## **Supplemental Label for Rice**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### **FOLIAR APPLICATION INSTRUCTIONS**

<b>Crop</b>	<b>Amount of product in fl. oz./10-gals water</b>	<b>Application timing</b>	<b>Remarks</b>
<b>Rice, Wild Rice</b>	0.6 – 1.2	1 <sup>st</sup> . Early tillering  2 <sup>nd</sup> . Flag leaf  3 <sup>rd</sup> . Anthesis	Helping even tiller and heading  Increase grain sets  To alleviate the heat stress and enhance pollination and set.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in possession of the user at the time of application.

Read the label affixed to the container for Sunergist<sup>®</sup> EBR before applying.

Use of Sunergist<sup>®</sup> EBR according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Sunergist<sup>®</sup> EBR.

# Sunergist® EBR

Plant growth regulator

EPA Reg. No. 92983-2

## Supplemental Label for Stone Fruits

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### FOLIAR APPLICATION INSTRUCTIONS

Crop	Amount of product in fl. oz./10-gals water	Application timing	Remarks
<b>Stone Fruits (Crop Group 12-12)</b> including Apricot, Apricot (Japanese), Capulin, Cherry (black, Nanking), Jujube (Chinese), Nectarine, Peach, Plum, Plum (American, beach, Canada, cherry, Chickasaw, Damson, Japanese, Klamath, prune), Plumcot, Sloe, Cultivars, varieties, and/or hybrids of these	1.2 – 6.4	1 <sup>st</sup> . White bud  2 <sup>nd</sup> . Petal fall  3 <sup>rd</sup> . During color change  4 <sup>th</sup> . Postharvest	Helping for even anthesis, recommend tank mix with cytokinin.  Enhance fruit set, recommend tank mix with cytokinin.  Uniform ripening.  To correct alternative bearing, recommend tank mix with cytokinin.
Tart & Sweet Cherry	6.4	1 <sup>st</sup> & 2 <sup>nd</sup> Tank mix with GA3 applications during straw color	Uniform ripening and even harvest

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in possession of the user at the time of application.

Read the label affixed to the container for Sunergist® EBR before applying.

Use of Sunergist® EBR according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Sunergist® EBR.

# ***Sunergist***<sup>®</sup> EBR

Plant growth regulator

EPA Reg. No. 92983-2

## **Supplemental Label for Table Grape**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### **FOLIAR APPLICATION INSTRUCTIONS**

<b>Crop</b>	<b>Amount of product in fl. oz./10-gals water</b>	<b>Application timing</b>	<b>Remarks</b>
Table grape	1.2-6.4	1 <sup>st</sup> . At bud break 2 <sup>nd</sup> At small fruit sizing 3 <sup>rd</sup> At berry softening	Tank-mix with GA3 Enhance ripening*

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in possession of the user at the time of application.

Read the label affixed to the container for Sunergist<sup>®</sup> EBR before applying.

Use of Sunergist<sup>®</sup> EBR according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Sunergist<sup>®</sup> EBR.

# **Sunergist<sup>®</sup> EBR**

Plant growth regulator

EPA Reg. No. 92983-2

## **Supplemental Label for Pome Fruits**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### **FOLIAR APPLICATION INSTRUCTIONS**

<b>Crop</b>	<b>Amount of product in fl. oz./10-gals water</b>	<b>Application timing</b>	<b>Remarks</b>
<b>Pome Fruits (Crop Group 11-10)</b> including Apple, Azarole, Crabapple, Loquat, Mahaw, Hook & Arn., Medlar, Pear, Pear (Asian), Pseudocydonia sinensis, Quince, Quince (Chinese, Japanese), Tejocote, and Cultivars, varieties and/or hybrids of these	0.64 – 6.4	1 <sup>st</sup> . Early flowering  2 <sup>nd</sup> . 7 – 10 days after petal fall  3 <sup>rd</sup> . Sizing stage  4 <sup>th</sup> . Ripening stage	

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in possession of the user at the time of application.

Read the label affixed to the container for Sunergist<sup>®</sup> EBR before applying.

Use of Sunergist<sup>®</sup> EBR according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Sunergist<sup>®</sup> EBR.