



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

**EPA Reg. Number:**

93236-2

**Date of Issuance:**

2/16/18

**NOTICE OF PESTICIDE:**

Registration  
 Reregistration  
(under FIFRA, as amended)

**Term of Issuance:**

Unconditional

**Name of Pesticide Product:**

Roundup Custom for Aquatic &  
Terrestrial Use

**Name and Address of Registrant (include ZIP Code):**

Stephen A. Adams  
Agent for Seamless Control LLC  
c/o Monsanto Company  
700 Chesterfield Parkway West  
Chesterfield, MO 63017

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Submit one copy of the revised final printed label for the record before you release the product for shipment.

*Continued on page 2*

**Signature of Approving Official:**

*Emily Schmid Jr.*

**Date:**

2/16/18

Reuben Baris, Product Manager 25  
Herbicide Branch, Registration Division (7505P)

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 09/27/2017
- Alternate CSF A dated 09/26/2017

If you have any questions, please contact Mindy Ondish by phone at 703-605-0723, or via email at [ondish.mindy@epa.gov](mailto:ondish.mindy@epa.gov).

Enclosure

# MASTER LABEL FOR EPA REG. NO. 93236-2

Primary Brand Name:

**Roundup Custom® for Aquatic & Terrestrial Use**

Alternate Brand Names:

(none)

## Master Label Table of Contents

I.	Container Labels	2 – 18
II.	Directions for Use for on Aquatic and Terrestrial Sites	19 – 72

\*\*See each label section for more detailed table of contents\*\*

**A C C E P T E D**

2/16/2018

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No.  
93236-2

## I. CONTAINER LABELS

### 1.0 EXTENDED CONTENT LABELS

[Extended content labels (ECL) have the complete directions for use attached as additional pages directly to a base label that is securely attached to the pesticide container.]

#### 1.1 Base Label for Extended Content Labels

[The following text represents the MINIMUM information that must be on the label that is securely attached to the primary pesticide container. Additional label information from the complete directions for use sections and the extended label contents included in this Master Label may be added to this base label as space allows.]

**[INSERT BRAND NAME]** [Logo optional]

GLYPHOSATE	GROUP	9	HERBICIDE
------------	-------	---	-----------

[Optional marketing claims and/or logos in this Master Label may be added to the Base Label, if space allows]

#### ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt..... 53.8%

OTHER INGREDIENTS: ..... 46.2%  
100.0%

\*Contains 648 grams of the active ingredient glyphosate, in the form of its isopropylamine salt, per liter or 5.4 pounds per U.S. gallon, which is equivalent to 480 grams of the acid, glyphosate, per liter or 4.0 pounds per U.S. gallon (39.9% by weight).

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

Keep out of reach of children

## CAUTION

[Optional base label text, if space allows: DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation could result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.]

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, socks and shoes.

[If room allows on container label, if not, reference to attached labeling: Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.]

[Optional base label text, as applicable: See attached labeling for additional [Optional text, as applicable: Precautions [and] information on Personal Protective Equipment].]

## I. CONTAINER LABELS

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to supplemental labeling under "AGRICULTURAL USE REQUIREMENTS" in the DIRECTIONS FOR USE section for information about this standard.

See Complete Directions for Use attached to this label for complete [Optional text, as applicable: Agricultural [and Non-Agricultural] Use Requirements of the Worker Protection Standard, Directions for Use [and] Limit of Warranty and Liability] [Optional label text, if applicable: in English and Spanish].

[Alternative label statement: See attached labeling booklet for complete [Optional label text, as applicable: Agricultural [and Non-Agricultural] Use Requirements of the Worker Protection Standard, Directions for Use and Limit of Warranty and Liability] [Optional label text, if applicable: in English and Spanish].]

### STORAGE AND DISPOSAL

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** STORE ABOVE 5°F (-15°C) TO KEEP PRODUCT FROM CRYSTALLIZING. Crystals will settle to the bottom. If allowed to crystallize, warm to 68°F (20°C) to redissolve and roll or shake container, or recirculate contents of larger containers, to mix well before using. Store pesticides away from food, pet food, feed, seed, fertilizers and veterinary supplies. Keep container closed to prevent spills and contamination.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

**CONTAINER HANDLING AND DISPOSAL:** [Insert appropriate Container Handling and Disposal Statement and Refilling Limitation from the following options]

[CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID CONTAINERS OF LESS THAN 1-GALLON CAPACITY]

Nonrefillable container. Do not reuse or refill this container.

[Alternative container statement: Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in this container. Contact your state regulatory agency to determine allowable practices in your state.]

Triple rinse this container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and

## I. CONTAINER LABELS

recap. Shake for 10 seconds. Pour rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Once properly rinsed, some plastic [*Optional text: agricultural*] pesticide containers can be taken to a container collection site or picked up for recycling. [*Alternative container disposal statement:* Then offer the container for recycling, if available.]

[*Optional container disposal statement:* To find the nearest collection site, contact your chemical dealer [*Optional text: or INSERT COMPANY NAME at TELEPHONE NUMBER*] [*Optional text: or call 1-800-768-6387*] [*Alternative telephone numbers:* 1-800-ROUNDUP (1-800-768-6387)].]

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

### [CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID PLASTIC 2.5-GALLON CONTAINER AND OTHER NONREFILLABLE CONTAINERS OF GREATER THAN 1-GALLON, BUT EQUAL TO OR LESS THAN 5-GALLON CAPACITY]

Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in this container. Contact your state regulatory agency to determine allowable practices in your state.

[*Alternative container statement:* Nonrefillable container. Do not reuse or refill this container.]

Triple rinse or pressure rinse (or equivalent) this container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic [*Optional text: agricultural*] pesticide containers can be taken to a container collection site or picked up for recycling. [*Alternative container disposal statement:* Then offer the container for recycling, if available.]

[*Optional container disposal statement:* To find the nearest collection site, contact your chemical dealer [*Optional text: or INSERT COMPANY NAME at TELEPHONE NUMBER*] [*Optional text: or call 1-800-768-6387*] [*Alternative telephone numbers:* 1-800-ROUNDUP (1-800-768-6387)].]

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

## I. CONTAINER LABELS

**[CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID PLASTIC 30-GALLON CONTAINER AND OTHER NONREFILLABLE CONTAINERS OF GREATER THAN 5-GALLON CAPACITY]**

Nonrefillable container. Do not reuse or refill this container.

*[Alternative container statement: Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in this container. Contact your state regulatory agency to determine allowable practices in your state.]*

Triple rinse or pressure rinse (or equivalent) this container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank. *[Optional label text: For containers not equipped with pumping systems,] Fill the container 1/4 full with water. Replace and tighten closures. Tip the container on its side and roll it back and forth for 30 seconds, ensuring at least one complete revolution. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Repeat this procedure two more times.*

*[Alternative or additional triple rinsing instructions for large containers equipped with pumping systems: [Optional label text: For large containers equipped with pumping systems,] Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.]*

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic *[Optional text: agricultural]* pesticide containers can be taken to a container collection site or picked up for recycling. *[Alternative container disposal statement: Then offer the container for recycling, if available.]*

*[Optional container disposal statement: Some container manufacturers offer container recycling. See additional information regarding manufacturer recycling programs attached to this container, if available. If no recycling information is available on this container, contact your chemical dealer [Optional text: or INSERT COMPANY NAME at TELEPHONE NUMBER] [Optional text: or call 1-800-768-6387] [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)] to find the nearest recycling location.]*

*[Optional container disposal statement: To find the nearest collection site, contact your chemical dealer [Optional text: or INSERT COMPANY NAME at TELEPHONE NUMBER] [Optional text: or call 1-800-768-6387] [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)].]*

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

*[Optional container disposal statement: Return Properly Rinsed Container for Recycling – Call 1-800-768-6387] [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)]*

*[Optional additional container disposal statement: IBC EMPTY? – FREE CALL – 1-888-SCHUETZ (1-888-724-8389) www.schuetz.net/ticket; Schuetz ticket service]*

## I. CONTAINER LABELS

[Optional additional container disposal statement: FREE IBC PICKUP] [For continental USA and Canada only]

[Optional additional container disposal statement: RETURNnet SYSTEM – To return empty IBC's Email or Call – [www.returnnetsystem.com](http://www.returnnetsystem.com) – 1-888-758-SHIP – United States and Canada (1-888-758-7447 – IBCNA – Clarkston, Michigan – USA)]

### **[CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR ALL REFILLABLE CONTAINERS, EXCEPT TRANSPORT VEHICLES]**

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning this container before refilling is the responsibility of the refiller. Cleaning this container before final disposal is the responsibility of the person disposing of the container.

To clean this container before final disposal, empty the remaining contents from the container into application equipment or mix-tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Then offer the container for recycling, if available. [Optional container disposal statement: Some container manufacturers offer container recycling. See additional information regarding manufacturer recycling programs attached to this container, if available. If no recycling information is available on this container, contact your chemical dealer [Optional text: or **INSERT COMPANY NAME** at **TELEPHONE NUMBER**] [Optional text: or call 1-800-768-6387] [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)] to find the nearest recycling location.]

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

[Optional additional container disposal statement: IBC EMPTY? – FREE CALL – 1-888-SCHUETZ (1-888-724-8389) [www.schuetz.net/ticket](http://www.schuetz.net/ticket); Schuetz ticket service]

[Optional additional container disposal statement: FREE IBC PICKUP] [For continental USA and Canada only]

[Optional additional container disposal statement: RETURNnet SYSTEM – To return empty IBC's Email or Call – [www.returnnetsystem.com](http://www.returnnetsystem.com) – 1-888-758-SHIP – United States and Canada (1-888-758-7447 – IBCNA – Clarkston, Michigan – USA)]

[Optional container disposal statement: To obtain information about recycling refillable containers, [Optional text: contact **INSERT COMPANY NAME** at **TELEPHONE NUMBER**] [Optional text: [or] call 1-800-768-6387] [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)].]

[Optional container disposal statement: Return Properly Rinsed Container for Recycling – Call 1-800-768-6387] [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387).]

[Optional additional container label statements for the CUBE refillable packaging system only:

**CUBE** Refillable Delivery System

FEATURES INCLUDE:

- Automatic Venting
- Heavy duty one-way 2-inch camloc ball valve with protective shield door
- Complete coated steel protective enclosure
- Durable 4-way plastic pallet

## I. CONTAINER LABELS

Lift door to access one-way valve]

[Optional label text: For product information or assistance using this product call [Optional text, if applicable: toll-free] 1-800-332-3111 [Or insert alternative telephone number, as appropriate].]

EPA Reg. No. 93236-2

EPA Est. [Insert appropriate EPA establishment number or include checkbox for options: 524-IA-1; 524-LA-1; or Other; or blank space] [Optional text: (Establishment Number when entered here supersedes all others)]

[Alternative EPA establishment text: EPA Est. (L) 524-LA-1 or (M) 524-IA-1 Lot number prefix (L) or (M) indicate appropriate establishment number.] [This lot number relationship with the establishment number can be expanded or changed to include additional or add new producing sites as appropriate]

[Insert net contents, as appropriate: NET [insert appropriate value or leave blank line for refillable container] GAL [or other appropriate unit of measure]] [Alternative label text: NET CONTENTS]

[Optional text, if applicable: [INSERT BRAND NAME and Design [and] TRUEBLUE ADVANTAGE PROVEN RELIABLE SUPPORT and Design] are [registered] trademarks of Seamless Control LLC.]

[Optional text, if applicable: [Roundup, Roundup Custom and Design] are [registered] trademarks of Monsanto Technology LLC and used under license.]

[Optional text, if applicable: All other trademarks are the property of their respective owners.]

Packed for:

SEAMLESS CONTROL LLC  
14111 SCOTTS LAWN ROAD  
MARYSVILLE, OH 43041

©[YEAR]

[Insert LOT number or LOT number will be printed directly on the container]

[Insert print plate number and barcodes as applicable]

[Optional text: ATTACH COMPLETE DIRECTIONS FOR USE HERE]

[Optional text that will be hidden behind the attached Extended Content Label and only become visible if the ECL is removed: COMPLETE DIRECTIONS FOR USE HAVE BEEN REMOVED]

[Optional text: LABEL CONTINUED ON BACK]

[Optional label text: FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC AT (800) 424-9300]

### 1.2 Extended Content

[The following represents the complete labeling text that must appear on the additional pages attached to the container base label as an Extended Content Label with the complete directions for use.]

[FRONT PANEL]

**[INSERT BRAND NAME]** [Logo optional]

GLYPHOSATE	GROUP	9	HERBICIDE
------------	-------	---	-----------

[Optional marketing claims and/or logos on this Master Label may be added to the Front Panel]

## I. CONTAINER LABELS

A broad-spectrum postemergence herbicide for [Optional text, as applicable to the final printed labeling: aquatic and industrial, turf, ornamental, forestry, roadside, utility rights-of-way, select crop, and other listed terrestrial] weed control.

(For a complete list of [Optional text, as applicable: aquatic and terrestrial uses], see the Directions for Use section inside this [Alternative text: the attached] labeling.)

[Optional statement for limited-geography product distribution: This product is not registered in all states.] [Optional text: See inside attached labeling for more details.]

[Additional optional statements for limited-geography product distribution: For control of annual and perennial weeds in] [list states and county-level information, as appropriate, where product is registered and distributed]. [Optional text: \*County Distribution:] [Optional text: see inside attached labeling for details.] [Optional text: In [list states] this product is distributed [Optional text: only] in the counties listed below:] [list counties by state]

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, [Optional text: GREEN] STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

---

### ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt..... 53.8%

OTHER INGREDIENTS: ..... 46.2%  
100.0%

\*Contains 648 grams of the active ingredient glyphosate, in the form of its isopropylamine salt, per liter or 5.4 pounds per U.S. gallon, which is equivalent to 480 grams of the acid, glyphosate, per liter or 4.0 pounds per U.S. gallon (39.9% by weight).

---

[Optional label text that will be updated at the time of printing, if necessary: This product is protected by [Optional text: one or more of the following] U.S. Patent No.'s [List appropriate patents].]

[Optional label text, if applicable: Other Patents Pending.]

[Optional label text, if applicable: No license granted under any non-U.S. patent(s).]

EPA Reg. No. 93236-2

EPA Est. [Insert appropriate EPA establishment number: 524-IA-1; 524-LA-1; or Other; or blank space] [Optional text: (Establishment Number when entered here supersedes all others)]

[Alternative EPA establishment text: EPA Est. (L) 524-LA-1 or (M) 524-IA-1 Lot number prefix (L) or (M) indicate appropriate establishment number.] [This lot number relationship with the establishment number can be expanded or changed to include additional or add new producing sites as appropriate]

---

Keep out of reach of children

## CAUTION

[Optional label statement, if applicable: See inside [Alternative text: back panel] for additional Precautions.]

See inside for Complete Directions for Use [Optional text: in English and Spanish].

[Alternative label statement: See attached labeling for Complete Directions for Use [Optional text: in English and Spanish].]

Read the entire label before using this product.

## I. CONTAINER LABELS

Use only according to label directions.

Read the LIMIT OF WARRANTY AND LIABILITY statement [Optional text, if statement not included on the container label: at the end of the attached labeling] before buying or using. If terms are not acceptable, return at once unopened.

[Insert net contents, as appropriate: NET [insert appropriate value or leave blank line for refillable container] GAL [or other appropriate unit of measure]] [Alternative text: NET CONTENTS]

Packed for:

SEAMLESS CONTROL LLC  
14111 SCOTTS LAWN ROAD  
MARYSVILLE, OH 43041

©[YEAR]

[Insert date]

[Insert print plate number and barcode(s) as applicable]

[Optional text: Open Here]

[EXTENDED CONTENT]

[For Extended Content Label, insert Complete Directions for Use, beginning with the Table of Contents from Part II of this Master Label, as appropriate]

## 2.0 SECONDARY PACKAGING FOR SHIPPING AND SALE

[This section defines labeling for secondary packaging of multiple units of this product that are themselves fully labeled for individual sale, but are distributed and can also be sold as a multi-pack unit.]

### 2.1 Secondary Packaging for Two 2.5-Gallon Nonrefillable Containers

[FRONT PANEL]

**[INSERT BRAND NAME]** [Logo optional]

GLYPHOSATE	GROUP	9	HERBICIDE
------------	-------	---	-----------

[Optional marketing claims and/or logos included on this Master Label may be added to this secondary packaging]

A broad-spectrum postemergence herbicide for [Optional text, as applicable to final printed labeling: aquatic and industrial, turf, ornamental, forestry, roadside, utility rights-of-way, select crop, and other listed terrestrial] weed control.

(For a complete list of [Optional text, as applicable to final printed labeling: aquatic and terrestrial uses], see the Directions for Use section of the attached labeling [Optional text: booklet] inside.)

[Optional label text: FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC AT (800) 424-9300]

[Optional statement for limited-geography product distribution: This product is not registered in all states.]  
[Optional text: See labeling [Optional text: booklet] inside for more details.]

[Additional optional statements for limited-geography product distribution: For control of annual and perennial weeds in [list states and county-level information as appropriate where product is registered and distributed]. [Optional text: \*County Distribution:] [Optional text: see labeling [Optional text: booklet]]

## I. CONTAINER LABELS

inside for details.] [Optional text: In [list states] this product is distributed [Optional text: only] in the counties listed below:] [list counties by state]

---

### ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt..... 53.8%

OTHER INGREDIENTS: ..... 46.2%

100.0%

\*Contains 648 grams of the active ingredient glyphosate, in the form of its isopropylamine salt, per liter or 5.4 pounds per U.S. gallon, which is equivalent to 480 grams of the acid, glyphosate, per liter or 4.0 pounds per U.S. gallon (39.9% by weight).

---

[Optional label text that will be updated at the time of printing, if necessary: This product is protected by [Optional text: one or more of the following] U.S. Patent No.'s [List appropriate patents].]

[Optional label text, if applicable: Other Patents Pending.]

[Optional label text, if applicable: No license granted under any non-U.S. patent(s).]

EPA Reg. No. 93236-2

EPA Est. [Insert appropriate pesticide establishment number: 524-IA-1; 524-LA-1; or Other]

[Alternative EPA establishment text: EPA Est. (L) 524-LA-1 or (M) 524-IA-1 Lot number prefix (L) or (M) indicate appropriate establishment number.] [This lot number relationship with the establishment number can be expanded or changed to include additional or add new producing sites as appropriate]

---

Keep out of reach of children

## CAUTION

[Optional label text, if applicable: See side panel for additional Precautions].

See labeling booklet inside for Complete Directions for Use [Optional text: in English and Spanish].

[Alternative label text: See attached labeling inside for Complete Directions for Use [Optional text: in English and Spanish].

Read the entire label before using this product.

Use only according to label directions.

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, [Optional text: GREEN] STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

Read the LIMIT OF WARRANTY AND LIABILITY statement [Optional text, if applicable: on side panel] before buying or using. If terms are not acceptable, return at once unopened.

Two 2.5 GAL Units

Packed for:

SEAMLESS CONTROL LLC  
14111 SCOTTS LAWN ROAD  
MARYSVILLE, OH 43041

©[YEAR]

[Insert print plate number and barcode(s) as applicable]

## I. CONTAINER LABELS

[ADDITIONAL CONTENT]

---

### PRECAUTIONARY STATEMENTS

---

Hazards to Humans and Domestic Animals

---

Keep out of reach of children

## CAUTION

**DOMESTIC ANIMALS:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation could result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

---

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, socks and shoes.

Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

---

### User Safety Recommendations

Users should:

- Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

---

### Environmental Hazards

---

*[For labels without aquatic uses: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters and rinsate.]*

*[Alternative text for labels with aquatic uses only: Killing aquatic weeds can result in depletion or loss of oxygen in the water due to decomposition of dead plant material. This oxygen loss can cause fish suffocation. Consult with your State agency with primary responsibility for regulating pesticides before applying to public waters to determine if a permit is required. Do not contaminate water when cleaning equipment or disposing of equipment wash waters and rinsate.]*

*[Alternative text for labels with both aquatic and terrestrial uses: Killing aquatic weeds can result in depletion or loss of oxygen in the water due to decomposition of dead plant material. This oxygen loss can cause fish suffocation. Consult with your State agency with primary responsibility for regulating pesticides before applying to public waters to determine if a permit is required. For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark [Optional text, if applicable: , except when applying this product by air over the forest canopy]. Do not contaminate water when cleaning equipment or disposing of equipment wash waters and rinsate.]*

## I. CONTAINER LABELS

### Physical or Chemical Hazards

Spray solutions of this product may be mixed, stored and applied using stainless steel, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas, which can form a highly combustible gas mixture. This gas mixture could flash or explode if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source and cause serious personal injury.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product may only be used in accordance with the Directions for Use on this label or in separately published supplemental labeling. Supplemental labeling for this product can be obtained from your Authorized Seamless Control Retailer or Representative.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### Agricultural Use Requirements

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to supplemental labeling under "AGRICULTURAL USE REQUIREMENTS" in the DIRECTIONS FOR USE section for information about this standard.

#### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

See labeling booklet inside for complete [Optional text, as applicable]: Agricultural [and] Non-Agricultural Use Requirements of the Worker Protection Standard and Directions for Use [and] Limit of Warranty and Liability [Optional label text: in English and Spanish].

[Alternative label statement: See attached labeling inside for complete] [Optional label text, applicable: Agricultural [and] Non-Agricultural Use Requirements of the Worker Protection Standard, Directions for Use, [and] Limit of Warranty and Liability] [Optional text: in English and Spanish].

#### STORAGE AND DISPOSAL

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** STORE ABOVE 5°F (-15°C) TO KEEP PRODUCT FROM CRYSTALLIZING. Crystals will settle to the bottom. If allowed to crystallize, warm to 68°F (20°C) to redissolve and roll or shake container, or recirculate contents of larger containers, to mix well before using. Store pesticides

## I. CONTAINER LABELS

away from food, pet food, feed, seed, fertilizers and veterinary supplies. Keep container closed to prevent spills and contamination.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in the container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

**CONTAINER HANDLING AND DISPOSAL:** [NOTE: Container Handling and Disposal Statement must match exactly the statement that is on the 2.5-gallon container inside the packaging.]

Nonrefillable container. Do not reuse the container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state.

[Alternative container statement: Nonrefillable container. Do not reuse or refill the container.]

Triple rinse or pressure rinse (or equivalent) the container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic [Optional text: agricultural] pesticide containers can be taken to a container collection site or picked up for recycling. [Alternative container disposal statement: Then offer the container for recycling, if available.]

[Optional container disposal statement: To find the nearest collection site, contact your chemical dealer [Optional text: or INSERT COMPANY NAME at TELEPHONE NUMBER] [Optional text: or call 1-800-768-6387] [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)].]

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

For product information or assistance using this product, call [Optional text, if applicable: toll-free] 1-800-332-3111 [Or insert alternative telephone number, as appropriate].

## LIMIT OF WARRANTY AND LIABILITY

Seamless Control LLC ("Company") warrants that this product conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

## I. CONTAINER LABELS

Buyer and all users shall use this product only for the purposes of and in accordance with the Complete Directions for Use label ("Directions") and shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss, injuries or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, crop injury or failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, use and/or application in any manner not explicitly set forth in or inconsistent with the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

---

*[Optional text, if applicable: [INSERT BRAND NAME and Design [and] TRUEBLUE ADVANTAGE PROVEN RELIABLE SUPPORT and Design] are [registered] trademarks of Seamless Control LLC.]*

*[Optional text, if applicable: [Roundup, Roundup Custom and Design] are [registered] trademarks of Monsanto Technology LLC and used under license.]*

*[Optional text, if applicable: All other trademarks are the property of their respective owners.]*

*[Insert UN packaging certification, if applicable]*

*[Optional packaging text: Pull at scored pullouts for easier opening]*

*[Optional packaging text: THIS SIDE UP] [Optional graphic of two arrows pointing upward]*

*[Optional packaging text: Liquid] [Optional graphic of a liquid droplet]*

## 3.0 TRANSPORT VEHICLE LABEL (AS DEFINED AT 40 CFR § 156.3)

**[INSERT BRAND NAME]** *[Logo optional]*

GLYPHOSATE	GROUP	9	HERBICIDE
------------	-------	---	-----------

*[Optional marketing claims and/or logos included on this Master Label may be added to this label]*

## I. CONTAINER LABELS

A broad-spectrum postemergence herbicide for [Optional text, as applicable to the final printed labeling: aquatic and industrial, turf, ornamental, forestry, roadside, utility rights-of-way, select crop, and other listed terrestrial] weed control.

[Optional label text: FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMREC AT (800) 424-9300]

[Optional statement for limited-geography product distribution: This product is not registered in all states.]

FOR BULK PESTICIDE TRANSPORT ONLY

---

### ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt..... 53.8%

OTHER INGREDIENTS: ..... 46.2%

100.0%

\*Contains 648 grams of the active ingredient glyphosate, in the form of its isopropylamine salt, per liter or 5.4 pounds per U.S. gallon, which is equivalent to 480 grams of the acid, glyphosate, per liter or 4.0 pounds per U.S. gallon (39.9% by weight).

[Optional label text that will be updated at the time of printing, if applicable: This product is protected by [Optional text: one or more of the following] U.S. Patent No.'s [List appropriate patents].]

[Optional label text, if applicable: Other Patents Pending.]

[Optional label text, if applicable: No license granted under any non-U.S. patent(s).]

EPA Reg. No. 93236-2

EPA Est. [Insert appropriate EPA establishment number: 524-IA-1; 524-LA-1; or Other; or blank space] [Optional text: (Establishment Number when entered here supersedes all others)]

[Alternative EPA establishment text: EPA Est. (L) 524-LA-1 or (M) 524-IA-1 Lot number prefix (L) or (M) indicate appropriate establishment number.] [This lot number relationship with the establishment number can be expanded or changed to include additional or add new producing sites as appropriate]

---

## PRECAUTIONARY STATEMENTS

---

### Hazards to Humans and Domestic Animals

---

Keep out of reach of children

## CAUTION

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation could result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

---

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, socks and shoes.

Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

## I. CONTAINER LABELS

### User Safety Recommendations

Users should:

- Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### Environmental Hazards

*[For labels without aquatic uses: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters and rinsate.]*

*[Alternative text for labels with aquatic uses only: Killing aquatic weeds can result in depletion or loss of oxygen in the water due to decomposition of dead plant material. This oxygen loss can cause fish suffocation. Consult with your State agency with primary responsibility for regulating pesticides before applying to public waters to determine if a permit is required. Do not contaminate water when cleaning equipment or disposing of equipment wash waters and rinsate.]*

*[Alternative text for labels with both aquatic and terrestrial uses: Killing aquatic weeds can result in depletion or loss of oxygen in the water due to decomposition of dead plant material. This oxygen loss can cause fish suffocation. Consult with your State agency with primary responsibility for regulating pesticides before applying to public waters to determine if a permit is required. For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark *[Optional text, if applicable: , except when applying this product by air over the forest canopy]*. Do not contaminate water when cleaning equipment or disposing of equipment wash waters and rinsate.]*

### Physical or Chemical Hazards

Spray solutions of this product may be mixed, stored and applied using stainless steel, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas, which can form a highly combustible gas mixture. This gas mixture could flash or explode if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source and cause serious personal injury.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and

## I. CONTAINER LABELS

exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear coveralls, socks and shoes, and waterproof gloves.

### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

### FOR BULK PESTICIDE TRANSPORT ONLY

#### STORAGE AND DISPOSAL

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** STORE ABOVE 5°F (-15°C) TO KEEP PRODUCT FROM CRYSTALLIZING. Crystals will settle to the bottom. If allowed to crystallize, warm to 68°F (20°C) to redissolve and roll or shake container, or recirculate contents of larger containers, to mix well before using. Store pesticides away from food, pet food, feed, seed, fertilizers and veterinary supplies. Keep container closed to prevent spills and contamination.

**PESTICIDE DISPOSAL:** To avoid wastes, empty as much product from this transport vehicle as possible for repackaging or use in accordance with label directions. If wastes cannot be avoided, offer remaining product or rinsate to a waste disposal facility or pesticide disposal program. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

**CONTAINER HANDLING AND DISPOSAL:** Emptied transport vehicle container retains vapor and product residue. Observe all precautions stated on this label until the container is cleaned, reconditioned or destroyed. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, and worn-out threads and closures. Clean thoroughly before reuse for transportation of a material of different composition or before retiring this transport vehicle container from service.

### THIS LABEL FOR USE WITH TRANSPORT VEHICLES ONLY

[Optional text, if applicable: [INSERT BRAND NAME and Design [and] TRUEBLUE ADVANTAGE PROVEN RELIABLE SUPPORT and Design] are [registered] trademarks of Seamless Control LLC.]

[Optional text, if applicable: [Roundup, Roundup Custom and Design] are [registered] trademarks of Monsanto Technology LLC and used under license.]

[Optional text, if applicable: All other trademarks are the property of their respective owners.]

Packed for:

SEAMLESS CONTROL LLC  
14111 SCOTTS LAWN ROAD  
MARYSVILLE, OH 43041

## I. CONTAINER LABELS

©[YEAR]

NET [*Insert net contents or blank space*] GAL [*or other unit of measure, as appropriate*]  
[Alternative label statement: NET CONTENTS:] [*Insert net contents or blank space*] GAL  
[Alternative label statement: NET CONTENTS: See Bill of Lading]

LOT [*Insert Lot number or blank space*]  
[Alternative label statement: LOT: See Bill of Lading]

[Alternative label statement: For Net Contents and Lot Number, see the Bill of Lading]

[*Insert print plate number and barcode(s) as applicable*]

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

[INSERT BRAND NAME] [Logo]

### Complete Directions for Use

GLYPHOSATE	GROUP	9	HERBICIDE
------------	-------	---	-----------

EPA Reg. No. 93236-2

A broad-spectrum postemergence herbicide for [Optional text, as applicable to the final printed labeling: aquatic and industrial, turf, ornamental, forestry, roadside, utility rights-of-way, select crop, and other listed terrestrial] weed control.

(For a complete list of [Optional text, as applicable to the final printed labeling: aquatic and terrestrial] use sites, see the Directions for Use section of this label.)

[Optional label text: FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC AT (800) 424-9300]

[Optional statement for limited-geography product distribution: This product is not registered in all states.]

[Additional optional statements for limited-geography product distribution: For control of annual and perennial weeds in [list states and or county-level information, as appropriate, where product is registered and distributed]. [Optional text: \*County Distribution:] [Optional text: see inside for details.] [Optional text: In [list states] this product is distributed [Optional text: only] in the counties listed below:] [list counties by state]

[Additional optional statement for limited-geography product distribution: This product is not registered for use in] [list states, if any, where this product is not registered for use]

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, [Optional text: GREEN] STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

[Optional label statement: TrueBlue Advantage – Proven – Reliable – Supported® (Logo)]

Read the entire label before using this product.

Use only according to label directions.

Not all products listed on this label are registered for use in California. Check the registration status of each product in California before using.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of this labeling before buying or using. If terms are not acceptable, return at once unopened.

### 1 1.0 INGREDIENTS

### 2 2.0 IMPORTANT PHONE NUMBERS

### 3 3.0 PRECAUTIONARY STATEMENTS

- 3.1 Hazards to Humans and Domestic Animals
- 3.2 Environmental Hazards
- 3.3 Physical or Chemical Hazards

### 4 4.0 STORAGE AND DISPOSAL

### 5 5.0 PRODUCT INFORMATION

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

### **6 6.0 WEED RESISTANCE MANAGEMENT**

- 6.1 Weed Management Practices
- 6.2 Management of Glyphosate-Resistant Biotypes

### **7 7.0 MIXING**

- 7.1 Mixing with Water
- 7.2 Surfactant
- 7.3 Tank Mixtures
- 7.4 Tank-Mixing Procedure
- 7.5 Mixing Spray Solution Concentrations
- 7.6 Ammonium Sulfate [*This section optional in final printed label*]
- 7.7 Colorants and Dyes
- 7.8 Drift Reduction Additives

### **8 8.0 APPLICATION EQUIPMENT AND TECHNIQUES**

- 8.1 Spray Drift Management
- 8.2 Aerial Application Equipment
- 8.3 Ground Application Equipment
- 8.4 Handheld Sprayers
- 8.5 Selective Application Equipment
- 8.6 Injection Systems
- 8.7 Controlled Droplet Applicator (CDA)

### **9 9.0 AQUATIC AND TERRESTRIAL USE SITES**

- 9.1 Aquatic Sites
- 9.2 Terrestrial Sites

### **10 10.0 ADDITIONAL SITE MANAGEMENT INFORMATION**

- 10.1 Fallow Cropland Management
- 10.2 Forestry, Hardwood and Christmas Tree Site Management
- 10.3 Grass Seed and Sod Production
- 10.4 Ornamental and Production Nursery Management
- 10.5 Commercial, Residential and Recreational Area Management
- 10.6 Railroad Management
- 10.7 Roadside Management
- 10.8 Utility Management

### **11 11.0 CITRUS TREE CROPS**

### **12 12.0 SUGARCANE**

### **13 13.0 WEEDS CONTROLLED**

- 13.1 Weed Control, Renovation and Chemical Mowing in Turf
- 13.2 Annual Weeds
- 13.3 Perennial Weeds
- 13.4 Woody Brush, Trees and Vines

### **14 14.0 LIMIT OF WARRANTY AND LIABILITY**

## **1.0 INGREDIENTS**

### ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt..... 53.8%

OTHER INGREDIENTS: ..... 46.2%

100.0%

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

\*Contains 648 grams of the active ingredient glyphosate, in the form of its isopropylamine salt per liter, or 5.4 pounds per U.S. gallon, which is equivalent to 480 grams of the acid, glyphosate, per liter or 4.0 pounds per U.S. gallon (39.9% by weight).

---

[Optional label text that will be updated at the time of printing, if applicable: This product is protected by [Optional text: one or more of the following] U.S. Patent No.'s [List appropriate patents].]

[Optional label text, if applicable: Other Patents Pending.]

[Optional label text, if applicable: No license granted under any non-U.S. patent(s).]

---

### 2.0 IMPORTANT PHONE NUMBERS

1. FOR PRODUCT INFORMATION OR ASSISTANCE USING THIS PRODUCT, CALL [Optional text, if applicable: TOLL-FREE, COLLECT, DAY OR NIGHT],  
(800) 332-3111 [Or insert alternative telephone number, as appropriate]
2. IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT OR FOR MEDICAL ASSISTANCE, CALL [Optional text, if applicable: TOLL-FREE, COLLECT, DAY OR NIGHT],  
(314) 694-4000 [Or insert alternative telephone number, as appropriate]

### 3.0 PRECAUTIONARY STATEMENTS

#### 3.1 Hazards to Humans and Domestic Animals

Keep out of reach of children

## CAUTION

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation could result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

---

#### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, socks and shoes.

Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations:

Users should:

- Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### 3.2 Environmental Hazards

Killing aquatic weeds can result in depletion or loss of oxygen in the water due to decomposition of dead plant material. This oxygen loss can cause fish suffocation. Consult with your State agency with primary

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

responsibility for regulating pesticides before applying to public waters to determine if a permit is required. For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark [Optional text, if applicable: , except if applying aerially over the forest canopy]. Do not contaminate water when cleaning equipment or disposing of equipment wash waters and rinsate.

[Alternative text for labels with aquatic uses only: Killing aquatic weeds can result in depletion or loss of oxygen in the water due to decomposition of dead plant material. This oxygen loss can cause fish suffocation. Consult with your State agency with primary responsibility for regulating pesticides before applying to public waters to determine if a permit is required. Do not contaminate water when cleaning equipment or disposing of equipment wash waters and rinsate.]

### 3.3 Physical or Chemical Hazards

Spray solutions of this product may be mixed, stored and applied using stainless steel, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas, which can form a highly combustible gas mixture. This gas mixture could flash or explode if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source and cause serious personal injury.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product may only be used in accordance with the Directions for Use on this label or on separately published supplemental labeling. Supplemental labeling for this product can be obtained from your Authorized Seamless Control Retailer or Representative.

*[The following paragraph is reserved and will only appear on final printed labeling for products under this registration if and when required by EPA:]*

**ENDANGERED SPECIES PROTECTION REQUIREMENTS:** This product may have effects on federally listed threatened or endangered species or their critical habitat in some locations. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.]

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear coveralls, socks and shoes, and waterproof gloves.

### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

## 4.0 STORAGE AND DISPOSAL

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage and disposal.

**PESTICIDE STORAGE:** STORE ABOVE 5°F (-15°C) TO KEEP PRODUCT FROM CRYSTALLIZING. Crystals will settle to the bottom. If allowed to crystallize, warm to 68°F (20°C) to redissolve and roll or shake container or recirculate contents of larger containers to mix well before using. Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination. See individual container label for additional storage conditions, if any.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in the container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

**CONTAINER HANDLING AND DISPOSAL:** [Optional label text, if applicable: See container label for container handling and disposal instructions and refilling limitations.] [Alternative optional text for ECL label: See base label attached to the container for container handling and disposal instructions and refilling limitation.]

[CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID CONTAINERS OF LESS THAN 1-GALLON CAPACITY]

Nonrefillable container. Do not reuse or refill the container.

[Alternative container statement: Nonrefillable container. Do not reuse the container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state.]

Triple rinse the container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix-tank, or store rinsate for

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Once properly rinsed, some plastic [Optional text: agricultural] pesticide containers can be taken to a container collection site or picked up for recycling. [Alternative container disposal statement: Then offer the container for recycling, if available.]

[Optional container disposal statement: To find the nearest collection site, contact your chemical dealer [Optional text: or INSERT COMPANY NAME at TELEPHONE NUMBER] [Optional text: or call 1-800-768-6387] [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)].]

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

### [CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID PLASTIC 2.5-GALLON CONTAINER AND OTHER NONREFILLABLE CONTAINERS OF GREATER THAN 1-GALLON, BUT EQUAL TO OR LESS THAN 5-GALLON CAPACITY]

Nonrefillable container. Do not reuse the container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state.

[Alternative container statement: Nonrefillable container. Do not reuse or refill the container.]

Triple rinse or pressure rinse (or equivalent) the container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic [Optional text: agricultural] pesticide containers can be taken to a container collection site or picked up for recycling. [Alternative container disposal statement: Then offer this container for recycling, if available.]

[Optional container disposal statement: To find the nearest collection site, contact your chemical dealer [Optional text: or INSERT COMPANY NAME at TELEPHONE NUMBER] [Optional text: or call 1-800-768-6387] [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)].]

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

**[CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID PLASTIC 30-GALLON CONTAINER AND OTHER NONREFILLABLE CONTAINERS OF GREATER THAN 5-GALLON CAPACITY]**

Nonrefillable container. Do not reuse or refill the container.

*[Alternative container statement: Nonrefillable container. Do not reuse the container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state.]*

Triple rinse or pressure rinse (or equivalent) the container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank. *[Optional label text: For containers not equipped with pumping systems,] Fill the container 1/4 full with water. Replace and tighten closures. Tip the container on its side and roll it back and forth for 30 seconds, ensuring at least one complete revolution. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Repeat this procedure two more times.*

*[Alternative or additional triple rinsing instructions for large containers equipped with pumping systems: [Optional label text: For large containers equipped with pumping systems,] Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.]*

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic *[Optional text: agricultural]* pesticide containers can be taken to a container collection site or picked up for recycling. *[Alternative container disposal statement: Then offer the container for recycling, if available.]*

*[Optional container disposal statement: Some container manufacturers offer container recycling. See additional information regarding manufacturer recycling programs attached to the container, if available. If no recycling information is available on the container, contact your chemical dealer [Optional text: or INSERT COMPANY NAME at TELEPHONE NUMBER] [Optional text: or call 1-800-768-6387] [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)] to find the nearest recycling location.]*

*[Optional container disposal statement: To find the nearest collection site, contact your chemical dealer [Optional text: or INSERT COMPANY NAME and TELEPHONE NUMBER] [Optional text: or call 1-800-768-6387] [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)].]*

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

*[Optional container disposal statement: Return Properly Rinsed Container for Recycling – Call 1-800-768-6387 [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)]*

*[Optional additional container disposal statement: IBC EMPTY? – FREE CALL – 1-888-SCHUETZ (1-888-724-8389) www.schuetz.net/ticket; Schuetz ticket service]*

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

[Optional additional container disposal statement: FREE IBC PICKUP] [For continental USA and Canada only]

[Optional additional container disposal statement: RETURNnet SYSTEM – To return empty IBC's Email or Call – [www.returnnetsystem.com](http://www.returnnetsystem.com) – 1-888-758-SHIP – United States and Canada (1-888-758-7447 – IBCNA – Clarkston, Michigan – USA)]

### [CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR ALL REFILLABLE CONTAINERS, EXCEPT TRANSPORT VEHICLES]

Refillable container. Refill the container with pesticide only. Do not reuse the container for any other purpose.

Cleaning the container before refilling is the responsibility of the refiller. Cleaning the container before final disposal is the responsibility of the person disposing of the container.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix-tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Then offer this container for recycling, if available. [Optional container disposal statement: Some container manufacturers offer container recycling. See additional information regarding manufacturer recycling programs attached to the container, if available. If no recycling information is available on the container, contact your chemical dealer [Optional text: or **INSERT COMPANY NAME** at **TELEPHONE NUMBER**] [Optional text: or call 1-800-768-6387] [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)] to find the nearest recycling location.]

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

[Optional additional container disposal statement: IBC EMPTY? – FREE CALL – 1-888-SCHUETZ (1-888-724-8389) [www.schuetz.net/ticket](http://www.schuetz.net/ticket); Schuetz ticket service]

[Optional additional container disposal statement: FREE IBC PICKUP] [For continental USA and Canada only]

[Optional additional container disposal statements: RETURNnet SYSTEM – To return empty IBC's Email or Call – [www.returnnetsystem.com](http://www.returnnetsystem.com) – 1-888-758-SHIP – United States and Canada (1-888-758-7447 – IBCNA – Clarkston, Michigan – USA)]

[Optional container disposal statement: To obtain information about recycling refillable containers, [Optional text: contact **INSERT COMPANY NAME** at **TELEPHONE NUMBER**] [Optional text: [or] call 1-800-768-6387 [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)].]

[Optional container disposal statement: Return Properly Rinsed Container for Recycling – Call 1-800-768-6387 [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)]

## 5.0 PRODUCT INFORMATION

**Product Description:** This product is a postemergence, systemic herbicide that, when mixed in the spray tank with a surfactant that is approved for aquatic use, may be used for both aquatic and terrestrial weed control. This product provides broad-spectrum control of many annual and perennial weeds, woody brush, trees and vines. This product does not control submerged weeds or provide residual weed control in soil. It is formulated as a water-soluble liquid that, unless otherwise directed, requires dilution with

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

water or another carrier and the addition of a surfactant according to label directions and intended use site before application using standard and specialized pesticide application equipment.

**Mechanism of Action:** The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to the formation of specific amino acids.

**No Soil Activity:** This product binds tightly to soil particles and does not provide residual weed control. Weeds must be emerged at the time of application to be controlled by foliar application of this product. Weed seeds in the soil will not be affected by this product and will continue to germinate. Unattached plant rhizomes and rootstocks beneath the soil surface will also not be affected by this product.

**Biological Degradation:** Degradation of this product is primarily a biological process carried out by soil microbes.

**Stage of Weeds:** Aquatic weeds must have foliage above the water surface in order to be controlled by this product. On terrestrial sites, annual and perennial weeds are easiest to control when they are small. See the "WEEDS CONTROLLED" section of this label for more information on the control of specific weeds.

**Cultural Considerations:** Reduced weed control could result when this product is applied to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to re-grow prior to application. Always use the highest application rate of this product within a given range when weed growth is heavy or dense, or when weeds are growing in an undisturbed (non-cultivated) area. Reduced weed control could result when this product is applied to weeds that show signs of disease or insect damage, are covered with dust, or are surviving under poor growing conditions.

**Spray Coverage:** For best results with this product, spray coverage must be uniform and complete. Do not spray foliage to the point of runoff.

**Rainfastness:** Rainfall or submersion of aquatic weeds by wave action within 4 hours of application could wash this product off of the foliage and a second application might be needed to achieve acceptable weed control. Refer to specific use sections of this label for additional information on minimum intervals required, if any, before re-application of this product.

**Time to Symptoms:** This product moves through the plant from the point of foliage contact to and into the root system. Visible effects are a gradual wilting and yellowing of the plant that advances to complete browning of aboveground growth and deterioration of underground plant parts. Effects are visible on most annual weeds within 2 to 4 days, but on most perennial weeds, effects might not be visible for 7 or more days after application. Extremely cool or cloudy weather following application could slow activity of this product and delay development of visual symptoms.

**Maximum Application Rates:** The maximum application or use rates stated throughout this label are given in units of volume (fluid ounces or quarts) of this product per acre. However, the maximum allowable application rates apply to this product combined with the use of any and all other herbicides containing the active ingredient glyphosate, whether applied separately or in a tank mixture, on a basis of total pounds of glyphosate (acid equivalents) per acre. If more than one glyphosate-containing product is applied to the same site within the same year, you must ensure that the total use of glyphosate (pounds acid equivalents) does not exceed the maximum allowed. See the "INGREDIENTS" section of this label for necessary product information.

Unless otherwise specified on this label, the combined total of all applications of this product on a site must not exceed 8 quarts (8 pounds of glyphosate acid) per acre per year.

## 6.0 WEED RESISTANCE MANAGEMENT

GLYPHOSATE	GROUP	9	HERBICIDE
------------	-------	---	-----------

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mechanism of action classification system of the Weed Science Society of America. Any weed population can contain plants that are naturally resistant to Group 9 herbicides. Weeds resistant to Group 9 herbicides can be effectively managed by using another herbicide from a different Group (either alone or in a mixture according to label directions), by using other cultural or mechanical methods of weed control, or a combination of the two. Consult your local company representative, state cooperative extension agent, professional consultant or other qualified authority to determine appropriate actions for controlling specific resistant weeds.

### **6.1      Weed Management Practices**

Resistant populations arise when rare individual plants are uncontrolled by a normal dose of a given herbicide under normal environmental conditions. In the absence of other control measures these individuals survive, produce seed, and eventually become the dominant biotype in the field through continuous selection. The best means of reducing this selection is to use diverse weed control practices such as multiple herbicides with different mechanisms of action, and often in combination with various mechanical and cultural practices.

Suspected herbicide resistance can be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially when control is achieved on adjacent weeds
- A spreading patch of non-controlled plants of a particular species
- Surviving plants mixed with controlled individuals of the same species

To minimize the occurrence of herbicide-resistant biotypes, including those resistant to glyphosate, implement the following weed management practice options that are practical to your situation. These management practices are applicable to reduce the spread of suspected and confirmed resistant biotypes (managing existing resistant biotypes) and to reduce the potential for selecting for resistance in new species (proactive resistance management).

- Use a diversified approach toward weed management focused on preventing weed seed production and reducing the number of weed seeds in the soil.
- Plant crops into fields that are as weed-free as possible and then keep them as weed-free as possible.
- Plant seed that is as weed-free as possible.
- Scout fields and application sites routinely, before and after herbicide application.
- Use multiple herbicide mechanisms of action that are effective against the most troublesome weeds at your application site and against those with known resistance.
- Apply herbicides at application rates listed on the label when weeds are within the size range indicated on the label.
- Emphasize cultural practices that suppress weeds by using crop competitiveness.
- Use mechanical and biological weed management practices, where appropriate.
- Prevent field-to-field and within-field movement of weed seed or vegetative propagules.
- Manage weed seed at harvest and after harvest to prevent a buildup of the weed seedbank.

### **6.2      Management of Glyphosate-Resistant Biotypes**

Appropriate testing is needed to determine if a weed is resistant to glyphosate. Call 1-800-768-6387 [Alternative telephone numbers: 1-800-ROUNDUP (1-800-768-6387)] [Optional alternative text: *INSERT COMPANY NAME AND/OR TELEPHONE NUMBER*] [Optional alternative text: [or] contact your

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Seamless Control Representative] to report any incidence of non-performance of this product against a particular weed species or to determine if resistance in any particular weed biotype has been confirmed in your area [Optional additional text: , or visit on the Internet at [www.weedresistancemanagement.com](http://www.weedresistancemanagement.com) or [www.weedscience.org](http://www.weedscience.org)].

Glyphosate-resistant weeds can be controlled or managed by applying this product in combination with residual preemergence herbicides and/or other postemergence herbicides labeled for control of the targeted weed in the crop being grown or on the site of application. For more information, see the "WEEDS CONTROLLED" section of this label.

Since the occurrence of resistant weeds is difficult to detect prior to use, Seamless Control LLC accepts no liability for any losses that result from the failure of this product to control resistant weeds.

### 7.0 MIXING

Spray solutions of this product may be mixed, stored and applied using clean stainless steel, fiberglass, plastic or plastic-lined steel containers.

**DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.**

Eliminate any risk of siphoning the contents of the tank back into the carrier source while mixing. Use approved anti-back-siphoning devices where required by state or local regulations.

A 50-mesh nozzle screen or line strainer on spray equipment is adequate.

Clean sprayer parts promptly after using this product by thoroughly flushing with water.

#### 7.1 Mixing with Water

PERFORMANCE OF THIS PRODUCT CAN BE SIGNIFICANTLY REDUCED IF WATER CONTAINING SOIL SEDIMENT IS USED AS CARRIER. DO NOT MIX THIS PRODUCT WITH WATER FROM PONDS OR DITCHES THAT IS VISIBLY MUDDY OR MURKY.

This product mixes readily with water. Mix spray solutions of this product as follows. Begin filling the mixing tank or spray tank with clean water. Add the required amount of this product near the end of the filling process and mix gently. Foaming of the spray solution can occur during mixing. To prevent or minimize foaming, mix gently, terminate by-pass and return lines at the bottom of the tank and, if necessary, add an appropriate anti-foam or defoaming agent to the spray solution.

#### 7.2 Surfactant

Unless otherwise directed, this product requires the addition of 2 or more quarts of a nonionic surfactant that is labeled for use with herbicides [Optional label text, if applicable: and approved for aquatic use] per 100 gallons of spray solution (0.5% or more by volume). Unless otherwise directed, use a higher concentration of surfactant when any of the following conditions apply to the use of this product:

- Adding surfactants that contain less than 70 percent active ingredient
- Making a broadcast application using a high carrier volume or using handheld spray equipment
- Applying under adverse growing conditions or anytime weeds are under stress
- Applying as a tank-mix with other products
- Applying to hard-to-control weeds, woody brush, trees and vines

[Optional label text to be used if surfactants that are not approved for aquatic use are allowed with this product: **NOTE:** For direct application of spray solutions of this product on emerged aquatic weeds or for use in intertidal areas below the mean high-water mark, or in application areas where a buffer that will

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

ensure no overspray of an adjacent body of water cannot be maintained, a surfactant that is also approved for aquatic use must be used. For terrestrial applications, surfactant is also needed in the spray solution, but does not have to be approved for aquatic use.]

*[Optional label text to be used if surfactants that are not approved for aquatic use are allowed with this product: RESTRICTION: If a surfactant that is NOT approved for aquatic use is added to the spray solution, DO NOT apply directly to or over water or use in intertidal areas below the mean high-water mark.]*

Check with your local State agency with primary responsibility for regulating pesticides for additional information about surfactants that are approved for aquatic use.

Read and follow all precautionary statements and directions for use on the surfactant label.

All reference throughout this label to concentration of surfactant in the spray solution is on a percentage-of-volume basis. Refer to the table below to achieve the appropriate concentration of surfactant in the spray solution.

Desired Volume of Spray Solution	Amount of Surfactant to Achieve Indicated Concentration in Spray Solution (percent by volume)					
	0.5%	0.75%	1%	1.5%	4%	8%
1 gallon	2/3 fl oz	1 fl oz	1.3 fl oz	2 fl oz	5 fl oz	10 fl oz
25 gallons	16 fl oz	24 fl oz	1 qt	1.5 qts	4 qts	2 gals
100 gallons	2 qts	3 qts	1 gal	1.5 gals	4 gals	8 gals

2 tablespoons = 1 fluid ounce (fl oz)

### 7.3 Tank Mixtures

This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control in the soil, a broader weed control spectrum or an alternate mechanism of action.

NOT ALL TANK-MIX PRODUCTS LISTED ON THIS LABEL ARE APPROVED FOR USE ON AQUATIC SITES. Refer to each individual label for all products in the tank mixture for approved use sites and application rates.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Seamless Control has not tested all tank-mix product formulations for compatibility, antagonism or reduction in product performance. Mixing this product with herbicides or other materials not specified on this label could result in reduced performance of this product. To the extent consistent with applicable law, buyer and all users are responsible for any loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly specified on this label, or on separate supplemental labeling or Fact Sheets published for this product.

This product may be applied at any rate listed on this label in a tank mixture with the following products to provide preemergence and/or improved postemergence control of weeds listed on the individual product labels.

---

*[Active ingredients and products to be added to the final printed labeling for tank-mixing may be selected from the following list:*

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

2,4-D; atrazine; dicamba; bromacil; diuron; imazapyr; metsulfuron methyl; oryzalin; pendimethalin, prodiamine; simazine; sulfosulfuron; trichlopyr

Arsenal; Arsenal Herbicide Applicators Concentrate; Banvel; Banvel 480; Barricade 4L; Barricade 65WG; Certainty Turf; Chopper Gen2; Crossbow; Endurance; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XLT Specialty; Gallery SC; Gallery 75 Dry Flowable Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Goal 2XL; GoalTender; Habitat; Hyvar X; Hyvar X-L; Karmex DF; Krenite S Brush Control Agent; Krovar I DF; Landmark; Landmark XP; Oust Extra; Oust XP; Outrider; Plateau; Poast; Poast Plus; Ronstar 50 WSP; Ronstar Flo; Ronstar G; Sahara DG; Spike 20P Specialty; Spike 80 DF Specialty; Stalker; Surflan AS Specialty; Surflan Flex; Surflan Flex T&O; Surflan XL 2G; Surflan Pro; Telar XP; Tordon 101 Mixture Specialty; Tordon 22K Specialty; Tordon K Herbicide Specialty; Transline Specialty; Vanquish; Velpar DF CU; Velpar DF VU; Velpar L CU; Velpar L; Velpar L VU]

When used in combination as described on this label and to the extent consistent with applicable law, the liability of Seamless Control LLC shall in no manner extend to any damage, loss or injury not solely and directly caused by the inclusion of the Seamless Control product in such combination use.

### 7.4 Tank-Mixing Procedure

Always predetermine the compatibility of all tank-mix products in the carrier by mixing small proportional quantities in advance.

Add individual tank-mix components to the tank as follows: wettable powders; flowables; emulsifiable concentrates; drift reduction additives; water soluble liquids (this product); nonionic surfactants. Ensure that the tank-mix products are well mixed in the spray solution before adding this product.

Mix only the quantity of spray solution that will be applied that day. Application of tank-mix solutions that are allowed to stand overnight could result in reduced weed control.

Maintain gentle agitation at all times until the contents of the tank are sprayed out. If the spray mixture is allowed to settle, agitate thoroughly to resuspend the mixture before resuming application.

Keep by-pass line on or near the bottom of the tank to minimize foaming.

### 7.5 Mixing Spray Solution Concentrations

All reference throughout this label to concentration of this product in a spray solution is on a percentage-of-volume basis.

Prepare the desired volume of spray solution at a given concentration by mixing the amount of this product indicated in the following table with water.

Desired Volume of Spray Solution	Amount of [INSERT BRAND NAME] to Achieve Indicated Concentration in Spray Solution (percent by volume)					
	0.5%	0.75%	1%	1.5%	4%	8%
1 gallon	2/3 fl oz	1 fl oz	1.3 fl oz	2 fl oz	5 fl oz	10 fl oz
25 gallons	16 fl oz	24 fl oz	1 qt	1.5 qts	4 qts	2 gals
100 gallons	2 qts	3 qts	1 gal	1.5 gals	4 gals	8 gals

2 tablespoons = 1 fluid ounce (fl oz)

For filling backpack and pump-up sprayers, consider mixing the appropriate amount of this product with water in a larger container and then filling the sprayer from the larger container.

### 7.6 Ammonium Sulfate [This section optional in final printed label]

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Unless otherwise directed, the addition of 1 to 2 percent dry ammonium sulfate by weight (8.5 to 17 pounds per 100 gallons of water), could increase the performance of this product on annual and perennial weeds, particularly under hard water conditions, drought conditions or when tank-mixed with certain residual herbicides. An equivalent amount of a liquid formulation of ammonium sulfate may also be used. Ensure that dry ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water promptly after use to reduce corrosion.

When using ammonium sulfate, apply this product at rates or spray solution concentrations as directed on this label; lower the application rate or concentration could result in reduced weed control.

### 7.7 Colorants and Dyes

Colorants and marking dyes may be added to spray solutions of this product; however, they could reduce the performance of this product. Use colorants and dyes according to the manufacturer's directions.

### 7.8 Drift Reduction Additives

Drift reduction additives may be used with all application equipment types, except wiper applicators, sponge bars and controlled droplet applicators (CDA). When a drift reduction additive is used, read and follow all directions for use, precautions, limitations and all other information on the product label. The use of drift reduction additives can affect spray coverage, which could result in reduced performance of this product.

## 8.0 APPLICATION EQUIPMENT AND TECHNIQUES

This product may be applied using the following equipment:

**Aerial Application Equipment**—fixed-wing and helicopter [Alternative text: helicopter only] [Optional text: (Aerial application allowed by helicopter only in) [list states where aerial application is allowed by helicopter only, if applicable.]]

**Ground Application Equipment**—boom or boomless systems, pull-type sprayers, floaters, pick-up sprayers, spray coupes and other ground broadcast application equipment

**Handheld Sprayers**—backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers\*, lances and other handheld and motorized spray equipment used to direct the spray onto unwanted foliage

\*This product is not registered in California or Arizona for use in mistblowers.

**Selective Application Equipment**—recirculating sprayer, shielded and hooded sprayers, wiper applicator, sponge bar, single or hollow stem injectors, tree injector, spray bottle

**Injection Systems**—aerial or ground injection sprayers

**Controlled Droplet Applicator (CDA)**—handheld or boom-mounted applicators that produce a spray consisting of a narrow range of droplet sizes

APPLY THIS PRODUCT USING PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF ACCURATELY DELIVERING DESIRED VOLUMES.

Do not apply this product through any type of irrigation system.

### 8.1 Spray Drift Management

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, [Optional text: GREEN] STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation, as even small quantities of this product can cause severe damage or destruction to the crop, plants or other vegetation on which application was not intended.

**AVOID DRIFT. USE EXTREME CARE TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHEN APPLYING THIS PRODUCT.**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator and grower are responsible for considering all these factors when making decisions regarding the application of this product.

The likelihood of injury occurring as the result of spray drift while applying this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or generation of fine particles (mist) that are likely to drift.

**TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFERS MUST BE MAINTAINED.**

**AVOID APPLYING THIS PRODUCT AT EXCESSIVE SPEED OR SPRAYER PRESSURE.**

### **8.2 Aerial Application Equipment**

Unless otherwise prohibited, all broadcast applications of this product described on this label may be made using aerial application equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this label and on separate supplemental labeling published for this product.

**DO NOT APPLY THIS PRODUCT USING AERIAL APPLICATION EQUIPMENT EXCEPT UNDER CONDITIONS SPECIFIED ON THIS LABEL OR ON SEPARATELY PUBLISHED SUPPLEMENTAL LABELING FOR THIS PRODUCT.**

*[Optional voluntary label restriction: Aerial application of this product may be made using helicopters only.]*

*[Optional voluntary label restriction: Aerial application of this product may be made using helicopters only in] [list states where aerial application is permitted by helicopter only, if applicable].*

**FOR SPECIFIC USE INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS RELATED TO THE AERIAL APPLICATION OF THIS PRODUCT IN CALIFORNIA OR SPECIFIC COUNTIES THEREIN, REFER TO THE LIMITATIONS ON AERIAL APPLICATION IN THAT STATE OR COUNTY PRESENTED IN THIS SECTION.**

Apply this product at a rate specified on this label in 3 to 25 gallons of water per acre when using aerial application equipment, unless otherwise directed. Use a larger spray volume within this range where weeds, brush, trees and vines are dense or form multiple canopy layers.

Avoid direct application to any body of water.

Drift control reduction additives may be used.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

### **Aircraft Maintenance**

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES COULD RESULT IN CORROSION AND POSSIBLE FAILURE OF THE

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

PART. LANDING GEAR IS MOST SUSCEPTIBLE. Maintaining an organic coating (paint) that meets aerospace specification MIL-C-38413 can help prevent corrosion.

### AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to minimize off-target drift movement during aerial application. [Optional label text, if applicable: These requirements do not apply to forestry applications.]

1. The distance of the outermost nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the [Optional text, if application using fixed-wing aircraft allowed: wingspan or] rotor.
2. Nozzles must always point backwards parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they must be followed.

#### Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if the application is made improperly or under unfavorable environmental conditions, such as in windy, high temperature with low humidity, and/or inversion conditions as described below.

#### Controlling Droplet Size

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure:** Operate the sprayer at a pressure towards the lower end of the range listed for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing the pressure.
- **Number of nozzles:** Use the minimum number of nozzles that provides uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the air stream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom length:** For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the [Optional text, if application using fixed-wing aircraft allowed: wingspan or] rotor length can further reduce drift without reducing swath width.
- **Application height:** Application must be made at a height of 10 feet or less above the tallest plants in the field, unless a greater height is needed for aircraft safety. Making the application at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

#### Swath Adjustment

When an application is made in the presence of a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Increase the swath adjustment distance with increasing drift potential (higher wind, smaller droplets, etc.).

#### Wind

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Drift potential is lowest at wind speeds of between 2 and 10 miles per hour. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. Avoid application when wind speeds are below 2 miles per hour due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Applicators must be familiar with local wind patterns and how they affect drift.

### Temperature and Humidity

When making an application in low relative humidity, set application equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### Temperature Inversions

Do not apply this product during a temperature inversion as drift potential is high under these conditions. Temperature inversions restrict vertical air mixing, which causes small droplets to remain suspended in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### Sensitive Areas

Apply this product only when the potential for drift to adjacent sensitive non-target areas (e.g., residential areas, known habitat for threatened or endangered species, emerged crops or other desirable vegetation) is minimal (e.g., when wind is blowing away from a sensitive area).

### State Specific Limitations on Aerial Application

---

#### LIMITATIONS ON AERIAL APPLICATION IN CALIFORNIA ONLY

---

*[Optional voluntary restriction: Aerial application of this product in California may be made by helicopter only.]*

DO NOT apply this product using aerial application equipment in residential areas.

AVOID DRIFT – DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT OF THIS PRODUCT ONTO ANY VEGETATION TO WHICH APPLICATION WAS NOT INTENDED CAN CAUSE DAMAGE. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, USE PROPER AERIAL APPLICATION EQUIPMENT FITTED WITH APPROPRIATE NOZZLES AND MAINTAIN ADEQUATE BUFFERS.

Follow the directions below when making an aerial application near emerged crops, desirable annual vegetation, or desirable perennial vegetation after bud break and before total leaf drop.

1. Do not apply this product within 100 feet of all desirable vegetation or non-target crops.
2. If winds are blowing up to 5 miles per hour TOWARD desirable vegetation or emerged crops, do not apply this product within 500 feet of the desirable vegetation or crops.
3. If winds are blowing between 5 and 10 miles per hour TOWARD desirable vegetation or emerged crops, a buffer zone greater than 500 feet might be needed to protect the desirable vegetation or crops.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

4. Do not apply this product using aerial application equipment when winds are blowing in excess of 10 miles per hour.
5. Do not apply this product using aerial application equipment when inversion conditions exist.

When tank-mixing this product with 2,4-D, only 2,4-D amine formulations may be applied in California using aerial application equipment. Tank mixtures of this product with 2,4-D amine formulations may be applied by air in California on fallow fields and in reduced tillage systems only.

This product, when tank-mixed with dicamba, may not be applied by air in California.

---

### ADDITIONAL LIMITATIONS ON AERIAL APPLICATION IN FRESNO COUNTY, CALIFORNIA ONLY

---

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

The following information applies only from February 15 through March 31 within the following boundaries of Fresno County, California:

North: Fresno County line  
South: Fresno County line  
East: State Highway 99  
West: Fresno County line

Observe the following directions to minimize off-site movement during aerial application of this product. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor and aerial applicator.

#### **Written Directions**

Written directions MUST be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. These written directions MUST state the proximity of surrounding crops and that conditions of each manufacturer's product label and this label have been satisfied.

#### **Aerial Applicator Training and Equipment**

Aerial application of this product is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-ins constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

**Applications at Night** – Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

For additional information on the proper aerial application of this product in Fresno County, call (800) 332-3111 [Or insert alternative telephone number as appropriate].

---

### 8.3     **Ground Application Equipment**

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Apply this product at an appropriate rate specified on this label in 3 to 40 gallons [*Alternative volume: 10 to 60 gallons*] of water per acre when making a broadcast application using ground application equipment, unless otherwise directed on this label or on separate supplemental labeling or Fact Sheets published for this product. As the weed density increases, increase the spray volume toward the upper end of this range to ensure complete coverage. Use nozzles that will avoid generating a fine mist. For best performance of this product when using ground application equipment, use flat-fan nozzles. Check spray pattern for uniform distribution of spray droplets.

### 8.4 Handheld Sprayers

When using a handheld sprayer, apply spray solutions of this product uniformly and completely to the foliage of target weeds using a coarse droplet spectrum and a spray-to-wet technique; do not spray to the point of runoff. For the appropriate concentration of this product in the spray solution and timing of application to control specific weeds, woody brush, trees and vines, refer to the "WEEDS CONTROLLED" section of this label.

For control of annual weeds, make application when weeds are small and prior to seedhead or bud formation. For control of perennial weeds, woody brush, trees and vines, make application after flowering and before fall color and leaf drop.

When making a low-volume directed spray application to annual and perennial weeds, woody brush, trees and vines using a handheld sprayer, ensure that at least 50 to 75 percent of the foliage or the top one-half of each unwanted plant is sprayed. If a straight stream nozzle is used, start the application at the top of the targeted plant and spray from top to bottom in a lateral zig-zag motion. To ensure uniform and complete coverage, spray both sides of large or tall woody brush, trees and vines, or when foliage is thick and dense, or where there are multiple sprouts. For best performance of this product on woody brush, trees and vines, apply to actively growing vegetation after full leaf expansion and flowering, prior to fall color and leaf drop.

The following table summarizes various methods of foliar application using a backpack sprayer with a spray-to-wet or low-volume directed spray technique and high-volume sprayer application using handheld application equipment for control or partial control of herbaceous weeds, woody brush, trees and vines listed in the "WEEDS CONTROLLED" section of this label.

Method of Application	Spray Solution Concentration	Spray Volume
Handgun or Backpack Sprayer	0.5 to 1.5% by volume	Spray-to-wet technique
Low-Volume Directed Spray (Backpack)	4 to 8% by volume	15 to 25 gallons/acre
Modified High-Volume Spray	1.5 to 3% by volume	40 to 60 gallons/acre

Low-volume directed spray application with a backpack sprayer works best when applying to weeds and brush less than 10 feet tall. For taller weeds and brush, a high-volume handgun can be modified by reducing the nozzle size and spray pressure to produce a modified high-volume directed spray application.

### 8.5 Selective Application Equipment

Selective application equipment allows this product to be applied to weeds growing near a crop or other desireable vegetation without killing the desireable vegetation. Selective application equipment must be capable of preventing all contact of the herbicide solution with the desirable vegetation and operated without spray mist escape, leakage or dripping of the herbicide solution.

**AVOID CONTACT OF THIS HERBICIDE WITH DESIRABLE VEGETATION.** Contact of this product with desirable vegetation could result in unwanted plant damage or destruction. To the extent consistent with applicable law, such damage shall be the sole responsibility of the applicator.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

This product may be diluted with water and applied using a recirculating sprayer, shielded sprayer, hooded sprayer, wiper applicator or sponge bar to weeds listed on this label growing in any aquatic or on any terrestrial non-food or non-feed crop site listed on this label, where feasible. This product may also be used with sprayers equipped with optical weed sensor technology. Other selective equipment that may be used to deliver or apply this product are single and hollow stem injectors, tree injectors, wiper applicators for cut stem and cut stump applications, and spray or squirt bottles for cut stem, cut stump and frill applications to control large stem weeds, brush, trees and vines listed on this label.

### **Recirculating Sprayer**

A recirculating sprayer directs the spray solution onto weeds growing above desirable vegetation, while spray solution that is not intercepted by weeds is collected and returned to the spray tank for reapplication. A recirculating sprayer may be used to apply spray solutions of this product to weeds listed on this label in any aquatic or on any terrestrial non-crop site described on this label.

### **Shielded and Hooded Sprayers**

A shielded sprayer directs the herbicide solution onto targeted weeds while using an impervious material, or shield, to protect desirable vegetation from coming into contact with the herbicide spray. To provide maximum protection for desirable vegetation, keep shields properly adjusted and use nozzles that provide uniform coverage within the application area.

A hooded sprayer is a type of shielded sprayer where the spray pattern is fully enclosed, including the top, sides, front and back, thereby shielding desirable vegetation from the spray solution.

This product may be diluted with water and, unless otherwise directed, mixed with a surfactant and applied using a shielded or hooded sprayer to control weeds listed on this label growing in any aquatic or on any terrestrial non-food and non-feed crop site described on this label, where feasible [Optional text, if applicable: , and between rows of plants or trees (row middles and strips) in any cropping system listed on this label].

Properly adjust the hood to protect desirable vegetation. Ensure that the hood is capable of completely enclosing the spray pattern. If necessary when applying around crops grown on raised beds, extend the front and rear flaps of the hooded sprayer downward to reach the ground in deep furrows.

A hooded sprayer must be configured and operated in a manner that minimizes bouncing and avoids raising the hood up off the ground surface at any time. If the hood is raised, spray particles can escape and come into contact with the crop or other desirable vegetation, causing damage to or destruction of the desirable vegetation. Avoid operating this equipment on rough or sloping terrain where the spray hood is likely to rise up off the ground surface.

Use hoods designed to minimize excessive dripping or runoff down the inside of the hood, such as a single, low pressure, low drift, flat-fan nozzle with an 80 to 95-degree spray angle positioned at the top center of the hood, with a spray volume of 20 to 30 gallons per acre.

The following procedures will help reduce the potential for injury to desirable vegetation when using a hooded sprayer:

- Operate the sprayer with the hood on the ground or skimming across the ground surface.
- Leave at least an 8-inch untreated strip over the drill row. (For example, if a crop row width is 38 inches, use a sprayer hood with a maximum width of 30 inches.)
- Operate at a ground speed no greater than 5 miles per hour to minimize bouncing of the hooded sprayer.
- Apply when wind speed is 10 miles per hour or less.
- Use low-drift nozzles that provide uniform coverage within the application area.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Injury to a crop or other desirable vegetation can occur when application is made to foliage of weeds that come into direct contact with the desirable vegetation. Do not apply this product when leaves of desirable vegetation are growing in direct contact with weeds. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation can result in discoloration, stunting or destruction.

### **Wiper Applicator**

A wiper applicator is a device that physically wipes this product or solutions of this product directly onto the unwanted vegetation or cut stump. Any handheld device that is capable of physically wiping this product or solutions of this product directly onto the target weed or stump, such as a paint brush, may be used.

A mechanical wiper applicator, such as a rope wick or sponge bar that can be driven through a field over the top of a crop or other desirable vegetation to control weeds that are taller than the desirable vegetation, must be designed, maintained and operated to prevent the herbicide solution from coming into contact with desirable vegetation. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation can result in discoloration, stunting or destruction. Avoid leakage or dripping onto desirable vegetation. To protect the desirable vegetation, adjust the height of the applicator to ensure that the wiper contact point is at least 2 inches above the desired vegetation.

Wiper applicators may be used over the top of food or feed crops ONLY if specifically permitted for use over that crop by this label or by separately published supplemental labeling for this product.

Weeds that do not come into contact with the herbicide solution will not be affected. The more weed foliage exposed above the desirable vegetation the better mechanical wiper applicators work. Better results can be obtained when weeds are a minimum of 6 inches above the desirable vegetation. Poor contact can also occur when weeds are growing in dense clumps, when operating in areas of severe weed infestation, or when weed height varies dramatically. In these situations, more than one application of this product might be necessary to achieve desired weed control.

For optimal results, operate wiper applicators at a ground speed of no greater than 5 miles per hour. Performance in areas of heavy weed infestation can be improved by reducing speed, which will provide more time for resaturation of the wiper with the herbicide solution and more contact time of the wiper with the weed. Better weed control using a wiper applicator can also be obtained when two applications are made traveling in opposite directions across the field.

Keep wiper surfaces clean.

Be aware that on sloping ground the herbicide solution can migrate to one side, causing dripping on the lower end and drying of the wiper on the upper end of the applicator.

Do not apply this product using a wiper applicator when weeds are wet.

Add a nonionic surfactant to a concentration of 10 percent by volume of the total applicator solution (one gallon of surfactant for every 10 gallons of solution) for use in a wiper applicator. See the "MIXING" section of this label for more information on the use of surfactants.

**For Rope Wick and Sponge Bar Applicators** – apply solutions ranging from 33 to 75 percent of this product by volume in water.

**For Panel Applicators** – apply solutions ranging from 33 to 90 percent of this product by volume in water.

Mix only the amount of this product that will be used during a 1-day period, as reduced product performance can result from the use of solutions held in storage.

Clean wiper parts promptly after using this product by thoroughly flushing with water.

### **Single and Hollow Stem Injectors**

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Control of certain weeds listed in the "WEEDS CONTROLLED" section can be obtained by injecting this concentrated product or solutions of this product directly in or onto the target weed. Ensure that the handheld injector being used for this application is capable of accurately delivering the volume specified on the label. When making stem injections, the combined total use of this product must not exceed 8 quarts per acre per year. At 5 milliliters of concentrated (undiluted) product per stem, 8 quarts will treat approximately 1500 stems per acre per year. The number of stems that can be treated per acre will vary depending on the injection volume and the concentration of this product in the application solution.

### 8.6 Injection Systems

This product may be used in aerial and ground injection spray systems as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this concentrated product with the undiluted concentrate of other products when using injection systems, unless otherwise directed. A non-ionic surfactant concentration of 0.5% or more in the spray stream is required for use of this product in injection systems.

### 8.7 Controlled Droplet Applicator (CDA)

The amount of this product applied per acre using a controlled droplet applicator (CDA) must be no less than the rate specified on this label for application using conventional broadcast application equipment.

A controlled droplet applicator produces a spray pattern that is not easily visible. Use extreme care to avoid spray or drift from coming into contact with the foliage or any other tissue of desirable vegetation, as plant damage or destruction could result.

## 9.0 AQUATIC AND TERRESTRIAL USE SITES

This product may be used according to the directions for use described on this label to control weeds, woody brush, trees and vines listed on this label growing in aquatic environments and on any terrestrial site described on this label.

### 9.1 Aquatic Sites

This product may be used to control emerged weeds, brush, trees and vines in all flowing, non-flowing or transient bodies of fresh and brackish surface water. These bodies of water include lakes, rivers, streams, ponds, estuaries, seeps, irrigation and drainage ditches, canals, reservoirs, wetlands and wastewater treatment facilities. This product may also be used to control weeds in intertidal areas below the mean high-water mark and on terrestrial sites where bodies of water may be present and a buffer that will ensure no overspray of the water cannot be maintained.

When applying spray solutions of this product in or near aquatic sites, a nonionic surfactant that is labeled for use with herbicides and approved for direct application to bodies of water must be used. See the "MIXING" section of this label for more information on the use of surfactants with this product.

Before using this product for aquatic weed control or for terrestrial weed control near aquatic sites, read the following information carefully.

- This product does not control plants that are completely submerged or have a majority of their foliage under water.
- There is no restriction on the use of water for irrigation, recreation or domestic purposes following direct application of this product to emerged aquatic plants.
- Consult your local State agency with primary responsibility for regulating pesticides, State fish and wildlife agency and/or water control authority before applying this product to vegetation growing in public waters to determine if a permit is required.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

- Do not apply this product directly to water within 0.5 mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within 0.5 mile of an active potable water intake in a standing body of water, such as a lake, pond or reservoir. To make aquatic applications around and within 0.5 mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after the application. The water intake may be turned on prior to 48 hours if the glyphosate level in the intake water is below 0.7 parts per million as determined by laboratory analysis. These aquatic applications may be made ONLY in those cases where there are alternative water sources or holding ponds that would permit the turning off of an active potable water intake for a minimum period of 48 hours after the application. This restriction does NOT apply to intermittent inadvertent overspray of water on terrestrial use sites.
- To achieve maximum weed control in dry ditches, apply this product within 1 day after water drawdown to ensure application to actively growing weeds and allow a minimum of 7 days after application before reintroduction of water
- Floating mats of vegetation could require more than one application of this product for control. Avoid washing this product off of foliage after application by boat backwash or rainfall within 4 hours of application. Wait a minimum of 24 hours before re-applying this product to the same vegetation.
- Application of this product to moving bodies of water must be made while traveling upstream to prevent concentration of this herbicide in the water.
- When making a bankside application, do not overlap more than 1 foot into open water.
- Do not apply this product to bodies of water where emerged weeds do not exist.
- If applying this product to more than 20 percent of the total area of a body of water, do not apply more than 3.75 quarts per acre in any single broadcast application. If applying to less than 20 percent of the total area of a body of water, any rate listed on this label may be applied. This single application rate restriction does not apply to stream crossings in utility rights-of-way.
- When emerged weed infestations cover the total surface area of an impounded waterbody, apply this product to the emerged vegetation in strips to help avoid oxygen depletion in the water due to decaying vegetation. Oxygen depletion in the water can result in increased fish mortality.

**TANK MIXTURES:** This product may be applied in a tank mixture with one or more of the following products for a broader spectrum of control of aquatic weeds, woody brush, trees and vines in aquatic sites, provided that the product used is labeled for aquatic use. Refer to the individual label of all products used in the tank mixture for approved uses and application rates. Always read and follow label directions for each product in the mix. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

---

*[Active ingredients and products to be added to the final printed labeling for tank-mixing may be selected from the following list:*

2,4-D amine; imazapyr; flumioxazin; triclopyr

Clipper; Garlon 3A Specialty; Habitat]

---

### 9.2 TERRESTRIAL SITES

This product may be used according to the directions described on this label to control annual and perennial weeds, woody brush, trees and vines listed on this label on any terrestrial site described on this label.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

This product may be used to control weeds, woody brush, trees and vines on maintained landscapes, on improved and unimproved land, on lawns and turf and around ornamentals on industrial, commercial and residential sites, including [Select all uses that are applicable to the final printed labeling: airports, apartment complexes, chaparrals, ditch banks, driveways, dry ditches, dry canals, farmsteads, fencerows, forestry sites, golf courses, greenhouses, lumber yards, manufacturing sites, municipal sites, natural areas, nurseries, office complexes, ornamental beds, parks, parking areas, petroleum tank farms, pumping installations, railroads, recreational areas, residential areas, roadsides, schools, shadehouses, sod and turfgrass seed farms, sports complexes, storage areas, substations, utility rights-of-way, utility sites, and warehouse areas.]

This product may be used for non-selective control of unwanted vegetation on any site listed on this label for trim-and-edge application around objects, including around building foundations, equipment storage areas and trees, and along and in fences, and to eliminate unwanted vegetation growing in and around established shrub beds and ornamental plantings. This product may also be used for complete elimination of vegetation from a terrestrial site prior to planting ornamentals, flowers, or turfgrass (sod or seed), and prior to land development, including prior to beginning construction projects or the laying of asphalt or other road material. Unless otherwise stated for a specific use, application of this product may be repeated, as needed, to maintain bare ground, up to a total application of 8 quarts per acre per year.

This product may be used for establishment and maintenance of fuel breaks, for establishing fire perimeters and black lines, along fire roads and to facilitate prescribed burning practices on any site described on this label.

[Optional label text: This product may also be used for weed control or growth regulation in or on] [Optional list of any terrestrial uses that are included on this Master Label, including, but not limited to: Christmas tree farms, citrus orchards, fallow cropland, farmsteads, production nurseries, sugarcane fields, and turfgrass seed and sod farms.]

This product requires the addition of a nonionic surfactant to the spray solution labeled for herbicide application. See the "MIXING" section of this label for more information on the use of surfactants with this product.

Unless otherwise directed, application of this product may be made according to the directions for use in the sections that follow on any of these sites using any method of application described on this label to control any weeds, woody brush, trees and vines listed in the "WEEDS CONTROLLED" section of this label.

### 10.0 ADDITIONAL SITE MANAGEMENT INFORMATION

The following sections contain additional use information specifically related to certain use sites. Unless otherwise directed, any application of this product described in the "WEEDS CONTROLLED" section or any other section of this label may be made on the use sites described in the sections that follow, where applicable, using any method of application described on this label that is appropriate.

#### 10.1 Fallow Cropland Management

This product may be applied to control or suppress annual and perennial weeds on cropland during fallow intervals between plantings. See the "WEEDS CONTROLLED" section of this label for application rates for specific weeds. Application of up to 48 fluid ounces of this product per acre may be made using aerial application equipment onto fallow fields where there is sufficient buffer available to prevent drift onto adjacent crops.

**RESTRICTIONS:** Application of this product on cropland must not exceed 6 quarts per acre per year. See the "PRODUCT INFORMATION" section of this label for information on Maximum Application Rates. Allow a minimum of 30 days between application of this product and the planting or transplanting of any food or feed crop.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

### 10.2 Forestry, Hardwood and Christmas Tree Site Management

This product may be used for control or partial control of woody brush, trees and herbaceous weeds on any tree site, including forestry settings, Christmas tree plantations, and silvicultural and production nursery sites, using any method of application described on this label. See the "WEEDS CONTROLLED" section of this label for application rates and specific use directions.

Unless otherwise directed, this product requires a nonionic surfactant that is labeled for the intended use on the site of application to be added to the spray mixture. Use of this product without a surfactant will result in reduced performance. See the "MIXING" section of this label for more information on the use of surfactants with this product.

**IMPORTANT: SOME SURFACTANTS CAN CAUSE TREE INJURY WHEN DIRECTLY APPLIED TO SOME SPECIES. READ AND FULLY UNDERSTAND ALL APPROVED USES, PRECAUTIONS AND LIMITATIONS OF THE SURFACTANT BEFORE USING.**

#### Weed Management, Site Preparation

This product may be used to control or partially control undesirable woody brush, trees, vines and herbaceous weeds listed on this label for preparing sites prior to planting any tree species, including Christmas trees, eucalyptus trees and hybrid tree cultivars, and for controlling weeds around established trees, [Optional text: for the release of conifer and hardwood trees,] establishing wildlife openings and maintaining roads on any tree site.

**TANK MIXTURES:** This product may be applied in a tank-mix with the products listed in this section to broaden the spectrum of vegetation controlled. Any application rate of this product listed on this label may be used in a tank-mix with the following products for tree site management, including site preparation, provided that the product is labeled for the use on the site of application and prior to planting the desired species. Refer to the individual label of all products used in the tank mixture for approved uses and application rates. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

---

*[Active ingredients and products to be added to the final printed labeling for tank-mixing may be selected from the following list:*

imazapyr; metsulfuron methyl; sulfometuron methyl; triclopyr

Arsenal; Arsenal Herbicide Applicators Concentrate; Chopper; Chopper GEN2; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XRT Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Landmark; Landmark XP; Oust Extra; Oust XP]

---

For control of herbaceous weeds, apply these tank-mix products at a rate toward the lower end of the range specified on the product label. For control or partial control of dense stands of vegetation or for hard-to-control woody brush, trees and vines, apply these products at a rate or spray solution concentration towards the higher end of the given range.

#### Conifer Release, Mid-Rotation Conifer Release, Hardwood Release, Timber Stand Improvement

This product may be applied as a directed spray using a handheld sprayer or using any selective application equipment described on this label to control woody and herbaceous weeds and other undesirable understory vegetation below the tree crop canopy in conifer plantations, hardwood sites, Christmas tree plantations and silvicultural and ornamental nurseries to facilitate the release and growth of conifer and hardwood trees.

This product may also be applied using ground broadcast equipment or as a directed spray application for mid-rotation release under the canopy of pines, other conifers and hardwoods.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

**PRECAUTIONS:** Avoid contact of spray drift, mist or drips with foliage, green bark or non-woody surface roots of desirable plant species. Use application techniques that prevent or minimize contact of this product with foliage of desired trees or other plants through direct contact or off-target spray movement.

*[Optional label text: RESTRICTIONS: Do not apply this product as an over-the-top broadcast spray for conifer or hardwood release, unless otherwise directed on this label or on separate supplemental labeling for this product.]*

### **Conifer Release – Broadcast Application** *[This section is optional in the final printed label]*

This product may be broadly applied over the top of conifer tree species listed in this section after formation of final conifer resting buds in the fall or prior to initial bud swelling in the spring for control, partial control or suppression of herbaceous weeds and hardwoods listed in the "WEEDS CONTROLLED" section of this label to facilitate the release of these desirable tree species in a forestry, plantation or nursery setting. Unless otherwise directed, make this application only where conifers have been established for a minimum of one growing season.

**PRECAUTIONS:** Conifer injury can occur when this product is applied at rates higher than prescribed on this label, where spray applications overlap, if application is made when conifers are actively growing, or when they are growing under stress from drought, flood, improper planting or insect, animal or disease damage.

#### Conifer Release Outside the Southeastern United States

For release of the following conifer species growing for a minimum of one growing season in most areas outside the southeastern United States, apply 24 to 48 fluid ounces of this product per acre as a broadcast application over the top of the conifer trees.

- Douglas fir
- Fir species
- Hemlock
- Pines\*
- California redwood
- Spruce

\* Includes all species except loblolly pine, longleaf pine, shortleaf pine or slash pine.

Apply 24 to 40 fluid ounces of this product for release of Douglas fir, pine and spruce that have been established for only one growing season (except in California).

For release of spruce (*Picea spp.*) in Maine, Michigan, Minnesota, New Hampshire and Wisconsin, up to 2.25 quarts of this product may be applied after formation of final resting buds in the fall for control of woody brush and tree species.

**PRECAUTIONS:** Ensure that the conifers are well hardened off before application of this product. Some nonionic surfactants can cause tree injury when broadly applied over the top of hemlock and California redwood and in mixed conifer stands. Test the nonionic surfactant to be used for tree safety before using.

#### Conifer Release in the Southeastern United States

For release of the following conifer species established for more than one growing season in the southeastern United States, apply 36 to 60 fluid ounces of this product per acre in the fall as a broadcast application over the top of the trees. For release of these species after only one growing season, apply only 24 fluid ounces of this product per acre.

- Eastern white pine
- Loblolly pine
- Longleaf pine
- Shortleaf pine
- Slash pine
- Virginia pine

**TANK MIXTURES:** This product may be applied for conifer release in a tank-mix with the following products to provide a broader spectrum of postemergence weed control and for residual control of weeds listed on the label of those products. Only apply these tank mixtures over the top of conifer species that are approved for this use for all products in the mix. Refer to the individual product labels for approved

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

uses and application rates. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

---

*[Active ingredients and products to be added to the final printed labeling for tank-mixing may be selected from the following list:*

atrazine; imazapyr; metsulfuron methyl; sulfometuron methyl

Arsenal; Arsenal Herbicide Applicators Concentrate; Oust Extra; Oust XP]

---

For release of Douglas fir established for a minimum of one growing season prior to bud swell in early spring, apply 24 fluid ounces of this product in a tank-mix with an appropriate rate of atrazine. Do not add surfactant for this application.

For herbaceous release of loblolly pine, Virginia pine and longleaf pine in the spring and early summer, apply 12 to 18 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Oust Extra or Oust XP.

### **Late-Summer and Fall after Resting Bud Formation**

For release of jack pine, white pine and white spruce, apply 24 to 48 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Oust Extra or Oust XP that will not harm these conifer species.

For release of Douglas fir, apply 24 to 36 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Arsenal or Arsenal Herbicide Applicators Concentrate.

For release of balsam fir and red spruce, apply 48 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Arsenal or Arsenal Herbicide Applicators Concentrate.

### **10.3 Grass Seed and Sod Production**

This product may be used for the production of grass seed and sod of any grass (*Gramineae* family) except barley, buckwheat, corn, millet (pearl, porso), oats, quinoa, rice, rye, sorghum, sugarcane, teff, teosinte, triticale, wheat (all types) and wild rice.

**TYPES OF APPLICATION:** Preplant; At-Planting; Preemergence; Renovation; Removal of Established Stands; Site Preparation; Shielded Sprayer; Wiper Applicator; Spot Treatment

**RESTRICTIONS:** Do not apply this product to grass fields grazed by domestic livestock. Do not harvest grasses for feed. Wait a minimum of 30 days after application of this product before planting any food or feed crops.

### **Preplant, At-Planting, Preemergence, Renovation, Removal of Established Stand, Site Preparation**

**USE INSTRUCTIONS:** This product controls most existing vegetation for purposes of renovating turf or forage grass seed production areas, or for establishing turfgrass grown for sod. This product may be used to destroy undesirable grass vegetation when production fields are converted to alternative species or crops. Do not disturb soil or underground plant parts before application and delay tillage or renovation techniques, including vertical mowing, coring and slicing, for a minimum of 7 days after application to allow for herbicide translocation into underground plant parts.

Apply before, during or after planting, or for renovation purposes. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the herbicide spray. For maximum control of existing vegetation, delay planting until determining if any re-growth of underground plant parts will occur. Where repeat applications are necessary, sufficient re-growth must be attained prior to application. For warm-season grasses, such as bermudagrass, summer or fall application can provide optimal control.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Broadcast application of this product may be used to control sod remnants or other unwanted vegetation after sod is harvested. Application rates of up to 3.75 quarts per acre may be used to totally remove an established stand of hard-to-kill grass species.

### **Shielded Sprayer**

USE INSTRUCTIONS: Apply 24 to 72 fluid ounces of this product in 10 to 20 gallons of water per acre using a shielded sprayer to control weeds between grass seed rows. Uniform planting in straight rows will aid shielded sprayer application. Best results can be obtained when the grass seed crop is small enough to easily pass by the protective shields. See additional instructions on the use of shielded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS: Contact of this product in any manner to any vegetation to which application is not intended could cause damage.

### **Wiper Applicator**

USE INSTRUCTIONS: This product may be applied over the top of desirable grasses using a wiper applicator for the control of tall weeds. See additional instructions on the use of wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS: Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation could result in discoloration, stunting or destruction.

### **Spot Treatment**

USE INSTRUCTIONS: Apply a 1-percent solution of this product using a handheld sprayer to control weeds within established vegetation prior to heading of grasses grown for seed or to control sod remnants or other unwanted vegetation after sod is harvested.

PRECAUTIONS: This product will kill the desirable grasses along with the weeds. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction.

### **10.4 Ornamental and Production Nursery Management**

All uses of this product described on this label may be used in a plant nursery setting using any method of application described.

This product may be used to clear an area of unwanted vegetation prior to planting any ornamental plant, tree, shrub or other plants.

This product may also be used to control weeds growing around established woody ornamental species, such as arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, Douglas fir, jojoba, hollies, lilac, magnolia, maple, oak, poplar, privet, pine, spruce and yew. This product may also be used to trim and edge around potted plants and other objects in a plant nursery.

PRECAUTIONS: Protect desirable plants from the spray solution using shields or coverings made of waterproof material. Take care to avoid contact of spray, drift or mist with foliage, green stems or immature bark of established ornamental species.

### **Greenhouse/Shadehouse**

This product may be used to control weeds growing in and around greenhouses and shadehouses.

RESTRICTIONS: Desirable vegetation must not be present during application in a greenhouse. Turn air circulation fans off before applying this product inside a greenhouse or shadehouse and leave them off until the application solution has dried.

### **10.5 Commercial, Residential and Recreational Area Management**

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

All applications of this product described on this label may be used on commercial, residential and recreational areas, including parks, schools and athletic fields, using any method of application described on this label, including spot treatment of unwanted vegetation, trim-and-edge application around trees, fences, walking paths, buildings, sidewalks, nature trails and other objects in these areas, to eliminate unwanted weeds growing in established shrub and ornamental beds, for turf management and renovation, and to eliminate vegetation from a site prior to development, including prior to planting an area to ornamentals, flowers or turfgrass (sod or seed), or beginning construction projects.

### 10.6 Railroad Management

All uses of this product described in the "WEEDS CONTROLLED" or any other section of this label may be used on railroad sites using any method of application described.

This product requires a nonionic surfactant that is labeled for the intended use on the site of application to be added to the spray mixture. If application is to be made where aquatic sites might be directly sprayed or inadvertently oversprayed, the surfactant must be labeled for aquatic use. Use of this product without a surfactant will result in reduced performance. See the "MIXING" section of this label for more information on the use of surfactants with this product.

Application of this product along railroad rights-of-way may be made in up to 80 gallons of spray solution per acre.

### Bare Ground, Ballast and Shoulders, Crossings, Spot Treatment

This product may be used to maintain bare ground on railroad ballast and shoulders and reduce the need for mowing and mechanical brush removal along railroad rights-of-way. Application of this product may be repeated as weeds continue to emerge in order to maintain bare ground, up to a maximum total application rate of 8 quarts of this product per acre per year.

**TANK MIXTURES:** This product may be applied in a tank mixture with the following products to broaden the spectrum of control of woody brush and trees for bare ground, ballast and shoulder, crossing and spot treatment applications, and other brush, tree and vine control on railroad sites, provided that the product used is labeled for these applications. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

---

*[Active ingredients and products to be added to the final printed labeling for tank-mixing may be selected from the following list:*

2,4-D; atrazine; bromacil; chlorsulfuron; clopyralid; dicamba; diquat; diuron; hexazinone; imazapyr; metsulfuron methyl; pelargonic acid; simazine; sulfometuron methyl; sulfosulfuron; tebuthiuron; triclopyr

Arsenal; Arsenal Herbicide Applicators Concentrate; Chopper; Chopper Gen2; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XRT Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Habitat; Hyvar X; Hyvar X-L; Krovar I DF; Oust Extra; Oust XP; Outrider; Princep 4L; Princep Caliber 90; Princep Liquid; Sahara DG; Scythe; Stalker; Spike 20P Specialty; Spike 80DF Specialty; Telar XP; Transline Specialty; Velpar DF CU; Velpar DF VU; Velpar L; Velpar L CU; Velpar L VU; Vastlan Specialty]

---

### Brush, Tree and Vine Control

This product may be used to control woody brush, trees and vines along railroad rights-of-way. Apply 3 to 8 quarts of this product per acre in up to 80 gallons of spray solution containing 0.5% or more by volume of a nonionic surfactant as a broadcast application using either a boom or boomless sprayer. Apply a 0.75 to 1.5-percent solution of this product when using high-volume application equipment with a

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

spray-to-wet technique, or a 4 to 8-percent solution when using a low-volume directed spray for spot treatment.

**TANK MIXTURES:** This product may be applied in a tank-mix with one or more of the following products to broaden the spectrum of control of woody brush, trees and vines along railroad rights-of-way, provided that the product is labeled for use on these sites. Refer to the individual product label for approved uses and application rates. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

---

*[Active ingredients and products to be added to the final printed labeling for tank-mixing may be selected from the following list:*

chlorsulfuron; clopyralid; dicamba; fosamine; hexazinone; imazapyr; metsulfuron methyl; picloram; triclopyr

Arsenal; Arsenal Herbicide Applicator's Concentrate; Chopper; Chopper Gen2; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XRT Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Habitat; Krenite S Brush Control Agent; Stalker; Telar XP; Tordon 101 Mixture Specialty; Tordon 22K Specialty; Tordon K Herbicide Specialty; Transline Specialty; Vanquish; Velpar DF CU; Velpar DF VU; Velpar L; Velpar L CU; Velpar L VU; Vastlan Specialty]

---

### Weed Control in Dormant and Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds in dormant and actively growing bermudagrass along railroad rights-of-way. See the "WEEDS CONTROLLED" section of this label for directions for use of this product for weed control in grasses.

### 10.7 Roadside Management

All uses of this product described on this label may be used for weed management along roadways, including weed control in dormant and active bermudagrass and bahiagrass, weed control along shoulders and under and around guardrails, signposts and other objects along the road, using any method of application described on this label. If applying this product in areas where the spray solution could inadvertently overspray a body of surface water, a non-ionic surfactant approved for aquatic use must be used. See the "MIXING" section of this label for more information on the use of surfactants with this product.

**TANK MIXTURES:** This product may be tank-mixed with the following products for shoulder, guardrail, spot treatment and maintaining bare ground applications, provided that the product used is labeled for use on these sites. Not all tank-mix products listed are labeled for aquatic use. Refer to the individual product labels for approved uses and application rates. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

---

*[Active ingredients and products to be added to the final printed labeling for tank-mixing may be selected from the following list:*

2,4-D; atrazine; bromacil; chlorsulfuron; clopyralid; dicamba; diuron; fosamine; hexazinone; imazapic; imazapyr; metsulfuron methyl; oryzalin; oxadiazon; pendimethalin; picloram; prodiamine; simazine; sulfometuron; sulfosulfuron; triclopyr

AAtrex 4L; AAtrex Nine-O; Banvel; Barricade 65WG; Chopper; Chopper Gen2; Crossbow; Direx 4L; Escort XP; Endurance; Formula 40; Gallery 75 Dry Flowable Specialty; Gallery SC; Garlon 4; Garlon XRT; Hyvar X; Karmex DF; Krenite S Brush Control Agent; Krovar I DF; Landmark;

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Landmark XP; Oust Extra; Oust XP; Outrider; Pendulum 3.3 EC; Pendulum AquaCap; Pendimax 3.3; Plateau; Poast; Poast Plus; Princep 4L; Ronstar 50 WSP; Ronstar Flo; Ronstar G; Sahara DG; Surflan AS Specialty; Surflan Flex; Surflan Flex T&O; Surflan Pro; Surflan XL 2G; Telar XP; Tordon K; Vanquish; Vastlan Specialty; Velpar DF CU; Velpar DF VU; Velpar L; Velpar L CU; Velpar L VU; Weedar 64]

---

### 10.8 Utility Management

This product may be used along electrical power, pipeline and telephone rights-of-way, and on all sites associated with these utility rights-of-way, including substations, access roads and rail lines, and along similar rights-of-way that run in conjunction with utilities, for spot treatment of unwanted vegetation, side-trimming, trim-and-edge application around objects, weed control prior to planting a utility site to ornamentals, flowers or turfgrass (sod or seed), turf management, to eliminate unwanted weeds growing in established shrub or ornamental beds, to prepare or establishing wildlife openings and for eliminating vegetation prior to beginning construction projects. Application of this product may be repeated as needed to maintain bare ground as weeds continue to emerge, up to a maximum application rate of 8 quarts per acre per year.

**TANK MIXTURES:** This product may be tank-mixed with the following products for use on utility sites, provided that the product is labeled for use on these sites. Refer to the individual product label for approved uses and application rates. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

---

*[Active ingredients and products to be added to the final printed labeling for tank-mixing may be selected from the following list:*

2,4-D; atrazine; bromacil; chlorsulfuron; clopyralid; dicamba; diuron; fosamine; hexazinone; imazapic; imazapyr; metsulfuron methyl; oryzalin; pendimethalin; prodiamine; simazine; sulfometuron methyl; sulfosulfuron; triclopyr

AAtrex 4L; AAtrex Nine-O; Arsenal Herbicide Applicators Concentrate; Endurance; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XLT Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Hyvar XL; Krenite S Brush Control Agent; Krovar I DF; Oust Extra; Oust XP; Outrider; Plateau; Sahara DG; Surflan AS Agricultural; Surflan AS Specialty; Surflan Flex; Surflan Flex T&O; Surflan XL 2G; Telar XP; Transline Specialty; Vanquish; Velpar DF CU; Velpar DF VU; Velpar L; Velpar L CU; Velpar L VU; Vastlan Specialty; Weedar 64]

---

Ensure that the Garlon product is thoroughly mixed with water according to label directions before adding this product to the spray mixture. Maintain continuous agitation when adding this product in order to avoid tank-mix compatibility problems.

To broaden the spectrum of weed control when side-trimming, apply this product in a tank-mix with one of the Garlon products listed above.

### 11.0 CITRUS TREE CROPS

**CROPS:** All cultivars, varieties and/or hybrids of Calamondin; Chironja; Citron; Citrus Hybrids; Grapefruit (including Japanese summer); Kumquat; Lemon; Lime (including Australian desert lime, Australian finger lime, Australian round lime, Brown river finger lime, Mount white, New Guinea wild, Russell river, sweet, and Tahiti); Mandarin (including Mediterranean, Satsuma); Orange (all); Pummelo; Tangelo (ugli); Tangerine (Mandarin); Tangor; Uniq Fruit (ugli)

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

**TYPES OF APPLICATION:** Preplant (site preparation); Broadcast Spray, Selective Equipment (shielded sprayer, wiper applicator), Directed Spray or Spot Treatment in Middles (between rows of trees) or Strips (within rows of trees); Perennial Grass Suppression; Cut Stump Application

**USE INSTRUCTIONS:** The following use instructions pertain to application of this product in Florida and Texas only.

This product may be applied using a boom sprayer, controlled droplet applicator (CDA), shielded sprayer, wiper applicator, handheld or backpack sprayer, lance or orchard gun, for site preparation prior to planting or transplanting citrus trees, in middles (between rows of citrus trees) and strips (within rows of citrus trees), and for weed control or perennial grass suppression in established citrus orchards.

Apply 12 fluid ounces to 4 quarts of this product per acre as directed in the "WEEDS CONTROLLED" section of this label. Use a higher application rate within a given range when weeds are stressed, growing in dense populations or greater than 12 inches tall. Application may be repeated as needed up to a maximum of 8 quarts of this product per acre per year. See the "PRODUCT INFORMATION" section of this label for more information on Maximum Application Rates.

To control goatweed, apply 48 to 72 fluid ounces of this product in 20 to 30 gallons of water per acre when plants are actively growing. Apply 48 fluid ounces per acre when plants are less than 8 inches tall and 72 fluid ounces per acre when plants are greater than 8 inches tall. If goatweed is greater than 8 inches tall, the use of this product in a tank-mix with Krovar I or Karmex could improve weed control. Refer to the individual product labels for specific crops, rates, geographic restrictions and precautionary statements.

**PRECAUTIONS:** Use extreme care to avoid contact of this herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of desirable trees, canes and vines. Avoid application when recent pruning wounds or other mechanical injury have occurred. Contact of this product with other than matured brown bark could result in serious crop damage or destruction. Only shielded or directed sprayers may be used where potential for crop contact is high, and then only where there is sufficient clearance. For application in strips (within rows of citrus trees), only selective equipment (directed sprayer, hooded sprayer, shielded sprayer or wiper applicator) may be used in order to minimize the potential for overspray or drift of this product onto the trees. See additional use instructions and precautions in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

**RESTRICTIONS:** Allow a minimum of 3 days between application of this product and transplanting. Allow a minimum of 1 day between application and harvest of citrus fruit crops. For citron groves, apply as a directed spray only.

### **Middles and Strips Management**

**USE INSTRUCTIONS:** This product will control or suppress annual and perennial weeds and ground covers growing between rows of citrus trees (middles) and within rows of citrus trees (strips). If weeds are under drought stress, irrigate prior to application. Reduced weed control could result if weeds have been recently mowed at the time of application.

**TANK MIXTURES:** A tank mixture of this product with Goal 2XL may be applied for annual weed control between and within rows of a variety of citrus tree crops when weeds are stressed or growing in dense populations. Application of 12 to 24 fluid ounces of this product per acre plus an appropriate rate of Goal 2XL will control annual weeds with a maximum height or length of 6 inches, including crabgrass, common groundsel, junglerice, common lambsquarters, redroot pigweed, London rocket, common ryegrass, shepherd's-purse, annual sowthistle, filaree (suppression), horseweed/marestail, stinging nettle and common purslane (suppression). This tank-mix will also control common cheeseweed (malva) or hairy fleabane with a maximum height or length of 3 inches.

This product may also be applied to row middles and strips in tank mixtures with the following products.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

*[Active ingredients and products to be added to the final printed labeling for tank-mixing may be selected from the following list:*

2,4-D; benefin; bromacil; clethodim; diuron; fluazifop-P-butyl; flumioxazin; glufosinate-ammonium; indaziflam; norflurazon; oryzalin; oxyfluorfen; pendimethalin; rimsulfuron; saflufenacil; sethoxydim; simazine; thiazopyr

Alion; Chateau Herbicide SW; Chateau Herbicide WDG; Direx 4L; Dri-Clean; Fusilade DX; Goal 2XL; GoalTender; Karmex DF; Krovar I DF; Matrix SG; Poast; Poast Plus; Prowl 3.3 EC; Prowl H2O; Princep 4L; Princep Caliber 90; Rely 280; Select 2 EC; Select Max Herbicide with Inside Technology; Simazine 4L; Simazine 90DF; Simazine 90 WDG; Sim-Trol 4L; Sim-Trol DF; Solicam DF; Surflan A.S. Agricultural; Surflan AS Specialty; Surflan Flex; Surflan Flex T&O; Surflan XL 2G; Treevix Powered by Kixor; Venue]

Ensure that the product used is labeled for application within the citrus crop being grown and at the maturity stage of the trees (bearing or non-bearing). Read and follow label directions for all products in the tank mixture. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**RESTRICTIONS:** Do not apply these tank mixtures in Puerto Rico.

### **Perennial Grass Suppression**

This product will suppress perennial grasses such as bahiagrass, bermudagrass, tall fescue, orchardgrass, Kentucky bluegrass and quackgrass that are grown as ground covers in citrus tree crops.

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 6 fluid ounces of this product in 10 to 20 gallons of water per acre.

For suppression of Kentucky bluegrass covers, apply 4.5 fluid ounces of this product per acre. Do not add ammonium sulfate to the spray mix.

To maximize performance of this product, mow cool-season grass covers in the spring to even their height and then apply this product 3 to 4 days after mowing.

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 4.5 fluid ounces of this product in 10 to 25 gallons of water per acre 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches prior to seedhead emergence. For suppression for up to 120 days, apply 3 fluid ounces of this product per acre, followed by an application of 1.5 to 3 fluid ounces per acre about 45 days later. Make no more than two applications per year.

For burndown of bermudagrass, apply 24 to 48 fluid ounces of this product in 3 to 20 gallons of water per acre. Make this application only if a reduction of the bermudagrass stand can be tolerated. When burndown is required prior to harvest, make the application a minimum of 21 days prior to harvest to allow sufficient time for burndown to occur.

For suppression of bermudagrass, apply 4.5 to 12 fluid ounces of this product in a spray volume of 3 to 20 gallons per acre no sooner than 1 to 2 weeks after full green-up. Under shaded conditions or where a lesser degree of suppression is desired, apply 4.5 to 7.5 fluid ounces of this product per acre. If the bermudagrass is mowed, ensure a minimum of 3 inches in height before applying this product. Sequential applications may be made when re-growth occurs and bermudagrass injury and stand reduction can be tolerated.

For burndown or control of the weeds listed below, apply this product at the specified rate in 3 to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.

Weed Species	Level of Perennial Weed Control at
--------------	------------------------------------

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

	Various Application Rates (amount of this product per acre)			
	24 fl oz	48 fl oz	2.25 quarts	3.75 quarts
Bermudagrass	B	–	PC	C
Guinea grass				
<i>Texas and Florida Ridge</i>	B	C	C	C
<i>Florida Flatwoods</i>	–	B	C	C
Para grass	B	C	C	C
Torpedograss	S	–	PC	C

S = Suppression, PC = Partial Control, B = Burndown, C = Control

### Cut Stump Application

Application of this product to a freshly cut tree stump may be made during site preparation or site renovation to control regrowth and re-sprouting of citrus trees, including calamondin, chironja, citron, citrus hybrids, grapefruit, kumquat, lemon, lime, mandarin (tangerine), orange (all), pummelo, tangelo, tangor. See the "WEEDS CONTROLLED" section of this label under "Woody Brush, Trees and Vines" for use instructions on controlling cut stumps.

## 12.0 SUGARCANE

CROPS: Sugarcane

**TYPES OF APPLICATION:** Chemical Fallow; Preplant; At-Planting; Preemergence; Hooded Sprayer in Row Middles; Shielded Sprayer in Row Middles; Wiper Applicator in Row Middles; Spot Treatment; Plant Growth Regulation; Post-Harvest

**USE INSTRUCTIONS:** Application of this product in sugarcane according to the directions for use that follow may be repeated as needed up to a total maximum of 6 quarts per acre per year.

### Preplant, At-Planting, Preemergence

**USE INSTRUCTIONS:** This product may be applied in or around sugarcane fields, or in fields prior to the emergence of plant cane.

**RESTRICTIONS:** Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation [Optional text, if this use included on label for aquatic use: unless the surfactant added to the spray solution is labeled for herbicide use and approved for aquatic application].

### Weed Control in Row Middles

**USE INSTRUCTIONS:** This product may be applied for weed control in between rows of sugarcane using a shielded/hooded sprayer or wiper applicator capable of preventing all contact of the herbicide solution with the crop. See additional instructions on the use of hooded sprayers and wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

**PRECAUTIONS:** Crop injury is possible with these methods of application. Refer to the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for information regarding the potential for crop injury using selective application equipment. Do not allow weeds within the application area to come into contact with the crop.

**RESTRICTIONS:** Allow a minimum of 14 days between application of this product in row middles and harvest.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

### **Spot Treatment**

**USE INSTRUCTIONS:** This product may be applied as a spot treatment in sugarcane. For control of volunteer or diseased sugarcane, apply a 1-percent solution of this product in water using a handheld sprayer and a spray-to-wet technique. Maximum control of volunteer and diseased sugarcane can be achieved when application is made when there are at least 7 new leaves. Avoid contact of this herbicide with healthy sugarcane plants as severe damage or destruction could result.

**RESTRICTIONS:** Do not apply this product as a spot treatment to more than 10 percent of the total sugarcane field to be harvested. Do not feed or graze foliage from within the application area.

### **Fallow Treatment**

**USE INSTRUCTIONS:** This product may be used as a replacement for tillage in fields that are lying fallow between sugarcane crops. This product may also be used to remove the last stubble of ratoon cane by applying 3 to 3.75 quarts of this product in 10 to 40 gallons of water per acre to new growth having at least 7 new leaves. Allow a minimum of 7 days after application before tillage. Aerial application of up to 72 fluid ounces per acre may be made onto fallow sites where there is sufficient buffer to prevent drift onto adjacent crops. Tank mixtures with 2,4-D or dicamba may be used. Ensure that the product used is labeled for this application in sugarcane. Read and follow label directions for all products in the tank mixture.

**RESTRICTIONS:** Do not plant subsequent crops within 30 days after application of this product other than the following: alfalfa or other forage legumes, beans (all types), corn (all types), cotton, melons (all types), pasture grasses, peanuts, potatoes (Irish or sweet), sorghum (milo), soybean, squash (all types), sugarcane or wheat.

### **Plant Growth Regulation**

**USE INSTRUCTIONS:** This product may be used as a foliar-applied plant growth regulator to hasten ripening and extend the period of high sucrose level in both low and high-tonnage sugarcane. Most of the sucrose increase is concentrated in the top nodes of the cane stalk. To maximize sugar recovery where topping is practiced at harvest, top at the base of the fourth leaf. Consult your state sugarcane authority or local Seamless Control representative regarding the degree of sucrose response that can be anticipated prior to application of this product.

As a result of leaf desiccation, improved trash burn can be expected.

For optimal results, apply this product at the following rates and timing according to the State in which the sugarcane is grown. Use a higher application rate within the given range when applying to sugarcane under adverse ripening conditions or to a less responsive variety.

**FLORIDA** – Apply 6 to 14 fluid ounces of this product per acre 3 to 5 weeks before harvest of LAST RATOON CANE ONLY.

**HAWAII** – Apply 10 to 24 fluid ounces of this product per acre 4 to 10 weeks before harvest.

**LOUISIANA** – Apply 4 to 14 fluid ounces of this product per acre 3 to 7 weeks before harvest of RATOON CANE ONLY.

**PUERTO RICO** – Apply 6 fluid ounces of this product per acre 3 to 5 weeks before harvest of RATOON CANE ONLY.

**TEXAS** – Apply 6 to 14 fluid ounces of this product per acre 3 to 5 weeks before harvest of RATOON CANE ONLY.

**PRECAUTIONS:** Application of this product can initiate development of shooting eyes. This product might not increase the sucrose content of sugarcane under conditions of good natural ripening. Within 2

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

to 3 weeks after application, this product could produce a slight yellowing to a pronounced browning and drying of leaves, and a shortening of upper internodes. Spindle death could occur.

Rainfall within 6 hours after application of this product for enhanced sugarcane ripening could reduce its effectiveness.

Application to sugarcane grown for seed could result in a reduction in germination or vigor. To the extent consistent with applicable law, buyer and all users are responsible for any and all loss or damage in connection with the preharvest use of this product on sugarcane grown for seed.

**RESTRICTIONS:** Do not feed or graze sugarcane forage following application of this product. Do not plant subsequent crops within 30 days after application for enhanced sugarcane ripening other than the following: alfalfa or other forage legumes, beans (all types), corn (all types), cotton, melons (all types), pasture grasses, peanuts, potatoes (Irish or sweet), sorghum (milo), soybean, squash (all types), sugarcane or wheat.

Do not apply for enhanced ripening to any crops other than sugarcane. Use of this product in any manner not consistent with this label could result in injury to persons, animals or crops, or other unintended consequences.

### 13.0 WEEDS CONTROLLED

Read the entire label before proceeding to use this product.

Unless otherwise directed, this product requires the addition of a nonionic surfactant that is labeled for use with herbicides to the spray solution. See the "MIXING" section of this label for more information on the use of surfactants with this product.

Always use a higher application rate or spray solution concentration of this product within a given range when weed growth is heavy or dense, or when weeds are growing in an undisturbed (non-cultivated) area.

Poor weed control could be realized if application is made to weeds covered with dust. For weeds that have been mowed, grazed or cut, allow re-growth to occur prior to application of this product.

Refer to the sections that follow for application rates and timing of application for the control of annual and perennial weeds and woody brush, trees and vines.

#### 13.1 Weed Control, Renovation and Chemical Mowing in Turf

The use of this product as described in this section may be applied to turfgrass growing on any terrestrial site described on this label. Ensure that any tank-mix product applied with this product is labeled for the intended use and on the site of application.

##### Weed Control in Dormant Bermudagrass and Bahiagrass

This product may be used to control or suppress many winter annual weeds and tall fescue for effective release of dormant bermudagrass and bahiagrass prior to spring green-up in areas where these turfgrasses are desirable ground covers and some temporary injury or discoloration can be tolerated.

Apply 6 to 48 fluid ounces of this product in 10 to 40 gallons of water per acre when bermudagrass and bahiagrass are dormant and prior to spring green-up.

Application of more than 12 fluid ounces of this product per acre on highly maintained bermudagrass and bahiagrass turf, such as golf courses and lawns, could result in injury or delayed green-up in the spring.

For residual weed control in dormant bermudagrass and bahiagrass, this product may be tank-mixed with Outrider, Oust Extra or Oust XP herbicides. Apply 6 to 48 fluid ounces of this product in a tank-mix with an appropriate rate of Outrider, Oust Extra or Oust XP herbicide in 10 to 40 gallons of water per acre. To

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

avoid delays in green-up and minimize injury, apply no more than 1 ounce of Oust Extra or Oust XP herbicide per acre on bermudagrass and no more than 0.5 ounce on bahiagrass, and avoid application when these grasses are in a semi-dormant condition.

DO NOT apply this product in a tank-mix with Outrider, Oust Extra or Oust XP herbicides on highly maintained bermudagrass and bahiagrass turf, such as on golf courses and lawns.

### Weed Control in Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds in actively growing bermudagrass. Some bermudagrass injury could result from the application of this product, but the bermudagrass will recover under moist conditions once the effects of the product wear off. Use only on well-established bermudagrass where some temporary injury or discoloration can be tolerated.

Apply 12 to 36 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Use a lower application rate within this range when controlling annual weeds less than 4 inches tall (or runner length) and increase the rate towards the upper end of the range as weeds increase in size or as they approach flower or seedhead formation. At these application rates, this product will provide partial control of the following perennial weeds in actively growing bermudagrass:

- Bahiagrass
- Bluestem, silver
- Fescue, tall
- Johnsongrass
- Trumpet creeper
- Vaseygrass

**PRECAUTIONS:** Applying more than 12 fluid ounces of this product per acre on highly maintained bermudagrass, such as on golf courses and lawns, could cause unacceptable turf injury and discoloration.

For a broader weed control spectrum in actively growing bermudagrass, this product may be tank-mixed with Outrider, Oust Extra or Oust XP herbicides. Apply these tank-mixtures only on well-established bermudagrass where some temporary injury or discoloration can be tolerated. Make no more than one application of this product in these tank mixtures in the same season, otherwise the bermudagrass could be severely injured.

Apply 6 to 24 fluid ounces of this product per acre in a tank-mix with Outrider herbicide for control or partial control of Johnsongrass and other weeds listed on the Outrider herbicide label. Use a higher application rate of both products within the given ranges for control of annual or perennial weeds greater than 6 inches tall.

Apply 12 to 24 fluid ounces of this product per acre in a tank-mix with Oust Extra or Oust XP herbicide to broaden the spectrum of weed control to include weeds listed on those labels. Use a lower application rate of each product within the given ranges to control annual weeds listed on the labels that are less than 4 inches tall (or runner length) and increase the rates toward the upper end of the ranges as annual weeds increase in size and approach the flower or seedhead stage. This tank-mix will provide partial control of the following perennial weeds in actively growing bermudagrass:

- Bahiagrass
- Bluestem, silver
- Broomsedge
- Dallisgrass
- Dock, curly
- Dogfennel
- Fescue, tall
- Johnsongrass
- Poorjoe
- Trumpet creeper
- Vaseygrass
- Vervain, blue

**PRECAUTIONS:** Apply these tank mixtures only on well-established bermudagrass where some temporary injury or discoloration can be tolerated. DO NOT apply this product in a tank mixture with Outrider or Oust herbicides on highly maintained bermudagrass, such as on golf courses and lawns.

### Weed Control in Actively Growing Bahiagrass

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 4 fluid ounces of this product in 10 to 40 gallons of water per acre 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches prior to seedhead emergence.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

For growth suppression of bahiagrass for up to 120 days, apply 3 fluid ounces of this product per acre, followed by an application of 2 to 3 fluid ounces per acre about 45 days later. Make no more than two growth suppression applications per year.

For broad spectrum weed control in actively growing bahiagrass, this product may be tank-mixed with Outrider, Oust Extra or Oust XP herbicides.

Apply 1.5 to 3.5 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Outrider herbicide per acre to control annual and perennial weeds greater than 4 inches in height.

Apply 4 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Oust Extra or Oust XP herbicide 1 to 2 weeks following an initial spring mowing to broaden the spectrum of weed control in actively growing bahiagrass to include weeds listed on the Oust herbicide label. Make this application only once per year.

**PRECAUTIONS:** Apply these tank mixtures only on well-established bahiagrass where some temporary injury or discoloration can be tolerated.

### **Turf Renovation**

This product controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding until after determining if any re-growth of underground plant parts will occur. Where repeat applications are necessary, sufficient re-growth must be attained prior to re-application of this product. Summer or fall application of this product is the optimal timing for control of warm-season grasses, such as bermudagrass. For managed turfgrass, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray solution.

This product has no residual soil activity and will not affect plants, seed or sod planted back into the area after application.

A handheld sprayer may be used for spot treatment of unwanted vegetation growing in existing turfgrass. Broadcast application or spot treatment using a handheld sprayer may be used to control sod remnants or other unwanted vegetation after sod is harvested.

**PRECAUTIONS:** Do not disturb soil or underground plant parts before application of this product. Delay tillage and renovation techniques, such as vertical mowing, coring or slicing, a minimum of 7 days after application to allow translocation of the herbicide into underground plant parts.

**RESTRICTIONS:** If application rates total 2.25 quarts of this product per acre or less, no waiting period between application and feeding or livestock grazing is required. If the rate is greater than 2.25 quarts per acre, remove domestic livestock before application and wait a minimum of 8 weeks after application before grazing or harvesting.

### **Chemical Mowing**

This product may be used to suppress growth of perennial and annual grasses to serve as a substitute for mowing.

Perennial Grasses – apply 5 fluid ounces of this product per acre to suppress growth of Kentucky bluegrass, or 6 fluid ounces to suppress tall fescue, fine fescue, orchardgrass, quackgrass or reed canarygrass in 10 to 40 gallons of spray solution per acre after grasses have greened up to at least 75 percent green color in the spring, or 7 to 10 days after mowing when sufficient re-growth has occurred to provide a desirable height for growth regulation. Use chemical mowing only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

Annual Grasses – apply 3 to 4 fluid ounces of this product in 10 to 40 gallons of spray solution per acre to suppress growth of some annual grasses, such as annual ryegrass, wild barley and wild oats when

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

actively growing in coarse turf on roadsides or other industrial areas and before the seedheads are in the boot stage of development. This application could injure the desired annual grasses.

**PRECAUTIONS:** Use this product for chemical mowing only in areas where some temporary injury or discoloration of perennial and annual grasses can be tolerated.

### 13.2 Annual Weeds

Annual weeds are easiest to control with this product when they are small and actively growing. New leaf development indicates active growth.

To control or partially control the annual weeds listed in this section when they are less than 6 inches in height or runner length and actively growing, apply 24 fluid ounces of this product per acre. If they are over 6 inches in height or runner length, or slowly growing under stressed conditions, increase the application rate to 1 to 4 quarts per acre, depending on weed height and the severity of the poor growing conditions.

For application using a handheld sprayer with a spray-to-wet technique, apply a 0.5-percent solution of this product to annual weeds less than 6 inches in height or runner length prior to seedhead formation in grasses or bud formation in broadleaf weeds. To control annual weeds over 6 inches tall, or even smaller weeds growing under stressed conditions, apply a 0.75 to 1.5-percent solution. Apply the maximum concentration of this product within this range for hard-to-control weeds or to control weeds over 24 inches tall.

For control of annual weeds using a handheld controlled droplet applicator (CDA), apply a 15-percent solution of this product (19 to 20 fluid ounces of this product per gallon of spray solution) at a flow rate of 2 fluid ounces of spray solution per minute and a walking speed of 1.5 miles per hour (1 quart of spray solution per acre). When using a vehicle-mounted CDA, apply the required amount of this product, as indicated in this section, in 2 to 15 gallons of water per acre.

For maximize performance of this product, do not mow, cut, till, burn or disturb vegetation in the application area for a minimum of 3 days after application.

This product has no residual soil activity and does not control emergence of new annual weeds from seed. Repeated application of this product will be needed to control weeds that continue to emerge.

### WEED SPECIES

---

Anoda, spurred  
Balsam apple<sup>1</sup>  
Barley  
Barley, little  
Barnyardgrass  
Bassia, fivehook  
Bittercress  
Bluegrass, annual\*  
Bluegrass, bulbous  
Brome, downy  
Brome, Japanese  
Broomsedge  
Buttercup  
Castor bean<sup>2</sup>  
Cheatgrass  
Cheeseweed (*Malva parviflora*)  
Chervil  
Chickweed

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Cocklebur  
Copperleaf, hophornbeam  
Copperleaf, Virginia  
Coreopsis, plains/tickseed  
Corn  
Crabgrass  
Cupgrass, woolly  
Dwarf dandelion  
Eclipta  
False dandelion  
Falseflax, smallseed  
Fiddleneck  
Filaree  
Fleabane, annual  
Fleabane, hairy\* (*Conyza bonariensis*)  
Fleabane, rough  
Foxtail  
Foxtail, Carolina  
Geranium, Carolina  
Goatgrass, jointed  
Goosegrass\*  
Groundsel, common  
Henbit  
Horseweed/Marestail (*Conyza canadensis*)  
Itchgrass  
Johnsongrass, seedling\*  
Junglerice\*  
Knotweed  
Kochia\*  
Lambsquarters  
Lettuce, prickly  
Mannagrass, eastern  
Mayweed  
Medusahead  
Morning glory (*Ipomoea* spp)  
Mustard, blue  
Mustard, tansy  
Mustard, tumble  
Mustard, wild  
Nightshade, black  
Oats  
Panicum, browntop  
Panicum, fall  
Panicum, Texas  
Pennycress, field  
Pepperweed, Virginia  
Pigweed\*  
Puncturevine  
Purslane, common  
Pusley, Florida  
Ragweed, common\*

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Ragweed, giant\*  
Rice, red  
Rocket, London  
Rocket, yellow  
Rye  
Ryegrass\*  
Sandbur, field  
Sesbania, hemp  
Shattercane  
Shepherd's-purse  
Sicklepod  
Signalgrass, broadleaf  
Smartweed, ladysthumb  
Smartweed, Pennsylvania  
Sorghum, grain (milo)  
Sowthistle, annual  
Spanish needles<sup>3</sup>  
Speedwell, corn  
Speedwell, purslane  
Sprangletop  
Spurge, annual  
Spurge, prostrate  
Spurge, spotted  
Spurry, umbrella  
Starthistle, yellow  
Stinkgrass  
Sunflower  
Teaweed / Prickly sida  
Thistle, Russian\*  
Velvetleaf  
Wheat  
Wild oats  
Witchgrass

---

<sup>1</sup> For control of balsam apple, apply this product using handheld equipment only.

<sup>2</sup> Control of castor bean can also be achieved by injecting 4 milliliters of this concentrated (undiluted) product per plant into the lower portion of the main stem.

<sup>3</sup> For control of Spanish needles, apply 48 fluid ounces of this product per acre.

\* A glyphosate-resistant biotype has been confirmed. For additional information, refer to the "WEED RESISTANCE MANAGEMENT" section of this label. You can also visit on the Internet, [www.weedscience.org](http://www.weedscience.org) or [www.weedresistancemanagement.com](http://www.weedresistancemanagement.com), or contact your Seamless Control representative.

### 13.3 Perennial Weeds

More effective control of perennial weeds with this product can be obtained when applied to target weeds that are small and actively growing. New leaf development indicates active growth. If application must be made to larger weeds or to weeds that are slowly growing under stressful conditions, apply this product at a rate or spray solution concentration towards the upper end of the specified range.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

If weeds have been mowed or tilled, do not apply this product until plants have resumed active growth and have reached the specified stage of growth or sufficient growth has been achieved to allow for good interception of the spray solution. To maximize performance of this product, do not mow, cut, till, burn or disturb vegetation in the application area for a minimum of 7 days after application.

For control of perennial weeds listed on this label using backpack or handheld equipment with a low-volume application technique, apply a 4 to 8-percent solution of this product over the crown of the target plant to cover 50 percent of the upper plant foliage.

For control of perennial weeds using a handheld controlled droplet applicator (CDA), apply a 15 to 30-percent solution of this product (19 to 38 fluid ounces of this product per gallon of spray solution) at a flow rate of 2 fluid ounces of spray solution per minute and a walking speed of 0.75 mile per hour (2 to 4 quarts of spray solution per acre). When using a vehicle-mounted CDA, apply the required amount of this product, as indicated in the following table, in 2 to 15 gallons of water per acre.

Application of this product in the fall must be made before a killing frost.

This product has no soil activity and does not control emergence of perennial weeds from seed and dormant underground roots, rhizomes or tubers present in the soil at the time of application. Repeated application of this product will be needed for continued control of weeds that emerge after application.

**PERENNIAL WEEDS RATE TABLE**

Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
<b>Alfalfa<sup>1</sup></b>	0.7	1.5
<b>Alligatorweed<sup>1</sup></b>	3	1.3
Apply this product when most of the target plants are in bloom. More than one application will be needed to achieve control.		
<b>Anise (fennel)</b>	1.5 – 3	1 – 1.5
<b>Bahiagrass</b>	2.3 – 3.75	1.5
<b>Beachgrass, European</b>	–	3.5
Apply a 3.5-percent solution of this product using a spray-to-wet technique or an 8-percent solution using a low-volume application technique. More effective control can be obtained when application is made onto target weeds that are actively growing at the boot through the full-heading stage of development. Make application prior to the loss of more than 50 percent of green leaf color in the fall. Monitor application site and re-apply this product to any target weeds that were missed, if necessary, before re-seeding the area with desirable vegetation.		
For selective control of European beachgrass, apply a 33.3-percent solution of this product containing 1 to 2.5 percent of a nonionic surfactant during period of active growth using a wiper applicator. Maximizing the amount of individual leaf tissue contacted by the wiper applicator or making a second pass in the opposite direction will improve control. Avoid contact of the herbicide solution with desirable vegetation.		
<b>Bentgrass<sup>1</sup></b>	1	1.5
<b>Bermudagrass</b>	4	1.5
Make application when seedheads are present.		
<b>Bermudagrass, water (knotgrass)</b>	1	1.5
<b>Bindweed, field</b>	2.3 – 3.75	1.5
For control, apply 3 to 3.75 quarts of this product per acre as a broadcast application west of the		

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
Mississippi River and 2.3 to 3 quarts per acre east of the Mississippi River when bindweed is at or beyond full bloom. For maximum performance, make application in late-summer or fall.		
<b>Bittersweet, Oriental</b>	2.25	2
For control of oriental bittersweet, apply this product as a broadcast spray in 3 to 40 gallons of spray solution containing 0.25 percent of a nonionic surfactant and 0.1 percent nonionic organosilicone per acre. Use a nonionic surfactant concentration of 0.5 to 2 percent by volume when using a handheld sprayer and a spray-to-wet application. For maximum performance, ensure complete coverage of the target plant with the spray solution.		
<b>Bluegrass, Kentucky</b>	1.5 – 2.3	0.75
Apply when most target plants have reached the boot to head stage of development. When application is made prior to the boot stage, reduced control can result. In the fall, make application before plants have turned brown.		
<b>Blueweed, Texas</b>	2.3 – 3.75	1.5
Apply 3 to 3.75 quarts of this product per acre west of the Mississippi River and 2.3 to 3 quarts per acre east of the Mississippi River when most target plants are at or beyond full bloom. For maximum performance, make application in late-summer or fall.		
<b>Brackenfern</b>	2.3 – 3	0.75 – 1
Apply to fully expanded fronds that are at least 18 inches long.		
<b>Bromegrass, smooth</b>	1.5 – 2.3	0.75
Apply this product when most target plants have reached the boot to head stage of development. When application is made prior to the boot stage, reduced control can result. In the fall, make application before plants have turned brown.		
<b>Bursage, woolly-leaf</b>	–	1.5
<b>Canarygrass, reed</b>	1.5 – 2.3	0.75
Apply this product when most target plants have reached the boot to head stage of development. When application is made prior to the boot stage, reduced control can result. In the fall, make application before plants have turned brown.		
<b>Cattail</b>	2.3 – 3.75	0.75
Apply this product when target plants are actively growing and are at or beyond the early to full bloom stage of development. For maximum performance, make application during the summer or fall months.		
<b>Clover; red, white</b>	2.3 – 3.75	1.5
<b>Cogongrass</b>	2.3 – 3.75	1.5
Apply this product in late-summer or fall when cogongrass is at least 18 inches tall and actively growing. Due to uneven stages of growth and the dense nature of cogongrass vegetation, more than one application might be necessary to achieve control.		
<b>Cordgrass</b>	2 – 8	5 – 8
Prior to application of this product for control of cordgrass, survey the area to determine if shellfish beds exist within the application area. If shellfish are intended to be harvested in the area, delay application of this product until after harvest or maintain a 50-foot buffer between the application area and commercial shellfish beds, or do not harvest shellfish for a minimum of 14 days following application of this product. See restrictions below.		
Ideal conditions for control of cordgrass are when target plants are free of silt and debris and actively growing, and good spray coverage is achievable. The presence of debris or silt on the surface of cordgrass will reduce the performance of this product. To improve herbicide uptake, wash targeted		

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
plants prior to application and allow a minimum of 4 hours for plants to dry before applying this product. Where cordgrass has been cut or mowed prior to application, allow for sufficient re-growth before applying this product to ensure adequate interception and uptake of this product. Rainfall or immersion of the plant in tidewater within 4 hours after application could reduce the effectiveness of this product.		
Apply 2 to 8 quarts of this product per acre using ground broadcast application or optical sensor equipment in 5 to 100 gallons of spray solution, or in 5 to 10 gallons of spray solution per acre when using aerial application equipment. Apply a 5 to 8-percent solution of this product when using a handheld backpack sprayer or high-volume sprayer. Make all applications of this product for the control of cordgrass in a spray solution containing 0.25% or more (1 or more quarts per 100 gallons of spray solution) of a nonionic surfactant or other adjuvant that is compatible with this product and labeled for use with herbicides and approved for use on aquatic sites. For maximum performance, ensure complete coverage of cordgrass clumps.		
RESTRICTIONS: If a minimum 50-foot buffer is maintained between the application area and commercial shellfish beds, there is no restriction on shellfish harvest. If application is made within 50 feet of commercial shellfish beds, DO NOT harvest shellfish for a minimum of 14 days following application of this product.		
<b>Cutgrass, giant<sup>1</sup></b>	3	1
More than one application of this product will be required to achieve control, especially where vegetation is partially submerged in water. Allow target weeds to re-grow to the 7 to 10-leaf stage before making next application.		
<b>Dallisgrass</b>	2.3 – 3.75	1.5
<b>Dandelion</b>	2.3 – 3.75	1.5
<b>Dock, curly</b>	2.3 – 3.75	1.5
<b>Dogbane, hemp</b>	3	1.5
Apply this product when most target plants have reached the late-bud to flower stage of growth. For maximum performance, make application in late-summer or fall.		
<b>Fescue (except tall)</b>	2.3 – 3.75	1.5
<b>Fescue, tall</b>	2.3	1
Apply this product when most target plants have reached the boot to head stage of growth. If applied prior to the boot stage, less than desirable control might be obtained.		
<b>Guinea grass</b>	2.3	0.75
Apply this product when most target plants have at least reached the 7-leaf growth stage.		
<b>Hemlock, poison</b>	1.5 – 3	0.75 – 1.5
Control can also be achieved by injecting 5 milliliters of a 5-percent solution of this product using a handheld injection device in one leaf cane per plant, 12 inches above the root crown. <sup>2</sup> No surfactant is required.		
<b>Hogweed, giant</b>	–	–
Inject 5 milliliters of a 5-percent solution of this product into one leaf cane per plant, 12 inches above the root crown. <sup>2</sup> No surfactant is required.		
<b>Horsenettle</b>	2.3 – 3.75	1.5
<b>Horseradish</b>	3	1.5
Apply this product when most target plants have reached the late-bud to flower stage of development.		

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
For maximum performance, make application in late-summer or fall.		
<b>Horsetail, field</b>	—	—
Inject 0.5 milliliter of this product per stem directly into the plant stem, one segment above the root crown. <sup>2</sup> No surfactant is required.		
<b>Iceplant</b>	1.5	1.5
<b>Iris, yellow flag</b>	—	—
Cut flower stems 8 to 9 inches above the root crown. Push a cavity needle into the stem center and then slowly remove it as you inject 0.5 milliliter of this product using a handheld injector. <sup>2</sup> No surfactant is required.		
<b>Ivy; cape, German</b>	1.5 – 3	0.75 – 1.5
<b>Jerusalem artichoke</b>	2.3 – 3.75	1.5
<b>Johnsongrass</b>	1.5 – 2.3	0.75
Apply this product when most target plants have reached the boot to head stage of development or before plants have turned brown in the fall. When applied prior to the boot stage, reduced control can result.		
<b>Kikuyu grass</b>	1.5 – 2.3	0.75
<b>Knapweed</b>	3	1.5
Apply this product when most target plants have reached the late-bud to flower stage of growth. For maximum performance, make application in late-summer or fall.		
<b>Knotweed; Bohemian, giant, Japanese</b>	3	2
Apply 3 quarts of this product per acre as a broadcast application in 3 to 40 gallons of spray solution with 0.5 to 1 percent by volume of a nonionic surfactant. For application using a backpack sprayer and a spray-to-wet technique, apply a 2-percent solution of this product containing 0.5 to 2 percent by volume of a nonionic surfactant. For maximum performance, do not disturb vegetation in the application area for a minimum of 7 days after application.		
Control can also be achieved by cutting stems cleanly just below the 2nd or 3rd node above the ground and immediately apply 0.36 fluid ounce (10 milliliters) of a 50-percent solution of this product in water into the "well" or remaining internode. Ensure that the upper plant material that was removed is gathered and properly discarded to prevent new plants from propagating from sprouting buds. Use of a bio-barrier, such as cardboard, plywood or plastic sheeting, will help guard against the spread of plant material. The combined total application rate of this product must not exceed 8 quarts per acre. <sup>2</sup>		
Control can also be achieved by injecting 5 milliliters of this product per stem into the second or third internode using a handheld injection device. <sup>2</sup> No surfactant is required.		
<b>Lantana</b>	—	0.75 – 1
Apply this product when most target plants are at or beyond the bloom stage of growth. Use the higher spray solution concentration on plants that have reached the woody stage of growth.		
<b>Lespedeza</b>	2.3 – 3.75	1.5
<b>Loosestrife, purple</b>	2	1 – 1.5
Apply this product when most target plants are at or beyond the bloom stage of growth. Maximum performance can be achieved when this product is applied during summer or fall months. Fall application must be made before a killing frost.		
<b>Lotus, American</b>	2	0.75
Apply this product when most target plants are at or beyond the bloom stage of growth. Maximum		

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
performance can be achieved when this product application is applied during summer or fall months. Fall application must be made before a killing frost. More than one application of this product might be necessary to control re-growth of underground plant parts and seeds.		
<b>Maidencane</b>	3	0.75
More than one application of this product will be needed for control, especially for vegetation partially submerged in water. Allow plants to re-grow to the 7 to 10-leaf stage before making next application.		
<b>Milkweed, common</b>	2.3	1.5
Apply this product when most target plants have reached the late-bud to flower stage of growth.		
<b>Muhly, wirestem</b>	1.5 – 2.3	0.75
Make application when most target plants are at least 8 inches in height (3 to 4-leaf stage of development) and actively growing.		
<b>Mullein, common</b>	2.3 – 3.75	1.5
<b>Napiergrass</b>	2.3 – 3.75	1.5
<b>Nightshade, silverleaf</b>	2.3 – 3.75	1.5
Apply 3 to 3.75 quarts of this product per acre as a broadcast application west of the Mississippi River and 2.3 to 3 quarts per acre east of the Mississippi River when most target plants are at or beyond full bloom. Maximum performance can be achieved when this product is applied in late-summer or fall after berries have formed.		
<b>Nutsedge; purple, yellow</b>	2.3	0.75
Apply this product to control existing nutsedge plants and attached immature nutlets when target plants are in flower or when new nutlets can be found at rhizome tips. Nutlets that have not germinated will not be controlled and will require repeated application of this product for long-term control.		
<b>Orchardgrass</b>	1.5 – 2.3	0.75
Make application when most target plants have reached the boot to head stage of development. When applied prior to the boot stage, less than acceptable control could be obtained. In the fall, make application before plants have turned brown.		
<b>Pampas grass</b>	2.3 – 3.75	1.5
<b>Para grass</b>	3	0.75
More than one application of this product will be needed to achieve complete control. Allow plants to re-grow to the 7 to 10-leaf stage before making next application.		
<b>Pepperweed, perennial</b>	3	1.5
<b>Phragmites<sup>1</sup></b>	2 – 3.75	0.75 – 1.5
For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 3.75 quarts of this product per acre as a broadcast application or a 1.5-percent solution using a handheld sprayer. In other areas of the U.S., apply 2 to 3 quarts per acre as a broadcast application or, for partial control, apply a 0.75-percent solution using a handheld sprayer. For maximum performance, make application in late-summer or fall when plants are actively growing and in full bloom. Due to the dense nature of this vegetation (which can prevent good spray coverage) and uneven stages of growth, more than one application of this product might be necessary to achieve control. Visual symptoms of control will be slow to develop.		
<b>Quackgrass</b>	1.5 – 2.3	0.75
Apply this product when most target plants are at least 8 inches in height (3- to 4-leaf stage of development) and actively growing.		

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
<b>Redvine<sup>1</sup></b>	1.5	1.5
<b>Reed; common, giant</b>	3 – 3.75	1.5
For maximum performance, apply this product in late-summer or fall. Control can also be achieved by injecting 5 milliliters of this concentrated product (undiluted) directly into the second or third internode using a handheld injection device. <sup>2</sup> No surfactant is required.		
<b>Ryegrass, perennial</b>	1.5 – 2.3	0.75
Apply this product when most target plants have reached the boot to head stage of growth. When applied prior to the boot stage, reduced control can result. In the fall, make application before ryegrass turns brown.		
<b>Salvinia, giant</b>	3 – 3.75	2
Apply a 2-percent solution of this product containing 0.5 to 2 percent by volume of a nonionic surfactant approved for aquatic use and containing at least 70 percent active ingredient using spray-to-wet technique. For broadcast application, apply 3 to 3.75 quarts of this product per acre in 3 to 40 gallons of spray solution containing 0.1 percent by volume nonionic organosilicone and 0.25 percent nonionic spreader sticker surfactant approved for aquatic use.  Allow a minimum of 3 days after application before disturbing vegetation in the application site. This product will not control plants that are completely submerged or have a majority of foliage under water.		
<b>Smartweed, swamp</b>	2.3 – 3.75	1.5
<b>Spatterdock</b>	3	0.75
Make application when most target plants are in full bloom. For maximum performance, make application in the summer or fall.		
<b>Spurge, leafy<sup>1</sup></b>	–	1.5
<b>Starthistle, yellow</b>	–	1.5
<b>Sweetpotato, wild<sup>1</sup></b>	–	1.5
Make application when most target plants are at or beyond the bloom stage of growth. More than one application will be needed to achieve control.		
<b>Thistle, artichoke</b>	1.5 – 2.3	2
Make application when target plants are at or beyond the bud stage of growth.		
<b>Thistle, Canada</b>	1.5 – 2.3	1.5
Make application when target plants are at or beyond the bud stage of growth.  Control can also be achieved by stem-injection. Cut 8 to 9 of tallest plants in a clump at bud stage. Push a cavity needle into the stem center and then slowly remove it as you inject 0.5 milliliter of this concentrated product into the stem. <sup>2</sup> No surfactant required.		
<b>Timothy</b>	1.5 – 2.3	1.5
Make application when most target plants have reached the boot to head stage of development. If application is made prior to the boot stage, reduced control can result. In the fall, make application before plants turn brown.		
<b>Torpedograss<sup>1</sup></b>	3 – 3.75	0.75 – 1.5
Apply this product at a lower rate or spray solution concentration within the specified range when torpedograss is growing on terrestrial sites and at a higher rate or concentration within the range when partially submerged under water or growing as a floating mat. Additional applications of this product will be needed to maintain control.		
<b>Trumpetcreeper<sup>1</sup></b>	1.5 – 2.3	1.5

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
<b>Tules, common</b> Make application to target plants at or beyond the seedhead stage of development. Visual symptoms will be slow to appear and might not appear for 3 or more weeks after application.	—	1.5
<b>Vaseygrass</b>	2.3 – 3.75	1.5
<b>Velvetgrass</b>	2.3 – 3.75	1.5
<b>Waterhyacinth</b> Make application when target plants are at or beyond the early bloom stage of development. Visual symptoms might require 3 or more weeks after application to appear, with complete necrosis and decomposition not occurring until 60 to 90 days after application. Use a higher application rate within the given range when more rapid visual effects are desired.	2.5 – 3	0.75 – 1
<b>Waterlettuce</b> Apply a 1-percent solution of this product in areas of heavy infestation. Maximum performance of this product can be obtained when applied from mid-summer through winter. Application in the spring could require more than one application to achieve control.	—	0.75 – 1
<b>Waterprimrose</b> Make application to target plants that are at or beyond the bloom stage of growth, but before fall color changes occur. Thorough coverage is necessary for maximum control.	—	0.75
<b>Wheatgrass, western</b> Make application when most target plants have reached the boot to head stage of development. Application made prior to the boot stage could result in reduced control. In the fall, make application before plants turn brown.	1.5 – 2.3	0.75

1 Partial control

2 When using stem injection, the combined total use of this product must not exceed 8 quarts per acre per year. At 5 milliliters of concentrated (undiluted) product per stem, 8 quarts will treat approximately 1500 stems per acre per year. The number of stems that can be treated per acre will vary depending on the injection volume and the concentration of this product in the application solution.

**Other perennials listed on this label** – Apply 2.3 to 3.75 quarts of this product per acre as a broadcast application or a 0.75 to 1.5-percent solution using a handheld sprayer.

### 13.4 Woody Brush, Trees and Vines

Apply this product to brush, trees and vines that are actively growing after full leaf expansion, unless otherwise directed. Apply this product at a rate or spray solution concentration within towards the upper end of a given range to control larger brush and trees and/or for application in areas of dense vegetative growth, for the control of vines that have reached the woody stage of growth. In most areas, maximum performance of this product on woody brush, trees and vines can be obtained when application is made in late-summer or fall after fruit formation. In arid areas, application of this product in the spring to early-summer when brush and trees are at high moisture content and flowering could provide better results. Poor control can be expected when this product is applied to drought-stressed brush, trees and vines.

Some autumn color on undesirable deciduous species of brush and trees is acceptable when applying this product in the fall, provided no major leaf drop has occurred. Reduced performance of this product could result if fall application is made after a frost. Symptoms might not appear prior to frost or senescence following fall application.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

For maximum performance, allow 7 or more days after application of this product before mowing, cutting, tilling, burning or removal of woody brush, trees or vines from the application site. Repeated application of this product will be necessary to control brush and trees regenerating from underground parts or seed.

**TANK MIXTURES:** This product may be applied at any rate stated on this label in a tank mixture with the following products to broaden the spectrum of control of herbaceous weeds, woody brush, trees and vines. For control of herbaceous weeds, apply the tank-mix product at the lower end of the given application rate or spray solution concentration range. For control of dense stands of vegetation or for hard-to-control woody brush, trees and vines, increase the application rate or spray solution concentration of the tank-mix product towards the higher end of the range. Refer to the individual product labels for approved uses and application rates. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

---

*[Active ingredients and products to be added to the final printed labeling for tank-mixing may be selected from the following list:*

imazapyr; metsulfuron methyl; triclopyr

Arsenal; Arsenal Herbicide Applicators Concentrate; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XRT Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Vastlan Specialty]

---

Ensure that the proper amount of the Garlon herbicide is thoroughly mixed with water in the spray tank before adding this product.

### **Cut Stump Application**

This product may be used to control re-growth and re-sprouting of woody brush and trees on any site listed on this label.

Cut the woody brush or tree close to the soil or water surface and immediately apply a 50 to 100-percent (undiluted) solution of this product to the freshly-cut surface using an applicator capable of applying this product to the entire cambium. A delay in application could result in reduced performance. For best results, cut the woody brush or tree during period of active growth and full leaf expansion and apply this product. No surfactant is needed for cut stump application.

For control of the tree of heaven (*Ailanthus altissima*), cut the tree close to the soil surface and immediately apply a 50-percent solution of this product (16 fluid ounces per quart of solution) and 10 percent Arsenal herbicide (3 to 4 fluid ounces per quart of solution) in water to the freshly-cut surface.

**DO NOT MAKE A CUT STUMP APPLICATION WHEN THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP, AS INJURY TO THE ADJACENT TREES COULD OCCUR.** Some sprouts, stems, or trees can share a common root system. Adjacent trees having a similar age, height and spacing could be an indicator of a shared root system. Whether grafted or shared, injury is likely to occur to adjacent stems or trees when this product is applied to one or more trees sharing common root system.

### **Woody Brush and Tree Injection and Frill Application**

This product may be used to control woody brush and trees listed in this section by injection or frill application on any aquatic and terrestrial site listed on this label.

Inject or apply the equivalent of 1 milliliter (0.04 fluid once) of this product for every 2 to 3 inches of trunk diameter at breast height (DBH). If injecting this product into the woody brush or tree, use equipment capable of penetrating into the living plant tissue under the bark. No surfactant is required for direct injection of this product into woody brush and trees.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

For frill application, apply a 50 to 95-percent solution of this product in water, with 0.5% or more by volume of a nonionic surfactant, to either a continuous frill around the tree or to cuts evenly spaced around the tree below all branches. As tree diameter increases, better results can be achieved by applying this product to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow this product to run out of the frill or cut areas. In species that freely exude sap, make the frill or cuts at an oblique angle to produce a cupping effect and apply a 95-percent solution of this product with a nonionic surfactant as described above. For maximum performance, make this application during period of active growth and after full leaf expansion.

### Modified High-Volume and Low-Volume Backpack Application

For control and partial control of woody bush, trees and vines listed on this label when using a backpack sprayer or other handheld equipment and a directed low-volume foliar application technique, apply a 4 to 8-percent solution of this product containing 0.5 to 1 percent by volume of a nonionic surfactant evenly over the plant crown to cover 50 percent of the upper foliage of the undesirable vegetation.

**WOODY BRUSH, TREES AND VINES RATE TABLE**

Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
Alder	2.3 – 3	0.75 – 1.2
Ash <sup>1</sup>	1.5 – 3.75	0.75 – 1.5
Aspen, quaking	1.5 – 2.3	0.75 – 1.2
Bearclover (Bearmat) <sup>1</sup>	1.5 – 3.75	0.75 – 1.5
Beech <sup>1</sup>	1.5 – 3.75	0.75 – 1.5
Birch	1.5	0.75
Blackberry	2.3 – 3	0.75 – 1.2
Blackgum	1.5 – 3.75	0.75 – 1.5
Bracken	1.5 – 3.75	0.75 – 1.5
Broom; French, Scotch	1.5 – 3.75	1.2 – 1.5
Buckwheat, California <sup>1</sup>	1.5 – 3	0.75 – 1.5
Cascara <sup>1</sup>	1.5 – 3.75	0.75 – 1.5
Castor bean	1.5 – 3.75	1.5
Also for control, inject 4 milliliters of this concentrated (undiluted) product per plant directly into the lower portion of the main stem using a handheld injection device. <sup>2</sup> No surfactant is required.		
Catsclaw <sup>1</sup>	–	1.2 – 1.5
For partial control, apply this product when at least 50 percent of the new leaves are fully developed.		
Ceanothus <sup>1</sup>	1.5 – 3.75	0.75 – 1.5
Chamise <sup>1</sup>	1.5 – 3.75	0.75
Cherry; bitter, black, pin	1.5 – 3.75	1 – 1.5
Cottonwood, eastern	1.5 – 3.75	0.75 – 1.5
Coyote brush	2.3 – 3	1.2 – 1.5
For control, apply this product when at least 50 percent of the new leaves are fully developed.		
Cypress; swamp, bald	1.5 – 3.75	0.75 – 1.5

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
<b>Deerweed</b>	1.5 – 3.75	0.75 – 1.5
<b>Dewberry</b>	2.3 – 3	0.75 – 1.2
<b>Dogwood<sup>1</sup></b>	3 – 3.75	1 – 2
<b>Elderberry</b>	1.5	0.75
<b>Elm<sup>1</sup></b>	1.5 – 3.75	0.75 – 1.5
<b>Eucalyptus, blue gum</b>	–	1.5
For control of eucalyptus re-sprouts, apply this product using a handheld sprayer when re-sprouts are 6 to 12 feet tall. Ensure complete coverage.		
<b>Gallberry</b>	1.5 – 3.75	0.75 – 1.5
<b>Gorse<sup>1</sup></b>	1.5 – 3.75	0.75 – 1.5
<b>Hackberry, western</b>	1.5 – 3.75	0.75 – 1.5
<b>Hasardia<sup>1</sup></b>	1.5 – 3	0.75 – 1.5
<b>Hawthorn</b>	1.5 – 2.3	0.75 – 1.2
<b>Hazel</b>	1.5	0.75
<b>Hickory<sup>1</sup></b>	3 – 3.75	1 – 2
<b>Honeysuckle</b>	2.3 – 3	0.75 – 1.2
<b>Hornbeam, American<sup>1</sup></b>	1.5 – 3.75	0.75 – 1.5
<b>Huckleberry</b>	1.5 – 3.75	0.75 – 1.5
<b>Ivy, poison</b>	3 – 3.75	1.5
<b>Kudzu</b>	3	1.5
<b>Locust, black<sup>1</sup></b>	1.5 – 3	0.75 – 1.5
<b>Madrone (re-sprouts)<sup>1</sup></b>	–	1.5
<b>Magnolia, sweetbay</b>	1.5 – 3.75	0.75 – 1.5
<b>Manzanita<sup>1</sup></b>	1.5 – 3.75	0.75 – 1.5
<b>Maple, red</b>	1 – 3.75	0.75 – 1.2
For control, apply a 0.75 to 1.2-percent solution of this product using a handheld sprayer when leaves are fully developed. For partial control, apply 1 to 3.75 quarts per acre as a broadcast application.		
<b>Maple, sugar</b>	–	0.75 – 1.2
For control, apply this product using a handheld sprayer when at least 50 percent of the new leaves are fully developed.		
<b>Maple, vine<sup>1</sup></b>	1.5 – 3.75	0.75 – 1.5
<b>Monkey flower<sup>1</sup></b>	1.5 – 3	0.75 – 1.5
<b>Oak; black, white<sup>1</sup></b>	1.5 – 3	0.75 – 1.5
<b>Oak; northern pin</b>	1.5 – 3	0.75 – 1.2
For control, apply this product when at least 50 percent of the new leaves are fully developed.		
<b>Oak, poison</b>	3 – 3.75	1.5
Repeat applications might be necessary to maintain control. Application in the fall must be made		

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
before leaves lose green color.		
<b>Oak, post</b>	2.3 – 3	0.75 – 1.2
<b>Oak, red</b>	–	0.75 – 1.2
For control, apply this product using a handheld sprayer when at least 50 percent of the new leaves are fully developed.		
<b>Oak, scrub<sup>1</sup></b>	1.5 – 3	0.75 – 1.5
<b>Oak, southern red</b>	1.5 – 3.75	1 – 1.5
<b>Orange, Osage</b>	1.5 – 3.75	0.75 – 1.5
<b>Peppertree, Brazilian (Florida holly)<sup>1</sup></b>	1.5 – 3.75	1.5
<b>Persimmon<sup>1</sup></b>	1.5 – 3.75	0.75 – 1.5
<b>Pine</b>	1.5 – 3.75	0.75 – 1.5
<b>Poplar, yellow<sup>1</sup></b>	1.5 – 3.75	0.75 – 1.5
<b>Prunus</b>	1.5 – 3.75	1 – 1.5
<b>Raspberry</b>	2.3 – 3	0.75 – 1.2
<b>Redbud, eastern</b>	1.5 – 3.75	0.75 – 1.5
<b>Redcedar, eastern</b>	1.5 – 3.75	0.75 – 1.5
<b>Rose, multiflora</b>	1.5	0.75
Make application prior to leaf deterioration by leaf-feeding insects.		
<b>Russian olive<sup>1</sup></b>	1.5 – 3.75	0.75 – 1.5
<b>Sage, black</b>	1.5 – 3	0.75
<b>Sage, white<sup>1</sup></b>	1.5 – 3	0.75 – 1.5
<b>Sagebrush, California</b>	1.5 – 3	0.75
<b>Salmonberry</b>	1.5	0.75
<b>Saltbush</b>	–	1
<b>Saltcedar<sup>1</sup></b>	3 – 3.75	1 – 2
For partial control, apply a 1 to 2-percent solution of this product using a handheld sprayer or 3 to 3.75 quarts per acre as a broadcast application. For control, apply a 1 to 1.5-percent solution of this product in a tank-mix with Arsenal herbicide or Arsenal Herbicide Applicators Concentrate using a handheld sprayer. For control using broadcast application, apply 1.5 quarts of this product per acre in a tank-mix with an appropriate rate of Arsenal herbicide or Arsenal Herbicide Applicators Concentrate to plants less than 6 feet tall. To control saltcedar greater than 6 feet tall using broadcast application, apply 3 quarts of this product per acre in a tank-mix with a higher rate of Arsenal herbicide or Arsenal Herbicide Applicators Concentrate.		
<b>Sassafras<sup>1</sup></b>	1.5 – 3.75	0.75 – 1.5
<b>Sea Myrtle</b>	–	1
<b>Sourwood<sup>1</sup></b>	1.5 – 3.75	0.75 – 1.5
<b>Sumac; laurel, poison, smooth, sugarbush, winged<sup>1</sup></b>	1.5 – 3	0.75 – 1.5
<b>Sweetgum</b>	1.5 – 2.3	0.75 – 1.5

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
<b>Swordfern<sup>1</sup></b>	1.5 – 3.75	0.75 – 1.5
<b>Tallowtree, Chinese</b>	–	0.75
<b>Tan oak (re-sprouts)<sup>1</sup></b>	–	1.5
<b>Thimbleberry</b>	1.5	0.75
<b>Tobacco, tree<sup>1</sup></b>	1.5 – 3	0.75 – 1.5
<b>Toyon<sup>1</sup></b>	–	1.5
<b>Trumpetcreeper</b>	1.5 – 2.3	0.75 – 1.2
<b>Virginia creeper</b>	1.5 – 3.75	0.75 – 1.5
<b>Waxmyrtle, southern<sup>1</sup></b>	1.5 – 3.75	1.5
<b>Willow</b>	2.3	0.75
<b>Yerba Santa, California<sup>1</sup></b>	–	1.5

<sup>1</sup> Partial control

<sup>2</sup> When using stem injection, the combined total use of this product must not exceed 8 quarts per acre per year. At 4 milliliters of concentrated (undiluted) product per stem, 8 quarts will treat approximately 1890 stems per acre per year. The number of stems that can be treated per acre will vary depending on the injection volume and the concentration of this product in the application solution.

**Other woody brush and trees listed on this label** – For partial control, apply 1.5 to 3.75 quarts of this product per acre as a broadcast application or a 0.75 to 1.5-percent solution using a handheld sprayer and a spray-to-wet application technique.

### 14.0 LIMIT OF WARRANTY AND LIABILITY

Seamless Control LLC (“Company”) warrants that this product conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall use this product only for the purposes of and in accordance with the Complete Directions for Use label (“Directions”) and shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss, injuries or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, crop injury or failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, use and/or application in any manner not explicitly set forth in or inconsistent with the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

## II. DIRECTIONS FOR USE ON AQUATIC AND TERRESTRIAL SITES

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

---

*[Optional text, if applicable: [INSERT BRAND NAME and Design [and] TRUEBLUE ADVANTAGE PROVEN RELIABLE SUPPORT and Design] are [registered] trademarks of Seamless Control LLC.]*

*[Optional text, if applicable: Roundup, Roundup Custom and Design,] are [registered] trademarks of Monsanto Technology LLC and used under license.]*

*[Optional text: All other trademarks are the property of their respective owners.]*

---

*In case of an emergency involving this product, or for medical assistance,  
call [Optional text, if applicable: toll-free, collect, day or night, (314) 694-4000 [Or insert alternative  
telephone number, as appropriate].*

---

Packed for:  
SEAMLESS CONTROL LLC  
14111 SCOTTS LAWN ROAD  
MARYSVILLE, OH 43041

©[YEAR]

*[Insert print plate number]*

*[Insert barcode]*