



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

November 10, 2021

Arianna Shorey  
Regulatory Consultant  
Infinicrop, LLC  
c/o Pyxis Regulatory Consulting Inc.  
4110 136th St. Ct. NW  
Gig Harbor, WA 98332

Subject: Registration Review Label Amendments for ATRAZINE Incorporating Mitigation Measures from the Interim Decision and the Technical Registrants' Commitments for the Endangered Species Act (ESA) Biological Evaluation  
*Product Name:* Atrazine 90DF  
*EPA Registration Number:* 94144-1  
*Application Date:* 11/6/2020  
*Decision Number:* 567676

Dear Arianna Shorey:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the ATRAZINE Interim Decision and with the technical registrants' commitments for the ESA Biological Evaluation. The Agency has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved

Page 2 of 2  
EPA Reg. No. 94144-1  
Decision No. 567676

labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Ben Tweed at [Tweed.  
Benjalmin@epa.gov](mailto:Benjalmin@epa.gov).

Sincerely,



Linda Arrington, Branch Chief  
Risk Management and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

Enclosure

{Note to reviewer: [Text] in brackets denotes optional text.}

{Note to reviewer: {Text} in braces denotes where in the final label text will appear and notes to reviewer.}

{BOOKLET FRONT PANEL}

## RESTRICTED USE PESTICIDE

### (Ground and Surface Water Concerns)

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification. This product is a restricted use herbicide due to ground and surface water concerns. Users must read and follow all precautionary statements and instructions for use in order to minimize potential for atrazine to reach ground and surface water.

ATRAZINE	GROUP	5	HERBICIDE
----------	-------	---	-----------

## Atrazine 90DF

For season-long weed control in Chemical fallow, Corn, Fallowland, Guava, Lawns, Macadamia nuts, Sorghum, Sugarcane and Turf.

#### ACTIVE INGREDIENTS:

Atrazine..... 88.4%  
Related compounds..... 1.6%

**OTHER INGREDIENTS:** ..... 10.0%

**TOTAL:** ..... 100.0%

### KEEP OUT OF REACH OF CHILDREN CAUTION

[See] [inside] [label] [booklet] [for] [additional] [Precautionary Statements][,] [and] [Directions for Use] [including] [Storage and Disposal] [instructions][.]


**EPA Reg. No.: 94144-1**

**EPA Est. No.:**

#### Manufactured for:

INFINICROP, LLC  
2364 East Southwood Road  
San Tan Valley, AZ 85240

**Net Weight:**



## {LANGUAGE INSIDE BOOKLET}

<b>FIRST AID</b>
<b>IF SWALLOWED:</b> <ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b> <ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF IN EYES:</b> <ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>HOTLINE NUMBER</b>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12 PM Pacific Standard Time. In the event of a medical emergency, call your poison control center at 1-800-222-1222.
<b>NOTE TO PHYSICIAN:</b> There is no specific antidote for atrazine. If this product is ingested, induce emesis or lavage stomach. The use of an aqueous slurry of activated charcoal may be considered.

### **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear protective eyewear.

#### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Mixers, loaders, cleaners of equipment or spills, and other handlers exposed to the concentrate must wear:** Coveralls over long-sleeved shirt and long pants, chemical-resistant gloves made of barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or Viton ≥ 14 mils, chemical-resistant footwear plus socks, chemical-resistant apron, and a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters..

**Applicators using spray equipment mounted on their backs must wear:** Coveralls over long-sleeved shirt and long pants, chemical-resistant gloves made of barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or Viton ≥ 14 mils, and chemical-resistant footwear plus socks.

**All other applicators and all other handlers exposed to the dilute must wear:** Long-sleeved shirt and long pants, chemical-resistant gloves made of barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or Viton ≥ 14 mils, and shoes plus socks.

**Aerial application is prohibited.**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

See engineering controls for additional requirements.

### **ENGINEERING CONTROLS**

When applicators use enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(5)), the handler PPE requirements may be reduced or modified as specified in the WPS.

## USER SAFETY RECOMMENDATIONS

### Users should:

Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to Sand and Loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

This product must not be mixed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes and reservoirs. This product must not be applied aerially or by ground within 66 feet of the points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 feet around natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66-foot buffer or setback from runoff points must be planted to crop or seeded with grass or other suitable crop.

Do not contaminate water when disposing of equipment washwaters. This product must not be mixed/loaded or used within 50 feet of all wells including abandoned wells, drainage wells and sinkholes. Operations that involve mixing, loading, rinsing or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spill or equipment leaks, container or equipment rinse or washwater and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading sites. Additional State imposed requirements regarding wellhead setbacks and operational area containment must be observed.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

**NON-TARGET ORGANISM ADVISORY STATEMENT:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

## TILE-OUTLETTED TERRACED FIELDS CONTAINING STANDPIPES

One of the following restrictions must be used in applying atrazine to tile-outletted terraced fields containing standpipes:

1. Do not apply this product within 66 feet of standpipes in tile-outletted terraced fields.
2. Apply this product to the entire tile-outletted terraced field and immediately incorporate it to a depth of 2 to 3 inches in the entire field.
3. Apply this product to the entire tile-outletted terraced field under a no-till practice only when high crop residue management practices are used. High crop residue management practice is described as a crop management practice where little or no crop residue is removed from the field during and after crop harvest.

## USE INFORMATION

Atrazine 90DF will control most annual broadleaf and grass weeds in corn, sorghum, sugarcane and certain other crops as specified on this label. This product may be applied before or after weeds emerge.

In each case where a range of rates is given, the lower rate should be used on coarse textured soil and soil low in organic matter and the higher rate should be used on fine textured soil and soil high in organic matter.

Since this product acts mainly through root absorption, its effectiveness depends on rainfall or irrigation to move it into the root zone. Should weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control.

This product is non-corrosive to equipment and metal surfaces, nonflammable and has low electrical conductivity. Take care to avoid using this product near adjacent desirable plants or in greenhouses, or injury may occur.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

FAILURE TO FOLLOW ALL USE PRECAUTIONS AND RESTRICTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY OR ILLEGAL RESIDUES.

### APPLICATION PROCEDURES

**GROUND APPLICATION:** Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and are the same size. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 10 gallons of spray mixture per acre for all pre-plant incorporated, pre-plant surface, pre-emergence and post-emergence applications (with or without oil or surfactant) with ground equipment. Use a pump with capacity to:

1. Maintain 35 to 40 psi at nozzles.
2. Provide sufficient agitation in tank to keep mixture in suspension.
3. Provide a minimum of 20% bypass at all times.

Use centrifugal pumps which provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gallons per minute per 100-gallon tank size circulated through a correctly positioned sparger tube or jets.

Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For band applications, calculate the amount to be applied per acre as follows:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \frac{\text{broadcast rate}}{\text{per acre}} = \frac{\text{amount needed}}{\text{per acre in field}}$$

### THIS PRODUCT IN WATER OR LIQUID FERTILIZER APPLICATIONS

Nitrogen solutions or complete liquid fertilizers may replace all or part of the water as a carrier for pre-emergence and preplant applications of this product on corn and sorghum. Check the compatibility of this product with liquid fertilizer and/or nitrogen solution as shown below before use. Do not apply in nitrogen solution or complete liquid fertilizer after corn or sorghum emerges, or crop injury may occur.

**Compatibility Test:** Since liquid fertilizers can vary, even within the same analysis, always check compatibility with herbicide(s) each time before use. Be especially careful when using complete suspension or fluid fertilizers as serious compatibility problems are more likely to occur. Commercial application equipment may improve compatibility in some instances. The following test assumes a spray volume of 25 gallons per acre. For other spray volumes, make appropriate changes in the ingredients.

#### Check compatibility using this procedure:

1. Add 1 pint of fertilizer to each of 2 one-quart jars with tight lids.
2. To one of the jars, add 0.25 teaspoon or 1.2 milliliters of a compatibility agent approved for this use, such as MIX™, Compex® or Unite® (0.25 teaspoon is equivalent to 2 pints per 100 gallons of spray). Shake or stir gently to mix.
3. To both jars add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowables next and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix. The appropriate amount of herbicides for this test follows:

**Dry Herbicides:** For each pound to be applied per acre, add 1.5 level teaspoons to each jar.

**Liquid Herbicides:** For each pint to be applied per acre, add 0.5 teaspoon or 2.5 milliliters to each jar.

- After adding all ingredients, put lids on and tighten and invert each jar ten times to mix. Let the mixtures stand 15 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: A) Slurry the dry herbicide(s) in water before addition, or B) Add half of the compatibility agent to the fertilizer and the other half of the emulsifiable concentrate or flowable herbicide before addition to the mixture. If incompatibility is still observed, do not use the mixture.

#### **THIS PRODUCT PLUS EMULSIFIABLE OIL OR OIL CONCENTRATE**

Adding emulsifiable oil (petroleum-derived, petroleum-derived oil concentrate or single or mixed crop-derived oil concentrate) to postemergence water-based sprays in corn and sorghum may improve weed control. However, under certain conditions, the use of either type of oil may seriously injure the crop. To minimize this possibility, observe the following directions. Use one of the following properly emulsified:

- A suitable oil concentrate containing at least 1% but not more than 20% suitable emulsifier or surfactant blend.
- Petroleum-derived oil containing at least 1% suitable emulsifier.

**Note:** In the event of a compatibility problem when mixing oil with this product and water, a compatibility agent such as MIX, Complex or Unite should be used. Any of the above oils contaminated with water or other materials can cause compatibility problems and/or crop injury.

## **DIRECTIONS FOR USE**

### **RESTRICTED USE PESTICIDE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A VIOLATION OF FEDERAL LAW. Before using this product, you must consult the Atrazine Watershed Information Center (AWIC) to determine whether the use of this product is prohibited in your watershed. AWIC can be accessed through [www.atrazinewatershed.info](http://www.atrazinewatershed.info) or 1-866-365-3014. If use of this product is prohibited in your watershed, you may return this product to your point of purchase for a refund.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **RESTRICTIONS**

- Use on roadsides; Conservation Reserve Program (CRP) land; conifers, including Christmas Tree plantings; timber; forestry; and, Miscanthus and other perennial bioenergy crops is prohibited.
- Do not use this product in combination with other herbicides or oils except as specifically described on the label or in literature published by the seller.
- Do not apply this product through any type of irrigation system.
- AERIAL APPLICATION IS PROHIBITED.**
- Not for use in the states of Hawaii or Alaska, or in the U.S. territories (Puerto Rico, Guam, American Samoa, the U.S. Virgin Islands, and the North Mariana Islands).
- Users must only apply to fallow land in the following states according to the prescribed rotation pattern in the table below:

<b>Fallow Rotation Pattern</b>	<b>Fallow Use Authorized in These States Only</b>
Wheat-Corn-Fallow	CO, KS, ND, NE, SD & WY
Wheat-Fallow-Wheat	CO, KS, ND, NE, SD & WY
Wheat-Sorghum-Fallow	AR, CO, GA, IL, KS, LA, MS, MO, NE, NM, NC, OK, SD & TX

### **Endangered Species**

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/espp/>, call 1-844-447-3813, or email [ESPP@epa.gov](mailto:ESPP@epa.gov). You must use the Bulletin valid for the month in which you will apply the product.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the REI of 12 hours. Exception: If the product is soil-injected or soil-incorporated, the WPS, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, chemical-resistant gloves made of barrier laminate, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, or Viton  $\geq$  14 mils, and shoes plus socks.

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the WPS for agricultural use pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter or allow others to enter treated area until sprays have dried.

### **MANDATORY SPRAY DRIFT MANAGEMENT**

#### **Ground Boom Applications:**

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a coarse or coarser droplet size (ASABE S572).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.
- User must maintain a 15 foot (4.6 meter) in-field downwind buffer (in the direction in which the wind is blowing) from the edge of streams and rivers, as well as high-tide line for all estuarine/marine environments.

#### **Boomless Ground Applications:**

- Applicators are required to use a coarse or coarser droplet size (ASABE S572).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.
- User must maintain a 15 foot (4.6 meter) in-field downwind buffer (in the direction in which the wind is blowing) from the edge of streams and rivers, as well as high-tide line for all estuarine/marine environments.

### **SPRAY DRIFT ADVISORIES**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.  
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### **Controlling Droplet Size – Ground Boom**

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **BOOM HEIGHT – Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.



## **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

## **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## **WIND**

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

### **Boomless Ground Applications:**

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

### **Handheld Technology Applications:**

- Take precautions to minimize spray drift.

## **WEED RESISTANCE MANAGEMENT**

For resistance management, Atrazine 90DF is a Group 5 herbicide. Any weed population may contain or develop plants naturally resistant to Atrazine 90DF and other Group 5 herbicides. Weed species with acquired resistance to Group 5 herbicides may eventually dominate the weed population if Group 5 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Atrazine 90DF or other Group 5 herbicides. Users should scout before and after application.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of Atrazine 90DF or other Group 5 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage ( or other mechanical control methods), cultural ( e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout prior to and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your INFINICROP, LLC representative. In addition to the guidance above, registrants are encouraged to incorporate the appropriate elements of Best Management Practices from HRAC and WSSA on the label.

### **MIXING PROCEDURES (All Uses)**

1. Be sure sprayer is clean and not contaminated with any other materials or crop injury or sprayer clogging may result.
2. Fill tank one-fourth full with clean water, nitrogen solution or complete liquid fertilizer.
3. Start agitation.
4. Be certain that the agitation system is working properly and creates a rippling or rolling action on the liquid surface.
5. Add this product directly into tank.
6. Continue filling tank until 90% full. Increase agitation, if necessary, to maintain surface action.
7. Add emulsifiable oil, oil concentrate or tank mix herbicide(s) after this product is thoroughly suspended.
8. Finish filling tank.
9. Empty tank as completely as possible before refilling to prevent buildup of oil or emulsifiable concentrate residue.  
Maintain agitation to avoid separation of materials.
10. If an oil or emulsifiable concentrate film starts to build up in tank, drain it and clean with strong detergent solution or solvent.
11. Clean sprayer thoroughly immediately after use by flushing system with water containing a detergent.

### **ROTATIONAL CROPS (All Uses)**

1. Land treated with this product should not be planted to any crop except corn or sorghum until the following year, or injury may occur.
2. If this product is applied after June 10, do not rotate with crops other than corn or sorghum the next year, or injury may occur.
3. In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use only when corn or sorghum is to follow corn or sorghum, or when a crop of untreated corn or sorghum is to precede other rotational crops.
4. In Western MN and Eastern parts of the Dakotas, KS, corn treated with this product should not be followed with soybeans if the broadcast rate applied was more than 2.2 lbs. per acre (2 lbs. a.i.), or comparable rate in a band, or injury to soybeans may occur.
5. Injury may occur to soybeans planted the year following an application on soils having a calcareous surface layer.
6. Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains or small-seeded legumes and grasses the year following application of this product, or injury may occur.

### **THIS PRODUCT APPLIED ALONE—CORN AND GRAIN SORGHUM**

#### **PRE-PLANT SURFACE APPLIED, PRE-PLANT INCORPORATED OR PRE-EMERGENCE (OR POST-EMERGENCE AT 2.2 LBS. OF THIS PRODUCT (2 LBS. A.I.) PER ACRE WITH OIL)**

**Broadleaf or Grass Weeds Controlled:** Annual morning glory, Barnyardgrass\* (Watergrass), Cocklebur\*\*, Giant foxtail\*\*, Green foxtail\*, Groundcherry, Jimsonweed, Kochia, Lambsquarters, Large (Hairy) crabgrass\*\*, Mustard, Nightshade, Pigweed, Purslane, Ragweed, Sicklepod\*\*, Velvetleaf\* (Buttonweed), Wild oats, Witchgrass\*\* (*Panicum capillare*) and Yellow foxtail\*.

#### **POST-EMERGENCE WITH EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER 1.3 LBS. OF THIS PRODUCT (1.2 LBS. A.I.) PER ACRE**

**Broadleaf Weeds Controlled:** Annual morningglory, Cocklebur, Jimsonweed, Lambsquarters, Mustards, Pigweed, Ragweed, Smartweed, Velvetleaf\* (Buttonweed) and Wild buckwheat.

\*Partial control only on Medium and Fine textured soils.

\*\*Partial control only.

Where there are state/local requirements regarding atrazine use (including lower maximum rates and/or greater setbacks) which are different from this label, the more restrictive/protective requirements must be followed. Certain states may have established rate limitations within specific geographical areas. Consult your State Lead Pesticide Control Agency for additional information. It is a violation of this label to deviate from State regulations.

## CORN

This product may be applied either before planting, at-planting or after planting as indicated.

### PRE-PLANT SURFACE APPLIED

**Broadleaf and Grass Control:** Use on medium and fine textured soils with minimum-tillage or no-tillage systems only in CO, IA, IL, IN, KS, KY, MM, MO, MT, ND, NE, SD, WI and WY. Apply the specified rate of this product shown in "TABLE 1" up to 45 days prior to planting. If unsatisfactory length of weed control results from adverse environmental conditions following early treatment, a follow-up application of an appropriately labeled herbicide may be used. If the follow-up treatment includes atrazine, do not exceed the labeled rate for corn indicated in "TABLE 1". On coarse textured soils, do not apply more than two weeks prior to planting.

If weeds are present at the time of treatment, apply in tank mix combination with a contact herbicide (for example, paraquat or glyphosate). Observe directions for use, use precautions and restrictions for use on the label of the contact herbicide.

**Precaution:** To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting or weed control will be diminished.

### PRE-PLANT INCORPORATED

**Broadleaf and Grass Control:** Broadcast in the Spring after plowing at the rate indicated in "TABLE 1". Application may be made before, during or after final seedbed preparation. Avoid deep incorporation of this product. For best results apply within two weeks prior to planting.

### PRE-EMERGENCE OR AT-PLANTING

**Broadleaf and Grass Control:** Apply during or shortly after planting, prior to weed emergence, at the rate indicated in "TABLE 1".

### POST-EMERGENCE

**Broadleaf and Grass Control:** Apply before weeds exceed 1.5 inches in height and before Corn exceeds 12 inches in height at the rate specified in "TABLE 1".

<b>TABLE 1</b>
<b>FOR CONTROL OF BROADLEAF AND GRASS WEEDS IN CORN*</b> <b>For All Soil Applications Prior To Crop Emergence</b>
<b>On Highly Erodible Soils**</b> If conservation tillage is practiced, leaving at least 30% of the soil covered with plant residues at-planting, apply a maximum of 2.2 lbs. of this product (2 lbs. a.i.) per acre as a single broadcast spray. If the soil coverage with plant residue is less than 30% at-planting, a maximum of 1.7 lbs. of this product (1.5 lbs. a.i.) per acre may be applied.
<b>On Soils Not Highly Erodible**</b> Apply a maximum of 2.2 lbs. of this product (2 lbs. a.i.) per acre as a single pre-emergence broadcast spray.
*Broadleaf control (Eastern CO, Western KS, Western NE, NM, OK Panhandle, West TX and Eastern WY): On sand, loamy sand, sandy loam, mild to strongly alkaline soil and all recently leveled soil, apply no more than 1.3 lbs. of this product (1.2 lbs. a.i.) per acre, either pre-plant or pre-emergence. On other soils in these areas, apply rate in this table for broadleaf and grass control. **As defined by the Natural Resource Conservation Service

### FOR POST-EMERGENCE APPLICATION

If no atrazine was applied prior to corn emergence, apply a maximum of 2.2 lbs. of this product (2 lbs. a.i.) per acre broadcast. If a post-emergence treatment is required following an earlier herbicide application, the total atrazine applied may not exceed 2.7 lbs. of this product (2.5 lbs. a.i.) per calendar year.

Where there are State/Local requirements regarding atrazine use (including lower maximum rates and/or greater set-backs) which are different from the label, the more restrictive/protective requirements must be followed. Certain states may have established rate limitations within specific geographical areas. Consult your State Lead Pesticide Control Agency for additional information. It is a violation of this label to deviate from State use regulations.

**POST-EMERGENCE WITH EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER:** Add the following volume of one of the type oils indicated for ground application unless the oil label specifies otherwise:

Oil Type	Ground Application Per Acre
Oil Concentrate (Crop or Petroleum-derived)	1 qt.
Petroleum -derived oil	1 gal.

**Note:** Crop-derived or petroleum-derived oil concentrates should contain at least 1%, but not more than 20%, suitable emulsifier or surfactant blend. Petroleum-derived oils should contain at least 1% suitable emulsifier.

**Broadleaf and Grass Control:** For post-emergence control of those weeds listed under "*THIS PRODUCT APPLIED ALONE — CORN AND GRAIN SORGHUM; PRE-PLANT SURFACE-APPLIED, PRE-PLANT INCORPORATED OR PREEMERGENCE*", broadcast 2.2 lbs. of this product (2 lbs. a.i.) per acre plus emulsifiable oil or oil concentrate after weed emergence, but before weeds reach 1.5 inches in height and before corn exceeds 12 inches in height.

**Broadleaf Control:** For post-emergence control of those weeds listed under "*THIS PRODUCT APPLIED ALONE — CORN AND GRAIN SORGHUM; POST-EMERGENCE WITH EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER*", broadcast 1.3 lbs. of this product (1.2 lbs. a.i.) per acre plus emulsifiable oil or oil concentrate before lambsquarters and pigweed reach 6 inches in height and before all other weeds reach 4 inches in height. A cultivation may be necessary if all weeds are not controlled or if weeds regrow.

**Use Precautions for Applications of This Product Plus Emulsifiable Oil or Oil Concentrate in Water to Corn**

- Adding other insecticides, herbicides, liquid fertilizers or other materials is not recommended, because they may cause compatibility problems or crop injury.
- Store and handle emulsifiable oil and oil concentrate carefully. Oil contaminated with even a small amount of water may not emulsify properly when added to the tank.

**Use Restrictions for Applications of This Product Plus Emulsifiable Oil or Oil Concentrate in Water to Corn**

- Do not use oil in sprays of this product when corn is under stress from prolonged cold, wet weather, poor fertility or other factors or when corn is wet and succulent from recent rainfall, as crop injury may occur.
- Do not use oils in sprays when treating inbred lines or any breeding stock as injury may occur.
- Do not exceed 2.7 lbs. of this product (2.5 lbs. a.i.) per acre per calendar year.
- Post-emergence application to corn must be made before corn exceeds 12 inches in height.

**TANK MIXTURE WITH THIS PRODUCT FOR CORN**

This product may be tank-mixed with these herbicides for control of certain broadleaf and grass weeds in corn:

Alachlor	Metolachlor + simazine
Alachlor + glyphosate	Metolachlor + simazine + glyphosate
Alachlor + paraquat	Metolachlor + simazine + paraquat
Butylate	Paraquat
Glyphosate	Propachlor
Metolachlor	Simazine
Metolachlor + glyphosate	Simazine + glyphosate
Metolachlor + paraquat	Simazine + paraquat

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

When tank-mixing or sequentially applying atrazine or products containing atrazine to corn, do not exceed an application rate of 2 lbs. atrazine active ingredient for any single application, and the total lbs. of atrazine applied (lbs. a.i./A) must not exceed 2.5 lbs. per acre per year.

When tank-mixing or sequentially applying atrazine or products containing atrazine to crops other than corn or sorghum, the total lbs. of atrazine applied (lbs. a.i./A) must not exceed the specific seasonal rate limits as noted in the use directions.

**Note:** When the labels of the above herbicides refer to Atrazine 80W, use equivalent rates of this product. One pound of Atrazine 80W equals 0.9 pound of Atrazine 90DF.

**THIS PRODUCT PLUS SIMAZINE**

In addition to the weeds listed under "THIS PRODUCT APPLIED ALONE — CORN AND GRAIN SORGHUM; PRE-PLANT SURFACE-APPLIED, PRE-PLANT INCORPORATED OR PRE-EMERGENCE" and "THIS PRODUCT APPLIED ALONE — CORN AND GRAIN SORGHUM; POST-EMERGENCE WITH EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER", this combination also controls carpetweed, crabgrass and Fall panicum. Broadcast tank mix before planting, at-planting or after planting, but before crop and weeds emerge, at rates in "TABLE 2". Use the 1:1 ratio for control of most weeds. Use the 1:2 ratio for expected heavy infestations of crabgrass and Fall panicum. Cultivate shallowly if weeds develop.

**PRE-PLANT SURFACE APPLIED**

Use on Medium and Fine textured soils with minimum-tillage or no tillage systems only in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, WI and WY.

Apply the labeled rate of this product and simazine shown in "TABLE 2" up to 45 days prior to planting Refer to "THIS PRODUCT ALONE" section for information if weeds should develop following the early treatment. On Coarse textured soils, do not apply more than 2 weeks prior to planting. Refer to "THIS PRODUCT APPLIED ALONE - PRE-PLANT SURFACE APPLIED" section of the corn label for additional details.

If weeds are present at time of treatment, apply in a tank mix combination with a contact herbicide (for example, paraquat or glyphosate).

It is the pesticide user’s responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Precaution:** To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

**PRE-PLANT INCORPORATED**

Apply to the soil and incorporate in Spring before, during or after final seedbed preparation. Avoid deep incorporation. For best results, apply within 2 weeks prior to planting.

**PRE-EMERGENCE**

Apply during or shortly after planting, but before crop and weeds emerge.

TABLE 2				
TANK MIXTURES WITH SIMAZINE ON CORN				
SOIL TEXTURE	BROADCAST RATE PER ACRE			
	1:1 Ratio*		1:2 Ratio**	
	This Product	Simazine	This Product	Simazine
<b>COURSE:</b> Sand, Loamy sand, Sandy loam	1.1 lbs.	Refer to product label	0.75 lb.	Refer to product label
<b>MEDIUM:</b> Loam, Silt loam, Silt, Clay loam, Sandy clay loam, Silty clay loam, Sandy clay or Silty clay with low organic matter	1.3 lbs.	Refer to product label	0.9 lb.	Refer to product label
<b>FINE:</b> Loam, Silt loam, Silt, Clay loam, Sandy clay loam, Silty clay loam, Sandy clay or Silty clay with medium to high organic matter and Clay (including dark prairie soils of the Corn Belt)	1.6 lbs.	Refer to product label	1.1 lbs.	Refer to product label
*For control of most weeds.				
**For control of expected heavy infestations of crabgrass and Fall panicum.				

Refer to "CORN" sections of this label and to Simazine label for further directions, limitations and precautions.

### THIS PRODUCT PLUS SIMAZINE PLUS PARAQUAT

For kill of existing vegetation and residual weed control where corn will be planted directly into a cover crop, established sod or in previous crop residues, apply 1.2 to 2.2 lbs. of this product (1.08 to 2 lbs. a.i.) plus the label specified rate of Simazine, or label specified rate of Simazine plus the label specified amount of paraquat per sprayed acre prior to, during or after planting, but before crop emergence. Use the lower rates of atrazine and simazine on Light soils and the higher rates on Heavy clay soils. Use 20 to 60 gallons of diluted spray per sprayed acre. Add Surf-Ac® 820 or another nonionic surfactant at the rate of 8 fluid ounces per 100 gallons of diluted spray. Add this product to the spray tank while agitating, then add the simazine, paraquat and Surf-Ac® 820. Continue agitation during application. Use the higher rate of paraquat if existing vegetation is 4 to 6 inches tall. This mixture will not control weeds taller than 6 inches. Refer to paraquat and simazine labels for further limitations and precautions.

### THIS PRODUCT PLUS SIMAZINE PLUS GLYPHOSATE

When applied as directed under the conditions described, this tank mixture controls many emerged weeds, and gives pre-emergence control of many annual weeds when corn will be planted directly into a cover crop, established or in previous crop residues.

Apply this tank mixture in 20 to 60 gallons of water per acre immediately before, during or after planting, but before crop emergence.

As density of stubble, crop residue or weeds increases, increase spray gallonage and rate within the specified ranges to ensure complete coverage.

**Annual Weeds:** Apply atrazine and simazine to actively growing grasses and broadleaf weeds that are 6 inches tall at the rates given in "TABLE 3" plus the label specified rate of glyphosate per acre. If the weeds are over 8 inches tall, increase the rate of glyphosate. Refer to the glyphosate label for specified rate.

**Perennial Weeds:** At normal application rates in minimum-tillage systems, perennial weeds may not be at the proper stage of growth for control. Refer to the glyphosate label for the proper stage of growth for perennial weeds. This combination will provide top kill and reduce competition from many emerged perennial grasses and broadleaf weeds.

#### Restrictions:

- Do not use this tank mix on sand or loamy sand soils.
- Do not exceed the label specified maximum rate of glyphosate per acre in this tank mixture.
- Do not use for control of bermudagrass or johnsongrass in minimum- tillage systems.

<b>Soil Texture</b>	<b>This Product*</b>	<b>Simazine</b>
<b>Coarse:</b> Sandy loam soil	1.1 lbs.	Refer to product label
<b>Medium:</b> Loam, Silt loam, Silt, Sandy clay loam	1.1 to 1.6 lbs.	Refer to product label
<b>Fine:</b> Silty clay, Loam, Clay loam, Sandy clay, Silty clay, Clay	1.3 to 1.7 lbs.	Refer to product label

\* Use the higher rate of atrazine and simazine in the specified range in soil with greater than 3% organic matter.

### FOR ALL SOIL APPLICATIONS PRIOR TO CROP EMERGENCE

**On Highly Erodible Soils\*:** If conservation tillage is practiced, leaving at least 30% of the soil covered with plant residues at planting, apply a maximum of 2.2 lbs. of this product (2 lbs. a.i.) per acre as a single broadcast spray. If the soil coverage with plant residue is less than 30% at planting, a maximum of 1.7 lbs. of this product (1.5 lbs. a.i.) per acre may be applied.

**On Soils Not Highly Erodible\*:** Apply a maximum of 2.2 lbs. of this product (2 lbs. a.i.) per acre as a single broadcast spray.

\*As defined by the Natural Resource Conservation Service.

### USE RESTRICTIONS FOR ALL APPLICATIONS TO CORN

- To avoid crop injury and illegal residues, do not apply more than 2.7 lbs. of this product (2.5 lbs. a.i.) per acre to corn in any one year.

- When tank-mixing or sequentially applying atrazine or products containing atrazine to corn, do not exceed an application rate of 2 lbs. atrazine active ingredient for any single application, and the total lbs. of atrazine applied (lbs. a.i./A) must not exceed 2.5 lbs. a.i. per acre per year.
- When tank-mixing or sequentially applying atrazine or products containing atrazine to crops other than corn or sorghum, the total lbs. of atrazine applied (lbs. a.i./A) must not exceed the specific seasonal rate limits as noted in the use directions.
- For best control of cocklebur and velvetleaf, the application rate cannot be less than 2 lbs. per acre of active ingredient, either alone or in a tank mix combination.
- Following harvest of a treated crop, plow (moldboard or disk-plow) and thoroughly till the soil in the Fall or Spring to minimize possible injury to rotational Spring-seeded crops, regardless of the rate used.
- For Field corn do not harvest for forage within 60 days of application. For Sweet corn, do not harvest forage within 45 days of application.
- For sweet corn, do not apply via mechanically pressurized handgun.

For "POST-EMERGENCE APPLICATIONS PLUS EMULSIFIABLE OIL OR OIL CONCENTRATE", see additional precautions under that section of this label.

## **SORGHUM AND SORGHUM-SUDAN HYBRIDS (Grain and Forage Types)**

Apply before planting, at-planting or after planting. See "CORN" section for weeds controlled.

### **PRE-PLANT SURFACE-APPLIED**

**Broadleaf and Grass Control:** Use on medium and fine textured soil with minimum-tillage or no-tillage systems only in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, WI and WY. Apply specified rate of this product shown in "TABLE 4" up to 45 days prior to planting. If an unsatisfactory length of weed control results from adverse environmental conditions following early treatment, a follow-up application of an appropriately labeled herbicide may be used.

If the follow-up treatment includes atrazine, do not exceed the labeled rate for corn indicated in Table 1.

Under dry conditions, irrigate after application to move atrazine into the soil.

If weeds are present at time of treatment, apply in a tank mix combination with a contact herbicide (for example paraquat or glyphosate).

Observe directions for use, precautions and restrictions on the label of the contact herbicide.

**Precaution:** To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting or weed control will be diminished.

### **PRE-PLANT INCORPORATED**

**Broadleaf and Grass Control:** Broadcast in Spring after plowing at rate in "TABLE 4". Apply before, during or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation. For best results, apply within 2 weeks prior to planting.

#### **Pre-Emergence or At-Planting (Broadleaf and Grass control):**

Apply during or shortly after planting, but prior to weed or crop emergence at rate shown in Table 4.

#### **FOR POST-EMERGENCE APPLICATION (Broadleaf and Grass control):**

Apply before weeds exceed 1.5 inches in height and before sorghum exceeds 12 inches in height. If no atrazine was applied prior to sorghum emergence, apply a maximum of 2.2 lbs. of this product (2 lbs. a.i.) per acre broadcast. If a post-emergence treatment is required following an earlier herbicide application, the total atrazine applied may not exceed 2.7 lbs. of this product (2.5 lbs. a.i.) per acre per calendar year.

Where there are State/Local requirements regarding atrazine use (including lower maximum rates and/or set-backs) which are different from the label, the more restrictive/protective requirements must be followed. Certain states may have established rate limitations within specific geographical areas. Consult your State Lead Pesticide Control Agency for additional information. It is a violation of this label to deviate from State use regulations.

<b>TABLE 4</b>	
<b>Pre-Plant Surface, Pre-Plant Incorporated and Pre-Emergence or At-Planting Applications* (For Broadleaf and Grass Weed Control in Sorghum)</b>	
<b>On Highly Erodible Soils**</b> If conservation tillage is practiced, leaving at least 30% of the soil covered with plant residues at the time of planting, apply a maximum of 2.2 lbs. of this product (2 lbs. a.i.) per acre as a single broadcast spray. If the soil coverage with plant residue is less than 30% at-planting, a maximum of 1.7 lbs. of this product (1.5 lbs. a.i.) per acre may be applied.	
<b>On Soils Not Highly Erodible**</b> Apply a maximum of 2.2 lbs. of this product (2 lbs. a.i.) per acre as a single broadcast spray.	
*Do not apply pre-plant, surface or pre-plant incorporated in AL, AR, FL, GA, LA, MS, NC, NM, OK, SC, TN or TX. Do not apply pre-emergence in NM, OK or TX, except in Northeast OK and the TX Gulf Coast and Blacklands areas. **As defined by the Natural Resource Conservation Service.	

In case of planting failures, sorghum or corn can be replanted. Do not make a second broadcast application or injury may occur. If this product is originally applied in a band and sorghum or corn is replanted in the untreated row middles, this product may be applied in a band to the second planting, provided the maximum application rate of 2.7 lbs. of this product (2.5 lbs. a.i.) per acre per calendar year is not exceeded.

**PRE-EMERGENCE BROADLEAF WEED CONTROL IN FURROW-IRRIGATED BEDDED SORGHUM (AZ and CA only):**

For pre-emergence control of many broadleaf weeds, broadcast 0.9 to 1.3 lbs. of this product (0.8 to 1.2 lbs. a.i.) per acre. Make application after bed preparation, during or after planting, but before sorghum and weeds have emerged and before the first furrow irrigation. Several regular irrigations should follow the application, making sure that all soil is thoroughly wet.

**USE PRECAUTIONS FOR PRE-EMERGENCE APPLICATIONS OF THIS PRODUCT TO FURROW-IRRIGATED BEDDED SORGHUM (AZ and CA):**

To avoid possible sorghum injury, do not use on sand or loamy sand soils or on sorghum planted in the furrow. Additionally, applications made to sorghum growing on alkali soils or where cuts, fills or erosions have exposed calcareous or alkali subsoils, may result in crop injury. In case of crop failure, do not replant sorghum for eight months following application. Corn may be planted immediately.

**POST-EMERGENCE**

**Broadleaf and Grass Weed Control:** Apply before weeds exceed 1 inch in height at the rate indicated in "TABLE 5". Application may be made to sorghum reaching up to 12 inches in height.

<b>TABLE 5</b>		
<b>POST-EMERGENCE BROADLEAF AND GRASS WEED CONTROL IN SORGHUM</b>		
<b>Soil Texture</b>	<b>Minimum Height of Sorghum At Treatment</b>	<b>Maximum Broadcast Rate of This Product Per Acre (lbs. a.i.)</b>
Sand or Loamy sand	DO NOT USE	
Sandy loam	See directions for Broadleaf and Weed control below.	
Silt loam to Sandy clay loams	Completely emerged	2.2 lbs. (2 lbs. a.i.)
Olton and Pullman clay soils	At least 6 inches high	2.2 lbs. (2 lbs. a.i.)
Silty clay loams and heavier soils	Completely emerged	2.2 lbs. (2 lbs. a.i.)
If no atrazine was applied prior to sorghum emergence, apply a maximum of 2.2 lbs. of this product (2 lbs. a.i.) per acre broadcast. If a postemergence treatment is required following an earlier herbicide application, the total atrazine applied may not exceed 2.5 lbs. active ingredient (2.7 lbs. of this product) per acre per calendar year.		



## **POST-EMERGENCE BROADLEAF WEED CONTROL WITH THIS PRODUCT PLUS EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER**

Broadcast 1.3 lbs. of this product (1.2 lbs. a.i.) per acre for control of many broadleaf weeds. Application should be made before lambsquarters and pigweed reach 6 inches in height and before all other weeds reach 4 inches in height. In CO, Western KS, NM, OK, TX and desert regions of AZ and CA, apply when sorghum is 6 to 12 inches in height, but before it reaches boot stage. In all other areas, apply after sorghum reaches the three-leaf stage, but before it exceeds 12 inches in height. Add emulsifiable oil at the rate of 1 gallon per acre or add 1 quart per acre of oil concentrate for ground application. A cultivation may be necessary if all weeds are not controlled or if regrowth of weeds occurs. For the list of weeds controlled, see "*THIS PRODUCT APPLIED ALONE - CORN OR GRAIN SORGHUM - POSTEMERGENCE WITH EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER.*"

**USE PRECAUTIONS FOR APPLICATION OF THIS PRODUCT PLUS EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER TO SORGHUM—** See "*USE PRECAUTIONS FOR APPLICATIONS WITH EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER*" in the "*CORN*" section.

**POST-EMERGENCE BROADLEAF WEED CONTROL WITH THIS PRODUCT PLUS SURFACTANT (CO, Western KS, NM, OK, TX and Desert regions of AZ and CA only):** Broadcast 1.3 lbs. of this product (1.2 lbs. a.i.) plus 0.75 to 1.5 pints of surfactant per acre after sorghum reaches 6 inches in height, but before weeds reach 1.5 inches in height. Apply only on sandy loams and finer textured soils.

### **USE PRECAUTIONS FOR ALL APPLICATIONS TO SORGHUM**

- Application made to Sorghum growing under stress caused by minor element deficiency or to Sorghum growing on highly calcareous soils may result in crop injury.
- Following harvest, plow (moldboard or disk-plow) and thoroughly till the soil in the Fall or Spring to minimize possible injury to Spring-seeded rotational crops, regardless of rate used.
- Injury may occur if both this herbicide, pre-plant surface, preplant incorporated or pre-emergence and an at-planting systemic insecticide applied in-furrow are used.
- For applications to furrow-irrigated bedded sorghum in AZ and CA and for post-emergence applications plus emulsifiable oil, see additional precautions under these sections of this label.

### **USE RESTRICTIONS FOR ALL APPLICATIONS TO SORGHUM**

- Heavy rains immediately following application tend to cause excessive concentrations of herbicide in seed furrow, resulting in possible crop injury. Do not make applications to furrow-planted sorghum until furrows are leveled (plow-in). Level deep planter marks or seed furrows before application.
- Do not apply more than 2.7 lbs. of this product (2.5 lbs. of a.i.) per acre per calendar year.
- For all soil applications prior to crop emergence (except for preemergence use on bedded sorghum in AZ and CA), do not apply to coarse textured soils, i.e., sand, loamy sand, sandy loam or to medium and fine-textured soils having less than 1% organic matter or injury may occur.
- For post-emergence applications, do not apply to sand or loamy sand or injury may occur.
- Do not harvest for forage within 60 days of a pre-emergence application and 45 days of a post-emergent application.
- Do not apply atrazine and propazine products to the same sorghum acre.

### **TANK MIXTURES FOR GRAIN SORGHUM**

**THIS PRODUCT + METOLACHLOR on Grain Sorghum (Cyometrinil-Treated Seed):** This product + metolachlor tank mixture provides control of weeds listed on the metolachlor label and this label.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

When tank-mixing or sequentially applying atrazine or products containing atrazine to sorghum, do not exceed an application rate of 2 lbs. atrazine active ingredient for any single application, and the total lbs. of atrazine applied (lbs a.i./A) must not exceed 2.5 lbs. per acre per year.

When tank-mixing or sequentially applying atrazine or products containing atrazine to crops other than corn or sorghum, the total lbs. of atrazine applied (lbs a.i./A) must not exceed the specific seasonal rate limits as noted in the use directions.

## **WINTER WEED CONTROL IN TEXAS**

For post-emergence control of Winter weeds only, such as annual thistle, henbit and seedling dock on Fall-bedded land in the Gulf Coast and Blacklands of TX, apply 0.9 to 1.1 lbs. of this product (0.8 to 1 lb. a.i.) per acre post-emergence to the weeds in November or December to land that will be planted to corn, grain sorghum or forage sorghum the following Spring. For best results add a suitable surfactant such as Surf-Ac 820 at the rate of 0.5% of the spray volume, an emulsifiable oil at the rate of 1% of the spray volume or an oil concentrate at the rate of 1 quart per acre.

Normal weed control programs may be used in the following: corn, grain sorghum or forage sorghum crops.

### **Use Restriction:**

- Do not plant any crops except corn, forage sorghum or grain sorghum the Spring following this treatment or illegal residues may result.

## **CHEMICAL FALLOW—THIS PRODUCT ALONE**

**WHEAT-SORGHUM-FALLOW (AR, CO, GA, IL, KS, LA, MS, MO, NE, NM, NC, OK, SD & TX only):** This treatment controls annual broadleaf and grass weeds following wheat harvest and in the following sorghum crop when grown under minimum-tillage. Apply 2.5 lbs. of this product (2.25 lbs. a.i.) per acre to wheat stubble immediately following wheat harvest. If weeds are present, remove them with a sweep plow or other suitable implement after application or use an approved contact herbicide before or after application of this product. Plant sorghum into wheat stubble the following Spring with minimum disturbance of the soil. Use a surface planter or a planter leaving a shallow furrow. If weeds are present at-planting, remove them with a sweep plow or other suitable implement before planting.

For the list of weeds controlled, see "*THIS PRODUCT APPLIED ALONE - CORN OR GRAIN SORGHUM - Pre-plant Surface-Applied, Pre-plant Incorporated, or Pre-emergence*".

### **Use Restrictions:**

- For soils in SD with a pH of 7.5 or greater, do not apply more than 1.6 lbs. of this product (1.5 lbs. A.i.) per acre for any application.
- For soils in SD with a pH of less than 7.5, do not apply more than 2.2 lbs. of this product (2 lbs. a.i.) per acre for any application.
- For all other locations, do not apply more than 2.5 lbs. of this product (2.25 lbs. a.i.) per acre for any application.
- Do not apply more than one application per cycle.
- Use only on silt loam or fine-textured soil, or crop injury may result.
- Wheat-Sorghum-Fallow cropping sequence must be followed.
- Do not apply following Sorghum harvest.
- To avoid illegal residues, do not graze or feed forage from treated areas to livestock.
- To avoid illegal residues and crop injury, do not plant any crop other than those on this label within 18 months following treatment.

**WHEAT-CORN-FALLOW (CO, KS, ND, NE, SD, AND WY ONLY):** This product controls cheatgrass (downy brome, chess), kochia, mustard, pigweed, Russian thistle, wild lettuce, wild sunflower and volunteer wheat during period after wheat harvest. Weed control may extend into following corn crop grown under minimum-tillage.

Follow directions for use and "*USE RESTRICTIONS*" in the "*WHEAT-SORGHUM-FALLOW*" section above, substituting corn for references to sorghum, and the Wheat-Corn-Fallow specific restrictions below.

### **Use Restrictions:**

- On soils in ND and SD with pH greater than 7.5, do not exceed 1.6 lbs. of this product (1.5 lbs. a.i.) per acre for any application.
- For soils with a pH less than 7.5 in ND and SD, apply 1.1 to 2.2 lbs. of this product (1 to 2 lbs. a.i.) per acre. Use the higher rate on fine-textured soils and where heavy weed infestations are expected; use the lower rate on coarse-textured soils and where light weed infestations are expected. Do not apply more than 2.2 lbs. of this product (2 lbs. a.i.) per acre for any application.
- For all other locations, do not apply more than 2.5 lbs. of this product (2.25 lbs. a.i.) per acre for any application. In the event grasses are present the following Spring, use a grass herbicide registered for use on corn.
- Do not apply more than one application per cycle.

**WHEAT-FALLOW-WHEAT (CO, KS, ND, NE, SD AND WY ONLY):** For pre-emergence and control of cheatgrass (downy brome, chess), common lambsquarters, field pennycress, kochia, mustard, Russian thistle, wild lettuce and suppression of volunteer wheat during fallow period of a Wheat-Fallow-Wheat rotation, broadcast 0.5 to 1.1 lbs. of this product (0.45 to 1 lb. a.i.) per acre on all soils except those listed under "*USE RESTRICTIONS*". For control of pigweed and wild sunflower, use the higher rate. Apply to stubble ground. Treat only once during same fallow period.

**Use Restrictions:**

- For soils in ND and SD with a pH of 7.5 or greater, do not apply more than 1.6 lbs. of this product (1.5 lbs. active ingredient) per acre for any application.
- For soils in ND and SD with a pH of less than 7.5, do not apply more than 2.2 lbs. of this product (2 lbs. active ingredient) per acre for any application.
- For all other locations, do not apply more than 2.5 lbs. of this product (2.25 lbs. active ingredient) per acre for any application.
- Do not apply more than one application per cycle.

**TANK MIXTURES FOR CHEMICAL FALLOW**

**WHEAT-SORGHUM-FALLOW OR WHEAT-CORN-FALLOW (KS, NE)**

**THIS PRODUCT PLUS PARAQUAT:** If weeds are present at application, a tank mix with paraquat may be used. Broadcast 2.5 lbs. of this product (2.25 lbs. a.i.) plus a suitable amount of paraquat in 20 to 60 gallons of water per acre by ground equipment. Add 0.5 to 1 pint of a non-ionic surfactant, such as Surf-Ac 820, per 100 gallons of spray mixture. Add this product to spray tank first and thoroughly mix with water. Then add paraquat followed by surfactant. Use the higher rate of paraquat if weeds are 4 to 6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground.

Refer to paraquat label for further directions, precautions and limitations.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Use Restrictions:**

- Do not apply more than 2.5 lbs. of this product (2.25 lbs. active ingredient) per acre for any application.
- Do not apply more than one application per cycle.

**WHEAT-FALLOW-WHEAT (CO, KS, ND, NE, SD AND WY ONLY)**

**THIS PRODUCT PLUS PARAQUAT:** If weeds are present at application, a tank mix with paraquat may be used. Broadcast 0.5 to 1.1 lbs. of this product (0.45 to 1 lb. a.i.) plus a suitable amount of paraquat in 20 to 60 gals, of water per acre by ground equipment. Add 0.5 to 1 pt. of a non-ionic surfactant, such as Surf-Ac 820 per 100 gals, of spray mixture. Add this product to spray tank first and thoroughly mix with water. Then add paraquat followed by surfactant. Use the higher rate of paraquat specified on the label if weeds are 4 to 6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground.

Refer to the paraquat label for further directions, precautions and limitations.

If weeds are present at application and this product is used alone, use either an approved contact herbicide before or after treatment or tillage after treatment.

Use tillage to control weeds which escape during fallow period. Till before planting. For this product applied alone or in tank mixture with paraquat, plant at least 2 inches deep and 12 months or more after application.

**Use Precautions:**

- Avoid spray overlap.

**Use Restrictions:**

- For soils in ND and SD with a pH of 7.5 or greater, do not apply more than 1.6 lbs. of this product (1.5 lbs. a.i.) per acre for any application.
- For soils in ND and SD with a pH of less than 7.5, do not apply more than 2.2 lbs. of this product (2 lbs. a.i.) per acre for any application.
- For all other locations, do not apply more than 2.5 lbs. of this product (2.25 lbs. a.i.) per acre for any application.
- Do not apply more than one application per cycle.
- Only one application is allowed per cycle for all chemical fallow applications.
- Do not use on sandy soil.
- Do not treat eroded hillsides, caliche and rocky outcroppings or exposed calcareous subsoil.

- Do not treat soils of the Rosebud and Canyon Series in Western NE and adjoining counties in CO and WY.
- Do not treat soils with calcareous surface layers.
- Do not graze treated areas within 6 months after application or illegal residues may result.

**All other locations restrictions:**

- Do not apply more than 2.5 lbs. of this product (2.25 lbs. a.i.) per acre for any application.
- Do not apply more than one application per year.

## SUGARCANE

For control of many broadleaf and grass weeds, such as amaranths, crabgrass, fireweed, flora's paintbrush, foxtail, junglerice and wiregrass, broadcast 2.2 to 4.4 lbs. of this product (2 to 4 lbs. a.i.) per acre at time of planting or ratooning, but before the sugarcane emerges.

Broadcast or band by ground equipment in a minimum of 20 gallons per acre, unless indicated otherwise. One additional application may be made over the cane as it emerges and two additional applications may be made interline after emergence, as a directed spray. Repeat treatments, where needed, may be applied broadcast, band or interline as suggested with the final application being prior to "close-in." Do not exceed the rate of herbicide listed for any one crop of sugarcane.

**Precaution:** Where high rates of this product are used, apply in a minimum of 1 gallon of water for each pound of this product applied per acre.

### USE RESTRICTIONS FOR ALL STATES AND USES

- Do not apply this product after "close-in".
- Do not apply more than 4.4 lbs. of this product (4.0 lbs. a.i.) per acre for any application. Do not apply more than 11.1 lbs. of this product (10 lbs. a.i.) per acre per crop.
- Injury to Sugarcane may occur when under moisture stress, when soil is of low absorptive capacity or when land is first cropped to Sugarcane.

For specific weed problems, the following may be used. Other rate and application timings may be used for other weed spectrums and cultural practices, provided they are consistent with the "*USE RESTRICTIONS FOR ALL STATES AND USES*".

### FLORIDA

For control of emerged pellitory weeds (artillery weed), apply 0.4 to 0.6 pound of this product (0.36 to 0.54 lb. a.i.) per acre in 40 gallons of water per acre as a directed spray by ground equipment prior to "close-in". Add 2 quarts of surfactant for each 50 gallons of spray and be sure Weed foliage is thoroughly covered.

For control of alexandergrass, large crabgrass, pellitory weed (artillery weed) and spiny amaranth, use one of the following methods at-planting or ratooning:

1. Apply 4.4 lbs. of this product (4 lbs. a.i.) per acre pre-emergence. Follow with one or two applications, as needed, post-emergence to sugarcane and weeds at 2.2 lbs. of product (2 lbs. a.i.) per acre. Treat before weeds exceed 1.5 inches in height.
2. Apply 1 to 3 times, as needed, at 2.2 lbs. of this product (2 lbs. a.i.) per acre post-emergence to sugarcane and weeds. Treat before weeds exceed 1.5 inches in height.

The total lbs. of active ingredient applied by pre-emergent PLUS post-emergent applications cannot exceed 11.1 lbs. of this product (10 lbs. a.i.) per acre per crop.

### LOUISIANA

For control of annual weeds during the Summer fallow period, apply 2.2 lbs. of this product (2 lbs. a.i.) per acre to weed-free beds immediately after bed formation. Follow normal weed control after planting.

#### **Additional Use Restrictions — To Avoid Crop Injury in LA:**

1. Do not apply more than 11.1 lbs. of this product (10 lbs. a.i.) per acre to any one crop of sugarcane.
2. If making 2.2 lbs. of this product (2 lbs. a.i.) per acre application during Summer fallow period, do not exceed 8.8 lbs. of this product (8 lbs. a.i.) per acre during the remainder of the growing season.

### TEXAS

Use this product for control of barnyardgrass, pigweed, purslane and sunflower in plant or ratoon sugarcane.

Apply 4.4 lbs. of this product (4 lbs. a.i.) per acre pre-emergence. Follow with one or two applications, as needed, at 3.3 lbs. of this product (3 lbs. a.i.) per acre post-emergence to sugarcane and weeds.

For best results, when weeds are emerged, add a non-ionic surfactant, such as Surf-Ac 820, at a concentration of 2 quarts per 100 gallons to the spray and apply before weeds exceed 1.5 inches in height.

## ST. AUGUSTINE GRASS, CENTIPEDEGRASS AND DORMANT BERMUDAGRASS

This product controls annual bluegrass, Florida betony, spurweed and many other weeds in home lawn, ornamental and recreational turf. This product may be applied with any pump-up or compressed air type sprayer or with a hose-on type sprayer. This product will control both emerged weeds and weeds from seeds. Rain or watering within 2 to 3 days of application may decrease the effectiveness on emerged weeds. However, for control of weeds from seeds, rainfall or watering is necessary within 7 to 10 days after treatment.

### WEEDS CONTROLLED OR SUPPRESSED

Annual bluegrass (*Poa annua*), Chickweed (Common and Mouseear), Crabgrass, Cransbill, Cudweed, Dichondra, Florida betony, Henbit, Knotweed, Lespedeza, Moneywort, Mustard, Narrowleaf vetch, Parsley-piert, Sandspur, Smutgrass, Spurge, Spurweed, Swinecrest, Woodsorrel and various annual Clovers.

### TIMING OF APPLICATION

The timing of applications to achieve maximum control may vary quite a bit with different weed species. The following application times are suggested for certain problem weeds.

**Spurweed:** Best control of spurweed can be obtained by applying this product when spurweed has emerged (December and January).

**Florida betony:** Apply this product when this weed emerges in the Fall.

**Dichondra, Moneywort:** Applications for Florida betony or spurweed generally will give control or suppression of the other weeds listed. This product will give the best control when applied to young, tender weeds or just prior to weed emergence. Do not apply to growing Bermudagrass.

### USE RESTRICTIONS

- Do not apply within the active root zone of vegetables or desirable ornamental plants such as azaleas, boxwoods, camellias, etc. However, treatments with this product should not normally cause injury to medium or large shrubs or trees in the landscape.
- Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result.
- Do not cut for hay or graze treated areas within 3 months following an application.
- Do not make more than one application per year.
- Do not seed or transplant species which are sensitive to this product within 18 months after application.
- Use of backpack applications to landscape turf is allowed for spot treatments only.

### ST. AUGUSTINE AND CENTIPEDEGRASSES

This product may be applied to established centipedegrasses and St. Augustine grasses during both the dormant and the growing season. Best results are usually obtained in the early Spring or dormant periods when weeds are small or have not emerged. Follow rates listed.

### DORMANT BERMUDAGRASS

This product may be applied to Bermudagrass during the dormant season only. Applications to Bermudagrass during the growing season will cause injury. Follow rates below, except for residential uses, for which the maximum rate is 1 lb. active ingredient per acre (0.6 tsp. of this product per 1,000 sq. ft.).

### RATE OF APPLICATION

Determine the total area to be sprayed and base rate of application on the chart below. Avoid overlapping spray pattern while treating.

Area To Be Treated	Amount of this product
500 sq. ft.	1.33 tbsps.
1,000 sq. ft.	2.75 tbsps. (1.33 ozs.)
3,000 sq. ft.	5 ozs.
5,450 sq. ft.	0.27 lb.
10,900 sq. ft. (0.25 acre)	0.55 lb.
1 acre	2.2 lbs.

## TURFGRASS FOR FAIRWAYS, LAWNS, SOD PRODUCTION\* AND RESIDENTIAL SITES (\*Except FL)

\*In states other than FL. For use on Turfgrass for sod in FL, see "TURFGRASSES FOR SOD (FL ONLY)" section below.

**Bermudagrass, Centipedegrass, St. Augustine grass and Zoysiagrass:** Apply this product after October 1 before emergence of Winter annual weeds for control of annual bluegrass, burclover, carpet burweed, chickweed, corn speedwell, henbit, hop clover and spurweed. This product will control annual bluegrass even if it is emerged at time of treatment. For control of Summer annual weeds listed in the pre-emergence portion of the "*THIS PRODUCT APPLIED ALONE—CORN OR GRAIN SORGHUM—Pre-plant Surface-Applied, or Pre-plant Incorporated, or Pre-emergence*" section of this label, also apply this product in late Winter before the weeds emerge. Apply in a minimum of 15 gallons of water per acre or 1 gallon per 1,000 square feet (0.4 fl. oz. per 1,000 sq. ft.).

Where annual bluegrass is the major weed, use 1.1 lbs. of this product (1 lb. a.i.) per acre. Use 2.2 lbs. of this product (2 lbs. a.i.) per acre for control of the other weeds named above. Do not exceed 1.1 lbs. of this product (1 lb. a.i.) per acre per treatment on newly sprigged Turfgrass or on hybrid Bermudagrass such as Tiflawn, Tifway and Ormond.

For continued Summer annual weed control, apply another 1.1 lbs. of this product (1 lb. a.i.) per acre at least 30 days after the previous application, but not after April 15. However, do not make more than 2 applications of this product per year. On newly sprigged Turfgrass and hybrid Bermudagrass, temporary slowing of growth and yellowing may occur following application.

### USE RESTRICTIONS

- For Turfgrass at residential sites including homes, day care facilities, schools, playgrounds, parks, recreational areas and sports fields, do not apply more than 1.1 lbs. of this product (1.0 lb. a.i.) per acre for any application. Do not apply more than 2.2 lbs. of this product (2.0 lbs. a.i.) per acre per year.
- Do not graze or feed turf clippings to animals or illegal residues may result.
- Use of backpack applications to landscape turf is allowed for spot treatments only.

To avoid turf injury: 1) Use only on Turfgrass reasonably free of infestations of insects, nematodes and diseases. 2) Do not use on golf greens. 3) Do not use North of NC (except may be used in VA Coastal Plains) or West of the high rainfall areas of Eastern OK and Eastern TX. 4) Do not use on muck or alkaline soils. 5) Do not apply over the rooting area of trees or ornamentals not listed on this label. 6) Do not overseed with desirable Turfgrass within 4 months before or 6 months after treatment. 7) Do not apply this product to newly seeded Bermudagrass until it has over-Wintered and has a well-developed rhizome system. Do not exceed 2.2 lbs. of this product (2 lbs. a.i.) per acre within 12 months of seeding Bermudagrass.

### TURFGRASSES FOR SOD (FL ONLY)

St. Augustine grass, Centipedegrass and Zoysiagrass: Broadcast 2.2 to 4.4 lbs. of this product (2 to 4 lbs. a.i.) per acre according to soil texture to control those weeds listed under "*THIS PRODUCT APPLIED ALONE—CORN OR GRAIN SORGHUM—Preplant Surface-Applied, or Pre-plant Incorporated, or Pre-emergence*".

Muck or Peat	4.4 lbs. of this product	Old Beds: Within 2 days after lifting sod.
		New Beds: 3 to 4 days after sprigging or plugging.
Sandy Soil	2.2 lbs. of this product	Old Beds: Within 2 days after lifting sod.
		New Beds: 7 to 10 days after sprigging or plugging.

If weeds regrow, apply an additional 2.2 lbs. of this product (2 lbs. a.i.) per acre on Muck or Peat, or 1.1 lbs. of this product (1 lb. a.i.) per acre on Sandy soil.

### Use Precautions — To avoid illegal residues and crop injury:

- Use only on Turfgrass reasonably free of infestations of insects, nematodes and diseases.
- On newly-sprigged Turfgrass, temporary slowing of growth may follow application.

**Use Restrictions — To avoid illegal residues and crop injury:**

- For Muck or Peat soils, do not apply more than 4.4 lbs. of this product (4 lbs. a.i.) per acre for any application. Do not apply more than 6.6 lbs. of this product (6 lbs. a.i.) per year. For sandy soil, do not apply more than 2.2 lbs. of this product (2 lbs. a.i.) per acre for any application. Do not apply more than 3.3 lbs. of this product (3 lbs. a.i.) per year.
- Do not apply within 30 days prior to cutting or lifting.
- Do not apply in combination with surfactants or other spray additives.

## **MACADAMIA NUTS**

For pre-emergence control of many broadleaf and grass weeds, including crabgrass, fireweed, Flora's paintbrush, foxtails, Spanishneedles and wiregrass, broadcast 2.2 to 4.4 lbs. of this product (2 to 4 lbs. a.i.) per acre before harvest and before weeds emerge. Repeat as necessary but do not exceed 8.8 lbs. of this product (8 lbs. a.i.) per acre per year.

### **USE RESTRICTIONS**

- Do not apply more than 4.4 lbs. of this product (4 lbs. a.i.) per acre for any application.
- Do not apply more than 8.8 lbs. of this product (8 lbs. a.i.) per acre per year.
- Do not spray when nuts are on ground during harvest period.
- Do not apply by air.
- Do not apply via mechanically pressurized handgun.

## **GUAVA**

Use only on established plantings which are at least 18 months old. Apply as a directed spray at 2.2 to 4.4 lbs. of this product (2 to 4 lbs. a.i.) per acre of this product in 20 to 50 gallons of spray mix pre-emergence or early post-emergence to weeds. When applying post-emergence, the use of a surfactant and greater spray volume (80 to 100 gallons of spray mix per acre) may enhance weed control.

This product controls many annual broadleaf and grass weeds, including fireweed, purslane, scarlet pimpernel, sowthistle and Spanishneedles.

### **USE RESTRICTIONS**

- Do not allow spray to contact foliage or fruit.
- Do not apply more frequently than at 4-month intervals.
- Do not apply more than 4.4 lbs. of this product (4 lbs. a.i.) per acre for any application.
- Do not apply more than 8.8 lbs. of this product (8 lbs. a.i.) per year.
- Do not apply via mechanically pressurized handgun.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

**PESTICIDE STORAGE:** Storage should be in a cool, dry area away from any heat or ignition source. Keep container tightly sealed when not in use. Do not allow bags to become wet or store in a damp, humid area. Keep away from any puncture source. Store in original container only.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING:

**[Nonrefillable Container (flexible-bag-all weights):]** [Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration.]

**[Nonrefillable Container (rigid-fifty lbs. or less):]** [Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration.]

**[Nonrefillable Container (rigid-greater than fifty lbs.):]** [Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration.]

**[Refillable Container:]** [Refillable container. Refill this container with this product only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.]

## WARRANTY—CONDITIONS OF SALE

OUR DIRECTIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically directed and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. To the extent consistent with applicable law, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

[EPA APPROVAL DATE]



**{LANGUAGE ON LABEL AFFIXED TO CONTAINER}**

**RESTRICTED USE PESTICIDE**

**(Ground and Surface Water Concerns)**

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification. This product is a restricted use herbicide due to ground and surface water concerns. Users must read and follow all precautionary statements and instructions for use in order to minimize potential for atrazine to reach ground and surface water.

ATRAZINE	GROUP	5	HERBICIDE
----------	-------	---	-----------

**Atrazine 90DF**

**ACTIVE INGREDIENTS:**

Atrazine.....	88.4%
Related compounds.....	1.6%
<b>OTHER INGREDIENTS:</b> .....	<b>10.0%</b>
<b>TOTAL:</b> .....	<b>100.0%</b>

**KEEP OUT OF REACH OF CHILDREN CAUTION**

**FIRST AID**

**IF SWALLOWED:**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

**IF IN EYES:**

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**HOTLINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12 PM Pacific Standard Time. In the event of a medical emergency, call your poison control center at 1-800-222-1222.

**NOTE TO PHYSICIAN:** There is no specific antidote for atrazine. If this product is ingested, induce emesis or lavage stomach. The use of an aqueous slurry of activated charcoal may be considered.

**PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear protective eyewear.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage and disposal.

**PESTICIDE STORAGE:** Storage should be in a cool, dry area away from any heat or ignition source. Keep container tightly sealed when not in use. Do not allow bags to become wet or store in a damp, humid area. Keep away from any puncture source. Store in original container only.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:**

**[Nonrefillable Container (flexible-bag-all weights):]** [Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration.]

**[Nonrefillable Container (rigid-fifty lbs. or less):]** [Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration.]

**[Nonrefillable Container (rigid-greater than fifty lbs.):]** [Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration.]

**[Refillable Container:]** [Refillable container. Refill this container with this product only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.]

[See] [inside] [label] [booklet] [for] [additional] [Precautionary Statements][,] [and] [Directions for Use] [including] [Storage and Disposal] [instructions][,]

**EPA Reg. No.: 94144-1**

**EPA Est. No.:**

**Manufactured for:**

INFINICROP, LLC  
266 Reservation Road, F#202  
Marina, CA 93933

**[Lot][Batch No.]** \_\_\_\_\_

**Net Weight:**