



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

March 4, 2024

Annette Marine
Agent for Aquatrols Corporation of America
c/o Wagner Regulatory Associates, Inc.
P.O. Box 640
7217 Lancaster Pike, Suite A
Hockessin, DE 19707

Subject: Approval of Label Amendment; Only Indicated Changes Reviewed – *Update New York state turfgrass restriction and add an ABN*
Product Name: Aquatrols Fluazinam UniTech
EPA Registration Number: 94396-32
Application Date: 11/04/2022; 01/27/2023; 02/27/2023
Case Number: 476567; 471911; 474473

Dear Annette Marine:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. However, EPA reviewed only the label changes highlighted, marked, or otherwise indicated on the submitted label. Any other changes to the previously approved label that were not clearly highlighted, marked, or otherwise indicated in your submission were not reviewed and may form the basis of regulatory and/or enforcement action if later discovered by the Agency. Further, submission of a label amendment application with unidentified changes may be considered a knowing submission of false information to the Agency. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

The label submitted with the application has been stamped "Accepted Only Indicated Revisions Reviewed" and is enclosed for your records.

The alternate brand name, "**Wingspan® Unitech® Fungicide**" has been added to the product record.

This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 C.F.R. § 152.130(c), you may distribute or sell this product under the

previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 C.F.R. § 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website contains any false or misleading statement, design, or graphic, the product may be misbranded and unlawful to sell or distribute under FIFRA Sections 2(q)(1)(A) and 12(a)(1)(E). 40 C.F.R. § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on the product label, claims made as part of the product's sale or distribution may not substantially differ from those claims approved through the registration process under FIFRA Section 12(a)(1)(B). Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the product will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Marc Sheahin at 202-566-2896 or at sheahin.marc@epa.gov.

A handwritten signature in black ink, appearing to read "Lydia Crawford". The signature is written in a cursive, flowing style.

Lydia Crawford, Acting Product Manager 25
Fungicide Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

FLUAZINAM	GROUP	29	FUNGICIDE
-----------	-------	----	-----------

Aquatrols Fluazinam UniTech

ABN: Wingspan® Unitech® Fungicide

A C C E P T E D

ONLY INDICATED

REVISIONS REVIEWED

03/04/2024

Controls certain diseases of turf grasses on golf course

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

94396-32

ACTIVE INGREDIENTS:

Fluazinam* No label revisions other than those indicated were reported to the Agency.

OTHER INGREDIENTS 60.0%

TOTAL: 100.00%

*3-chloro-N-[3-chloro-2,6-dinitro-4-(trifluoromethyl)phenyl]-5-trifluoromethyl-2-pyridinamine (CA)
Equivalent to 4.17 pounds Fluazinam per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

WHERE TO USE: For use on turfgrass on golf courses only. Aquatrols Fluazinam Unitech can be utilized on cool-season and warm-season turfgrass varieties. See label booklet for more detailed information.

WHEN TO USE: Aquatrols Fluazinam Unitech may be utilized at any point throughout the growing season but is most effective as a preventative application when conditions become favorable for turfgrass diseases.

See [side] [back] [panel] [label booklet] for [First Aid,] [additional] [Precautionary Statements,] [and] [complete] [Directions For Use,] including [Storage and Disposal.]

Net Contents: _____ Gallons
EPA Reg. No. 94396-32

EPA Est. No. _____

Aquatrols Corporation of America
1273 Imperial Way
Paulsboro, NJ 08066
1-800-257-7797
Aquatrols.com

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF ON SKIN	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
<p>For chemical emergency spill, leak, fire, exposure or accident, call CHEMTEL day or night. Domestic North America 800-255-3924. International call 813-248-0585 (collect calls accepted). You may also call the poison control center at 1-800-222-1222.</p>	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before use. **DO NOT** take internally.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Coveralls worn over long-sleeved shirt, long pants
- Socks and chemical resistant footwear
- Chemical resistant gloves made of any waterproof material including polyethylene or polyvinyl chloride
- Protective eyewear

When mixing and loading, or when cleaning equipment, also wear a chemical resistant apron.

USER SAFETY REQUIREMENTS

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, or enclosed cabs, in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for “applicators and other handlers” and have such PPE immediately available for use in an emergency, including a spill or equipment break-down. **DO NOT** allow contact between contaminated sprayer parts and unprotected skin. Ensure sprayer is washed down daily.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when cleaning equipment or disposing of equipment wash waters or rinsate. **DO NOT** apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas.

PHYSICAL-CHEMICAL HAZARDS

Do not mix or allow coming in contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE USE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN TURF INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DO NOT allow people (other than the applicator) or pets on the treatment area during application. **DO NOT** enter or allow others to enter the treated area until sprays have dried.

PRODUCT INFORMATION

Aquatrols Fluazinam UniTech is a unique multi-site contact fungicide which controls certain diseases of turf grasses on golf courses.

Aquatrols Fluazinam Unitech is effective for strategic use in fungicide disease management programs that attempt to minimize disease resistance to fungicides. Aquatrols Fluazinam Unitech has a multi-site mode of action that disrupts the energy production in the fungus. It is listed in FRAC code 29 as an uncoupler of oxidative phosphorylation. Some other fungicides, which are at risk from disease resistance, exhibit a single-site mode of fungicidal action.

Aquatrols Fluazinam Unitech, with its multi-site mode of action, may be used to prevent or delay the development of resistance to single-site fungicides when used as a tank-mix partner or alternated in a disease management program.

Aquatrols Fluazinam Unitech has demonstrated excellent control of *Sclerotinia homeocarpa*, *Lanzia* and *Moellerodiscus* strains (causing Dollar Spot) that are insensitive to several classes of fungicides.

Benefits of Using Aquatrols Fluazinam Unitech on Turfgrass on Golf Courses

- Excellent control of dollar spot and 10 other turf diseases
- Controls dollar spot insensitive to other chemistries
- Fantastic tank mix partner with triadimefon or other systemic Aquatrols fungicides to offer complete control

TURFGRASS USE RESTRICTIONS:

- **DO NOT** apply Aquatrols Fluazinam Unitech to home lawns (turf) or any other residential turf.
- **DO NOT** apply Aquatrols Fluazinam Unitech by aerial application.
- **DO NOT** apply Aquatrols Fluazinam Unitech through any irrigation system.
- **DO NOT** apply Aquatrols Fluazinam Unitech within 25 feet of permanent water bodies (lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, and estuaries).
- ~~In the State of New York,~~ **DO NOT** apply Aquatrols Fluazinam Unitech within 100 feet of any water bodies that ~~are connected to other~~ with any off-off-site surface waters bodies such as a freshwater lake, pond, river, stream or wetland in the State of New York.
- **DO NOT** apply Aquatrols Fluazinam Unitech when the wind direction is toward aquatic areas to minimize drift to these areas.

Integrated Pest (Disease) Management (IPM)

Aquatrols Fluazinam Unitech must be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Sound pest management resulting in healthy, vigorous turf and ornamentals is the foundation of a good IPM program. Cultural practices including proper choice of varieties, nutrient management, proper cutting height, thatch management, and proper watering, drainage, and moisture stress management must be integrated with the use of fungicides to increase vigor and reduce the susceptibility to disease. Immunoassay detection kits and extension service diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

Resistance Management

For resistance management, Aquatrols Fluazinam Unitech contains a Group 29 fungicide. Any fungal population may contain individuals naturally resistant to Aquatrols Fluazinam Unitech and other Group 29 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Aquatrols Fluazinam UniTech or other Group 29 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Aquatrols Corporation of America at 1-800-257-7797 or at www.aquatrols.com. You can also contact your pesticide distributor or university extension specialist to report resistance.

PLANT RESISTANCE

Note to User: Although Aquatrols Fluazinam Unitech has been evaluated on several turf species with no indication of phytotoxicity, neither the manufacturer nor seller has determined whether or not Aquatrols Fluazinam Unitech can be used safely on all turf species. The professional user must determine if Aquatrols Fluazinam Unitech can be used safely prior to commercial use by testing on the type of turf to be treated at specified rates for phytotoxicity.

TURFGRASS APPLICATION INFORMATION

Aquatrols Fluazinam Unitech can be used effectively in dilute sprays. Thorough, uniform coverage is essential for disease control.

NOTE: Slowly invert container several times to ensure uniform mixture.

Prepare only the amount of spray mixture required for immediate use. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. **DO NOT** allow spray mixture to stand overnight or for prolonged periods in the spray tank. Spraying equipment must be thoroughly cleaned immediately after the application.

Aquatrols Fluazinam Unitech Alone: Fill the spray tank with half of the required amount of water. With the agitator running, add Aquatrols Fluazinam Unitech to the spray tank. Continue agitation while adding the remainder of the water. Begin application of the solution after Aquatrols Fluazinam Unitech has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Tank Mixtures: Aquatrols Fluazinam Unitech is physically compatible with many products used for control of diseases, liquid fertilizers, growth regulators, insecticides, and biological control products. If tank mixes are desired, read and follow all directions, precautions, and limitations on labeling of all products used. This product must not be mixed with any product which prohibits such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

Tank mix combinations containing Aquatrols Fluazinam Unitech have not been tested on all varieties of every species or under all possible growing conditions. If a user is unfamiliar with the performance of Aquatrols Fluazinam Unitech in tank mixes under user's growing conditions, a limited area must be tested prior to large-scale application. The user must always exercise reasonable judgment and caution when using this and all other products.

Aquatrols Fluazinam Unitech is compatible with the products mentioned above, provided sufficient free water is available for dispersion of all the tank mix products. However, the physical compatibility of Aquatrols Fluazinam Unitech with tank mix partners must be evaluated before use. A jar test must be conducted with intended tank mix products prior to preparation of large volumes.

Use the following procedure: 1) Pour the specified proportions of the products into a suitable container of water, 2) Mix thoroughly and 3) Allow to stand for 5 minutes.

If the combination remains mixed or can be re-mixed readily, it is considered physically compatible. Any physical incompatibility in the jar test indicates that Aquatrols Fluazinam Unitech must not be used in the tank mix.

Consult compatibility charts or your local or state agricultural authorities for additional compatibility information.

To prepare spray solution, add ½ of the required amount of water to the mix/sprayer tank. Start the agitator running before adding any tank mix partners. In general, add tank mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry fowables), liquid fowables (including Aquatrols Fluazinam Unitech), liquids, and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

When using Aquatrols Fluazinam Unitech in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including Aquatrols Fluazinam Unitech. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

Use a pump with capacity to maintain the correct rated pressure for the nozzles selected. Maintain sufficient agitation to keep mixture in suspension. Use a jet agitator, liquid sparge tube, or mechanical paddle for agitation.

Use screens to prevent nozzles from clogging. Use 50-mesh or coarser screens placed after the tank and before the nozzles. Check nozzle manufacturer's directions. For more information on spray equipment and calibration, consult sprayer manufacturer's and state recommendations.

Apply Aquatrols Fluazinam Unitech in sufficient water to obtain adequate uniform coverage of the turf. Apply in a minimum spray volume of 40 gal of water per acre on fairways and 90 gal of water per acre on tees and greens.

Turfgrass Disease Control

1. Use Aquatrols Fluazinam Unitech in a Preventative Disease Control Program.
2. Apply in sufficient water to ensure thorough coverage.
3. Apply after mowing or allow sprayed area to completely dry before mowing.
4. For control of foliar diseases, allow sprayed area to completely dry before irrigation.
5. For optimum turf quality and disease control, use Aquatrols Fluazinam Unitech in conjunction with turf management practices that promote good plant health and optimum disease control.
6. Evaluate spray additives prior to use. Label directions are based on data obtained with no additives.
7. Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

Restrictions:

- **DO NOT** exceed 258 f oz of Aquatrols Fluazinam Unitech (8.4 pounds active ingredient) per acre per year.
- **DO NOT** apply more than twelve (12) applications per year.
- **DO NOT** apply more than 21.5 f oz (0.7 pounds active ingredient) per acre (0.5 f oz/1000 ft²) per application.
- **Retreatment Interval:** 14 days

Turfgrass Application Rates

Aquatrols Fluazinam Unitech Use Rates			
Diseases Controlled Pathogen(s)	Application Interval (days)	Rate fl. oz. product/ 1000 ft ²	Instructions for Use on Golf Course Tees, Greens, Fairways and Roughs
Dollar Spot <i>Sclerotinia homeocarpa</i> <i>Lanzia</i> spp. <i>Moellerodiscus</i> spp.	14	0.5 (0.7 lb. AI)	Apply as a preventative disease program, starting when conditions become favorable for disease. For broad spectrum disease control or for control greater than 14 days, tank mix with [another fungicide labeled for control of this disease] [including] [one of the following fungicides:] [Aquatrols Triadimefon + Azoxystrobin; Aquatrols Triadimefon; another Aquatrols fungicide]
Leaf spot, Melting-out, Brown blight <i>Drechslera</i> spp. (including <i>D. poae</i> , <i>D. siccans</i>)	14	0.5 (0.7 lb. AI)	Apply when conditions become favorable for disease. For broad spectrum disease control tank mix with [another fungicide labeled for control of this disease] [including] [one of the following fungicides:] [Aquatrols Triadimefon + Azoxystrobin; Aquatrols Triadimefon; another Aquatrols fungicide]
Brown patch Large patch <i>Rhizoctonia solani</i>	14	0.5 (0.7 lb. AI)	Begin applications in May or June before disease is present for Brown patch. To control Large patch, apply in fall when temperature drops below 70°F. For broad spectrum control or for control greater than 14 days, tank mix with [another fungicide labeled for control of this disease] [including] [one of the following fungicides:] [Aquatrols Triadimefon + Azoxystrobin; Aquatrols Triadimefon; another Aquatrols fungicide]
Red Thread <i>Laetisaria fuciformis</i>	14	0.5 (0.7 lb. AI)	Apply when conditions become favorable for disease.
Anthracnose <i>Colletotrichum cereale</i>	14	0.5 (0.7 lb. AI)	Applications must be made prior to infection. Alternate or mix Aquatrols Fluazinam Unitech with another fungicide for anthracnose disease prevention. For optimum disease control, tank mix with [another fungicide labeled for control of this disease] [including] [one of the following fungicides:] [Aquatrols Triadimefon + Azoxystrobin; Aquatrols Triadimefon; another Aquatrols fungicide]
Gray Snow Mold <i>Typhula</i> spp. Pink Snow Mold <i>Microdochium nivale</i>	Late Fall	0.5 (0.7 lb. AI)	Apply one to two applications in the late fall before snow cover. Restriction: DO NOT apply on top of the snow. For optimum disease control tank mix with [another fungicide labeled for control of this disease] [including] [one of the following fungicides:] [Aquatrols Triadimefon + Azoxystrobin; Aquatrols Triadimefon; another Aquatrols fungicide]
Pink Patch <i>Microdochium nivale</i>	14	0.5 (0.7 lb. AI)	Apply in the fall or early spring when conditions are favorable for disease development. For optimum disease control, tank mix with [another fungicide labeled for control of this disease] [including] [one of the following fungicides:] [Aquatrols Triadimefon + Azoxystrobin; Aquatrols Triadimefon; another Aquatrols fungicide]
Stem Rust <i>Puccinia graminis</i> var. <i>graminicola</i>	14	0.5 (0.7 lb. AI)	Apply on a preventative schedule when disease conditions are favorable (cool and cloudy weather).
Algal Scum (filamentous blue-green algae/cyanobacteria)	14	0.5 (0.7 lb. AI)	This product must be tank mixed with another fungicide labeled for control of this disease, including mancozeb.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container, in a secured, cool, dry place separate from food and feed. **DO NOT** store near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: [Nonrefillable Containers 5 Gallons or Less] Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip.

Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration.

[Nonrefillable containers larger than 5 gallons] Non-refillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT INFORMATION READ BEFORE USING PRODUCT

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If the terms are not acceptable, **DO NOT** use the product and instead, return the unopened product container immediately. By using this product, you accept the following Conditions, Disclaimer of Warranties and Limitation of Liability.

For technical information, contact Aquatrols Corporation of America at 800-257-7797. Information regarding the contents and levels of metals in this product is available on the internet at www.aapfco.org/metals.html

CONDITIONS: The directions for use on this label are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Insufficient performance or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions, the failure to follow the label directions or good application practices, all of which are beyond the control of Aquatrols Corporation of America. In addition, failure to follow label directions may cause poor performance, injury to crop, animals, humans, or the environment. You assumed all such risks by using this product.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, AQUATROLS CORPORATION OF AMERICA MAKES NO OTHER WARRANTIES OR REPRESENTATIONS OF ANY KIND, EXPRESSED OR IMPLIED, CONCERNING THE PRODUCT, INCLUDING NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR ANY PARTICULAR PURPOSE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL AND NO SUCH WARRANTY SHALL BE IMPLIED BY LAW. No agent of Aquatrols Corporation of America is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, Aquatrols Corporation of America disclaims any and all claims are waived of any liability whatsoever for special, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profit or income, resulting from the use or handling of this product.

LIMITATION OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, the remedy for any losses or cause of action relating to injuries, damages or the handling or use of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or, at Aquatrols Corporation of America's election, the replacement of the product. Aquatrols shall not be liable and any and all claims against Aquatrols are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profit or income.

Optional Marketing Claims

- *Unique multi-site contact active ingredient*
- *Excellent control of dollar spot and 10 additional turf diseases*
- *Controls dollar spot that is insensitive to other chemistries*
- *Unique chemical class aids resistance management*
- *Perfect rotation partner with Daconil Action (EPA Reg. No. 100-1364 containing Chlorothalonil + Acibenzolar-S-methyl) for season-long contact protection*
- *Perfect partner with systemics for inside out protection*
- *Formulation holds fungicide to the surface of grass blades even after irrigation and heavy rains*
- *One convenient rate for 12 applications per year on all turfgrass at any height*
- *Provides superior protection against dollar spot*