



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Biopesticides and Pollution Prevention Division (7511P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

94610-1

Date of Issuance:

2/7/2020

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Concentric SS

Name and Address of Registrant (include ZIP Code):

Concentric Ag Corporation
 7304 South Joliet Street, Suite 200
 Centennial, CO 80112

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.
2. Submit storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) data as these data requirements are not satisfied. A one-year study is required to satisfy these data requirements. You have 18 months from the date of this registration to provide these data to the EPA.

Signature of Approving Official:

Andrew Bryceland, Team Leader
 Biochemical Pesticides Branch
 Biopesticides and Pollution Prevention Division (7511P)
 Office of Pesticide Programs

Date:

2/7/2020

3. Make the following labeling change before you release this product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 94610-1."
4. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated 10/11/2019

Any CSFs other than those listed above are superseded.

If you have any questions, please contact James Parker by phone at (703) 306-0469 or via email at parker.james@epa.gov.

Sincerely,




Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure

{Note to reviewer: [Text] in brackets denotes optional text.}

{Note to reviewer: {Text} in braces denotes where in the final label text will appear.}



Concentric SS Plant Growth Regulator

For field or greenhouse use on vegetables, fruits, nuts, vine crops, field crops, ornamentals, turf and seed treatment to stimulate root and shoot growth when used alone or in mixtures with other pesticide products.

ACTIVE INGREDIENTS:

Complex polymeric polyhydroxy acids.....	00.108%
Cytokinin (as kinetin)	00.004%
OTHER INGREDIENTS	<u>99.888%</u>
TOTAL	100.000%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
<p>HOTLINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For 24-hour medical emergency assistance (human or animal) call 1-877- 424-7452.</p>	

SEE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS, COMPLETE DIRECTIONS FOR USE, WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY

EPA Reg. No. 94610-[xx]

EPA Est. No. _____

Manufactured By:
Concentric Ag Corporation
7304 South Joliet Street, Suite 200
Centennial, CO 80112

Net Contents: _____

[EPA Approval Date]

Precautionary Statements

Hazards to Humans and Domestic Animals

Caution: Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse. Might cause slight eye irritation. Do not get in eyes or on clothing. Wear goggles or face shield when handling concentrate. After product is diluted in accordance with the directions for use, goggles or face shield are not required. Avoid contact with skin, eyes or clothing. Wear appropriate personal protective equipment (PPE).

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes and socks.
- Waterproof gloves
- Protective eyewear. (Wear goggles or face shield when handling concentrate. After product is diluted in accordance with the directions for use, goggles or face shield are not required.)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

Users should:

- Remove clothing immediately if pesticide gets inside.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

Environmental Hazards

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation

GENERAL USE INFORMATION

Concentric SS is a broad-spectrum plant growth regulator for use on field or greenhouse crops, fruits, nuts, vines and ornamentals. Concentric SS may be applied in liquid sprays, with irrigation, impregnated on fertilizer granules, or used as a seed treatment. Optimum growth effects are achieved when the product is applied on a regular scheduled application program.

Since all combinations or sequences of foliar pesticide and nutrient applications including surfactants and adjuvants have not been tested, before wide spread application, test a small area to be sprayed first to make certain that no phytotoxicity occurs.

Read the entire label before using Concentric SS. Consult your State Agricultural Experimental Station or Extension Service Specialist for additional information on application timing, rates and any additional requirements or restrictions.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes and socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated area until sprays have dried.

FOLIAR AND SOIL APPLICATION INSTRUCTIONS

Concentric SS is a broad spectrum biopesticide to improve germination and seedling development, stimulate root and shoot growth, increase chlorophyll content, improve plant ability to withstand stress, and increase yields when used alone or in mixtures with nutrients and other pesticide products on field crops, vegetables, fruits, nuts, vine crops, turf and ornamentals.

For all crops, apply Concentric SS at a rate of 5-20 fluid ounces per acre unless otherwise specified on this label. Refer to the Foliar and Soil Application Use Rates for Outdoor and Greenhouse Crops section of this label for specific soil and foliar application use rate and timing information.

Applications can begin prior to planting for soil applications and continue as long as plants are still in the vegetative stage. Foliar applications may begin when plants reach or surpass the 4- leaf stage of development. For best results, apply Concentric SS before leaf hardening.

Repeat applications no sooner than every 7 days. Concentric SS is effective for use in programs where repeated applications throughout the growing season are necessary. If not applied on a routine spray schedule, apply Concentric SS alone or in combination with appropriate fertilizers and pesticides.

Specific Use Restrictions and Precautions: Do not make foliar applications when conditions favor drift from target area or wind speed is greater than 10 mph. Spray equipment must be cleaned thoroughly before and after applications.

MIXING INSTRUCTIONS:

For foliar applications, be sure the sprayer is clean and not contaminated with other materials prior to use. When using an agitated spray tank fill tank 1/2 to 3/4 full with clean water and start agitation. Be certain that the agitation system is working properly. With the agitator running add the required amount of Concentric SS to the tank. If tank mixing with other materials, add them to the tank and continue agitation. Continue filling tank with the remainder of the water. Agitate until mixed thoroughly and avoid excessive

foaming. Mix as needed. Do not store diluted material overnight.

For granular applications, Concentric SS is applied to provide uniform coverage of all granules. In cases where the amount of fertilizer per acre is less than 100 pounds, Concentric SS should be applied to a small amount of fertilizer to ensure that it will not cause the fertilizer to cake.

COMPATIBILITY

Concentric SS is compatible with most commonly used agricultural pesticides. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Dilute Concentric SS to its use rate and then with stirring add the other components in the appropriate amounts. If precipitation, gelation, or sedimentation occurs, do not use the combination' of pesticides. Because of the wide variety of possible combinations that can be encountered, observe all precautions and limitations on the label of all products used in mixtures.

SPRINKLER IRRIGATION SYSTEM APPLICATION

Apply this product only through drip, microjet, lateral move, end tow, side (wheel) roll, hand move, solid set and center pivot irrigation systems. Do not apply this product through any other type of irrigation system. Use standard tank agitation when mixing Concentric SS alone or with other pesticides. Fill the tank halfway with water, begin agitation and add Concentric SS, other tank mix pesticides or fertilizers and fill with water.

Preparation of Injection Equipment: Remove pesticide, scale residues and other foreign matter from the chemical tank and entire injection system. Flush with clean water.

Set the sprinkler system to deliver 0.1 to 0.3 inch of water per acre. Start the sprinkler system and uniformly inject the solution of Concentric SS into the irrigation water line. Inject the Concentric SS solution with a positive displacement pump into the main line before a right angle turn to ensure adequate mixing. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Crop injury or lack of effectiveness in the crop can result from non-uniform distribution of treated water. A person with knowledge of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut down and make necessary adjustments should the need arise. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

When applying Concentric SS using microjet and drip irrigation systems avoid further irrigation after the treatment has been completed for 24 to 48 hours.

When applying Concentric SS using solid set, hand move and center pivot irrigation systems avoid further irrigation of the treated area until the foliage is dry to prevent washing the product from the crop.

When applying Concentric SS using a continuously moving system, such as lateral move, or side (wheel) roll system, inject this product-water mixture continuously, applying the labeled rate per acre for that crop.

When applying Concentric SS through stationary or non-continuous moving systems, inject the product-water mixture in the last 15-30 minutes of each set allowing sufficient time for all the required pesticide to be applied by all the sprinkler heads and applying the labeled rate per acre for that crop. Apply Concentric SS continuously for the duration of the water application.

General Information and Instructions for Irrigation Systems: Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Public system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Irrigation systems connected to public water systems must contain a functional, reduced- pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticides distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. For drip (trickle) irrigation: The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

FOLIAR AND SOIL APPLICATION USE RATES FOR OUTDOOR AND GREENHOUSE CROPS

For all crops, Concentric SS can be applied to the soil or to the foliage. Use rates range from 5-20 fluid ounces per acre for soil and foliar applications depending upon the application method, product placement and timing. Refer to the Foliar applications and Soil applications directions hereafter for use rate and timing information.

Foliar applications: Concentric SS may be applied directly to the foliage of actively growing plants at a minimum rate of 10 fluid ounces per acre in a minimum spray volume of 5 gallons per acre. Concentric SS may be applied with all nutrient products and pesticides, but provides best results when applied with nutrients.

Soil applications (including irrigation): Concentric SS works best when soil applications are made with fertilizer or nutrient products and can be applied with liquid or granular formulations. Rates will depend upon the formulation and placement of the product. Rates are as follows:

1. Impregnated on granules - apply Concentric SS at a rate of 13-20 fluid ounces per acre.
2. Formulated with liquid fertilizer - apply Concentric SS at 8 to 13 fluid ounces per acre when applied in furrow at planting. For broadcast, side dress, or top dress applications, apply at 13-20 fluid ounces per acre.
3. For all applications through an irrigation system, apply at 13 fluid ounces per acre.

FOR CEREAL GRAINS, VEGETABLE CROPS, AND OTHER AGRONOMIC CROPS: apply a minimum rate of 10 fluid ounces of Concentric SS in a minimum of 5 gallons finished spray per acre when applied as a foliar spray. For soil applications, a rate of 8-20 fluid ounces per acre is recommended, depending upon the application method, product placement and timing.

Specific Use Restrictions: Do not apply more than 20 fluid ounces per acre per application. Do not apply more than 40 fluid ounces per acre per season. Do not make post-harvest applications. PHI: 0 days.

Cereal Grains, including:

Barley, Buckwheat, Corn (sweet and field), Millet, Oats, Popcorn, Rice, Rye, Sorghum, Triticale, Wheat, Wild rice

Cucurbit and Fruiting Vegetables, including:

Cucumber, Eggplant, Gherkin, Muskmelon, Okra, Pepper, Pumpkin, Squash, Tomato, Watermelon,

Leafy Vegetables, including:

Arugula, Cardoon Celery, Chinese Celery, Chervil, Cilantro, Endive, Fennel, Lettuce, Orach, Parsley, Purslane, Radicchio, Rhubarb, Spinach, Swiss Chard

Brassica (Cole) Leafy Vegetables, including:

Broccoli. Broccoli Raab. Brussel Sprouts. Cabbage. Chinese Cabbage (Boc Choy), Chinese Cabbage (Napa), Chinese Mustard Cabbage (Gai Choy), Cauliflower, Collards, Kale, Kohlrabi, Mustard Greens, Mustard Spinach

Legume Vegetables, including:

Beans, Broad bean, Chickpea, Lentil, Peas, Soybean

Root, Tuber and Bulb Vegetables, including:

Beet (garden and sugar), Carrot, Cassava, Celeriac, Chayote, Chervil, Chicory, Garlic, Ginger, Ginseng, Horseradish, Leek, Onion, Parsley root, Potato, Radish, Rutabaga, Salsify, Shallot, Sweet potato, Turnip, Yam

Other Agronomic Crops, including:

Artichoke, Asparagus, Coffee, Cotton, Grass grown for seed, Hops, Industrial hemp (non-food use), Jojoba, Papaya, Pistachio, Sesame, Sunflower, Tea, Mint, Spearmint

FOR FRUIT, NUT, BERRY AND VINE CROPS: apply a minimum rate of 10 fluid ounces of Concentric SS in a minimum of 25 gallons finished spray per acre when applied as a foliar spray. For soil applications, a rate of 8-20 fluid ounces per acre is recommended, depending upon the application method, product placement and timing.

Specific Use Restrictions: Do not apply more than 20 fluid ounces per acre per application. PHI: 0 days.

Berries and Vine Crops, including:

Blackberry, Blueberry, Cranberry, Elderberry, Gooseberry, Grape, Huckleberry, Loganberry, Raspberry, Strawberry

Citrus, including:

Grapefruit, Lemon, Lime, Mandarin, Orange, Pummelo, Tangerine

Nut Crop, including:

Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory, Macadamia, Pecan, Pistachio, Walnut

Pome Fruit, including:

Apple, Crabapple, Loquat, Mayhaw, Pear, Quince Stone Fruit, including:
Apricot, Cherry, Nectarine, Peach, Plum, Prune, Plumcot

FOR ORNAMENTAL CROPS (including broadleaf shrubs and trees, flowering plants and bulbs, and foliage plants): apply a minimum rate of 10 fluid ounces of Concentric SS in a minimum of 25 gallons finished spray per acre when applied as a foliar spray. For soil applications, a rate of 8 to 20 fluid ounces per acre is recommended, depending upon the application method, product placement and timing.

Specific Use Restrictions: Do not apply more than 20 fluid ounces per acre per application. Do not apply more than 40 fluid ounces per acre per season.

IMPORTANT NOTE: Plant sensitivities to Concentric SS have been found to be acceptable for plants listed on this label; however, it is impossible to know sensitivities under all conditions and phytotoxicity may occur. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for sensitivity to Concentric SS. Neither the manufacturer nor seller, endorses use upon species not listed on the label, nor has it been determined that Concentric SS can be safely used on ornamental or nursery plants not listed on this label. The user must determine if Concentric SS can be used safely prior to commercial use. In a small area, apply the listed rates to the plants in question, i.e. foliage, fruit, etc., and observe for 7-10 days for symptoms of phytotoxicity prior to commercial use. Do not apply foliar sprays to open blooms of Geranium, Marigold, Pansy, and Petunia.

Broadleaf Shrubs and Trees:

Andromeda, Ash, Aspen, Azalea, Buckeye, Camellia, Cherry Laurel, Crabapple, Dogwood, Eucalyptus, Euonymus, Firethorn, Flowering almond, Flowering cherry, Flowering peach, Flowering plum, Flowering quince, Hawthorn, Holly, Laurel, Lilac, Magnolia, Maple, Oak, Poplar, Privet, Red-tip, Rhododendron, Sequoia, Spirea, Sycamore, Viburnum, Walnut

Flowering Plants and Bulbs:

African violet, Begonia, Carnation, Chrysanthemum, Crocus, Daffodil, Daisy, Geranium*, Gladiolus, Hollyhock, Hydrangea, Iris, Lily, Marigold*, Narcissus, Pansy*, Petunia*, Phlox, Rose, Statice, Zinnia

*NOTE: Do not apply foliar sprays of Concentric SS to open blooms of these species.

Foliage plants:

Aglaonema, Arnesenia, Boston fern, Dracaena, Dumbcane, Fatsia, Ficus, Leatherleaf fern, Lipstick plant, Ming aralias, Oyster plant, Pachysandra, Palm, Parlor palm, Peperomia, Philodendron, Prayer plant, Ruffle fern, Syngonium, Zebra plant

SEED TREATMENT APPLICATION INSTRUCTIONS

Concentric SS is a water-based liquid plant growth regulator for use as a seed treatment on a wide range of crops including field crops, vegetables, fruits, grains, and turf seed. Concentric SS is used to provide better and faster seedling emergence through improved seed germination and seedling development, to stimulate root and shoot growth, increase chlorophyll content, improve the plants ability to withstand stress and increase yields. Concentric SS can be used alone, but should be applied in mixtures with fungicide and insecticide seed treatments in situations where the seed and seedlings are at risk from disease or insect attack. Concentric SS can also be applied in mixtures with a wide range of other seed amendments such as bio-stimulants, fertilizers, polymers, fungicides, insecticides, and inoculants.

Concentric SS is designed for easy use as a seed treatment, and should not form sediment during normal storage. It may be applied using standard commercial slurry or mist-type seed treatment equipment which can be calibrated and which provides good mixing of seed and treatment products to provide even distribution and coverage of the seed. Uniform application to seed will give optimum performance. The seed should be sound, well-conditioned, and free from dust and chaff.

Concentric SS seed treatment can be pre-diluted with clean water, and other seed amendments can also be mixed with it as required, and the total volume of slurry adjusted according to the application equipment and seed type. Apply gentle agitation during mixing and use, and do not store the diluted product overnight.

Read the entire label before using Concentric SS for seed treatment. Consult your State Agricultural Experimental Station or Extension Service Specialist for additional information on rates of use, application and any additional requirements or restrictions.

SEED TREATMENT APPLICATION RATES

Apply Concentric SS at rates of 0.8 to 3.3 fl. oz per 100 lbs of seed. Rate of use will depend on seed type, seed size and on use of mixtures.

Specific Use Restrictions and Precautions:

- Do not use treated seed for food, feed or oil processing purposes
- Ensure that the treated seed is colored with a suitable dye to prevent its accidental use as food or feed
- Do not use the same equipment including augers for treating seed that will handle commercial or feed grains
- Store treated seed in cool dry conditions
- Any treated seed which is stored until the following season must be re-tested for germination and vigour prior to sowing

Compatibility: Concentric SS is compatible with most commonly used seed amendment products, and it is advisable not to tank-mix Concentric SS with any other product unless adequate physical compatibility and phytotoxicity (germination) tests have been conducted and proven to be satisfactory. Consult your supplier or advisor for further information.

Treated Seed Must Be Labelled: Seed that is treated with Concentric SS and then packaged or bagged for future use must carry appropriate labelling on the outside of the package or bag

"This seed has been treated with Concentric SS (plus any other pre-mixed seed amendments) - Do not use for food, feed or oil purposes."

Also include the date and establishment where treated, name and address on the label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Keep pesticide in original container. Keep container tightly closed when not in use. Store product above 400F. Do not store in aluminum, fiberglass, copper, brass, zinc, or galvanized containers. Protect from excessive heat. Store in a cool, dry place.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: {for non-refillable containers \leq 5 gallons} Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident,
call CHEMTREC 1-800-424-9300**

WARRANTY DISCLAIMER

The Directions for Use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY THE MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

[Batch Code/Lot Number: ____]

[EPA APPROVAL DATE]