



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

94730-30

Date of Issuance:

3/4/22

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

GCS Sulfentrazone 62.1% +
Cloransulam 7.9% DF

Name and Address of Registrant (include ZIP Code):

Generic Crop Science, LLC
c/o Wagner Regulatory Associates, Inc.
P.O. Box 640
Hockessin, DE 19707

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Continues page 2

Signature of Approving Official:

Mindy Ondish, Product Manager 23
Herbicide Branch, Registration Division (7505P)

Date:

3/4/22

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Please note the alternate brand name, "**GCS Sulfen Cloran**" has been added to the product record.

Please also note that the record for this product currently contains the following CSF:

- Basic CSF dated 10/14/2021

If you have any questions, please contact Julia Kerr by phone at 202-566-2810, or via email at kerr.julia@epa.gov

Enclosure

SULFENTRAZONE	GROUP	14	HERBICIDE
CLORANSULAM-METHYL	GROUP	2	HERBICIDE

GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF

ABN: GCS Sulfen Cloran

Active Ingredients:	By Wt.
Sulfentrazone*	62.1%
Cloransulam-methyl*	7.9%
Other Ingredients:	30.0%
Total:	100.0%

*GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF contains 0.7 pound of active ingredient per pound of product (0.62 pound a.i. of sulfentrazone and 0.08 pound of a.i. of cloransulam-methyl).

KEEP OUT OF REACH OF CHILDREN CAUTION


FIRST AID	
If Swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on Skin or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For 24-Hour Medical Emergency Assistance (Human or Animal), call: 1-800-222-1222 . For Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), call CHEMTREC: 1-800-424-9300 .	

[Optional referral statements when booklets and container labels are used:]

[See label booklet for [complete] [additional] [First Aid,] [Precautionary Statements,] [Directions For Use,] and [Storage and Disposal.]]

EPA Reg. No. 94730-30
EPA Est. No. _____

Manufactured [For] [By]:
Generic Crop Science, LLC
1887 Whitney Mesa Drive, #9740
Henderson, NV 89014



Net Contents: _____ [Lbs./Kgs.]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators, mixers, loaders, and other pesticide handlers must wear:

- Protective eyewear
- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protections Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-f)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.607(e-f)]. Aerial applicators must be in enclosed cockpits.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling the product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to marine/estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Groundwater Advisory:

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not use on coarse soils classified as sand which have less than 1% organic matter.

Surface Water Advisory:

Sulfentrazone can contaminate surface water through spray drift. Under some conditions, sulfentrazone may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

Non-Target Organism Advisory Statement:

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

PHYSICAL/CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Restrictions:

- **Engineering Controls** - When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-f)], the handler PPE requirements may be reduced or modified as specified in the WPS. Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.607(e-f)]. Aerial applicators must be in enclosed cockpits.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

WEED RESISTANCE MANAGEMENT

GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF, which contains the active ingredients sulfentrazone and cloransulam-methyl is a group 14 and 2 herbicide based on the mode of action (MOA) classification system of the Weed Science Society of America. Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different sites of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small (less than 4 inches).
- Apply full rates of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** for the most difficult to control weed in the field at the specified time (correct weed size) to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your local retailer or county extension agent.
- Contact your crop advisor or extension agent to find out if suspected resistant weeds to these MOAs have been found in your region. Do not assume that each listed weed is being controlled by multiple sites of action. Products with multiple active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredient in this product.
- If resistance is suspected, treat weed escapes with an herbicide having a site of action other than Group 14 and 2 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species; and
 - Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad-spectrum soil-applied herbicide with other sites of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative sites of action.
- Rotate the use of this product with non-Group 14 and 2 herbicides.
- Avoid making more than two applications of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** and any other Group 14 and 2 herbicides within a single growing season unless mixed with an herbicide with a different site of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in an around fields, during and after harvest to reduce weed seed production.

PRODUCT INFORMATION

GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF is for preemergence control of broadleaf and grass weeds in soybeans only.

The mode of action of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** involves uptake by weed roots and shoots. Preemergence and preplant incorporated applications of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** require rainfall or irrigation to activate the herbicide. The amount of rainfall or irrigation required for activation following application depends on existing soil moisture, organic matter content and soil texture. If adequate moisture (1/2" to 1") is not received within 7 to 10 days after the **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** treatment, a shallow cultivation may be needed to obtain desired weed control. When sufficient moisture is received after dry conditions, **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** will provide control of susceptible germinating weeds.

GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF exhibits excellent crop safety. Poor growing conditions, such as excessive moisture, cool temperatures, and soil compaction or the presence of various pathogens may impact seedling vigor. Under these conditions, the active ingredients in **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF**, like other soil-applied herbicides, can contribute to crop response.

It is the pesticide user's responsibility to ensure that all products in tank mixtures with **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

IMPORTANT RESTRICTIONS

1. Do not apply **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** if there are visible signs of cracking due to soybean emergence, or serious crop injury such as but not limited to stand loss may result.

IMPORTANT PRECAUTIONS

1. Back-to-back application of ALS or ALS containing herbicides can occasionally result in residual herbicide stacking and potential crop injury. Applicator and grower are responsible and should be aware of previous herbicide use and potential interaction it may have with **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** application.
2. Ensure the seed furrow is closed and the seed covered on acres treated with **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF**.
3. Soybean stunting may occur if excessive rainfall occurs after application but before soybeans emerge. Injury is more prevalent under poor drainage or compacted conditions or when soil is saturated for long periods of time. Soybeans outgrow stunting once favorable growing conditions return.
4. Seedling disease, nematodes, cold weather, deep planting (more than 2"), excessive moisture, high salt concentration, or drought may weaken soybean seedlings and increase the possibility of crop injury.
5. When tank mixing, follow the most restrictive use rates and precautions of the mixing partners.

APPLICATION INFORMATION

DO NOT APPLY TO CROPS OTHER THAN SOYBEANS.

Ground Application

Use a standard low pressure herbicide boom sprayer equipped with suitable nozzles and screens. Apply uniformly using properly calibrated nozzles and screens and strainers no finer than 50 mesh. Use 10 to 40 gallons of spray solution per acre. A minimum of 15 gallons of spray solution per acre must be used for use in a tank mix with one or more contact, burndown herbicides. Do not exceed 40 psi spray pressure unless required by the spray nozzle manufacturer.

Continuous agitation during application is required. Avoid swath overlaps. Shut off spray booms while turning, slowing or stopping, as over application may result. Do not allow **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** spray mixtures to sit overnight as settling of product and difficulty of re-suspending may occur.

To avoid injury to sensitive crops, spray equipment used for **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** applications must be drained and thoroughly cleaned with water plus ammonia before being used to apply other products. See Spray Clean-out Section.

Avoid all direct, and/or indirect spray contact with non-target plants. Do not apply near desirable vegetation. Allow adequate distance between target area and desirable plants to minimize exposure.

Aerial Application

Use Restrictions

Aerial application is allowed only when environmental conditions prohibit ground application. When this product is applied by air, applicator must use a minimum finished spray volume of 5 gallons per acre. The maximum release height must be 10 feet from the top of the canopy, unless a greater application height is required for pilot safety.

Do not apply when wind speed favors drift beyond the area intended for treatment.

These requirements must be followed to avoid off-target movement from aerial applications. These requirements do not apply to forestry applications, public health uses or to applications of dry materials.

1. The distance of the outermost nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

3. Observe the regulations of the State where applications are made.
4. Applicators must observe and abide by the requirements of the **SPRAY DRIFT ADVISORIES**.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage for pesticide performance. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (see information on **Wind**, **Temperature and Humidity**, and **Temperature Inversions** in subsequent sections). Select nozzles and application pressure that deliver medium to coarse or larger spray droplets as indicated in the nozzle manufacturer's recommendations and in accordance with ASABE Standard S-572. Select coarse to very coarse droplet size when used as a preemergent/preplant application. Select medium to very coarse droplet size when used postemergence with a contact burndown herbicide. Do not apply as spray droplets smaller than medium to coarse (defined by the ASAE standard).

Controlling Spray Droplet Size

Volume – Use high flow rate nozzles to apply the greatest practical spray volume. Nozzles with higher rated flow generally produce larger droplets.

Pressure – When higher flow rates are needed, use higher flow rate nozzles rather than increasing spray pressure. Do not exceed the nozzle manufacturer's recommended pressures. Lower pressure produces larger droplets in many types of nozzles.

Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation – For aerial application, the recommended practice is to orient nozzles so that the spray is released parallel to the airstream. This orientation usually produces larger droplets as compared to other nozzle orientations. Significant nozzle deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low drift nozzles for both ground and aerial applications. Solid stream nozzles oriented straight back usually produce the largest droplets and the lowest drift potential in aerial applications.

Boom Length – For some aerial use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height – Aerial applications should not be made at a height greater than 10 feet above the top of the target plant canopy unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment – When aerial applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by the path of the aircraft upwind. Swath adjustment or offset distance should increase when conditions favor increased drift potential (higher winds, smaller droplets, etc).

Wind – Drift potential is lowest between wind speeds of 3-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. Application should be avoided below 3 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they may potentially affect spray drift.

Temperature and Humidity – When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions – Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small-suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the low speed and variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common during conditions of limited cloud cover and little to no wind. They often begin to form as the sun sets and may often continue into the morning. The presence of a temperature inversion may be indicated by ground fog. However if fog is not present, the movement of smoke from a ground source or an aircraft smoke generator can also identify inversions. Smoke that remains in layers and moves laterally in a concentrated cloud (under low-speed wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas – The pesticide should only be applied when the wind is blowing away from sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops).

Off-Target Movement of GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF

Drift of dilute spray mixtures containing **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** must be prevented. Observation of the preceding environmental conditions, correct application equipment design, calibration and application practices will significantly diminish the risk of off-target spray drift. **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** can cause significant symptomology by drift on to sensitive crops and other plants. This symptomology may manifest initially as discreet, localized spots where contacted by **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** drift mixtures. Depending on concentration of the spray solution and droplets size (effectively determining the dosage of sulfentrazone) and also depending on the inherent sensitivity of the plants involved, these spots or lesions may or may not coalesce. These effects will usually not have lasting effects on plant growth, but will likely reduce the value of affected fruit or foliage where grade or quality is associated with appearance. In severe drift instances with particularly sensitive crops, defoliation of affected foliage could result. Failure to follow these guidelines and environmental prohibitions that then result in off-target movement or drift of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** on to unintended crops or plants, irrespective of severity, constitutes misapplication of this product. Generic Crop Science, LLC accepts no responsibility or liability for potential crop effects that may result from such misapplication of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF**.

MIXING INSTRUCTIONS AND LOADING INSTRUCTIONS**Restrictions**

- Do not apply this product through any type of irrigation system.
- Do not use flood irrigation to apply or incorporate this product.

Proper handling instructions

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Product must be used in a manner which will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF Applied Alone

Select the proper **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** application rate from the following **TIMING AND METHOD OF APPLICATION** section of this label. Fill the spray tank with approximately one-half of the volume of water needed for the acreage being treated. With agitator operating, add the required amount of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** for acreage being treated by opening the bottle(s) and measuring directly into the spray tank. Allow the product to fully disperse. Complete the addition of spray water. Maintain agitation during filling, mixing and application. Apply the **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** spray mixture immediately after mixing. Do not store mixture.

GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF Applied in Tank Mix Combination

Select the proper **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** application rate from **TIMING AND METHOD OF APPLICATION** section of label. It is the pesticide user's responsibility to ensure that all products in tank mixtures with **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. To ensure product compatibility, a jar test should be conducted before large volume mixing. Provided the jar test indicates the mixture is compatible, prepare the tank mixture as follows.

Fill the spray tank with approximately one-half of the volume of water needed for the acreage being treated. With agitator operating, add the required amount of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** for the acreage being treated by opening the bottle(s) and measuring directly into the spray tank. Allow the product to fully disperse. Next add the specified amount(s) of the additional tank mix product(s) in the following order: first dry formulations (e.g., wettable powders, dry flowables), next liquid suspensions (e.g., flowables) and finally liquids (e.g., EC's). Allow time for complete mixing and dispersion after each addition, adding water as necessary. Complete the addition of spray water. Maintain agitation during filling, mixing and application. Use **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** tank mixtures immediately after mixing. Do not store tank mixtures.

Fertilizer Spray Mixtures

Applications of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** alone, or with recommended tank mixtures, in conjunction with fertilizer solutions may be used unless use directions specifically state otherwise. Small quantities should be tested for compatibility by the following procedure before mixing in full spray tank quantities.

- Put 1 pint of fertilizer solution in a quart jar.
- Add the appropriate amount of herbicide based on the table below. If more than one product is to be used, add each separately using the following sequence: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's).

Herbicide Type	Herbicide Field Use Rate	Amount Herbicide Added Per Pint*
Wettable Powder Or Dry Flowable	0.5 lb.	0.75 teaspoon
	1.0 lb.	1.50 teaspoons
	2.0 lbs.	3.00 teaspoons
	3.0 lbs.	4.50 teaspoons
Emulsified Concentrates	1.0 pt.	0.5 teaspoon
Liquid Flowables	1.0 qt.	1.0 teaspoon
	2.0 qts.	2.0 teaspoons
	3.0 qts.	3.0 teaspoons

*Based on a spray volume of 25 gal/A. For lower or higher spray volumes, adjust fluid fertilizer quantity accordingly.

- Close jar and shake well.
- Watch mixture for several seconds, again after 5 minutes and again after 30 minutes. If herbicide/fertilizer combination remains mixed or can be remixed readily (i.e., does not permanently separate, foam, gel or become lumpy), the mixture is compatible and can be mixed in full volumes and sprayed. If the mixture is compatible, prepare spray by adding fertilizer solution to the tank first, then follow directions noted below.

GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF Applied Alone with Liquid Fertilizer

In order to add **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** to a liquid fertilizer carrier, **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** must be premixed in a slurry of product and clear water.

Fill the spray tank one-half full with fertilizer solution. With agitator operating, add the **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** slurry to the spray tank.

Use a minimum of one gallon of water for each container of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF**. Stir until completely dissolved. Then add slurry to the spray tank through a 20-35 mesh screen. Rinse container used for pre-mixing and add rinsate to the spray tank. Complete filling the sprayer tank with fertilizer. Maintain agitation during filling, mixing and application. Use **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** spray mixture immediately after mixing. Do not store mixture.

GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF Applied in Tank Mix Combinations with Fertilizer

Fill the spray tank one-half full with fertilizer solution. With the agitator operating, add a slurry of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** as described in the preceding paragraph. Next dilute the individual tank mix partners with sufficient water to form a free-flowing dispersion, then add to the spray tank of fertilizer. While maintaining agitation, add the other products using the following order: slurry of dry formulations (wetable powders, dry flowables) first, diluted liquid formulations (EC's, flowables) second. Complete filling the sprayer tank with fertilizer. Maintain agitation during filling, mixing and application. Use **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** tank mixtures immediately after mixing. Do not store tank mixtures.

SPRAYER EQUIPMENT CLEAN-OUT

After spraying GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF and before using sprayer equipment for any other applications, the sprayer must be thoroughly cleaned using the following procedure:

- Drain sprayer tank, hoses, and spray boom and thoroughly rinse the inside of the sprayer tank with clean water to remove sediment and residues. Thoroughly flush sprayer hoses, boom, and nozzles with clean water.
- Fill the tank 1/2 full with clean water, and add appropriate detergent or ammonia (follow manufacturer's directions for use). Fill the tank to capacity and operate the sprayer for 15 minutes to flush hoses, boom, and nozzles.
- Convenient and thorough cleaning of the sprayer can be achieved if the cleaning solution is left in the spray tank, hoses, spray booms and spray nozzles overnight or during storage.
- Before using the sprayer, drain the sprayer system. Rinse the tank with clean water and flush through the hoses, boom, and nozzles. Remove and clean spray tips and screens separately with the detergent or ammonia solution.
- Properly dispose of all cleaning solution and rinsate in accordance with Federal, State, and local regulations and guidelines.

Do not drain or flush equipment on or near desirable trees or plants. Do not contaminate any body of water including irrigation water that may be used on other crops.

Should small quantities of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to certain crops and other vegetation. Generic Crop Science, LLC accepts no liability for any effects due to inadequately cleaned equipment.

CROP ROTATIONAL RESTRICTIONS

Shown below are the minimum intervals in months from the time of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** application until **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** treated soil may be replanted with the crops listed. Cover crops for soil health

and erosion control can be planted at any time after an application of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF**, but do not use for food or feed. Residual activity of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** may result in injury to some cover crop species if planted too soon following application. Consult your local University extension service for cover crop sensitivity to **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF**. For crops not listed the interval is 30 months and a successful field bioassay. It is the pesticide user's responsibility to ensure that all products in tank mixtures with **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

CROP ROTATIONAL RESTRICTIONS TABLE

Crop	Interval (Months)
Alfalfa	12
Barley	12
Canola	24
Corn, Field*	18 or 10
Corn, Pop*	18 or 10
Corn, Seed*	18 or 10
Corn, Sweet*	18 or 12
Cotton	18 or 12 [†]
Dry Shelled Beans and Peas	9
Lima Beans	12
Oats	12
Peanuts	12
Potatoes	18
Rice	10
Rye	12
Snap Beans	12
Sorghum	12
Soybeans	Anytime
Succulent Peas	9
Sugar Beets**	30
Sunflower**	30
Tobacco***	30
Wheat	4

*Corn (including field corn, popcorn, and seed corn): Observe an 18-month rotational interval if 6.45 - 8.0 oz./A of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** is applied to soils of 1.5% organic matter or less, and pH is above 7.

Hybrid Seed Production: Corn inbred lines grown for hybrid seed production may be injured the growing season following an application of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF**. Inbred lines should be thoroughly tested for crop tolerance before rotating to production scale acreages. Generic Crop Science, LLC will not accept responsibility for any crop injury on field corn grown for seed following an application of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF**.

**These crops require a 30-month rotational interval and a successful field bioassay.

***Transplanted tobacco may be planted 10 months after application of a maximum application rate of 3.0 oz./A of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF**. Tobacco in seedbed nurseries may be replanted 18 months after applications of 3.0 oz./A of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** and following a successful field bioassay. A rotational interval of 30 months and a successful field bioassay is required for all applications of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** greater than 3.0 oz./A.

[†]Cotton may be planted after 12 months where **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** was applied at rates 5 oz./A or less and meets the following conditions:

- Medium and fine soils
- pH <7.2
- Rainfall or irrigation must exceed 15" after application of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF**

REPLANTING INSTRUCTIONS

If the initial planting of soybeans fails to produce a uniform stand, soybeans may be replanted in fields treated with **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** alone. Do not retreat fields with a second application of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF**. When tank mixing with a labeled product, refer to the replant instructions for that product. Do not replant treated fields with any crop at intervals that are inconsistent with the Rotational Crop Guidelines on the **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** label. It is the pesticide user's responsibility to ensure that all products in tank mixtures with **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

SOYBEANS (Conventional and Glyphosate-Resistant)

TIMING AND METHOD OF APPLICATION

GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF may be used alone or in tank mixture combinations for the control of the weeds listed in conventional or glyphosate-resistant soybean varieties.

Standard Rate Table 1:

Soil Organic Matter*	GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF (Oz./A)	Product Use Rates (Lb. a.i./A)
----------------------	---	--------------------------------

		Sulfentrazone	Cloransulam-methyl
3% or less	6.45	0.25	0.032
Greater than 3%	8.00	0.31	0.040

*Do not apply GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF to soils classified as sand with less than 1% organic matter.

Preplant Incorporated Application

Apply GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF alone or in tank mix combination with other herbicides registered for preplant incorporated application to soybeans. Incorporate the herbicide(s) into the top 1 to 3 inches of the final seedbed using equipment that provides thorough soil mixing. When GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF is applied in tank mix combination with other herbicide(s), follow the incorporation directions for the tank mix partner(s). It is the pesticide user's responsibility to ensure that all products in tank mixtures with GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Preplant Surface Application

Apply GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF alone or in tank mix combination with other herbicides registered for preplant soil surface application to soybeans. If applied in tank mix combination, follow use instructions, including application rates (**Note:** Apply 1/2 of the maximum application rate for suppression of weeds in glyphosate-resistant soybeans, maintaining control with sequential application(s) of registered postemergence herbicides), precautions and restrictions of each product used in the tank mixture.

Preemergence Application

Apply at planting time or within 3 days after planting. GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF may be applied alone or in tank mix combination with other herbicides registered for preemergence application to soybeans. When applied in tank mix combination, follow applicable use instructions, including application rates (**Note:** Apply 1/2 of the maximum application rate for suppression of weeds in glyphosate-resistant soybeans, maintaining control with sequential application(s) of registered postemergence herbicides). Observe the precautions and restrictions of each product used in the tank mixture. Apply before planting, at planting time or prior to seed germination. Properly closed seed furrows are necessary when applying at planting time of before seed germination. Do not apply later than 3 days after planting (or after seed germination), as crop injury may result.

Weeds Controlled

When used as directed above, GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF will provide control or suppression of the following broadleaf weeds and grasses.

Common Name	Scientific Name
Broadleaves	
Amaranth, Palmer	<i>Amaranthus, Palmer</i>
Amaranth, spiny	<i>Amaranthus, spinosus</i>
Anoda, spurred	<i>Anoda cristata</i>
Beggarweed, Florida	<i>Desmodium tortuosum</i>
Carpetweed	<i>Mollugo verticillata</i>
Cocklebur, common	<i>Xanthium Pensylvanicum</i>
Copperleaf, Hophornbeam	<i>Acalypha ostryaefolia</i>
Croton, tropic	<i>Croton glandulosus</i>
Daisy, American	<i>Eclipta alba</i>
Dayflower, common	<i>Commelina communis</i>
Galinsoga, hairy	<i>Galinsoga ciliata</i>
Groundcherry, clammy	<i>Physalis heterophylla</i>
Groundcherry, cutleaf	<i>Physalis angulata</i>
Horseweed (Marestail)* *	<i>Hippuris vulgaris</i>
Jimsonweed	<i>Datura stramonium</i>
Kochia	<i>Kochia scoparia</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, common	<i>Chenopodium album</i>
Mallow, Venice	<i>Hibiscus trionum</i>
Mexicanweed	<i>Caperonia castaneifolia</i>
Morningglory, entireleaf	<i>Ipomea hederacea integriuscula</i>
Morningglory, ivyleaf	<i>Ipomea hederacea</i>
Morningglory, palmleaf	<i>Ipomea Wrightii</i>
Morningglory, pitted *	<i>Ipomea, lacunosa</i>
Morningglory, purple	<i>Ipomea turbinata</i>
Morningglory, red	<i>Ipomea coccinea</i>
Morningglory, smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomea, purpurea</i>
Mustard, wild	<i>Brassica kaber</i>
Nightshade, Eastern black	<i>Solanum americanum</i>
Nightshade, hairy	<i>Solanum sarrachoides</i>
Nightshade, silverleaf	<i>Solanum elaeagnifolium</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>

Pigweed, smooth	<i>Amaranthus hybridus</i>
Pigweed, tumble	<i>Amaranthus albus</i>
Poorjoe	<i>Diodia teres</i>
Purslane, common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scabra</i>
Ragweed, common **	<i>Ambrosia artemisiifolia</i>
Ragweed, giant **	<i>Ambrosia trifida</i>
Senna, coffee	<i>Cassia occidentalis</i>
Teaweed	<i>Sida, prickly</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Smellmelon	<i>Cucumis melo</i>
Spurge, spotted	<i>Euphorbia maculata</i>
Starbur, bristly	<i>Acanthospermum hispidum</i>
Sunflower, common	<i>Helianthus annuus</i>
Thistle, Russian	<i>Salsola kali</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatus</i>
Grasses	
Barnyardgrass*	<i>Echinochloa crus-galli</i>
Broadleaf signalgrass	<i>Brachiaria platyphylla</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>
Crabgrass, southern*	<i>Digitaria ciliaris</i>
Crowfootgrass*	<i>Dactyloctenium aegyptium</i>
Foxtail, giant*	<i>Setaria faberi</i>
Foxtail, Green*	<i>Setaria viridis</i>
Foxtail, yellow*	<i>Setaria lutescens</i>
Goosegrass	<i>Eleusine indica</i>
Johnsongrass, seedling *	<i>Sorghum halepense</i>
Orchardgrass	<i>Dactylis glomerata</i>
Panicum fall*	<i>Panicum dichotomiflorum</i>
Panicum, Texas	<i>Panicum texanum</i>
Sedges	
Nutsedge, purple	<i>Cyperus rotundus</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Sedge, annual	<i>Cares spp.</i>

*Provides suppression or partial control only.

**Will not control ALS resistant biotypes of these weed species.

Limited Residual Rates for Planned Sequential Application Program in Soybeans

Use rates in Table 2 are to be used in conjunction with an effective planned post herbicide program; **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** at these reduced rates will provide early season control or suppression to reduce early season weed competition. If resistance with the post herbicide is documented in your area, use rates in Table 1.

Apply before planting, at planting time or prior to seed germination. Properly closed seed furrows are necessary when applying at planting time of before seed germination. Postemergence treatments may include any product or combination of products labeled for use.

Limited Residual Rate Table for Planned Sequential Application Program in Soybeans**Table 2:**

Soil Organic Matter*	GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF (Oz./A)	Product Use Rates (Lb. a.i./A)	
		Sulfentrazone	Cloransulam-methyl
3% or less	3.00 - 5.00	0.116 - 0.193	0.015 - 0.025
Greater than 3%	4.00 - 6.00	0.155 - 0.233	0.020 - 0.030

*Do not apply **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** to soils classified as sand with less than 1% organic matter.

Preplant Burndown Application GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF, used at 6.45 - 8.0 oz./A as in Full Rate Table 1 above, aids in the burndown of weeds listed below, when applied as follows. **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** can provide for increased burndown activity on emerged weeds in no-till applications, but is not intended to replace part or all of an appropriate preplant burndown program. For control of the weeds in the Weeds Controlled table in no-till / minimum till fields, **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** must be tank-mixed or used in combination with a full burndown program. This may include 2,4-D alone or in combination with carfentrazone-ethyl, dicamba, glyphosate, glufosinate, paraquat, or other appropriate burndown herbicides in tank-mixes at their appropriate rate for the size and species of weeds present. Reduced rates of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** and/or the corresponding burndown partner herbicides can result in weed escapes and unsatisfactory performance.

Apply a minimum of ten gallons per acre finished spray volume. Thorough coverage is essential. Use a non-ionic surfactant (NIS) having at least 80% active ingredient strength at 0.125 - 0.25% v/v (1 - 2 pints per 100 gallons of spray solution) plus ammonium sulfate (AMS) at 2.5% v/v. Crop oil concentrate (COC) and Methylated Seed Oil (MSO) at 1.2% v/v plus ammonium sulfate may be used. Burndown results may be slowed or reduced when the growth of the weeds is affected by unusual environmental factors just prior to or after application such as especially cool or widely fluctuating day and night air temperatures, drought, heat stress, or waterlogged soils.

Weeds Controlled

When used as directed for burndown, **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** will aid in the control or suppression of the following broadleaf weeds up to 3" tall.

Common Name	Scientific Name
Broadleaves	
Cocklebur, common	<i>Xanthium Pensylvanicum</i>
Horseweed (Marestail)**	<i>Hippuris vulgaris</i>
Jimsonweed	<i>Datura stramonium</i>
Mallow, Venice	<i>Hibiscus trionum</i>
Morningglory, entireleaf	<i>Ipomea hederacea integruscula</i>
Morningglory, ivyleaf	<i>Ipomea hederacea</i>
Morningglory, palmleaf	<i>Ipomea Wrightii</i>
Morningglory, pitted*	<i>Ipomea lacunosa</i>
Morningglory, purple	<i>Ipomea turbinata</i>
Morningglory, red	<i>Ipomea coccinea</i>
Morningglory, smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomea purpurea</i>
Ragweed, common**	<i>Ambrosia artemisiifolia</i>
Ragweed, giant**	<i>Ambrosia trifida</i>
Sicklepod	<i>Cassia obtusifolia</i>
Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Sunflower, common	<i>Helianthus annuus</i>
Velvetleaf *	<i>Abutilon theophrasti</i>

*For Velvetleaf control, use 28% nitrogen (UAN) or AMS with NIS or COC.

****GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** will not control ALS resistant biotypes of these weed species.

Precautions

- Properly closed seed furrows are necessary when applying at planting time of before seed germination.
- Maintain spray tank agitation until the spray mixture is applied.

Restrictions

- Do not use flood irrigation to apply or incorporate this product.
- Do not apply this product through any type of irrigation system.
- The maximum single application rate for **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** is 8.0 oz./A (0.31 lb. sulfentrazone a.i./A and 0.04 lb. cloransulam-methyl a.i./A).
- The maximum amount of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** that can be applied per year is 8.0 oz./A (0.31 lb. sulfentrazone a.i./A and 0.04 lb. cloransulam-methyl a.i./A).
- Do not apply more than 0.375 lb. a.i./A of sulfentrazone from any source per year.
- Do not apply more than 0.055 lb. a.i./A of cloransulam-methyl from any source per year.
- Do not feed treated soybean forage or soybean hay to livestock.
- Do not harvest soybeans for 65 days after application of **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF**.
- Do not apply **GCS Sulfentrazone 62.1% + Cloransulam 7.9% DF** to soils classified as sand containing less than 1% organic matter.
- Do not drain or flush equipment on or near desirable trees or plants. Do not contaminate any body of water including irrigation water that may be used on other crops.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store product in original container only, away from other pesticides, fertilizer, food, or feed.

In Case of Spill

Avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call **CHEMTREC (Transportation and spills): (800) 424-9300**.

To Confine Spill

Dike surrounding area, sweep up spillage. Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in an approved manner. Place damaged package in a holding container. Identify contents per required hazardous waste labeling regulations.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

Container Handling

[Nonrefillable Container (flexible-bag-all weights):] [Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by any other procedure allowed by state and local authorities.]

[Nonrefillable Container (rigid 50 lbs. or less):] [Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration.]

[Nonrefillable Container (rigid-greater than 50 lbs.):] [Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water.

Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by any other procedure allowed by state and local authorities.]

[Refillable Container:] [Refillable container. Refill this container with this product only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Dispose of empty container in a sanitary landfill or by any other procedure allowed by state and local authorities.]

IMPORTANT: READ BEFORE USE

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the unopened product container at once. By using the product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Generic Crop Science, LLC. To the extent consistent with applicable law, such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GENERIC CROP SCIENCE, LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. To the extent consistent with applicable law, no agent of Generic Crop Science, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GENERIC CROP SCIENCE, LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID OR AT GENERIC CROP SCIENCE, LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

All trademarks are the property of their respective owners.