



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

95009-1

Date of Issuance:

10/27/20

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Maxunitech Carfentrazone-ethyl  
240EC Herbicide

Name and Address of Registrant (include ZIP Code):

Maxunitech North America, LLC  
c/o Pyxis Regulatory Consulting, Inc.  
4110 136<sup>th</sup> St. Ct. NW  
Gig Harbor, WA 98332

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 95009-1."
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Signature of Approving Official:

Shaja B. Joyner, Product Manager 20  
Fungicide-Herbicide Branch, Registration Division 7505P

Date:

10/27/20

EPA Form 8570-6

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 1/16/2020

If you have any questions, please contact Nathan Mellor by phone at 703-347-8562, or via email at [mellor.nathan@epa.gov](mailto:mellor.nathan@epa.gov)

Enclosure

[Note to reviewer: [Text] in brackets denotes optional text].  
[Note to reviewer: {Text} in braces denotes where in the final label text will appear.]

{BOOKLET FRONT PANEL}

CARFENTRAZONE-ETHYL	GROUP	14	HERBICIDE
---------------------	-------	----	-----------

## Maxunitech Carfentrazone-ethyl 240EC Herbicide

ACTIVE INGREDIENT:	By Wt.
Carfentrazone-ethyl .....	23.93%
<b>OTHER INGREDIENTS:</b> .....	<u>76.07%</u>
<b>TOTAL:</b> .....	100.00%

This product contains 2.0 lbs. active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

[See] [inside] [label] [booklet] [for] [additional] [Precautionary Statements],[.] [and] [Directions for Use]  
[including] [Storage and Disposal] [instructions][.]

**EPA Reg. No.: 95009-XX**


**EPA Est. No.:**

**Manufactured for:**

Maxunitech North America LLC  
853 Valparaiso Ave.  
Menlo Park, CA 94025

**[Lot][Batch]No:**

**Net Contents:**



## {LANGUAGE INSIDE BOOKLET}

<b>FIRST AID</b>	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12 PM Pacific Standard Time. In the event of a medical emergency, call your poison control center at 1-800-222-1222.	

### **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION**

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

#### **Personal Protective Equipment (PPE)**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

#### **User Safety Requirements:**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### **User Safety Recommendations**

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling the product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### **ENVIRONMENTAL HAZARDS**

Carfentrazone-ethyl is very toxic to algae and moderately toxic to fish. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the high-water mark, except as specified on this label. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

#### **For Groundwater:**

Residues of this chemical have properties and characteristics associated with chemicals detected in ground water. Residues of this chemical may leach into ground water if the chemical is used in areas where soils are permeable, particularly where the water table is shallow.

#### **For Surface Water:**

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface

water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of carfentrazone-ethyl residues from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT** apply this product through any type of irrigation system.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is: Coveralls, waterproof gloves, and shoes plus socks.

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Re-entry Statement: DO NOT** allow people (other than applicator) or pets on treatment area during application. **DO NOT** enter treatment area until spray has dried.

### **WEED RESISTANCE MANAGEMENT**

For resistance management, Maxunitech Carfentrazone-ethyl 240EC Herbicide is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to Maxunitech Carfentrazone-ethyl 240EC Herbicide and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of Maxunitech Carfentrazone-ethyl 240EC Herbicide or other Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage ( or other mechanical control methods), cultural ( e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled

individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method, for example, hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.

- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Maxunitech North America LLC retailer or representative.

Report any incidence of non-performance of this product against a particular weed species to your Maxunitech North America LLC retailer or representative. If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemical means to remove escapes, as practical, with the goal of preventing further seed production.

## **PRODUCT INFORMATION**

Maxunitech Carfentrazone-ethyl 240EC Herbicide is an emulsifiable concentrate formulation. Maxunitech Carfentrazone-ethyl 240EC Herbicide is to be mixed with water, liquid fertilizer or mixtures of water and liquid fertilizer and adjuvants and applied to labeled crops and non-crop areas for selective postemergence control of broadleaf weeds, for sucker control, for burndown prior to planting, as a harvest aid and to defoliate/desiccate labeled crops.

Weed control is optimized when the product is applied to actively growing weeds. This product is a contact herbicide. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation.

Extremes in environmental conditions including temperature, moisture, soil conditions, and cultural practices may affect the activity of this product. Herbicide symptoms may be accelerated under moist conditions. Weed control may be reduced when weeds are hardened off by drought and become less susceptible to Maxunitech Carfentrazone-ethyl 240EC Herbicide.

This product is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications should not be made within 6 to 8 hours of either rain or irrigation or when heavy dew is present on the crop. Environmental conditions and with certain spray tank additives may increase herbicidal symptoms on the crop.

## **TANK MIXTURES**

This product may be tank-mixed with other registered herbicides for controlling broader spectrum weeds. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. When preparing a new tank mix conduct an appropriate compatibility test by mixing proportional amounts of all spray ingredients in a test vessel (jar) prior to tank mixing with other products. Shake the mixture vigorously and allow it to stand for five to ten minutes. Rapid precipitation of the ingredients and failure to re-suspend when shaken indicates that the mixture is incompatible and should not be applied. Provided the jar test indicates the mixture to be compatible, prepare the tank mixture as follows: Fill the tank one fourth full with water. With the agitator operating, add the directed amounts of ingredients using the following order: dry granules first and liquid suspensions (flowables) second. As the agitation continues and the tank is filled with water add emulsifiable concentrate products third followed by the addition of water-soluble products.

## **ADJUVANT USE REQUIREMENTS**

The use of a quality spray adjuvant is required for optimum performance. Refer to the individual crop sections of this label for specific adjuvant type and use rates.

## **ON-FARM TESTING**

Not all varieties or cultivars of labeled crops have been fully evaluated under all environmental and soil conditions. Consult with your local seed company for additional information.

It may also be beneficial to conduct small on-farm trials under actual conditions with specific varieties or cultivars before treating large acreage.

## **APPLICATION INFORMATION**

### **Mixing and Loading Instructions**

Start by filling the tank with  $\frac{3}{4}$  of the desired volume of clean water and, with agitation, add the proper amount of this product. Complete filling the spray tank to the desired volume. Maintain sufficient agitation to keep materials in solution during both mixing and application and until the spray tank has been emptied. For tank mixtures, follow your local extension guidelines for mixing order. Guidelines are: add dry materials first and agitate until mixed; then EW or

water soluble liquids; then EC formulations; then, add adjuvants last. Ensure the compatibility of other products and/or liquid fertilizers with this product before mixing them together in the spray tank.

### **Mixing Precautions**

Avoid the overnight storage of Maxunitech Carfentrazone-ethyl 240EC Herbicide spray mixtures. If spray solution is stored overnight or longer, thoroughly agitate spray mixture before applying the solution. **DO NOT** premix Maxunitech Carfentrazone-ethyl 240EC Herbicide spray solutions in nurse tanks. Maintain continuous and adequate spray solution agitation until all the spray solution has been used. **DO NOT** use with tank additives that alter the pH of the spray solution below pH 5 or above pH 8. Buffer spray solutions if necessary to maintain the pH between 5-8.

### **SPRAY EQUIPMENT CLEAN-OUT**

**Many new pesticides are very active at low rates, especially to sensitive crops.** Residues left in mixing equipment, spray tanks, hoses, spray booms and nozzles can cause crop effects if they are not properly cleaned. As soon as possible after spraying this product and before using the sprayer equipment for any other applications, the sprayer equipment must be thoroughly cleaned using the following procedure. In addition, users must take appropriate steps to ensure proper equipment clean-out for any other products mixed with this product as required on the other product labels. More complete cleaning can be achieved if the spray system is cleaned immediately following the application.

1. Drain sprayer tank, hoses, spray boom and spray nozzles. Use a high-pressure detergent wash to remove physical sediment and residues from the inside of the sprayer tank and thoroughly rinse. Then, thoroughly flush sprayer hoses, spray boom and spray nozzles with a clean water rinse. Remove and clean spray tips and all filters and screens (tank, spray hose and spray tips) separately in the ammonia solution of Step 2.
2. Next, prepare a sprayer cleaning solution by adding three gallons of ammonia (containing at least 3% active) per 100 gallons of clean water. Prepare sufficient cleaning solution to allow the operation of the spray system for a minimum of 15 minutes to thoroughly flush hoses, spray boom and spray nozzles.
3. Convenient and thorough cleaning of the sprayer can be achieved if the ammonia solution or fresh water is left in the spray tank, hoses, spray booms and spray nozzles overnight or during storage.
4. Before using the sprayer, completely drain the sprayer system. Rinse the tank with clean water and flush through the hoses, spray boom, and spray nozzles with clean water.
5. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State, and local regulations and guidelines.

**DO NOT** apply sprayer cleaning solutions or rinsate to sensitive crops.

**DO NOT** store the sprayer overnight or for any extended period of time with Maxunitech Carfentrazone-ethyl 240EC Herbicide spray solution remaining in the tank, spray lines, spray boom plumbing, spray nozzles or strainers.

If the sprayer has been stored or idle, purge the spray boom and nozzles with clean water before beginning any application.

Should small quantities of Maxunitech Carfentrazone-ethyl 240EC Herbicide remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to certain crops and other vegetation. Maxunitech North America LLC accepts no liability for any effects due to inadequately cleaned equipment.

### **APPLICATION METHODS**

#### **GROUND APPLICATION**

Use ground sprayers designed, calibrated and operated to deliver uniform spray droplets to the targeted plant or plant parts. Adjust sprayer nozzles to achieve uniform plant coverage. Overlaps and slower ground speeds (caused by continuing to spray while starting, stopping or turning) may result in higher application rates and possible crop response.

#### **Spray Buffer for Ground Application**

Spray buffer zones for ground applications, listed in chart below, are required near desirable perennial vegetation or crops before blossom and after total leaf drop, and/or near other desirable or annual crops.

<b>Buffers for Ground Application</b>		
<b>MAXUNITECH CARFENTRAZONE-ETHYL 240EC HERBICIDE USE RATE (lb. ai per acre)</b>	<b>Low Spray Boom Buffer (ft.)</b>	<b>High Spray Boom Buffer (ft.)</b>
0.024	20	33
0.031	26	46

### **Broadcast Boom Sprayers**

Use a broadcast boom sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Use nozzles that produce minimal amounts of fine spray droplets. **DO NOT** exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Apply a minimum of 10 gallons of finished spray per acre. Use higher spray volumes when there is a dense weed population or crop canopy. Adjust sprayers to position spray tips no lower than 12-18 inches above the crop or weed canopy depending on the nozzle specification. Operate the sprayer to avoid the application of high herbicide rates directly over the rows or into the whorl of treated crop plants.

### **Directed Sprayers**

For directed sprayers apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with drop nozzles or other post directed spray equipment.

**Post directed** applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. **DO NOT** apply when conditions favor drift or when wind speed is above 10MPH.

Use drop nozzles or other spray equipment capable of directing the spray to target weeds and away from sensitive plant parts. Apply when labeled crops have reached minimum growth stages described in specific crop sections of this label and when spray will not be deposited on green stems, foliage, blossoms or fruit.

### **Hooded Sprayers**

To apply Maxunitech Carfentrazone-ethyl 240EC Herbicide using a hooded sprayer, refer to the Hooded Sprayer Section for specific adjustment and operation instructions. For additional information, refer to the individual crop sections of this label.

### **Hand-held or high-volume orchard gun sprayers**

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be applied to certain labeled crops and non-crop areas with hand operated sprayers including backpack sprayers, compression sprayers, knapsack sprayers, or high-volume orchard gun sprayers. Directed applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. Refer to individual crop sections of this label.

### **AERIAL APPLICATION**

Use nozzle types and arrangements that will provide optimum coverage while producing a minimal amount of fine droplets. Apply at a minimum of 3 gallons of finished spray per acre. Spray volumes greater than 3 GPA may be needed for harvest aid and defoliation treatments, or for dense weed populations or with heavy crop canopies.

#### **For Aerial Application in California:**

**Refer to individual crop sections to see if application is permitted by air**

For applications near desirable perennial vegetation or crops before blossom and after total leaf drop, and/or near other desirable or annual crops:

-**DO NOT** apply within 100 feet of all desirable vegetation or crops.

-If wind up to 10 miles per hour is blowing toward desirable vegetation or crops, **DO NOT** apply within 500 feet of the desirable vegetation or crops.

-**DO NOT** apply when winds are in excess of 10 mph or when inversion conditions exist.



## MANDATORY SPRAY DRIFT

### Aerial Applications

- **DO NOT** release spray at a height greater than 10 ft above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

### Ground Applications

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

### Boom-less Ground Applications:

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

## SPRAY DRIFT ADVISORIES

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- IMPORTANCE OF DROPLET SIZE  
An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.  
Controlling Droplet Size - Ground Boom
  - Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
  - Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
  - Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.Controlling Droplet Size - Aircraft
  - Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.
- BOOM HEIGHT - Ground Boom  
Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.
- RELEASE HEIGHT - Aircraft  
Higher release heights increase the potential for spray drift. When applying aerially to crops, **DO NOT** release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.
- SHIELDED SPRAYERS  
Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- TEMPERATURE AND HUMIDITY  
When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS  
Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence

of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

- WIND  
Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.  
Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.
- Boom-less Ground Applications:  
Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.
- Handheld Technology Applications:  
Take precautions to minimize spray drift.

## ALLOWABLE MAXUNITECH CARFENTRAZONE-ETHYL 240EC HERBICIDE USE INFORMATION

Refer to the crop section of this label for specific product use directions.

Table 1:

Maximum Allowable Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Per Acre Per Year* for Crop or Subgroup		
Crop Group/Subgroup	Maximum Rate Maxunitech Carfentrazone-ethyl 240EC Herbicide (fl oz/acre) Per Year	Maximum Rate Maxunitech Carfentrazone-ethyl 240EC Herbicide (lb. ai/acre) Per Year
Alfalfa and Clover (Group 18)	2.5	0.04
Alfalfa and Clover (Group 18), harvest aid only	3.8	0.06
Asparagus	3.8	0.06
Banana	7.9	0.124
Berry, low growing (Subgroup 13-07G)	6.1	0.096
Bushberry (Subgroup 13-07B)	6.1	0.096
Cacao	7.9	0.124
Caneberry (Subgroup 13-07A)	25.6	0.4
Citrus fruit (Group 10-10)	7.9	0.124
Coconut	7.9	0.124
Coffee	7.9	0.124
Corn	2.0	0.031
Cotton	7.9	0.124
Cotton, harvest aid only	3.2	0.05
Date	7.9	0.124
Fig	7.9	0.124
Fruit, small vine climbing – except fuzzy kiwifruit (Subgroup 13-07F)	7.9	0.124
Globe Artichoke	6.1	0.096
Grass (Group 17)	5.9	0.093
Guayule	7.9	0.124
Herbs and Spices (Group 19)	6.1	0.096
Hops	7.7	0.12
Horseradish	6.1	0.096
Indian Mulberry	7.9	0.124
Kiwifruit	7.9	0.124
Mint	1.92	0.030
Nut, Tree (Group 14-12)	7.9	0.124
Oil Seed – except cottonseed (Group 20)	6.1	0.096
Olive	7.9	0.124
Palm Heart	7.9	0.124
Peanut	6.1	0.096
Peanut (harvest aid)	2.0	0.031
Persimmon	7.9	0.124
Pome fruit (Group 11-10)	7.9	0.124
Pomegranate	7.9	0.124

**Maximum Allowable Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Per Acre Per Year\* for Crop or Subgroup**

<b>Crop Group/Subgroup</b>	<b>Maximum Rate Maxunitech Carfentrazone-ethyl 240EC Herbicide (fl oz/acre) Per Year</b>	<b>Maximum Rate Maxunitech Carfentrazone-ethyl 240EC Herbicide (lb. ai/acre) Per Year</b>
Rice (in California only)	19.2	0.3
Rice (Southern use only)	8.8	0.138
Rice, harvest aid only, (not permitted in California)	1.5	0.023
Small Grains	1.0	0.016
Small Grains (except winter wheat)	2.0	0.031
Small Grains (winter wheat)	2.0	0.031
Sorghum (harvest aid)	1.0	0.016
Sorghum (grown for seed and grain)	1.0	0.016
Soybeans (preplant, in-season and harvest aid)	1.5	0.023
Stone fruit (Group 12-12)	7.9	0.124
Sugarcane	6.1	0.096
Sugarcane (harvest aid)	2.0	0.031
Tea	7.9	0.124
Teff	2.0	0.031
Tobacco	3.2	0.05
Tropical fruit Trees	6.1	0.096
Vanilla	7.9	0.124
Vegetable, brassica (Group 5)	6.1	0.096
Vegetable, bulb (Group 3-07)	6.1	0.096
Vegetable, cucurbit (Group 9)	6.1	0.096
Vegetable, foliage of legume (Group 7)	6.1	0.096
Vegetable, fruiting (Group 8-10)	6.1	0.096
Vegetable, leafy (except Brassica) (Group 4)	6.1	0.096
Vegetable, leaves of root and tuber (Group 2)	6.1	0.096
Vegetable, legume (Group 6 – except soybean)	6.1	0.096
Vegetable, root (Subgroups 1A and 1B)	6.1	0.096
Vegetable, tuberous and corm (Subgroups 1C and 1D)	11.6	0.181
Wild Rice	19.2	0.3

**PREHARVEST INTERVALS**

Refer to the crop section of this label for specific product use directions.

**Table 2:**

**Preharvest Intervals (PHI) or Maximum Growth Stage for Maxunitech Carfentrazone-ethyl 240EC Herbicide Applications**

<b>Crop Group/Subgroup</b>	<b>PHI (Days Before Harvest) or Growth Stage</b>
Alfalfa and Clover (Group 18) grown for Forage and/or Hay	21
Alfalfa and Clover (Group 18), grown for Seed	3
Asparagus	5
Banana	3
Berry, low growing (Subgroup 13-07G)	0
Bushberry (Subgroup 13-07B)	0
Cacao	3
Caneberry (Subgroup 13-07A)	15
Citrus fruit (Group 10-10)	3
Coconut	3
Coffee	3
Corn	14 Leaf Collars
Corn, Sweet corn grown for seed, popcorn, field corn (harvest aid)	3
Cotton (harvest aid)	7

**Preharvest Intervals (PHI) or Maximum Growth Stage for Maxunitech Carfentrazone-ethyl 240EC Herbicide Applications**

<b>Crop Group/Subgroup</b>	<b>PHI (Days Before Harvest) or Growth Stage</b>
Cotton (preplant and in-season)	7
Date	3
Fig	3
Fruit, small vine climbing – except fuzzy kiwifruit (Subgroup 13-07F)	3
Globe Artichoke	0
Grass (Group 17)	0
Guayule	3
Herbs and Spices (Group 19)	0
Hops	7
Horseradish	0
Indian Mulberry	3
Kiwifruit	3
Mint	5
Nut, Tree (Group 14-12)	3
Oil Seed – except cottonseed (Group 20)	0
Olive	3
Palm Heart	3
Peanut	7
Persimmon	3
Pome fruit (Group 11-10)	3
Pomegranate	3
Rice (in California only)	60
Rice (Southern use only)	60
Rice, harvest aid only, (not permitted in California)	3
Small Grains (except winter wheat)	Jointing Stage
Small Grains (harvest aid) – include Winter Wheat	7
Sorghum (harvest aid)	3
Sorghum (grown for seed and grain)	14 Leaf Collars Stage
Soybean (harvest aid)	3
Soybeans (preplant, in-season and harvest aid)	V10
Stone fruit (Group 12-12)	3
Sugarcane	7
Tea	3
Teff	Jointing Stage
Teff (forage – harvest aid)	7
Teff (grain – harvest aid)	3
Tobacco	6
Tropical fruit	0
Vanilla	3
Vegetable, brassica (Group 5)	0
Vegetable, bulb (Group 3-07)	0
Vegetable, cucurbit (Group 9)	0
Vegetable, foliage of legume (Group 7)	0
Vegetable, fruiting (Group 8-10)	0
Vegetable, leafy (except Brassica) (Group 4)	0
Vegetable, leaves of root and tuber (Group 2)	0
Vegetable, legume (Group 6 – except soybean)	0
Vegetable, root (Subgroups 1A and 1B)	0
Vegetable, tuberous and corm (Subgroups 1C and 1D)	7
Wild Rice	60

### CROP ROTATIONAL RESTRICTIONS

Following an application of Maxunitech Carfentrazone-ethyl 240EC Herbicide, a treated field may be rotated to a registered crop at any time, subject to specific crop restrictions that may be found in the individual crop sections. All other crops may be planted after 12 months.

### WEED CONTROL

When used as directed, Maxunitech Carfentrazone-ethyl 240EC Herbicide will provide control of the listed weeds up to four (4) inches in height, or as specified.

Table 3:

<b>Weeds Controlled</b>	<b>Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rate fl oz (lb. ai) per acre</b>
Lambsquarters, common (up to 3 inches tall)	0.5 fl oz (0.008 lb. ai) per acre
Morningglory, ivyleaf (up to 3 leaves)	
Morningglory, pitted (up to 3 leaves)	
Nightshade, Eastern black	
Pigweed, redroot	
Velvetleaf	
Waterhemp (up to 2 inches tall)	
<b>Weeds Controlled</b>	<b>Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rate fl oz (lb. ai) per acre</b>
All the weeds controlled at 0.5 fl oz (0.008 lb. ai) per acre plus the weeds listed below:	0.8 fl oz (0.013 lb. ai) per acre
Cheeseweed	
Filaree, redstem	
Flixweed	
Lambsquarters, common	
Mallow, common	
Morningglory, entireleaf	
Morningglory, ivyleaf	
Morningglory, pitted	
Morningglory, scarlet	
Nightshade, hairy	
Pennycress, field	
Pigweed, prostrate	
Pigweed, smooth	
Pigweed, tumble	
Purslane, common	
Sesbania, hemp	
Smartweed, PA (seedling)	
Spurge, prostrate	
Tansymustard	
Velvetleaf (24")	
Waterhemp, common & tall	
<b>Weeds Controlled</b>	<b>Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rate fl oz (lb. ai) per acre</b>
All the weeds controlled at 0.8 fl oz (0.013 lb. ai) per acre plus the weeds listed below:	1.0 fl oz (0.016 lb. ai) per acre
Amaranth, spiny	
Anoda, spurred	
Bedstraw, catchweed	
Buffalobur	
Carpetweed	
Cocklebur	
Copperleaf, hophornbeam	
Cotton, GMO Varieties	

<b>Weeds Controlled</b>	<b>Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rate fl oz (lb. ai) per acre</b>
Cotton, volunteer	
Eclipta	
Fiddleneck, coast	
Groundcherry, smooth (seedling)	
Groundcherry, Wright's	
Jimsonweed	
Kochia	
Lettuce, Prickly 2-3 leaf	
Nettle, burning	
Nightshade, American black	
Nightshade, black	
Rocket, London	
Shepherdspurse	
Speedwell, Virginia	
Spiderwort, tropical	
Thistle, Russian (up to 2 inches tall)	
Wallflower, bushy	
<b>Weeds Controlled</b>	<b>Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rate fl oz (lb. ai) per acre</b>
All the weeds controlled at 1.1 fl oz (0.016 lb. ai) per acre plus the weeds listed below:	1.6 fl oz (0.025 lb. ai) per acre
Amaranth, Palmer	
Corn Spurry	
Filaree, broadleaf	
Filaree, white	
Lettuce, prickly	
Mallow, Venice (up to 2 inches tall)	
Meadowfoam	
Redmaids	

### **Burndown of top growth**

<b>Weed List</b>	<b>Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rate fl oz (lb. ai) per acre</b>
Bindweed, field	1.0 - 2.0 fl oz (0.016 – 0.032 lb. ai) per acre
Burclover	
Dayflower	
Sage, lanceleaf	
Sowthistle	

### **AGRICULTURE FARM AND FARMSTEAD USE – NON-CROP**

#### **Timing and Method of Application**

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be used for broadleaf weed control on farms and farmsteads in areas outside of crop growing areas. See the rate and weed table to determine the proper rate for areas including grass waterways, field edges, terraces, equipment storage areas, shelter belts, fence lines, farm buildings, dry ditch, canal banks etc. Maxunitech Carfentrazone-ethyl 240EC Herbicide is a contact herbicide and coverage is essential for good weed control. Maxunitech Carfentrazone-ethyl 240EC Herbicide will control emerged weeds only. Weeds that germinate after application will require repeat treatments.

#### **BOOM EQUIPMENT**

##### **Maxunitech Carfentrazone-ethyl 240EC Herbicide use Rate – Boom Equipment:**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at up to 2.0 fl oz (0.031 lb. ai) per acre.

##### **Adjuvant Requirements – Boom Equipment**

A nonionic surfactant crop oil concentrate or methylated seed oil is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient or a petroleum or oil seed

based crop oil concentrate (COC) at 1.5 to 2 % v/v (1.5 to 2.0 gallons per 100 gallons of spray solution) or a methylated seed oil (MSO). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons) or ammonium sulfate at 2 to 4 pounds per acre in addition to the selected NIS, MSO or COC is allowed.

### Tank Mixes – Boom Equipment

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be mixed with other herbicides labeled for this method of application in non-crops areas for broader spectrum weed control. See Mixing and Loading Instructions under the PRODUCT INFORMATION section of this label for specific mixing instructions. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank-mix partner.

### SPOT TREATMENTS (Applications with hand operated sprayer including backpack sprayers, compression sprayers, knapsack sprayers.)

Mix the amount of Maxunitech Carfentrazone-ethyl 240EC Herbicide for the desired percent spray solution from the table below. These mixtures are based on 1 gallon of solution evenly covering 1000 square feet. Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. **DO NOT** spray to runoff. See Table 4 for weeds controlled at specific concentrations. Use lower concentrations for small seedling weeds at the 2-3 leaf stage. Higher concentrations are needed for larger weeds up to the 6-leaf stage. Applications beyond the 6-leaf stage may result in only partial control. Maxunitech Carfentrazone-ethyl 240EC Herbicide may be mixed with other labeled herbicides including glyphosate, glufosinate, and paraquat for broader spectrum weed control.

**Table 4:**

Desired Volume	Amount Maxunitech Carfentrazone-ethyl 240EC Herbicide				
	0.5 fl oz/acre (0.008 lb. ai)	0.8 fl oz/acre (0.013 lb. ai)	1.0 fl oz/acre (0.016 lb. ai)	1.6 fl oz/acre (0.026 lb. ai)	2.0 fl oz/acre (0.032 lb. ai)
1 Gal	0.4 ml	0.6 ml	0.7 ml	1.1 ml	1.4 ml
5 Gal	1.7 ml	2.7 ml	3.4 ml	5.4 ml	6.8 ml
25 Gal	8.5 ml	13.6 ml	17.0 ml	27.2 ml	34.0 ml

### Adjuvant Requirements

A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v having at least 80% active ingredient, or a methylated seed oil (MSO), or crop oil concentrate (COC) (petroleum or seed oil) at 1 to 2% v/v. A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v or ammonium sulfate (AMS) at the rate of 0.75 to 1.5 ounces per gallon in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.

**Table 5:**

Desired Volume	Adjuvants				
	NIS	COC or MSO		Liquid Nitrogen	
	0.25% v/v	1.5 % v/v	2.0% v/v	2.0% v/v	4.0% v/v
1 Gal	0.35 fl oz	1.9 fl oz	2.5 fl oz	2.5 fl oz	5.0 fl oz
5 Gal	1.6 fl oz	9.6 fl oz	12.8 fl oz	12.8 fl oz	25.6 fl oz
25 Gal	8.0 fl oz	47 fl oz	2 qt	2 qt	4 qt

### PREPLANT BURNDOWN

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or with other herbicides or liquid fertilizers as a burndown treatment to control or suppress weeds. Maxunitech Carfentrazone-ethyl 240EC Herbicide is effective as a burndown treatment for crops prior to new plantings. Apply up to 2.0 fl oz Maxunitech Carfentrazone-ethyl 240EC Herbicide (0.031 lb. ai) per acre. **DO NOT** exceed the applicable amounts as listed for the specific crop in the MAXIMUM ALLOWABLE MAXUNITECH CARFENTRAZONE-ETHYL 240EC HERBICIDE USE TABLE 1. For optimum performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across. **Coverage is essential for good control.** Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a labeled burndown herbicides e.g. glyphosate, glufosinate, paraquat, 2, 4-D, or dicamba.

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a burndown treatment no later than one (1) day after planting by seed to any of the following crops. (See specific crop section for other timings)
Alfalfa and Clover (Crop Group 18)
Cereal grains (Crop Group 15)
Grasses (Crop Group 17)
Oil Seed (Crop Group 20 – except cottonseed)
Peanut
Soybean

<b>Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a burndown treatment no later than one (1) day after planting by seed to any of the following crops. (See specific crop section for other timings)</b>
Sugarcane
Vegetables, legume (succulent or dried) (Crop Group 6)
Vegetable, tuberous and corm (Subgroup 1C)
<b>Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a burndown treatment no later than one (1) day before transplanting any of the following crops.</b>
Avocado
Banana
Berry, low growing subgroup 13-07G
Cacao
Coconut
Coffee
Date
Fig
Fruit, citrus (Crop Group 10-10)
Fruit, pome (Crop Group 11-10)
Fruit, stone (Crop Group 12-12)
Globe Artichoke
Guayule
Hops
Horseradish
Indian Mulberry
Kiwifruit
Nuts, Tree (Group 14-12)
Olive
Palm Heart
Persimmon
Pomegranate
Small Fruit Vine, Climbing – except fuzzy kiwifruit (Subgroup 13-07F)
Tea
Tobacco
Vanilla
<b>For transplants (not seeded) of the following crops</b>
Vegetable, brassica (Crop Group 5)
Vegetable, cucurbit (Crop Group 9)
Vegetable, fruiting (Crop Group 8-10)
Vegetable, leafy except Brassica (Crop Group 4)
<b>Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a burndown treatment no later than seven (7) days before planting by seed any of the following crops.</b>
Vegetable, brassica (Crop Group 5)
Vegetable, cucurbit (Crop Group 9)
Vegetable, fruiting (Crop Group 8-10)
Vegetable, leafy except Brassica (Crop Group 4)
Vegetable, tuberous and corm (Crop Subgroups 1C and 1D)
<b>Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a burndown treatment no later than thirty (30) days before planting by seed any of the following crops.</b>
Sugarbeet
Vegetable, bulb (Group 3-07)

#### **Adjuvant Requirements for Preplant Burndown**

A nonionic surfactant crop oil concentrate or methylated seed oil is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient or a petroleum or oil seed based crop oil concentrate (COC) at 1.0 to 2 % v/v (1.0 to 2.0 gallons per 100 gallons of spray solution) or a methylated seed oil (MSO). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons) or ammonium sulfate at 2 to 4 pounds per acre in addition to the selected NIS, MSO or COC is allowed.

#### **Maxunitech Carfentrazone-ethyl 240EC Herbicide Plus Glyphosate or Glufosinate**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at 0.5 to 1.0 fl oz (0.008 to 0.16 lb. ai) per acre in combination with glyphosate or glufosinate products at their labeled rates for increased speed of activity and improved control of weeds e.g. those listed below.



When applied as directed, Maxunitech Carfentrazone-ethyl 240EC Herbicide plus labeled herbicides including glyphosate, glufosinate, or paraquat will provide increased speed of activity and improved control of weeds listed below in Table 6 plus the weeds listed in Table 3 for the rate of Maxunitech Carfentrazone-ethyl 240EC Herbicide used.

**Table 6:**

Buttercup, smallflower	Morningglory spp.
Chickweed	Pennycress, field
Curled Dock	Prostrate knotweed
Cutleaf Evening Primrose	Purslane, common
Bindweed, field	Smartweed, PA
Dandelion, common	Star-of-Bethlehem
*Fleabane	Shepherdspurse
Groundsel	Tansymustard
Henbit	Thistle, Russian
Kochia	Thistles, annual & biennial
Lambsquarters, common	Wild buckwheat
*Marestail	Wild hemp

\*glyphosate susceptible marestail and fleabane

**When tank mixing with fertilizer solutions, be sure to prepare a premixture of Maxunitech Carfentrazone-ethyl 240EC Herbicide and clean water.**

For other specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section.

### **HOODED SPRAYER APPLICATIONS**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide to the row middles of the following emerged crops using hooded sprayers in accordance with specific use information in the following Directions for Use section.

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with hooded sprayers to control labeled weeds between the rows of the below listed emerged crops. This treatment is for crops grown in rows, and includes crops grown in rows where mulch or plastic barriers are used as a weed control tool in the drill or plant line.

**Hooded sprayers must be designed, adjusted and operated in such a manner to totally enclose the spray pattern and to prevent any spray deposition to green stem tissue, foliage, blooms or fruit of the crop.**

Sprayers shall not be operated at more than five (5) miles per hour in order to minimize vertical movement of the sprayer during application, including the bouncing or raising of the equipment. Use extreme care in applying to fields where the soil surface is uneven, has deep furrows, drains or other contours that would disturb the adjustment and positioning of the spray equipment and/or the spray pattern. Applications must not be made when wind conditions may disturb the spray patterns and result in spray deposition to sensitive plants or plant parts.

For optimum performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. **Coverage is essential for good control.**

#### **Crops Labeled for Use with Hooded Sprayers:**

Hooded Spray application can be used for all crops listed on this Maxunitech Carfentrazone-ethyl 240EC Herbicide label

**Note: Crop injury will occur when spray is allowed to come in contact with the green stem tissue, leaves, blooms or fruit of the crop.**

#### **See listing for individual commodities contained within the respective Crop groups:**

Vegetable, Root and Tuber (Group 1) including: Arracacha, Arrowroot, Chinese artichoke, Jerusalem artichoke, Garden Beet, Sugar beet, Edible Burdock, Edible Canna, Carrot; Bitter and Sweet Cassava, Celeriac, Chayote (root), Turnip-rooted Chervil, Chicory, Chufa, Dasheen (taro), Ginger, Ginseng, Horseradish, Leren, Turnip-rooted Parsley, Parsnip, Potato, Radish, Oriental (daikon) Radish, Rutabaga, Salsify, Black Salsify, Spanish Salsify, Skirret, Sweet Potato, Tanier, Turmeric, Turnip, Yam bean; True Yam

Vegetable, leaves of root and tuber (Group 2) including: Garden Beet, Sugar Beet, Edible Burdock, Carrot, Bitter and Sweet Cassava, Celeriac, Chervil, Turnip-rooted, Chicory, Dasheen (taro), Parsnip, Radish, Oriental (daikon) Radish, Rutabaga, Black Salsify, Sweet Potato, Tanier, Turnip, True Yam

Vegetable, bulb (Group 3-07) including: Fresh Leaves Chive, Chinese Fresh Leaves Chive, Bulb Daylily, Elegans Hosta; Bulb Fritillaria, Leaves Fritillaria, Bulb Garlic, Great-headed Garlic, Serpent Bulb Garlic, Kurrat; Lady's Leek, Leek, Wild Leek, Bulb Lily, Beltsville Bunching Onion, Bulb Onion, Chinese Bulb Onion, Fresh Onion, Green Onion, Macrostem onion, Pearl onion, Potato Bulb Onion, Tree Tops Onion, Welsh Tops Onion, Bulb Shallot, Fresh Leaves shallot, and cultivars, varieties, and/or hybrids of these

Vegetable, leafy except brassica (Group 4) including: Amaranth (Chinese Spinach), Arugula (Roquette), Cardoon, Celery, Chinese Celery, Celtuce, Chervil, Edible-Leaved Chrysanthemum, Garland Chrysanthemum, Corn Salad, Cress, Garden, Upland Cress, Dandelion, Dock (Sorrel), Endive (Escarole), Florence Fennel, Head And Leaf Lettuce, Orach, Parsley, Garden Purslane, Winter Purslane, Radicchio (Red Chicory), Rhubarb, Spinach, New Zealand Spinach, Vine Spinach, Swiss Chard

Vegetable, brassica (Group 5) including: Broccoli; Chinese Broccoli, (gai lon), Broccoli Raab (rapini), Brussels Sprouts, Cabbage, Chinese Cabbage, (bok choy); Chinese Cabbage (napa), Chinese Mustard Cabbage (gai choy), Cauliflower, Cavalo Broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens

Vegetable, legume, except soybean (succulent or dried) (Group 6) including: Bean (Lupinus spp.) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (Phaseolus spp.) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); bean (Vigna spp.) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); broad bean (fava); chickpea (garbanzo); guar; jackbean; lablab bean (hyacinth bean); lentil; pea (Pisum spp.) (includes dwarf pea, ediblepodded pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea); pigeon pea; soybean (immature seed); sword bean

Vegetable, foliage of legume (Group 7) including: Plant parts of any legume vegetable included in the legume vegetables group that will be used as animal feed

Vegetable, fruiting (Group 8-10) including: African eggplant, Bush Tomato, Bell Pepper, Cocona, Currant Tomato, Eggplant, Garden Huckleberry, Goji Berry, Groundcherry, Martynia, Naranjilla, Okra, Pea Eggplant, Pepino, Non-Bell Pepper, Roselle, Scarlet Eggplant, Sunberry, Tomatillo, Tomato, Tree Tomato, and cultivars, varieties, and/or hybrids of these

Vegetable, cucurbit (Group 9) including: Chayote (fruit), Chinese Waxgourd (Chinese Preserving Melon), Citron Melon, Cucumber, Gherkin, Edible Gourd (includes Hyotan, Cucuzza, Hechima, Chinese Okra), Momordica spp. (includes Balsam Apple, Balsam Pear, Bittermelon, Chinese Cucumber), Muskmelon (includes Cantaloupe), Pumpkin, Summer Squash, Winter Squash (includes Butternut Squash, Calabaza, Hubbard Squash, Acorn Squash, Spaghetti Squash), Watermelon

Citrus Fruit (Group 10-10) including: Australian Desert Lime, Australian Finger-Lime, Australian Round Lime, Brown River Finger Lime, Calamondin, Citron, Citrus hybrids, Grapefruit, Japanese Summer Grapefruit, Kumquat, Lemon, Lime, Mediterranean Mandarin, Mount White Lime, New Guinea Wild Lime, Sour Orange; Sweet Orange, Pummelo, Russell River Lime, Satsuma Mandarin, Sweet Lime, Tachibana Orange, Tahiti Lime, Tangelo, Tangerine (mandarin), Tangor, Trifoliolate Orange; Uniq Fruit, and cultivars, varieties, and/or hybrids of these

Pome Fruit (Group 11-10) including: Apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these

Stone Fruit (Group 12-12) including: Apricot, Japanese Apricot, Capulin, Black Cherry, Nanking Cherry, Sweet Cherry, Tart Cherry, Chinese Jujube, Nectarine, Peach, Plum, American Plum, Beach Plum, Canada Plum, Cherry Plum, Chickasaw Plum, Damson Plum, Japanese Plum, Klamath Plum, Prune Plum, Plumcot, Sloe and cultivars, varieties, and/or hybrids of these

Caneberry (subgroup 13-07A) including: Blackberry, Loganberry, Black and Red Raspberry, Wild Raspberry, and cultivars, varieties, and/or hybrids of these

Bushberry (subgroup 13-07B) including Aronia Berry, Highbush Blueberry, Lowbush Blueberry, Buffalo Currant, Chilean Guava, Highbush Cranberry, Black Currant, Red Currant, Elderberry, European Barberry, Gooseberry, Edible Honeysuckle, Huckleberry, Jostaberry, Juneberry (Saskatoon Berry), Lingonberry, Native Currant, Salal, Sea Buckthorn and cultivars, varieties, and/or hybrids of these

Fruit, small vine climbing – except fuzzy kiwifruit (subgroup 13-07F) including: Amur River Grape, Gooseberry, Grape, Hardy Kiwifruit, Maypop, Schisandra Berry and cultivars, varieties, and/or hybrids of these

Berry, low growing (subgroup 13-07G) including: Bearberry, Bilberry, Lowbush Blueberry, Cloudberry, Cranberry, Lingonberry, Muntries, Partridgeberry, Strawberry, and cultivars, varieties, and/or hybrids of these

Tree Nuts (Group 14-12) including: African Nut-Tree, Almond, Beechnut, Brazil Nut; Brazilian Pine, Bunya, Bur Oak, Butternut, Cajou Nut, Candlenut, Cashew, Chestnut, Chinquapin, Coconut, Coquito nut, Dika Nut, Ginkgo, Guiana

Chestnut, Hazelnut (filbert); Heartnut, Hickory Nut, Japanese Horse-Chestnut, Macadamia Nut, Mongongo Nut, Monkey-Pot, Monkey Puzzle Nut, Okari Nut, Pachira Nut, Peach Palm Nut, Pecan, Pequi, Pili Nut, Pine Nut, Pistachio, Sapucaia Nut, Tropical Almond, Black Walnut, English Walnut, Yellowhorn and cultivars, varieties, and/or hybrids of these

Cereal Grains (Group 15) including: Barley, Buckwheat, Corn, Millet (Pearl and proso), Oats, Popcorn, Rice, Rye, Sorghum (milo), Teosinte, Triticale, Wheat, and Wild Rice

Forage, fodder and straw of Cereal Grains (Group 16) including forage fodder and straw of all commodities included in the cereal grains (Group 15)

Grasses (Group 17) including: Any grass, Gramineae family (either green or cured) except sugarcane and those included in the cereal grains group, that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage

Non-grass Animal Feed (Group 18) including: Alfalfa, Velvet Bean, Clover (*Trifolium* spp., *Melilotus* spp.), Kudzu, Lespedeza, Lupin, Sainfoin, Trefoil, Vetch, Crown Vetch, Milk Vetch

Herbs and Spices (Group 19) including: Allspice, Angelica, Anise (seed), Star Anise, Annatto (seed), Balm (Lemon Balm), Basil (Fresh and Dried), Borage, Burnet, Camomile, Caper Buds, Caraway, Black Caraway, Cardamom, Cassia Bark, Cassia Buds, Catnip, Celery Seed, Chervil (dried), Chive, Chinese Chive, Cinnamon, Clary, Clove Buds, Coriander Leaf (Cilantro or Chinese Parsley), Coriander Seed (Cilantro), Costmary, Culantro (Leaf), Culantro (Seed), Cumin, Curry (Leaf), Dill (Dillweed), Dill (Seed), Fennel (Common), Florence Fennel (seed), Fenugreek, Grains of Paradise, Horehound, Hyssop, Juniper Berry, Lavender, Lemongrass, Lovage (leaf), Lovage (seed), Mace, Marigold, Marjoram (includes Sweet or Annual Marjoram, Wild Marjoram or Oregano, and Pot Marjoram), Mustard (Seed), Nasturtium, Nutmeg, Parsley (Dried), Pennyroyal, Black Pepper, White Pepper, Poppy (Seed), Rosemary, Rue, Saffron, Sage, Summer and Winter Savory, Sweet Bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood

Oil Seeds (Subgroups 20A & 20B, except Cottonseed) including: Borage, Crambe, Cuphea, Echium, Flax Seed, Gold of Pleasure, Hare's Ear Mustard, Lesquerella, Lunaria, Meadowfoam, Milkweed, Mustard Seed, Oil Radish, Poppy Seed, Rapeseed, Sesame, Sweet Rocket, Calendula, Castor Oil Plant, Chinese Tallowtree, Euphorbia, Evening Primrose, Jojoba, Niger Seed, Rose Hip, Safflower, Stokes Aster, Sunflower, Tallowwood, Tea Oil Plant, Vernonia and cultivars, varieties, and/or hybrids of these

Tropical fruit: including Acerola, Atemoya, Avocado, Biriba, Black Sapote, Canistel, Cherimoya, Custard apple, Feijoa, Guava, Jaboticaba, Llama, Longan, Lychee, Mamey Sapote, Mango, Papaya, Passionfruit, Pawpaw, Pulasan, Rambutan, Sapodilla, Soursop Spanish lime, Star apple, Starfruit, Sugar apple, Wax jambu, Aloe vera, Cactus

## APPLICATION INSTRUCTIONS

### ALFALFA AND CLOVER (Established Stands Only) Crop group 18

Methods and Timing	Target Weeds	Application Rates
<b>Postemergence Weed Control (Dormant, In-crop, and Stubble)</b>	Refer to table 3	0.5-2.5 fl oz (0.008 – 0.04 lb. ai) per acre
<b>Harvest Aid</b>	Refer to table 3	2.0 to 3.8 fl oz/A (0.03 – 0.06 lb. ai) per acre

#### DIRECTIONS FOR USE:

##### Postemergence Weed Control Treatment

##### Dormant Season (Fall or Winter Application Postemergence on Weeds)

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be applied on dormant crop stubble alone or in combination with other registered herbicides for the post emergence control of weeds in established nongrass animal feed stands during the dormant season (between growing seasons). To control insect pests, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with insecticides, including Mustang Max.

##### Between Cutting In-Season Application (Spring/Summer Applications Postemergence on Weeds)

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be applied alone or in combination with other registered herbicides between cuttings (in-season) for the post emergence control of weeds in established crop stands. In-

season applications should be made as soon as possible after removal of the previous hay crop and prior to significant regrowth on stems and crowns. Applications may be made from hay removal up to 6 inches of new growth. To control insect pests, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with insecticides, including Mustang Maxx.

#### **Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rates**

For optimum results, weeds should be treated when small. Applications should be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 10 gallons of finished spray per acre for ground application equipment, and a minimum of 3 gallons per acre of finished spray for aerial equipment. For optimum results, apply Maxunitech Carfentrazone-ethyl 240EC Herbicide to weeds up to 4 inches tall and rosettes less than 3 inches across. Use a quality nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. For more active treatments, use a Crop Oil Concentrate (COC) at 0.5 to 1.0% v/v (one half to one gallon per 100 gallons). Some temporary leaf speckling and necrosis may occur on green alfalfa or clover tissue present with between cutting applications, which should be rapidly outgrown under good growing conditions. Adjuvant selection and high moisture environmental conditions will enhance this effect. A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed. Coverage is essential for satisfactory performance. Repeat application if necessary. **DO NOT** irrigate just prior to or just after application. Weed control under dry and hot conditions will be improved with COC or similar products.

#### **Tank Mix**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with other labeled herbicides to control weeds not listed on this label. Read and follow all manufacturers' label directions and label restrictions for the companion herbicide. When tank mixing Maxunitech Carfentrazone-ethyl 240EC Herbicide with other products, be sure Maxunitech Carfentrazone-ethyl 240EC Herbicide is mixed in the spray tank water first.

#### **Harvest Aid Treatment**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide to crops grown for forage, hay or seed alone or as a tank mixture with other harvest aids. Applications shall be made when the crop is mature, or according to Extension Service guidelines in the use area. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at 2.0 to 3.8 fl oz (0.031 to 0.06 lb. ai) per acre, but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE MAXUNITECH CARFENTRAZONE-ETHYL 240EC HERBICIDE USE RATE CHART and the PREHARVEST INTERVAL charts for additional application information. If treatments of Maxunitech Carfentrazone-ethyl 240EC Herbicide have been made to the crop earlier, that volume must be considered in determining the maximum use rate as a harvest aid treatment

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 10 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed. Coverage is essential for satisfactory performance. Repeat application if necessary.

#### **RESTRICTIONS:**

**DO NOT** apply more than 6.3 fl oz (0.1 lb. ai) per acre per year inclusive of all applications.

**DO NOT** make more than 4 applications per year inclusive of Postemergence Weed Control and Harvest Aid applications.

**DO NOT** apply more than 2.5 fl oz (0.04 lb. ai) per acre per application for Postemergence Weed Control applications.

**DO NOT** apply more than 2.5 fl oz (0.04 lb. ai) per acre per year for Postemergence Weed Control applications in nongrass animal feeds.

**DO NOT** apply more than 3.8 fl oz (0.06 lb. ai) per acre per application for Harvest Aid applications.

**DO NOT** apply more than 3.8 fl oz (0.06 lb. ai) per acre per year for Harvest Aid applications.

**DO NOT** make more than 2 Postemergence Weed Control applications per year at reduced rates.

**DO NOT** make more than 2 Harvest Aid applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply within 21 days of harvest for stands grown for forage and hay.

**DO NOT** apply within 3 days of harvest for stands grown for seed as a Harvest Aid treatment.

After an application of this product to crop group 18 (nongrass animal feed crops), you may only rotate the field to a carfentrazone-ethyl registered crop.

**Note**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**ASPARAGUS**

Methods and Timing	Target Weeds	Application Rates
Postemergence Weed Control	Refer to table 3	Apply one to two applications of Maxunitech Carfentrazone-ethyl 240EC Herbicide at 0.5 to 1.92 fl oz (0.008 to 0.03 lb. ai) per acre.  Use higher rates when Asparagus tissues and weeds are under stress or are larger.

**DIRECTIONS FOR USE:**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a broadcast application after harvest of Asparagus spears for control of broadleaf weeds and new existing Asparagus tissues.

**Coverage is essential for good control.**

**Adjuvant Requirements**

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 10 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed. Repeat application if necessary.

For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section.

**RESTRICTIONS:**

**DO NOT** apply more than 1.92 fl oz (0.03 lb. ai) per acre per application.

**DO NOT** apply more than 3.8 fl oz (0.06 lb ai) per acre per year.

**DO NOT** make more than 2 applications per year.

**DO NOT** make applications less than 20 days apart.

**DO NOT** apply within 5 days of harvest.

## BUSHBERRY Subgroup 13-07B

Methods and Timing	Target Weeds	Application Rates
Postemergence Weed Control	Refer to table 3	Up to 2.0 fl oz (0.031 lb. ai) per acre.
<p><b>DIRECTIONS FOR USE</b></p> <p>Maxunitech Carfentrazone-ethyl 240EC Herbicide applications will control susceptible emerged broadleaf weeds. Repeat applications may be necessary for weeds that emerge after an Maxunitech Carfentrazone-ethyl 240EC Herbicide treatment.</p> <p><b>Equipment and Application</b> Apply only by ground equipment including boom sprayers, shielded or hooded sprayers, hand-held or high-volume wands or orchard guns. Use a minimum of 20 gallons finished spray solution per broadcast acre.</p> <p><b>Dormant Applications</b> Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a broadcast application to the base of the trunk to control emerged and actively growing weeds during the dormant stage of the crop.</p> <p><b>Post-directed Applications for Broadleaf Weed Control</b> Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a directed spray avoiding contact with the berry plant but directed at actively growing weeds. Maxunitech Carfentrazone-ethyl 240EC Herbicide is a contact herbicide and coverage is essential for good weed control. <b>DO NOT</b> allow Maxunitech Carfentrazone-ethyl 240EC Herbicide spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage.</p> <p>Newly planted bush berries should only be treated with shielded sprayers or hooded sprayers.</p> <p><b>Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rates</b> Apply up to 2 fl oz (0.031 lb. ai) Maxunitech Carfentrazone-ethyl 240EC Herbicide per broadcast acre. For best control, apply to seedling weeds in the 2 to 3-leaf stage. Use higher labeled rates of Maxunitech Carfentrazone-ethyl 240EC Herbicide for larger weeds up to 6 leaves. Weeds greater than 6 leaves may be only partially controlled. See Table 3 for Maxunitech Carfentrazone-ethyl 240EC Herbicide use rates and weeds controlled.</p> <p><b>Adjuvant Requirements</b> A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.</p> <p><b>Tank Mix</b> Maxunitech Carfentrazone-ethyl 240EC Herbicide may be mixed with other registered herbicides for broader spectrum weed control. When tank mixing with fertilizer solutions, be sure to prepare a premixture of Maxunitech Carfentrazone-ethyl 240EC Herbicide and clean water.</p> <p>See Mixing and Loading Instructions under the PRODUCT INFORMATION section of this label for specific mixing instructions. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p><b>Precautions</b> Extreme caution must be taken during applications when desirable fruit, foliage and/or blooms are present in order to avoid spotting or necrosis. <b>DO NOT</b> allow Maxunitech Carfentrazone-ethyl 240EC Herbicide spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage.</p> <p>For seedling or newly transplanted bushes, <b>DO NOT</b> allow spray to contact green bark of trunk area. Use shielded sprayers only.</p> <p><b>Band Treatment Application</b> For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:</p> $\frac{\text{Band Width}}{\text{Row Width}} \times \text{Broadcast Rate Per Acre} = \text{Band Rate}$ <p>Inches Inches</p>		

Band Width Inches	X	Broadcast Volume Per Acre	=	Band Volume
Row Width Inches				
<b>RESTRICTIONS:</b>				
<b>DO NOT</b> apply more than 2 fl oz (0.031 lb. ai) per acre per application.				
<b>DO NOT</b> apply more than 6.1 fl oz (0.096 lb. ai) per acre per year.				
<b>DO NOT</b> apply more than 2 fl oz (0.031 lb. ai) during the dormant season.				
<b>DO NOT</b> make more than 4 applications per year at reduced rates.				
<b>DO NOT</b> make applications less than 14 days apart.				
Can be applied up to harvest.				

### CANEBERRY Subgroup 13-07A

Methods and Timing	Target Weeds	Application Rates
<b>Postemergence Weed Control</b>	Refer to table 3	Apply 6.4 fl oz Maxunitech Carfentrazone-ethyl 240EC Herbicide (0.1 lb. ai) per broadcast acre as a directed spray when weeds and primocanes are approximately 6 inches tall.  Apply up to 2 fl oz (0.031 lb. ai) Maxunitech Carfentrazone-ethyl 240EC Herbicide per broadcast acre. For best control, apply to actively growing weeds up to 4 inches tall or rosettes less than 3 inches across.

#### DIRECTIONS FOR USE:

##### Equipment and Application

Apply only by ground equipment including boom sprayers, shielded or hooded sprayers, hand-held or high-volume wands or orchard guns. **DO NOT** allow Maxunitech Carfentrazone-ethyl 240EC Herbicide spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage. **DO NOT** apply when conditions favor drift or when wind is above 10 mph.

##### Post-Directed Application for Primocane and Weed Control

Maxunitech Carfentrazone-ethyl 240EC Herbicide is a contact herbicide for directed application for the control of primocanes and weeds.

Use a minimum of 20 gallons finished spray per broadcast acre at intervals of 14 to 21 days. Direct spray to the bottom 18 inches of the canes and to the soil 24 inches from each side of the plant row. Refer to weed control list in Table 3 for appropriate weed control information.

##### Adjuvant Requirements

An adjuvant is required. See Adjuvant Requirements below under weed control.

##### Post-directed Application for Weed Control

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide to actively growing weeds. Maxunitech Carfentrazone-ethyl 240EC Herbicide is a contact herbicide and coverage is essential for good weed control. Use a minimum of 20 gallons finished spray solution per acre.

##### Band Treatment Application

For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Rate Per Acre} = \text{Band Rate}$$

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Volume Per Acre} = \text{Band Volume}$$

**Coverage is essential for good control.**

**Adjuvant Requirements**

A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be mixed with other herbicides registered in caneberries for broader spectrum weed control.

Maxunitech Carfentrazone-ethyl 240EC Herbicide should be the first product added to the spray tank water. See Mixing and Loading Instructions under the PRODUCT INFORMATION section of this label for specific mixing instructions. Refer to this and the other product's labels for mixing instructions, precautions, and restrictions. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Precautions**

Extreme caution must be taken during applications when desirable fruit, foliage and/or blooms are present in order to avoid spotting or necrosis. **DO NOT** allow Maxunitech Carfentrazone-ethyl 240EC Herbicide spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage.

Newly planted caneberries should only be treated with shielded sprayers or hooded sprayers.

**RESTRICTIONS:**

**DO NOT** apply more than 6.4 fl oz (0.1 lb. ai) per acre per application as a directed spray when weeds and promocanes are approximately 6 inches tall.

**DO NOT** apply more than 2 fl oz (0.031 lb. ai) per acre per application to actively growing weeds up to 4 inches tall or rosettes less than 3 inches across.

**DO NOT** make more than 5 applications per year at reduced rates.

**DO NOT** apply more than 25.6 fl oz (0.4 lb. ai)per acre per year.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply within 15 days of harvest.



**CORN (Field, Seed, Silage, Popcorn, Sweet Corn – Processing and Fresh Market)**

Methods and Timing	Target Weeds	Application Rates
Preplant Burndown	Refer to table 3	Up to 2.0 fl oz (0.031 lb. ai) per acre
Postemergence (Broadcast)	Refer to table 3	Up to 1.0 fl oz (0.016 lb. ai) per acre
Postemergence (Hooded Sprayer and Directed Applications)	Refer to table 3	Up to 2.0 fl oz (0.031 lb. ai) per acre
Harvest Aid	Refer to table 3	1.0 to 2.0 fl oz (0.016 – 0.031 lb. ai) per acre

**Directions for Use:**

**Postemergence Weed Control Treatment**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to corn in all tillage systems from prior to planting up to 14-leaf collar growth stage. When applying Maxunitech Carfentrazone-ethyl 240EC Herbicide to corn greater than V8 stage, utilize drop nozzles aligned between the rows with directed application to reduce contact with the corn foliage and improve contact with the weeds. For optimum performance, make application to actively growing weeds up to 4 inches high and rosettes less than 3 inches across. **Coverage is essential for good control.**

**Broadcast Applications:**

Use Maxunitech Carfentrazone-ethyl 240EC Herbicide at 0.5 to 1.0 fl oz (0.008 – 0.016 lb. ai) as a broadcast application using a minimum of 10 gallons per acre of spray volume by ground or 3 gallons per acre by air. Broadcast applications may be applied through V8 stage corn.

**Tank Mix**

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank-mixed with other corn herbicides to control weeds not listed on this label. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Maxunitech Carfentrazone-ethyl 240EC Herbicide plus Atrazine**

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed at a rate of 0.5 fl oz (0.008 lb. ai) per acre with Atrazine 4L (16 fl oz per acre) or Atrazine 90DF (0.6 -1.6 lbs. per acre) to control the following weeds:

**When used as directed, Maxunitech Carfentrazone-ethyl 240EC Herbicide + atrazine will provide control of listed weeds up to 4 inches tall.**

Amaranth, Palmer (not triazine resistant)	Copperleaf, hophornbeam	Mallow, Venice	Purslane, common
Amaranth, spiny	Croton, wooly	Morningglory spp.	Sesbania, hemp
Anoda, spurred	Devilsclaw	Nightshade, Eastern black	Thistle, Russian
Buckwheat, wild	Eveningprimrose, cutleaf	Nightshade, hairy	Velvetleaf
Buffalobur	Jimsonweed	Pigweed, redroot	Waterhemp, common
Carpetweed	Kochia *	Pigweed, smooth	Waterhemp, tall
Cocklebur	Lambsquarters, common	Potato, volunteer	

\* Kochia control up to 2 inches tall with Maxunitech Carfentrazone-ethyl 240EC Herbicide + Atrazine + COC only. Refer to the Atrazine labels for additional weed listings and for higher use rates.

**Maxunitech Carfentrazone-ethyl 240EC Herbicide plus Dicamba**

Maxunitech Carfentrazone-ethyl 240EC Herbicide at 0.5 fl oz (0.008 lb. ai) per acre plus 0.25% v/v nonionic surfactant (2 pints per 100 gallons) can be tank mixed with dicamba herbicides (8 -16 fluid ounces per acre) for control of broadleaf weeds including the following:

**When used as directed, Maxunitech Carfentrazone-ethyl 240EC Herbicide + dicamba will provide control of listed weeds up to 4 inches tall.**

Buckwheat, wild	Morningglory spp.	Potato, volunteer	Thistle, Russian
Cocklebur, common	Nightshade, black	Ragweed, common	Velvetleaf
Jimsonweed	Pigweed, redroot	Ragweed, giant	Waterhemp, common
Kochia	Pigweed, smooth	Smartweed, PA (seedling)	Waterhemp, tall
Lambsquarters	Pigweed, Triazine resistant	Sunflower, common	

Refer to the dicamba labels for additional weed listings and for higher use rates.

Refer to the Tank Mixture Section for information on potential leaf injury.

**Maxunitech Carfentrazone-ethyl 240EC Herbicide Plus Atrazine Plus Dicamba or 2,4-D**

For the control of additional or certain larger weeds up to 6 inches tall, Atrazine may be added to the tank mixtures of Maxunitech Carfentrazone-ethyl 240EC Herbicide plus dicamba or Maxunitech Carfentrazone-ethyl 240EC Herbicide plus 2,4-D (amine).

Add 2,4-D (amine) to the tank mix at 0.125 to 0.25 lb. ai per acre or dicamba at 3 to 8 fluid ounces per acre. Higher rates of atrazine and dicamba herbicides are allowed, but **DO NOT** exceed the specific label use rates allowed by these labels. Add a 0.25% v/v nonionic surfactant (2 pints per 100 gallons) to the tank mixture. Under very dry soil moisture conditions, the use of crop oil concentrate at 1% v/v (1 gallon per 100-gallon spray solution) may improve weed control. The use of crop oil concentrate may increase leaf speckling. Refer to the Tank Mixture section for information on potential leaf injury.

**For control of the following weeds up to 6 inches in height, or as specified, add dicamba at 3 to 8 ounces per acre to Maxunitech Carfentrazone-ethyl 240EC Herbicide tank mixes with atrazine or to Maxunitech Carfentrazone-ethyl 240EC Herbicide tank mixes with other products that allow the use of dicamba on their labels.**

Amaranth, Palmer (up to 4 inches)	Nightshade, Eastern black	Smartweeds, annual (seedling)
Amaranth, spiny (up to 4 inches)	Nightshade, hairy	Sunflower, common (up to 4 inches tall)
Cocklebur, common	Pigweed, redroot	Velvetleaf (up to 24 inches)
Kochia (up to 4 inches)	Pigweed, smooth	Waterhemp, common
Lambsquarters, common	Ragweed, common	Waterhemp, tall
Morningglory spp.	Ragweed, giant (up to 4 inches tall)	

**Adjuvant Requirements:**

Use a non-ionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution). Under dry conditions, the use of a crop oil concentrate (COC) at 1.0% v/v may improve weed control. The use of crop oil concentrate can increase leaf speckling and crop response on treated corn leaves.

**Directed Spray Applications:**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with drop nozzles between the rows to the target weeds and away from the whorl of the corn plant. Directed spray applications should be used when corn is V8 to V14 stage. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide up to 2.0 fl oz (0.031 lb. ai) per acre. Be aware that weeds growing in and under the dense canopies may not receive adequate spray coverage and may require the use of higher spray volumes for acceptable control. Use appropriate rates of adjuvants including non-ionic surfactant (NIS), crop oil concentrate (COC), or methylated seed oil (MSO).

**Hooded Sprayer Applications:**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

**Harvest Aid:**

Apply 1.0 to 2 fl oz (0.016 – 0.032 lb. ai) Maxunitech Carfentrazone-ethyl 240EC Herbicide per acre, but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE MAXUNITECH CARFENTRAZONE-ETHYL 240EC HERBICIDE USE RATE and the PREHARVEST INTERVAL Table (Table 2) for additional application information. If treatments of Maxunitech Carfentrazone-ethyl 240EC Herbicide have been made to the crop earlier, that volume must be considered in determining the maximum use rate as a harvest aid treatment

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 15 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. A methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use methylated seed oil, or crop oil concentrate (COC) (petroleum or seed oil) at 1 to 2% v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the methylated seed oil or crop oil is allowed.

**Coverage is essential for satisfactory performance****Seed Corn Production:**

For seed production fields, apply Maxunitech Carfentrazone-ethyl 240EC Herbicide using drop nozzles or other equipment to make a directed spray treatment. Avoid directing spray solution into the whorl.

Seed corn inbred lines have shown good tolerance to Maxunitech Carfentrazone-ethyl 240EC Herbicide. However, all inbred lines have not been tested. Broadcast applications may result in spray being concentrated into the whorl of the plant that will increase leaf response. To minimize application into the whorl of the plants, drop nozzles or other type directed sprayers must be used to direct the spray to the targeted weeds.

**Sweet Corn Precaution:**

When applying Maxunitech Carfentrazone-ethyl 240EC Herbicide to sweet corn; broadcast applications may result in spray being concentrated into the whorl of the plant that will increase leaf response. To minimize application into the whorl of the plants, drop nozzles or other type directed sprayers must be used to direct the spray to the targeted weeds.

Use only NIS as the spray adjuvant in sweet corn applications.

**Application Precautions:**

Leaf speckling can occur when Maxunitech Carfentrazone-ethyl 240EC Herbicide is used with certain crop protection products and adjuvants. Refer to the Tank Mixtures and Adjuvants requirements sections under PRODUCT INFORMATION. Bromoxynil mixtures and bentazon mixtures may cause significant crop response when in contact with crop foliage.

**Crop Response**

The application of Maxunitech Carfentrazone-ethyl 240EC Herbicide to corn may result in temporary crop response, for example, speckling or necrosis of the leaves. Grain yields will not be affected. **DO NOT** make applications when air temperatures are abnormally cool or humidity is high or if the corn foliage is wet from dew, rainfall or irrigation. Users should be aware of these inherent risks and accept these risks prior to application of Maxunitech Carfentrazone-ethyl 240EC Herbicide.

For additional information regarding potential crop response, refer to the PRODUCT INFORMATION section of the Maxunitech Carfentrazone-ethyl 240EC Herbicide label.

**Coverage is essential for satisfactory performance.**

**RESTRICTIONS:**

**DO NOT** apply more than 2 fl oz (0.031 lb. ai) per acre per application for Preplant Burnout, Postemergence (Hooded Sprayer and Directed Applications) and Harvest Aid.

**DO NOT** apply more than 1 fl oz (0.016 lb. ai) per acre per application for Postemergence (Broadcast).

**DO NOT** apply more than 2.0 fl oz. (0.031 lb. ai) per acre per year including all preplant, in-crop, and harvest aid applications.

**DO NOT** make more than 3 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply after 14 Leaf Collar for Postemergence applications.

**DO NOT** apply within 3 days of harvest for Harvest Aid treatments.

**COTTON**

Methods and Timing	Target Weeds	Application Rates
Removal of Failed Cotton Stands	Failed Cotton (up to 3 leaf cotton)	1.0 to 1.6 fl oz (0.016 – 0.025 lb. ai) per acre
Pre Plant Burndown	Refer to table 3	Up to 1.6 fl oz (0.025 lb. ai) per acre
Hooded Sprayer	Refer to table 3	Up to 1.6 fl oz (0.025 lb. ai) per acre
Post-directed and Lay-by	Refer to table 3	Up to 1.6 fl oz (0.025 lb. ai) per acre
Defoliation/Harvest Aid	Defoliate and desiccate cotton and troublesome weeds	Up to 1.6 fl oz (0.025 lb. ai) per acre

**DIRECTIONS FOR USE:****Removal of Failed Cotton Stands**

Apply 1.0 to 1.6 fl oz Maxunitech Carfentrazone-ethyl 240EC Herbicide (0.016 to 0.025 lb. ai) per acre broadcast as a foliar spray over the top of the remaining cotton plants with sufficient spray volume to provide adequate coverage of the cotton plant, particularly the terminal area. Use higher rates on larger failed cotton. For best results **DO NOT** exceed 3 leaf cotton. **Coverage is essential for good control.**

**PREPLANT BURNDOWN**

See instructions under the Preplant Burndown section of this label.

### **Hooded Sprayer Applications**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

### **Post-directed and Lay-by Applications**

Maxunitech Carfentrazone-ethyl 240EC Herbicide is a contact herbicide for postemergence directed sprayer or hooded/shielded sprayer applications for the control of broadleaf weeds in cotton. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other herbicides to emerged and actively growing weeds. For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section. Applications of Maxunitech Carfentrazone-ethyl 240EC Herbicide or Maxunitech Carfentrazone-ethyl 240EC Herbicide tank mixes must be made with directed sprayers or hooded sprayers to prevent contact of spray solution with the cotton plant. **DO NOT** allow spray solution to contact cotton foliage, green stem tissue, or blooms. Directed spray equipment must position nozzles a minimum 3 to 4 inches above the soil, with nozzles directed beneath the crop canopy. Maxunitech Carfentrazone-ethyl 240EC Herbicide or Maxunitech Carfentrazone-ethyl 240EC Herbicide tank mix applications shall be made to cotton that is a minimum of 6 inches in height. Applications to cotton at 5 to 6 nodes or less must be made with hooded or shielded sprayer equipment to completely avoid contact with cotton plants. Apply lay-by applications of Maxunitech Carfentrazone-ethyl 240EC Herbicide or Maxunitech Carfentrazone-ethyl 240EC Herbicide tank mixtures at later growth stages of cotton when cotton plants have achieved a height of 12 inches or more with sufficient bark development and height differential between crop bottom leaves and the soil. Spray solution shall be directed at the base of cotton plants for minimal contact with green stem tissue or foliage while maintaining maximum contact with broadleaf weeds that are at appropriate treatment size.

For optimum performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. **Coverage is essential for good control.**

### **Adjuvant Requirements**

A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2% v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.

### **Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rates and Weeds Controlled**

Apply up to 1.6 fl oz (0.025 lb. ai/A) Maxunitech Carfentrazone-ethyl 240EC Herbicide as a post-directed treatment using a directed sprayer a hooded sprayer or lay-by sprayer delivering a minimum finished spray volume of 10 gallons per acre. **DO NOT** apply more than 3.2 fl oz (0.05 lb.ai) Maxunitech Carfentrazone-ethyl 240EC Herbicide per year by post-directed and lay-by applications. Refer to weed control list in Table 3 for appropriate weed control information.

For control of additional broadleaf weeds and grasses, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with other herbicides registered for cotton post-directed and/or lay-by applications. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions and rotational cropping restrictions.

### **Defoliation / Harvest Aid Application**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a harvest aid to defoliate and desiccate cotton and troublesome weeds that may be present at harvest. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other cotton harvest aids.

Use a quality spray adjuvant e.g. nonionic surfactant (NIS) or crop oil concentrate (COC) at the directed rates. Use the adjuvant, NIS during warmer periods with COC being the better choice for applications during cooler periods. Make application when 60 to 70 percent of the bolls are open, or according to the State Agricultural Extension Service guidelines in the use area.

Apply up to 1.6 fl oz Maxunitech Carfentrazone-ethyl 240EC Herbicide (up to 0.025 lb. ai per acre) in spray volume sufficient to provide complete coverage of cotton foliage. Use a minimum of 10 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. **Coverage is essential for good defoliation.** Repeat application if necessary to remove remaining foliage. **DO NOT** apply more than 3.2 fl oz (0.05 lb. ai) per acre total as a harvest aid. Dense cotton canopy, large plant size, and environmental conditions not conducive to complete plant coverage may reduce initial application performance and increase the need for a second application.

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone, as a tank mix, or as a sequential application alone or tank mixed with other registered cotton harvest aid products.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Restrictions:**

**DO NOT** apply more than 1.6 fl oz (0.025 lb. ai) per acre per application.

**DO NOT** apply more than 7.9 fl oz (0.124 lb. ai) per acre per year for preplant, in-season weed control and harvest aid.

**DO NOT** apply more than 3.2 fl oz (0.05 lb. ai) per acre per year for Harvest Aid applications.

**DO NOT** make more than 5 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply within 7 days of harvest.

**DRIED SHELLED BEANS, PEAS Crop subgroup 6-C  
FLAX (except soybean) and VEGETABLE FOLIAGE OF LEGUME Crop group 7.**

Methods and Timing	Target Weeds	Application Rates
Preplant Burndown	Refer to table 3	Up to 2.0 fl oz (0.031 lb. ai) per acre.
Harvest Aid Applications	Refer to table 3	1.0 to 6.1 fl oz (0.016 to 0.096 lb. ai) per acre.

**Directions for Use:**

**Preplant Burndown:**

Refer to the preplant burn down section of this label.

**Harvest AID Treatment:**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a harvest aid to dry beans and dry peas at maturity when 80 to 90% of seed pods are yellow or buck skin in color and only 30% of green leaves remain on the plant. Apply to flax when 75% of the bolls have turned brown. Thorough coverage is essential for harvest aid and multiple applications may be needed. For optimum performance use 15 to 30 gallons per acre finished sprayed with a methylated seed oil (MSO) type adjuvant to ensure thorough coverage and retention for harvest aid.

**Maxunitech Carfentrazone-ethyl 240EC Herbicide Use rates:**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other harvest aids. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at 1.0 to 6.1 fl oz (0.016 to 0.096 lb. ai) per acre, but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE MAXUNITECH CARFENTRAZONE-ETHYL 240EC HERBICIDE USE RATE CHART and the PREHARVEST INTERVAL charts for additional application information.

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 15 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. A methylated seed oil (MSO) or crop oil concentrate (COC) is required at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). The addition of a high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the methylated seed oil or crop oil may enhance performance. If spraying dry beans before full maturity and pods are not all mature and turning color, a repeat application may be necessary.

**Restrictions:**

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per application for Preplant Burnout.

**DO NOT** apply more than 6.1 fl oz (0.096 lb. ai) per acre per application for Harvest Aid.

**DO NOT** apply more than 6.1 fl oz (0.096 lb. ai) per acre per year.

**DO NOT** make more than 4 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

Can be applied up to 0 days before harvest as a Harvest Aid application. **Note**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**FRUIT, SMALL VINE CLIMBING (except fuzzy kiwifruit) Crop subgroup 13-07F**

Methods and Timing	Target Weeds	Application Rates
<b>Postemergence Weed Control</b>	Refer to table 3	Up to 2.0 fl oz (0.031 lb. ai) per acre.

**DIRECTIONS FOR USE**

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be applied for postemergence weed control or for sucker control.

**Weed Control**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other herbicides as a postemergence directed spray treatment or as a hooded spray treatment to control emerged and actively growing weeds. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide to middles (between rows of plants) and in strips (in row of plants).

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at any time during the year (see precautions). Maxunitech Carfentrazone-ethyl 240EC Herbicide may be mixed with other herbicides that have pre-emergence or post-emergence activity. Any preemergence activity must rely on activity from other herbicides as directed on their labels.

Herbicides including glyphosate may be tank mixed with Maxunitech Carfentrazone-ethyl 240EC Herbicide for broader spectrum weed control. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Sucker Management**

Maxunitech Carfentrazone-ethyl 240EC Herbicide is effective as an aid in the management of undesirable sucker growth from the base of vine trunks or root sprouts. Suckers and other undesirable growth must be treated when the tissue is young and not mature and/or hardened off. Care must be taken not to allow spray mist to contact desirable fruit or foliage or green stem tissue (see precautions). Application of Maxunitech Carfentrazone-ethyl 240EC Herbicide with other sucker control herbicides is allowed.

**Hooded Sprayer Applications**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

**Equipment and Application**

Coverage is essential for good control. Use a spray volume adequate to obtain thorough coverage with a minimum of 10 gallons of finished spray per acre. Apply only with ground equipment. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with hooded sprayers, boom equipment, shielded sprayers, hand-held and high-volume wands or orchard guns. Always add Maxunitech Carfentrazone-ethyl 240EC Herbicide to the spray tank first. See "Mixing and Loading Instructions" under PRODUCT INFORMATION.

**Adjuvant Requirements**

Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a quality nonionic surfactant (NIS) containing at least 80% active at 0.25% v/v (2 pints NIS per 100 gallons) or a crop oil concentrate (COC) at 1% v/v (one-gallon COC per 100 gallons), or a methylated seed oil (MSO). The use of a high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v or ammonium sulfate (AMS) used at 2 to 4 pounds per acre in addition to the NIS, or MSO or COC is allowed.

**Application Precautions: Extreme caution must be used during applications when desirable fruit or foliage is present in order to avoid fruit spotting or leaf necrosis.**

**RESTRICTIONS:**

**DO NOT** allow Maxunitech Carfentrazone-ethyl 240EC Herbicide spray mist to come in contact with desirable fruit, green stem tissue, foliage or blooms.

**DO NOT** use on seedling or newly transplanted vines

**DO NOT** allow spray to contact green bark of trunk area.

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per application (including preplant site preparation treatments).

**DO NOT** apply more than 7.9 fl oz (0.124 lb. ai) per acre per year.

**DO NOT** make more than 5 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply within 3 days of harvest.

**FRUIT TREE**

**TREE NUT**

**OTHER CROPS** (banana, cacao, coconut, coffee, date, fig, guayule, Indian mulberry, olive, palm heart, persimmon, pomegranate, tea, vanilla)

<b>CROP GROUP</b>	<b>Target Weeds</b>	<b>Application Rates</b>
<b>Citrus Fruits</b> including Calamondin, Citrus Citron, Chironja, Tangelo, Tangor, Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Orange (sour), Orange (Sweet), Pummelo, Satsuma and Mandarin	Refer to table 3	Apply up to 2.0 fl oz (0.031 lb. ai) per acre.
<b>Pome Fruits:</b> including Apple, Crabapple, Loquat, Mayhaw, Pear, Pear (Oriental) and Quince	Refer to table 3	Apply up to 2.0 fl oz (0.031 lb. ai) per acre.
<b>Stone Fruits:</b> including Apricot, Cherry (Sweet), Cherry (Tart), Nectarine, Peach, Plum, Plum (Chickasaw), Plum (Damson), Plum (Japanese), Prune and Plumcot	Refer to table 3	Apply up to 2.0 fl oz (0.031 lb. ai) per acre.
<b>Tree Nuts:</b> including Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut (Bush Nut), Pecan, Pistachio and Walnut (Black and English)	Refer to table 3	Apply up to 2.0 fl oz (0.031 lb. ai) per acre.
<b>Tropical fruit:</b> including Papaya, Avocado, Black Sapote, Canistel, Mamey Sapote, Mango, Sapodilla, Star apple, Guava, Feijoa, Jaboticaba, Wax jambu, Starfruit, Passionfruit, Acerola, Lychee, Longan, Spanish lime, Rambutan, Pulasan, Sugar apple, Atemoya, Custard apple, Cherimoya, Llama, Soursop, and Biriba	Refer to table 3	Apply up to 2.0 fl oz (0.031 lb. ai) per acre.
<b>Other Crops:</b> including Banana, Cacao, Coconut, Coffee, Date, Fig, Guayule, Indian Mulberry, Olive, Palm Heart, Persimmon, Pomegranate, Tea, and Vanilla	Refer to table 3	Apply up to 2.0 fl oz (0.031 lb. ai) per acre.

**DIRECTIONS FOR USE**

**PRODUCTION SYSTEMS**

Different production systems dictate different application techniques. Skirted trees are those allowing the lower branches of the trees to grow to the ground line. Non-skirted trees are grown in production systems where branches are pruned allowing access to the trunk area.

**Equipment and Application**

**Skirted Orchards and Groves**

Hooded sprayers are required for Maxunitech Carfentrazone-ethyl 240EC Herbicide applications in skirted trees. Refer to the HOODED SPRAYER APPLICATIONS section of this label.

**Non-Skirted Orchards and Groves**

Apply only by ground equipment including boom sprayers, shielded or hooded sprayers, hand-held or high-volume wands or orchard guns. Use a minimum of 20 gallons finished spray solution per broadcast acre.

**Weed Control**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mix with other registered herbicides to actively growing weeds. Maxunitech Carfentrazone-ethyl 240EC Herbicide is a contact herbicide and coverage is essential for good weed control. Use a minimum of 20 gallons finished spray solution per broadcast acre.

**DO NOT** allow Maxunitech Carfentrazone-ethyl 240EC Herbicide spray solution to contact green stem tissue, leaves, fruit or blooms of trees.

**Maxunitech Carfentrazone-ethyl 240EC Herbicide Application Rates**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide up to 2 fl oz (0.031 lb. ai) per acre for postemergence control of susceptible broadleaf weeds. Refer to weed control list in Table 3 for appropriate weed control information. For best control, apply to seedling weeds in the 2 to 3-leaf stage. For larger weeds up to 6 leaves, use higher labeled rates of Maxunitech Carfentrazone-ethyl 240EC Herbicide. Weeds greater than 6 leaves may be only partially controlled.

**Adjuvant Requirements**

Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints NIS per 100 gallons) or a crop oil concentrate at 1% v/v (one-gallon COC per 100 gallons). Maxunitech Carfentrazone-ethyl 240EC Herbicide may also be applied with labeled rates of MSO or silicone adjuvants.

**Tank Mix**

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be mixed with other herbicides that have preemergence or postemergence activity. Maxunitech Carfentrazone-ethyl 240EC Herbicide only controls emerged vegetation. Any preemergence activity must rely on activity from registered preemergence herbicides mixed with instructions. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Sucker Management**

Maxunitech Carfentrazone-ethyl 240EC Herbicide is effective as an aid in the management of undesirable sucker growth from the base of the trunks or root sprouts. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at 2.0 fl oz (0.031 lb. ai) per acre. Suckers and other undesirable growth must be treated when the tissue is young and not mature and/or hardened off. Care must be taken not to allow spray mist to contact desirable fruit, foliage or green stem tissue (see Precautions).

**Adjuvant Requirements**

Refer to adjuvant section of this label.

**Chemical Mowing**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or in tank mixtures with other herbicides in chemical mowing practices for orchard vegetation management.

**Hooded Sprayer Application**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the **Hooded Sprayer Applications** section of this label for additional specific use directions.

**Precautions**

Extreme caution must be used during applications when desirable fruit and/or foliage are present in order to avoid fruit spotting and/or leaf necrosis.

**RESTRICTIONS:**

**DO NOT** allow spray mist of Maxunitech Carfentrazone-ethyl 240EC Herbicide to come in contact with green stem tissue, foliage, blooms or desirable fruit.

On seedling or newly transplanted trees **DO NOT** allow spray to contact green bark of trunk area. For new seedlings up to 2-year old trees, the trunk base should be wrapped to help prevent chemical contact with the bark.

**DO NOT** make applications with air-blast sprayers.

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per application.

**DO NOT** apply more than 6.1 fl oz (0.096 lb. ai) per acre per year for Tropical Fruits.



**DO NOT** apply more than 7.9 fl oz (0.124 lb. ai) per acre per year for Citrus Fruits, Pome Fruits, Stone Fruits, Tree Nuts, and Other Crops.

**DO NOT** make more than 5 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply within 3 days of harvest except Tropical Fruits which can be applied up to harvest.

**Tank Mix**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**GRASSES (Forage, Fodder, Hay, Seed and Sod)**

Methods and Timing	Target Weeds	Application Rates
Postemergence Weed Control	Refer to table 3	Up to 2.0 fl oz (0.031 lb. ai) per acre.

**Directions for Use:**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or in combination with other registered pesticides for the control of weeds in rangeland, pastures, hay, grasses grown for hay or silage and grass seed production and grass grown in Conservation Reserve Programs (CRP). Note that CRP usage must be in compliance with Federal, State, and local use guidelines.

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at use rates up to 2.0 fl oz (0.031 lb. ai) per broadcast acre. For optimum results, weeds should be treated when small. Applications shall be made with ground equipment delivering a minimum of 10 gallons of finished spray per acre or aerial delivering a minimum of 3 gal/acre of finished spray. Adjust sprayers to provide optimum coverage of the target weeds. Refer to weed control list in Table 3 for appropriate weed control information.

**Adjuvant Requirements**

Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a quality nonionic surfactant (NIS) containing at least 80% active at 0.25% v/v (2 pints NIS per 100 gallons) or a crop oil concentrate (COC) at 1% v/v (one-gallon COC per 100 gallons), or a methylated seed oil (MSO). The use of a high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v or ammonium sulfate (AMS) used at 2 to 4 pounds per acre in addition to the NIS, or MSO or COC is allowed.

When Maxunitech Carfentrazone-ethyl 240EC Herbicide is applied alone, grazing and hay operations may proceed with no restrictions.

**Tank Mix**

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with other labeled herbicides to control weeds not listed on this label. Read and follow all manufacturers' label directions for the companion herbicide.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**RESTRICTIONS:**

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per application.

**DO NOT** apply more than 5.9 fl oz (0.093 lb. ai) per acre per year.

**DO NOT** make more than 3 applications per year at reduced rates.

**DO NOT** make applications less than 7 days apart.

When Maxunitech Carfentrazone-ethyl 240EC Herbicide is applied alone, grazing and hay operations may proceed with no restrictions.

## HOPS

Methods and Timing	Target Weeds	Application Rates
Post-Directed for Sucker Management	Refer to table 3	2.0 fl oz (0.031 lb. ai) per acre.
Postemergence Weed Control	Refer to table 3	Up to 2.0 fl oz (0.031 lb. ai) per acre.

### DIRECTIONS FOR USE

#### Post-Directed Application for Sucker Management.

Maxunitech Carfentrazone-ethyl 240EC Herbicide is a contact herbicide for directed spray application to the basal portion of the hop plant for the management of sucker growth. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at 2.0 fl oz (0.031 lb. ai) per acre per application in a minimum of 20 gallons of spray solution by boom-type ground application equipment only to the basal portion of the hop plant (approximately the lower 1.5 feet) and to the sucker mat which extends from the base of the plant to approximately 1.5 to 2 feet into the row.

An alternate row treatment program may be followed to avoid the removal of excessive photosynthetic capacity from the crown area by treating alternate rows on different days. Applications timing and techniques may vary from region to region. Please consult local university extension personnel for local management practices.

#### Postemergent Control of Broadleaf Weeds

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide using shielded sprayers or hooded sprayers to control emerged and actively growing broadleaf weeds within or between the rows of the crop. Refer to Table 3 for appropriate weed control information.

#### Adjuvant Requirements

Coverage is essential to obtain good basal growth management. Use a nonionic surfactant (NIS) having at least 80 percent active ingredient at 0.25 % v/v (2 pints of NIS per 100 gallons of spray volume) or a quality crop oil concentrate (COC) at labeled rates.

If Maxunitech Carfentrazone-ethyl 240EC Herbicide is used in a tank mixture, refer to the other product labels for all restrictions on tank mixing and observe all label precautions, instructions and rotational cropping restrictions.

For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Rate Per Acre} = \text{Band Rate}$$

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Volume Per Acre} = \text{Band Volume}$$

#### Application Precautions

Extreme caution must be taken during application to avoid upward drift of the spray solution and contact with the highly susceptible new growth. Avoid applications until newly trained vines have developed sufficient barking to avoid damage to the stem and are high enough up the string to avoid contact with the apical bud.

#### RESTRICTIONS:

**DO NOT** apply Maxunitech Carfentrazone-ethyl 240EC Herbicide using air blast or air assisted sprayers.

**DO NOT** apply through any type of irrigation system.

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per application.

**DO NOT** apply more than 7.6 fl oz (0.12 lb. ai) per acre per year.

**DO NOT** make more than 4 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply within 7 days of harvest.

**LOW GROWING BERRY Crop subgroup 13-07G**

Methods and Timing	Target Weeds	Application Rates
Postemergence Weed Control	Refer to table 3	Up to 2.0 fl oz (0.031 lb. ai) per acre.

**DIRECTIONS FOR USE**

Maxunitech Carfentrazone-ethyl 240EC Herbicide applications will control susceptible emerged broadleaf weeds. Repeat applications may be necessary for weeds that emerge after an Maxunitech Carfentrazone-ethyl 240EC Herbicide treatment.

**Equipment and Application**

Apply only by ground equipment including boom sprayers, shielded or hooded sprayers, hand-held or high-volume wands or orchard guns. Use a minimum of 20 gallons finished spray solution per broadcast acre.

**Dormant Applications**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a broadcast application to the base of the trunk to control emerged and actively growing weeds during the dormant stage of the crop.

**Post-directed Applications for Broadleaf Weed Control**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a directed spray avoiding contact with the berry plant but directed at actively growing weeds. Maxunitech Carfentrazone-ethyl 240EC Herbicide is a contact herbicide and coverage is essential for good weed control. **DO NOT** allow Maxunitech Carfentrazone-ethyl 240EC Herbicide spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage.

Newly planted bush berries should only be treated with shielded sprayers or hooded sprayers.

**Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rates**

Apply up to 2.0 fl oz (0.031 lb. ai) Maxunitech Carfentrazone-ethyl 240EC Herbicide per broadcast acre. For best control, apply to seedling weeds in the 2 to 3-leaf stage. Use higher labeled rates of Maxunitech Carfentrazone-ethyl 240EC Herbicide for larger weeds up to 6 leaves. Weeds greater than 6 leaves may be only partially controlled. See Table 3 for Maxunitech Carfentrazone-ethyl 240EC Herbicide use rates and weeds controlled.

**Adjuvant Requirements**

A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be mixed with other registered herbicides for broader spectrum weed control. When tank mixing with fertilizer solutions, be sure to prepare an Maxunitech Carfentrazone-ethyl 240EC Herbicide premixture of Maxunitech Carfentrazone-ethyl 240EC Herbicide and clean water.

See Mixing and Loading Instructions under the PRODUCT INFORMATION section of this label for specific mixing instructions. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Precautions**

Extreme caution must be taken during applications when desirable fruit, foliage and/or blooms are present in order to avoid spotting or necrosis.

**RESTRICTIONS:**

**DO NOT** allow Maxunitech Carfentrazone-ethyl 240EC Herbicide spray mist to come in contact with green stem tissue, desirable fruit, blooms or foliage.

For seedling or newly transplanted bushes, **DO NOT** allow spray to contact green bark of trunk area. Use shielded sprayers only.

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per application.

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre during the dormant season.

**DO NOT** apply more than 6.15 fl oz (0.096 lb. ai) per acre per year.

**DO NOT** make more than 4 applications per year at reduced rates.

**DO NOT** make more than one dormant season application per year.

**DO NOT** make applications less than 14 days apart.

Can make applications up to harvest.

**Band Treatment Application**

For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Rate Per Acre} = \text{Band Rate}$$

$$\frac{\text{Band Width Inches}}{\text{Row Width Inches}} \times \text{Broadcast Volume Per Acre} = \text{Band Volume}$$

**MINT**

Methods and Timing	Target Weeds	Application Rates
<b>Broadcast</b>	Refer to table 3	Apply one application of Maxunitech Carfentrazone-ethyl 240EC Herbicide at 0.5 to 1.92 fl oz (0.008 to 0.030 lb. ai) per acre.  Use higher rates when weeds are under stress or are larger.

**DIRECTIONS FOR USE**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a broadcast application before Mint break dormancy for control of existing broadleaf weeds.

**Coverage is essential for good control.**

**Adjuvant Requirements**

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 10 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application. A nonionic surfactant (NIS), methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed. Repeat application if necessary.

For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section.

**RESTRICTIONS:**

**DO NOT** apply to actively growing crop.

**DO NOT** apply more than 1.92 fl oz (0.030 lb. ai) per acre per application.

**DO NOT** apply more than 1.92 fl oz (0.030 lb. ai) per acre per year.

**DO NOT** make more than 2 application per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply within 5 days of harvest.

## PEANUT

Methods and Timing	Target Weeds	Application Rates
Postemergence Weed Control	Refer to table 3	Up to 2.0 fl oz (0.031 lb. ai) per acre.
Harvest Aid	Refer to table 3	

### DIRECTIONS FOR USE

#### Weed Control

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other herbicides as a postemergence treatment or as a hooded/directed spray treatment to control emerged and actively growing weeds. Apply hooded/directed applications of Maxunitech Carfentrazone-ethyl 240EC Herbicide to middles (between rows of plants) and in strips (in row of plants). Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at any time during the year (see precautions). Maxunitech Carfentrazone-ethyl 240EC Herbicide may be mixed with other herbicides that have pre-emergence or post-emergence activity. Any pre-emergence activity must rely on activity from other herbicides as directed on their labels. Herbicides including glyphosate may be tank mixed with Maxunitech Carfentrazone-ethyl 240EC Herbicide for broader spectrum weed control. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### Harvest Aid Application

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a harvest aid to defoliate and desiccate troublesome weeds that may be present at harvest. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other peanut harvest aids.

#### Adjuvant Requirements

Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a quality nonionic surfactant (NIS) containing at least 80% active at 0.25% v/v (2 pints NIS per 100 gallons) or a crop oil concentrate (COC) at 1% v/v (one-gallon COC per 100 gallons), or a methylated seed oil (MSO). The use of a high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v or ammonium sulfate (AMS) used at 2 to 4 pounds per acre in addition to the NIS, or MSO or COC is allowed.

#### Harvest Aid

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a harvest aid to defoliate and desiccate troublesome weeds that may be present at harvest. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other peanut harvest aids.

**Coverage is essential for satisfactory performance.**

#### Crop Rotation Restriction:

After an application of this product to peanuts, you may only rotate the field to a carfentrazone-ethyl registered crop.

#### RESTRICTIONS:

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per application.

**DO NOT** apply more than 6.1 fl oz (0.096 lb. ai) per acre per year.

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per year as a Harvest Aid treatment.

**DO NOT** make more than 5 applications per year inclusive of all applications.

**DO NOT** make more than 1 Harvest Aid application per year.

**DO NOT** make more than 4 applications Postemergence Weed applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply within 7 days of harvest.

**DO NOT** feed immature peanut plant or peanut hay to livestock.

**RICE (Southern US Only)**

Methods and Timing	Target Weeds	Application Rates
<b>Pre-flood Applications to Dry Seeded Rice</b>	See weed list in table 10 below.	1.25 to 3.2 fl oz (0.0195 to 0.05 lb. ai) per acre
<b>Post Flood Applications to Exposed Weed</b>	See weed list in table 11 below.	1.6 to 6.4 fl oz (0.025 to 0.10 lb. ai) per acre
<b>Harvest Aid</b> (not permitted in California)	Desiccate troublesome broadleaf weeds e.g. hemp sesbania, Indian and northern jointvetch, morningglories, and pigweeds	1.25 to 1.5 fl oz (0.0195 to 0.023 lb. ai) per acre

**DIRECTIONS FOR USE**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other rice herbicides to emerged and actively growing weeds. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with either ground or aerial spray equipment. **DO NOT** apply when conditions favor drift.

To control weeds not listed on this label, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with other herbicides registered for use on rice. For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Postemergence Pre-flood Applications to Dry Seeded Rice**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at 1.25 to 3.2 fl oz (0.0195 to 0.05 lb. ai) per acre. Use a minimum of 10 gallons of finished spray per acre for ground application equipment, and a minimum of 3 gallons per acre of finished spray for aerial equipment. For optimum results, apply Maxunitech Carfentrazone-ethyl 240EC Herbicide to weeds up to 4 inches tall Use a quality nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. For more active treatments, use a Crop Oil Concentrate (COC) at 0.5 to 1.0% v/v (one half to one gallon per 100 gallons). Apply when the rice is at the 2 leaf stage or larger, but prior to flooding. Some leaf speckling may occur. Once field is flooded, water must be held for at least 23 days following treatment before release.

**When used as directed Maxunitech Carfentrazone-ethyl 240EC Herbicide will provide Control of listed weeds up to 4 inches tall.**

**Table 10:**

Cocklebur, common	Morningglory spp.
Copperleaf, hophornbeam	Pigweed spp.
Dayflower, spreading	Purslane, common
Groundcherry, cutleaf	Redweed
Hyssop, water	Sesbania, hemp
Jointvetch, Indian	Smartweed, PA (seedling)
Jointvetch, northern	

**Suppression of listed weeds**

Alligatorweed	Flatsedge, rice
Ducksalad	Redstem
Eclipta	Texasweed

**Tank Mix**

For control of weeds listed as suppressed or not listed on this label, apply Maxunitech Carfentrazone-ethyl 240EC Herbicide following a preemergence grass herbicide or tank with other rice herbicides for broad spectrum weed control. Use tank mix applications when rice is well established and in the appropriate stage of growth for treatment with Maxunitech Carfentrazone-ethyl 240EC Herbicide and the tank mix partner. For optimum results, weed species should also be in the proper stage of growth as specified on the Maxunitech Carfentrazone-ethyl 240EC Herbicide and tank mix partner label. Read and follow all manufacturers' label directions for the companion herbicide except for specific directions on this label. **DO NOT** add a surfactant or crop oil concentrate when tank mixing herbicides

formulated as emulsifiable concentrates unless required by the tank mix partners label. For other herbicide tank mix partners that are not Emulsifiable concentrates refer to their label for specific adjuvant directions. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. **Post Flood**

**Applications to Exposed Weeds**

For post flood applications apply Maxunitech Carfentrazone-ethyl 240EC Herbicide to rice and weeds after the establishment of the permanent flood and when 80% of the foliage of the weeds are exposed. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at 1.25 to 6.4 fl oz per acre (0.0195 to 0.10 lb. ai) per acre to actively growing weeds. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. For more active treatments, use a Crop Oil Concentrate (COC) at 1.0% v/v (one gallon per 100 gallons). Apply when the rice is at the 2- leaf stage or later. Use a minimum of 10 gallons of finished spray per acre for ground application equipment and a minimum of 3 gallons of finished spray per acre for aerial application equipment. For optimum results, make applications to small rather than larger weeds. If water level has been lowered to allow this treatment, it should be returned to normal levels 24 hours following treatment. Users of Maxunitech Carfentrazone-ethyl 240EC Herbicide must hold the water on the rice fields for 23 days following treatment.

**When used as directed, Maxunitech Carfentrazone-ethyl 240EC Herbicide will provide control of listed weeds.**

**Table 11:**

Arrowhead, annual	Morningglory spp.
Jointvetch, Indian	Sesbania, hemp
Jointvetch, northern	

**Suppression of listed weeds up to 4 inches.**

Alligatorweed	Ducksalad
Ammannia, purple	Flatsedge, rice
Dayflower, spreading	Texasweed

**Harvest Aid Application:**

Maxunitech Carfentrazone-ethyl 240EC Herbicide is effective as a harvest aid to defoliate and desiccate troublesome weeds that may be present at harvest. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other rice harvest aids. Harvest aid treatment applications may be made no earlier than soft dough up to the 3-day PHI. Refer to Table 1 for maximum use rate as harvest aid.

**Crop Rotation Restriction:**

After an application of this product to rice, you may only rotate the field to a carfentrazone-ethyl registered crop.

**RESTRICTIONS:**

**DO NOT** apply when conditions favor drift or when wind is above 10 mph.

**DO NOT** apply more than 3.2 fl oz (0.05 lb. ai) per acre per application for Pre-flood Applications to Dry Seeded Rice.

**DO NOT** apply more than 6.4 fl oz (0.10 lb. ai) per acre per application for Post Flood Applications to Exposed Weed.

**DO NOT** apply more than 1.5 fl oz (0.023 lb. ai) per acre per application for Harvest Aid.

**DO NOT** apply more than 8.8 fl oz (0.138 lb. ai) per acre per year including fallow/preplant burndown and other labeled crop applications for Pre-flood Applications to Dry Seeded Rice and Post Flood Applications to Exposed Weed.

**DO NOT** apply more than 1.5 fl oz (0.023 lb. ai) per acre per year as a Harvest Aid treatment.

**DO NOT** make more than 4 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply within 60 days of harvest once field is flooded for Pre-flood Applications to Dry Seeded Rice and Post Flood Applications to Exposed Weed.

**DO NOT** apply earlier than soft dough up to within 3 days of harvest for Harvest Aid applications.

Once field is flooded, water must be held for at least 23 days following treatment before release for Pre-flood Applications to Dry Seeded Rice.

**DO NOT** release water for at least 23 days following a Post Flood treatment in the water.

Harvest Aid Restriction: not permitted in California.

## RICE (For Rice Grown in California)

Methods and Timing	Target Weeds	Application Rates
<b>Early Post Seeding Applications to Submerged Weeds</b>	See weed list in table 12 below.	12.0 fl oz (0.19 lb. ai) per acre
<b>Foliar Applications to Emerged Weeds Above the Water Surface</b>	See weed list in table 12 below.	Up to 6.4 fl oz (0.10 lb. ai) per acre

### DIRECTIONS FOR USE

Apply alone or as a tank mixture with other rice herbicides to emerged and actively growing weeds. Applications shall be made by ground equipment only using a minimum finished spray volume of 10 gallons of spray per acre.

To control weeds not listed on this label, this product may be tank mixed with other herbicides registered for use on rice. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### Early Post Seeding Applications to Submerged Weeds

Apply at 12 fl oz (0.19 lb. ai) per acre. Evenly distribute the spray solution over the flooded rice. The floodwater must be 3 to 6 inches deep. Apply at 1.5 leaf stage of rice. Earlier applications may cause unacceptable crop response. Rice must be well rooted and actively growing at the time of application. Hold the floodwater at a static depth for at least five days after application. Once field is flooded, water must be held for at least 23 days following treatment before release.

**When used as directed this product will provide control of listed weeds at the 2-leaf stage or less.**

**Table 12:**

Arrowhead, California
Ammannia, purple (suppression only)
Ammannia, redstem (suppression only)
Bulrush, ricefield
Umbrellaplant, smallflower (suppression only)

#### Tank Mixtures

This product may be tank mixed with other herbicides to control weeds not listed on this label. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Apply before, after, or with an application of a herbicide with the following active ingredient: bensulfuron-methyl, molinate and thiobencarb. Observe all applicable directions, restrictions (including water holding requirements) and precautions on the bensulfuron-methyl, molinate and thiobencarb herbicide labels.

**DO NOT** apply as a tank mixture with a herbicide containing bispyribac-sodium.

#### Foliar Applications to Emerged Weeds Above the Water Surface

Apply up to 6.4 fl oz (0.10 lb. ai) per acre to the foliage of exposed weeds. At least 80% of the weed foliage must be exposed before spraying. For optimum results, apply to actively growing weeds 20 to 45 days postseeding or the earliest practical opportunity to spray. Weed control is enhanced with greater weed exposure. If the field was drained at application, relood twenty-four hours after application to the normal flood depth.

**When used as directed this product will provide control or suppression of the following weeds.**

**Table 13:**

Bulrush, ricefield
Arrowhead, California
Ammannia, purple (suppression only)
Ammannia, redstem (suppression only)
Umbrellaplant, smallflower (suppression only)

#### Crop Response

Some temporary leaf speckling may occur shortly after application.



**Tank Mix**

This product may be tank mixed with other herbicides to control weeds not listed on this label. This product may be tank mixed with propanil-containing herbicides. Not all combinations of this product and other formulated herbicides have been tested. The EC formulations, nonionic and silicone based surfactants and crop oil concentrates, when mixed with this product will increase leaf speckling on the rice leaves. These tank mixtures should be tested on a small portion of the field to ensure crop safety prior to use.

**Crop Rotation Restriction:**

After an application of this product to rice, you may only rotate the field to a carfentrazone-ethyl registered crop.

**RESTRICTIONS:**

**DO NOT** apply by air.

**DO NOT** apply within ½ mile of sensitive crops.

**DO NOT** apply when conditions favoring drift exist.

**DO NOT** apply more than 12.0 fl oz (0.19 lb. ai) per acre per application for Early Post Seeding Applications to Submerged Weeds.

**DO NOT** apply more than 6.4 fl oz (0.10 lb. ai) per acre per application for Foliar Applications to Emerged Weeds Above the Water Surface.

**DO NOT** apply more than 19.2 fl oz (0.3 lb. ai) per acre per year, including fallow, preplant, burndown, and labeled crop applications.

**DO NOT** make more than 4 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply within 60 days of harvest.

Once field is flooded, water must be held for at least 23 days following treatment before release.

**DO NOT** release water for at least 23 days following a Post Flood treatment in the water.

**RICE, WILD (Wild rice grown in cultivated fields where the water discharge/release can be controlled)**

Methods and Timing	Target Weeds	Application Rates
<b>Postemergence Weed Control</b>	See weed list in table 14.	6.4 to 12.0 fl oz (0.1 to 0.19 lb. ai) per acre

**DIRECTIONS FOR USE**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other rice herbicides to emerged and actively growing weeds. Wild rice should be well rooted and vigorously growing at the time of application. Earlier applications may cause unacceptable crop response. Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons per acre.

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide to weeds at the rate of 6.4 to 12.0 fl oz (0.1 to 0.19 lb. ai) per acre to the foliage of exposed weeds above the water surface. Make applications after the floating leaf stage through tillering. The water in paddies may be lowered if practical. Smaller weeds with more leaf area exposed will give better control. If water is lowered for application, it may be reflooded to normal depths 24 hours after the application.

**When used as directed Maxunitech Carfentrazone-ethyl 240EC Herbicide will provide control or suppression of the following weeds.**

**Table 14:**

Ammannia, purple (suppression only)
Ammannia, redstem (suppression only)
Arrowhead, California
Bulrush, ricefield
Burrweed, giant (Suppression only)
Umbrellaplant, smallflower (suppression only)
Waterplantain, common (Suppression only)

**Crop Response**

Some temporary leaf speckling may occur following application.

**Tank Mix**

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with other herbicides to control weeds not listed on this label. Not all combinations of Maxunitech Carfentrazone-ethyl 240EC Herbicide and other formulated herbicides and adjuvants have been tested. EC formulations, nonionic and silicone based surfactants, and crop oil concentrates, will increase leaf speckling on the wild rice leaves. These tank mixes should be tested on a small portion of the field to ensure crop safety prior to use.

**Precautions**

Wet leaf surfaces at the time of application can cause unacceptable injury.

**Crop Rotation Restriction**

After an application of this product to wild rice, you may only rotate the field to a carfentrazone-ethyl registered crop.

**RESTRICTIONS:**

**DO NOT** apply when conditions favoring drift exist.

**DO NOT** apply when winds exceed 10 mph.

**DO NOT** apply more than 12.0 fl oz (0.19 lb. ai) per acre per application.

**DO NOT** apply more than 19.2 fl oz (0.3 lb. ai) per acre per year including fallow/preplant, burndown, and other labeled crop applications.

**DO NOT** make more than 3 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply within 60 days of harvest.

**DO NOT** apply during the floating leaf stage when exposed wild rice leaves are most susceptible to injury.

**DO NOT** apply to wild rice when there is heavy dew on the leaves or under high humidity conditions.

**DO NOT** release flood water off wild rice field(s) for a minimum of 23 days after application of this product.

In California, **DO NOT** apply within 0.5 mile of sensitive crops.

In California, **DO NOT** apply to wild rice by air.

In California, **DO NOT** release flood water off wild rice field(s) for a minimum of 23 days after an application of this product.

**SMALL GRAINS**

Methods and Timing	Target Weeds	Application Rates
Preplant Burndown	Refer to Table 3	Up to 1.0 fl oz (0.031 lb. ai) per acre.
Postemergence	Refer to Table 3	0.5 to 1.0 fl oz (0.008 to 0.016 lb. ai) per acre.
Harvest Aid Applications	Refer to Table 3	Up to 2.0 fl oz (0.031 lb. ai) per acre.

**Directions for Use:****Timing and method of application:**

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be applied preplant (up to 1 day before seeding), postemergence or harvest aid. For optimum performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. For dense weed pressure, use the higher labeled application rate plus tank mix combinations. **Coverage is essential for good control.** Refer to Table 3 for weeds controlled at labeled rates of Maxunitech Carfentrazone-ethyl 240EC Herbicide. For broader spectrum weed control, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with other herbicides registered for use in small grains.

**Preplant Burndown:**

Refer to the pre plant burndown section of this label.

**Postemergence Application:**

In-season application may be made from 4-inches tall to just prior to the boot stage, **DO NOT** apply more than 0.016 lb. ai/acre including preplant and postemergent application (not including harvest aid). **DO NOT** apply more than 0.016 lb. ai/acre as a harvest aid treatment.

### Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rate

Apply from 0.5 to 1.0 fl oz Maxunitech Carfentrazone-ethyl 240EC Herbicide (0.008 – 0.016 lb. ai) per acre. Use a minimum finished spray solution of 10 gallons per acre by ground or 3 gallons per acre by air. Up to half of the spray volume (by air or ground) may be liquid nitrogen fertilizer.

### Adjuvant Requirements

Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. The use of a high quality sprayable liquid nitrogen fertilizer (2 to 4% v/v or 2 to 4 gallons per 100-gallon spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant is allowed. **DO NOT** use this product with crop oil concentrates (COC), methylated seed oils (MSO) or silicone based adjuvants for postemergence applications.

### Tank Mix

To control weeds not listed on this label, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with other registered herbicides.

For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Use aerial or ground equipment for Maxunitech Carfentrazone-ethyl 240EC Herbicide applications. **Coverage is essential for good control.** Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons of spray per acre. Applications made by air shall utilize a minimum finished spray volume of 3 gallons per acre. Up to half of the spray volume (by air or ground) may be liquid nitrogen fertilizer. Refer to Table 3 for appropriate weed control information.

### Maxunitech Carfentrazone-ethyl 240EC Herbicide Plus 2,4-D (amine or ester) or MCPA (amine or ester)

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed at a rate of 0.5 to 1.0 fl oz (0.008-0.016 lb. ai) per acre with 2,4-D (amine or ester) or MCPA (amine or ester) for use on small grains. For optimum results add 2,4-D (amine or ester) to the tank at 0.25 lb. acid equivalent per acre or MCPA (amine or ester) at 0.375 lb. acid equivalent per acre. Higher rates of these herbicides are allowed, but **DO NOT** exceed the label use rates allowed by these labels. Add nitrogen fertilizer (2 to 4% v/v) 2 to 4 gallons per 100 gallons or ammonium sulfate 4 lbs. per acre) to the tank mixture.

**When applied as directed, Maxunitech Carfentrazone-ethyl 240EC Herbicide in tank mixtures with 2,4-D (amine or ester) or MCPA (amine or ester) herbicides will provide control of listed weeds up to 4 inches tall.**

**Table 15:**

Amaranthus spp.	Nightshade, black
Bedstraw, catchweed	Pennycress, field **
Buckwheat, wild	Pepperweed, greenflower**
Cocklebur	Pigweed, prostrate
Croton, woolly	Pigweed, redroot
Fiddleneck	Pigweed, smooth
Filaree, redstem	Primrose, cutleaf
Flixweed**	Primrose, tumble
Gromwell, common	Radish, wild
Groundsel, common	Ragweed, common
Knotweed, prostrate*	Ragweed, giant
Kochia	Rocket, London
Lambsquarters, common	Sowthistle, annual
Lettuce, miners	Speedwell, ivyleaf
Lettuce, prickly	Sunflower, wild
Mustard, blue***	Tarweed, coast
Mustard, tansy***	Thistle, Russian
Mustard, tumble**	Wallflower, bushy
Mustard, wild**	Waterhemp, tall

\*For Knotweed control, use Maxunitech Carfentrazone-ethyl 240EC Herbicide + 2,4-D (amine or ester) only.

\*\*These weeds can be treated from the rosette through bolting growth stages.

\*\*\*Apply to rosette growth stage (before bolting) of blue mustard.

**Harvest Aid**

Apply up to 2.0 fl oz (0.032 lb. ai) Maxunitech Carfentrazone-ethyl 240EC Herbicide per acre, but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE MAXUNITECH CARFENTRAZONE-ETHYL 240EC HERBICIDE USE RATE and the PREHARVEST INTERVAL Table (Table 2) for additional application information. If treatments of Maxunitech Carfentrazone-ethyl 240EC Herbicide have been made to the crop earlier, that volume must be considered in determining the maximum use rate as a harvest aid treatment.

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 15 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application.

**Adjuvant Requirements – Harvest Aid**

A methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use methylated seed oil, or crop oil concentrate (COC) (petroleum or seed oil) at 1 to 2% v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the methylated seed oil or crop oil is allowed.

**Coverage is essential for satisfactory performance.**

**Restrictions:**

**DO NOT** apply when conditions favor drift.

**DO NOT** apply more than 1.0 fl oz (0.016 lb. ai) per acre per application for Preplant Burndown and Postemergence.

**DO NOT** apply more than 1.0 fl oz (0.016 lb. ai) per acre per application for Harvest Aid Applications.

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per year.

**DO NOT** apply more than 1.0 fl oz (0.016 lb. ai) per acre per year including fallow, preplant burndown or labeled crop applications.

**DO NOT** make more than 3 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** harvest for forage within 7 days of application.

**PHI:** Except Winter Wheat – jointing state. Winter Wheat – boot stage for Postemergence applications.

**SORGHUM (Grown for Grain and Seed)**

Methods and Timing	Target Weeds	Application Rates
<b>Preplant Burndown</b>	Refer to Table 3	Up to 1.0 fl oz (0.016 lb. ai) per acre.
<b>Foliar Broadcast Application (Grain Sorghum Only)</b>	Refer to Table 3 for weeds controlled at 0.5 fl oz per acre rate.	Up to 0.5 fl oz (0.008 lb. ai) per acre.
<b>Directed or Shielded Spray Applications</b>	Refer to Table 3	Up to 1.0 fl oz (0.016 lb. ai) per acre.
<b>Harvest Aid</b>	Desiccate troublesome broadleaf weeds e.g. morningglories, pigweeds and velvetleaf.	Up to 1.0 fl oz (0.016 lb. ai) per acre.

**DIRECTIONS FOR USE**

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be applied to grain and forage sorghum as a pre plant burndown; a hooded or shielded spray; and a post directed spray. In addition to these applications methods, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be applied to grain sorghum (sorghum grown for grain but not for seed production) as a foliar broadcast and harvest aid treatment. See Table 1 for Maximum Yearly Maxunitech Carfentrazone-ethyl 240EC Herbicide Use and Table 3 for weeds controlled at labeled rates of Maxunitech Carfentrazone-ethyl 240EC Herbicide on sorghum.

**PRE PLANT BURNDOWN**

See instructions under the Preplant Burndown section of this label.

**FOLIAR BROADCAST (Grain Sorghum Only)**

Apply to grain sorghum from 4 inches tall to just prior to the boot stage. Maxunitech Carfentrazone-ethyl 240EC Herbicide may be applied alone or as a tank mixture with other herbicides labeled for use on sorghum. Broadcast applications of Maxunitech Carfentrazone-ethyl 240EC Herbicide to sorghum with wet foliage or application during periods of adverse environmental conditions including cool, cloudy, wet, or high humidity may cause increased crop

response. Directed sprays are suggested under these conditions. For additional information on crop response, refer to the PRODUCT INFORMATION section of the Maxunitech Carfentrazone-ethyl 240EC Herbicide label.

#### **Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rates – Foliar Grain Only**

**DO NOT** exceed 0.5 fl oz (0.008 lb. ai) Maxunitech Carfentrazone-ethyl 240EC Herbicide per acre. See Table 3 for weeds controlled at 0.5 fl oz of Maxunitech Carfentrazone-ethyl 240EC Herbicide. Rates below 0.5 fl oz may not fully control weeds.

#### **Adjuvant Requirements – Foliar Grain Only**

Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. **DO NOT** use crop oil concentrates or methylated seed oils for broadcast applications on emerged sorghum.

#### **Tank Mix – Foliar Grain Only**

For control of additional broadleaf weeds and grasses, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with 2,4-D (amine); atrazine; dicamba; atrazine and sodium bentazon; halosulfuron-methyl; fluroxypyr-meptyl; or dicamba, dimethylamine salt. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Leaf speckling can occur when Maxunitech Carfentrazone-ethyl 240EC Herbicide is used with certain formulations of crop protection products and adjuvants.

#### **DIRECTED OR SHIELDED SPRAY APPLICATIONS**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide when the sorghum is at least 4 inches tall to prior to the boot stage. Use drop nozzles or other sprayers capable of directing the spray to the target weeds and away from the whorl and leaves of the sorghum plant. Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons per acre. Refer to Table 3 for weeds controlled at labeled rates of Maxunitech Carfentrazone-ethyl 240EC Herbicide. **Coverage is essential for good control.** Directed, shielded, or hooded sprayers are required for post emergence treatments to forage sorghum and sorghum grown for seed.

#### **Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rates – Directed or Shielded Spray**

Apply up to 1.0 fl oz Maxunitech Carfentrazone-ethyl 240EC Herbicide (0.016 lb. ai) per acre using directed or shielded sprayers.

#### **Adjuvant Requirements – Directed or Shielded Spray**

Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. Crop oil concentrates or methylated seed oils may increase crop injury on sorghum.

#### **Tank Mix – Directed or Shielded Spray**

For control of additional broadleaf weeds and grasses, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with 2,4-D (amine); atrazine; dicamba; atrazine and sodium bentazon; halosulfuron-methyl; fluroxypyr-meptyl; or dicamba, dimethylamine salt. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### **HOODED SPRAYER APPLICATION**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

#### **HARVEST AID (WEED CONTROL)**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide to defoliate and/or desiccate troublesome broadleaf weeds e.g. morningglories, pigweeds and velvetleaf that may be present at harvest. Apply up to 1.0 fl oz (0.016 lb. ai) Maxunitech Carfentrazone-ethyl 240EC Herbicide per acre, but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE MAXUNITECH CARFENTRAZONE-ETHYL 240EC HERBICIDE USE RATE and the PREHARVEST INTERVAL Table (Table 2) for additional application information. If treatments of Maxunitech Carfentrazone-ethyl 240EC Herbicide have been made to the crop earlier, that volume must be considered in determining the maximum use rate as a harvest aid treatment.

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 15 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application.

#### **Adjuvant Requirements – Harvest Aid**

A methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use methylated seed oil, or crop oil concentrate (COC) (petroleum or seed oil) at 1 to 2% v/v (1 to 2 gallons per 100 gallons of spray solution). A high

quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the methylated seed oil or crop oil is allowed.

**Coverage is essential for satisfactory performance.**

**PRECAUTIONS**

Leaf speckling can occur when Maxunitech Carfentrazone-ethyl 240EC Herbicide is used with certain formulations of crop protection products and adjuvants.

**RESTRICTIONS:**

**DO NOT** use crop oil concentrates or methylated seed oils for broadcast applications on emerged sorghum.

**DO NOT** apply more than 1.0 fl oz (0.016 lb. ai) per acre per application for Preplant Burndown.

**DO NOT** apply more than 0.5 fl oz (0.008 lb. ai) per acre per application for Foliar Broadcast Application (Grain Sorghum Only).

**DO NOT** apply more than 1.0 fl oz (0.016 lb. ai) for Directed or Shielded Spray Applications

**DO NOT** apply more than 1.0 fl oz (0.016 lb. ai) for Harvest Aid Applications.

**DO NOT** apply more than 1.0 fl oz (0.016 lb. ai) per acre per year including fallow, preplant burndown, labeled applications to the growing crop, and Harvest Aid treatment.

**DO NOT** make more than 3 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply past 14 leaf collar stage for Foliar Broadcast Application (Grain Sorghum Only).

**DO NOT** apply past preboot stage (forage), 14 collar (grain) for Directed or Shielded Spray Applications.

**DO NOT** apply within 3 days of harvest for Harvest Aid treatment.

**DO NOT** make foliar broadcast applications to forage sorghum or sorghum grown for seed.

**SOYBEANS**

Methods and Timing	Target Weeds	Application Rates
<b>Preplant Burndown</b>	Refer to table 3	Up to 1.5 fl oz (0.023 lb. ai) per acre
<b>Postemergence</b> (Broadcast)	Refer to table 3	0.25 to 0.5 fl oz (0.004 to 0.008 lb. ai) per acre. (See Directions for Use below for details).
<b>Postemergence</b> (Directed Spray and Hooded Sprayer Applications)	Refer to table 3	Up to 1.5 fl oz (0.023 lb. ai) per acre
<b>Harvest Aid</b>	Refer to table 3	Up to 1.5 fl oz (0.023 lb. ai) per acre

**Directions for Use:**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to soybeans in all tillage systems from prior to planting up to prior to emergence. **DO NOT** apply Maxunitech Carfentrazone-ethyl 240EC Herbicide during a period from emergence to V2. After plants have reached V3, applications are allowed up to V10.

For optimum performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. Use the higher rates when treating more mature weeds or dense vegetative growth. **Coverage is essential for good control.** Refer to weed control list in Table 3 for appropriate weed control information.

**Broadcast Postemergence Application**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at 1.5 fl oz (0.023 lb. ai) per acre for the control of velvetleaf. **DO NOT** apply Maxunitech Carfentrazone-ethyl 240EC Herbicide to soybeans with maturities less than

Group 2.0. For soybeans of maturity Group 2.1 to 3.4, apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at rates up to 1.5 fl oz (0.023 lb. ai) per acre. Use caution when making applications when making these treatments.

For soybeans maturing later than Group 3.5, apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at rates up to 1.5 fl oz (0.023 lb. ai) per acre.

#### Adjuvant Requirements

Use NIS only as the adjuvant for this treatment at the rate of 0.25% v/v (2 pints per 100 gallons of spray solution).

#### Broadcast Application Precaution

The application of Maxunitech Carfentrazone-ethyl 240EC Herbicide to soybeans may result in crop response. Soybeans may show some burn, speckling or necrosis of crop leaves. Soybeans quickly outgrow initial herbicide effects and yields are not affected. **DO NOT** make applications during conditions of abnormal cool, high humidity or if foliage is wet from dew, rainfall or irrigation. Users should be aware of these potential effects prior to making applications. If the user is not willing to accept these risks, applications should not be made.

For additional information on crop response, refer to the PRODUCT INFORMATION section of this label.

#### Tank Mix

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank-mixed with other herbicides to control weeds not listed on this label. **DO NOT use with diphenylether herbicides.** It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section. For control of additional broadleaf weeds and grasses, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank-mixed with glyphosate or glufosinate products for use on GMO soybeans. Leaf injury can occur when Maxunitech Carfentrazone-ethyl 240EC Herbicide is used with certain formulations of crop protection products and adjuvants. Refer to the Tank Mixtures and Required Adjuvants sections under PRODUCT INFORMATION.

**When used as directed Maxunitech Carfentrazone-ethyl 240EC Herbicide at 0.25 fl oz (0.004 lb. ai) per acre will provide:**

**Control of listed weeds up to 4 inches tall.**

Velvetleaf	
------------	--

**When used as directed, Maxunitech Carfentrazone-ethyl 240EC Herbicide at 0.5 fl oz (0.008 lb. ai) per acre will provide:**

**Control of weeds up to 4 inches tall, or as specified.**

Lambsquarters, common	Nightshade, black
Morningglory, Pitted (up to 3 true leaves)	Pigweed, redroot
Morningglory, Ivyleaf (up to 3 true leaves)	Waterhemp, spp. (up to 3 inches tall)

#### Hooded Sprayer Application

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications of this label for additional specific use directions.

#### Directed Sprayer Application

Use Maxunitech Carfentrazone-ethyl 240EC Herbicide at 0.5 to 1.5 fl oz (0.008 to 0.023 lb. ai) per acre. Applications shall be made by ground equipment using a finished volume of 10 to 20 gallons of spray per acre. When soybeans are grown under very dry soil moisture conditions, the use of a high quality sprayable liquid nitrogen fertilizer (2 to 4% v/v) or 2 to 4 gallons per 100-gallon spray solution) used in addition to the nonionic surfactant is allowed. Apply as a post-directed treatment with spray directed toward the base of the plant and avoid contact with soybean foliage. The use of spray shields may reduce spray contact with soybean foliage. Maxunitech Carfentrazone-ethyl 240EC Herbicide contact with soybean foliage can result in significant crop response.

#### RESTRICTIONS:

**DO NOT** apply more than 1.5 fl oz (0.023 lb. ai) per acre per application for Preplant Burndown, Postemergence (Directed Spray and Hooded Sprayer Applications), and Harvest Aid treatments.

**DO NOT** apply more than 0.5 fl oz (0.008 lb. ai) for Postemergence (Broadcast)

**DO NOT** apply more than 1.5 fl oz (0.023 lb. ai) per acre per year.

**DO NOT** make more than 5 applications per year at reduced rates.  
**DO NOT** make applications less than 14 days apart.  
**DO NOT** apply when conditions favoring drift exist.  
**DO NOT** feed treated soybean forage or hay to livestock.  
**DO NOT** use with diphenylether herbicides.  
**DO NOT** apply when crop foliage is wet from dew, rainfall or irrigation.  
**DO NOT** apply within 3 days of harvest for Harvest Aid applications.  
**DO NOT** apply past V10 for Postemergence applications.

## SUGARCANE

Methods and Timing	Target Weeds	Application Rates
Postemergence Treatment or Hooded/directed Spray	Refer to table 3	Up to 2.0 fl oz (0.031 lb. ai) per acre.
Harvest Aid	Desiccate troublesome broadleaf weeds e.g. morningglories, pigweeds and velvetleaf.	1.0 - 2.0 fl oz (0.016 - 0.031 lb. ai) per acre

### DIRECTIONS FOR USE

#### Postemergence/Hood Spray Application

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other herbicides as a postemergence treatment or as a hooded/directed spray treatment to control emerged and actively growing weeds. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide up to 2.0 fl oz (0.031 lb. ai) per acre. Apply hooded/directed applications of Maxunitech Carfentrazone-ethyl 240EC Herbicide to middles (between rows of plants) and in strips (in row of plants). Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide at any time during the year (see precautions). Maxunitech Carfentrazone-ethyl 240EC Herbicide may be mixed with other herbicides that have pre-emergence or postemergence activity. Any pre-emergence activity must rely on activity from other herbicides as directed on their labels. Herbicides including glyphosate may be tank mixed with Maxunitech Carfentrazone-ethyl 240EC Herbicide for broader spectrum weed control. If Maxunitech Carfentrazone-ethyl 240EC Herbicide is used in a tank mixture, observe the other product's label for restrictions, precautions and rotational cropping instructions.

#### Harvest Aid Application

Maxunitech Carfentrazone-ethyl 240EC Herbicide is effective as a harvest aid to defoliate and desiccate troublesome weeds that may be present at harvest. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other sugarcane harvest aids.

#### Adjuvant Requirements

Control is enhanced with the addition of a nonionic surfactant (NIS) or crop oil concentrate (COC). Use a quality nonionic surfactant (NIS) containing at least 80% active at 0.25% v/v (2 pints NIS per 100 gallons) or a crop oil concentrate (COC) at 1% v/v (one-gallon COC per 100 gallons), or a methylated seed oil (MSO). The use of a high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v or ammonium sulfate (AMS) used at 2 to 4 pounds per acre in addition to the NIS, or MSO or COC is allowed.

#### Crop Rotation

After an application of Maxunitech Carfentrazone-ethyl 240EC Herbicide to sugarcane, you may only rotate the field to a carfentrazone-ethyl registered crop.

#### RESTRICTIONS:

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per application.  
**DO NOT** apply more than 6.1 fl oz (0.096 lb. ai) per acre per year.  
**DO NOT** apply more than one Harvest Aid treatment per year.  
**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per year as a Harvest Aid treatment.  
**DO NOT** make more than 4 applications per year at reduced rates.  
**DO NOT** make applications less than 14 days apart.  
**DO NOT** apply within 7 days of harvest.



## TEFF (Grain and Forage)

Methods and Timing	Target Weeds	Application Rates
<b>Pre Plant Burndown</b>	Refer to table 3	Up to 1.0 fl oz (0.016 lb. ai) per acre
<b>Foliar Broadcast Application (Grain Teff Only)</b>	Refer to table 3 for weeds controlled at 0.5 fl oz (0.008 lb. ai) per acre rate.	Up to 0.5 fl oz (0.008 lb. ai) per acre
<b>Directed or Shielded Spray Applications</b>	Refer to table 3	Up to 1.0 fl oz (0.016 lb. ai) per acre
<b>Harvest Aid - Forage</b>	Desiccate troublesome broadleaf weeds e.g. morningglories, pigweeds and velvetleaf.	Up to 2.0 fl oz (0.031 lb. ai) per acre
<b>Harvest Aid - Grain</b>	Desiccate troublesome broadleaf weeds e.g. morningglories, pigweeds and velvetleaf.	Up to 2.0 fl oz (0.031 lb. ai) per acre

### DIRECTIONS FOR USE

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be applied to grain and forage teff as a pre-plant burndown; a hooded or shielded spray; and a post directed spray. In addition to these applications methods, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be applied to grain teff (teff grown for grain but not for seed production) as a foliar broadcast and harvest aid treatment. See Table 1 for Maximum Yearly Maxunitech Carfentrazone-ethyl 240EC Herbicide Use and Table 3 for weeds controlled at labeled rates of Maxunitech Carfentrazone-ethyl 240EC Herbicide on teff.

#### PRE PLANT BURNDOWN

See instructions under the Pre Plant Burndown section of this label.

#### FOLIAR BROADCAST (Grain Teff Only)

Apply to grain teff from 4 inches tall to just prior to the boot stage. Maxunitech Carfentrazone-ethyl 240EC Herbicide may be applied alone or as a tank mixture with other herbicides labeled for use on teff. Broadcast applications of Maxunitech Carfentrazone-ethyl 240EC Herbicide to teff with wet foliage or application during periods of adverse environmental conditions including cool, cloudy, wet, or high humidity may cause increased crop response. Directed sprays are suggested under these conditions. For additional information on crop response, refer to the PRODUCT INFORMATION section of the Maxunitech Carfentrazone-ethyl 240EC Herbicide label.

#### Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rates – Foliar Grain Only

**DO NOT** exceed 0.5 fl oz (0.008 lb. ai) Maxunitech Carfentrazone-ethyl 240EC Herbicide per acre. See Table 3 for weeds controlled at 0.5 fl oz (0.008 lb. ai) of Maxunitech Carfentrazone-ethyl 240EC Herbicide. Rates below 0.5 fl oz (0.008 lb. ai) may not fully control weeds.

#### Adjuvant Requirements – Foliar Grain Only

Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. **DO NOT** use crop oil concentrates or methylated seed oils for broadcast applications on emerged teff.

#### Tank Mix – Foliar Grain Only

For control of additional broadleaf weeds and grasses, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with 2,4-D (amine), atrazine; dicamba; atrazine and sodium bentazon; halosulfuron-methyl; fluroxypyr-meptyl; or dicamba, dimethylamine salt. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Leaf speckling can occur when Maxunitech Carfentrazone-ethyl 240EC Herbicide is used with certain formulations of crop protection products and adjuvants.

#### DIRECTED OR SHIELDED SPRAY APPLICATIONS

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide when the teff is at least 4 inches tall to prior to the boot stage. Use drop nozzles or other sprayers capable of directing the spray to the target weeds and away from the whorl and leaves of the teff plant. Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons per acre. Refer to Table 3 for weeds controlled at labeled rates of Maxunitech Carfentrazone-ethyl 240EC Herbicide. **Coverage is essential for good control.** Directed, shielded, or hooded sprayers are required for post emergence treatments to forage teff and teff grown for seed.

#### Maxunitech Carfentrazone-ethyl 240EC Herbicide Use Rates – Directed or Shielded Spray

Apply up to 1.0 fl oz Maxunitech Carfentrazone-ethyl 240EC Herbicide (0.016 lb. ai) per acre using directed or shielded sprayers.

**Adjuvant Requirements – Directed or Shielded Spray**

Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. Crop oil concentrates or methylated seed oils may increase crop injury on teff.

**Tank Mix – Directed or Shield Spray**

For control of additional broadleaf weeds and grasses, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with 2,4- D (amine); atrazine; dicamba; atrazine and sodium bentazon; halosulfuron-methyl; fluroxypyr-meptyl; or dicamba, dimethylamine salt. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**HOODED SPRAYER APPLICATION**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with hooded sprayers to control labeled weeds between the rows of the crop. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.

**HARVEST AID (WEED CONTROL)**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide to defoliate and/or desiccate troublesome broadleaf weeds e.g. morningglories, pigweeds and velvetleaf that may be present at harvest. Apply up to 2.0 fl oz (0.031 lb. ai) Maxunitech Carfentrazone-ethyl 240EC Herbicide per acre, but not to exceed maximum labeled rates. Refer to the MAXIMUM ALLOWABLE MAXUNITECH CARFENTRAZONE-ETHYL 240EC HERBICIDE USE RATE and the PREHARVEST INTERVAL Table (Table 2) for additional application information. If treatments of Maxunitech Carfentrazone-ethyl 240EC Herbicide have been made to the crop earlier, that volume must be considered in determining the maximum use rate as a harvest aid treatment.

Applications shall be made in spray volumes sufficient to provide complete coverage of foliage. Use a minimum of 15 gallons of finished spray per acre for ground application and 5 gallons per acre for aerial application.

**Adjuvant Requirements – Harvest Aid**

A methylated seed oil (MSO) or crop oil concentrate (COC) is required. Use methylated seed oil, or crop oil concentrate (COC) (petroleum or seed oil) at 1 to 2% v/v (1 to 2 gallons per 100 gallons of spray solution). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the methylated seed oil or crop oil is allowed.

**Coverage is essential for satisfactory performance.**

**PRECAUTIONS:**

Leaf speckling can occur when Maxunitech Carfentrazone-ethyl 240EC Herbicide is used with certain formulations of crop protection products and adjuvants.

**RESTRICTIONS:**

**DO NOT** apply more than 1.0 fl oz (0.016 lb. ai) per acre per application for Preplant Burndown and Directed or Shielded Spray Applications.

**DO NOT** apply more than 0.5 fl oz (0.008 lb. ai) per acre per application for Foliar Broadcast Application (Grain Teff Only).

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per application for Harvest Aid Applications.

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per year for all applications.

**DO NOT** make more than 3 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

**DO NOT** apply after the Jointing Stage for Foliar Broadcast Application (Grain Teff Only) and Directed or Shielded Spray Applications.

**DO NOT** apply within 7 days of harvest for Harvest Aid treatments (Forage).

**DO NOT** apply within 3 days of harvest for Harvest Aid treatments (Grain).

**DO NOT** make foliar broadcast applications to forage Teff or Teff grown for seed.

**DO NOT** use crop oil concentrates or methylated seed oils for broadcast applications on emerged teff.

## TOBACCO

Methods and Timing	Target Weeds	Application Rates
<b>Postemergence Weed Control</b> (pre-transplant, shielded/hooded spray, directed spray)	Refer to table 3	Up to 1.5 fl oz (0.024 lb. ai) per acre.
<p><b>DIRECTIONS FOR USE:</b></p> <p>Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or as a tank mixture with other registered herbicides to emerged and actively growing weeds. For optimum performance, make applications to weeds up to 4 inches tall and rosettes less than 3 inches across. Use higher rates when treating more mature weeds or dense vegetative growth.</p> <p><b>Coverage is essential for good control.</b></p> <p><b>Adjuvant Requirements</b>            Use adequate spray volume to achieve thorough coverage, but a minimum of 10 gallons of finished spray per acre is required. Use a quality crop oil concentrate (COC) at 1% v/v (1 gallon of COC per 100 gallons of spray solution).            Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with other herbicides registered for use on tobacco to provide additional weed control. For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. For additional information refer to the PRODUCT INFORMATION section of the Maxunitech Carfentrazone-ethyl 240EC Herbicide label.</p> <p><b>Pre-transplant burndown</b>            Maxunitech Carfentrazone-ethyl 240EC Herbicide is a contact herbicide for pre-transplant burndown control of broadleaf weeds in tobacco. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a broadcast application alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide up to one (1) day prior to transplanting.</p> <p><b>Shielded spray or Hooded spray</b>            Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide using shielded sprayers or hooded sprayers to emerged and actively growing broadleaf weeds in tobacco from transplanting until layby. Shielded spray or hooded spray applications of Maxunitech Carfentrazone-ethyl 240EC Herbicide or Maxunitech Carfentrazone-ethyl 240EC Herbicide tank mixtures should utilize application equipment that must prevent contact of spray solution with the tobacco plant. <b>DO NOT</b> allow spray solution to contact tobacco foliage or green stem tissue. Refer to the Hooded Sprayer Applications section of this label for additional specific use directions.</p> <p><b>Directed spray after first priming (Flue Cured Tobacco Only)</b>            Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a directed spray application after the first priming in only flue cured tobacco only for the control of emerged and actively growing broadleaf weeds. Directed spray equipment should position nozzles a minimum of 3 to 4 inches above the soil, with nozzles directed underneath the crop canopy. Spray solution should be directed at the base of tobacco plants for minimal contact with foliage while maintaining maximum contact with broadleaf weeds that are at appropriate treatment size. <b>DO NOT</b> apply when conditions favor drift or wind is above 10 mph.</p> <p>For control of additional broadleaf weeds and grasses, Maxunitech Carfentrazone-ethyl 240EC Herbicide may be tank mixed with other herbicides registered for use in tobacco at the appropriate timing. Refer to weed control list in Table 3 for appropriate weed control information. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions and rotational cropping restrictions.</p> <p><b>Restrictions:</b></p> <p><b>DO NOT</b> apply more than 1.5 fl oz (0.024 lb. ai) per acre per application.</p> <p><b>DO NOT</b> apply more than 3.2 fl oz (0.05 lb. ai) per acre per year.</p> <p><b>DO NOT</b> make more than 3 applications per year at reduced rates.</p> <p><b>DO NOT</b> make applications less than 14 days apart.</p> <p><b>DO NOT</b> apply within 6 days of harvest.</p>		

**TUBEROUS AND CORM VEGETABLES (except Potato) Crop subgroup 1C & 1D**

Methods and Timing	Target Weeds	Application Rates
<p><b>Fallow Systems</b> See the <b>Fallow Systems</b> section for directions for application.</p> <p><b>Preplant Burndown</b> See the <b>Preplant Burndown</b> section for directions for application.</p>	Refer to table 3	Up to 2.0 fl oz Maxunitech Carfentrazone-ethyl 240EC Herbicide (0.031 lb. ai) per acre.
<p><b>Harvest Aid</b></p>	Refer to table 3	3.2 to 5.8 fl oz (0.05 to 0.09 lb. ai) per acre. 2.0 – 5.8 fl oz (0.031 to 0.09 lb. ai) with other registered potato desiccants.

**DIRECTIONS FOR USE**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or in a tank mix combination with other herbicides and insecticides as a fallow systems treatment, as a preplant burndown treatment and/or as a harvest aid to desiccate potatoes and those susceptible weeds that may be present.

**Fallow Systems**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide by ground or air alone or with other herbicides in the fallow period prior to planting or the emergence of any crop listed on this label to control or suppress weeds. For optimum performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across.

**Coverage is essential for good weed control.**

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be utilized in Fallow Cropping Systems for chemical weed control to aid in moisture conservation between cropping periods.

**Preplant Burndown**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide alone or with other herbicides or liquid fertilizers as a burn-down treatment to control or suppress weeds. Maxunitech Carfentrazone-ethyl 240EC Herbicide is effective as a burndown treatment for crops prior to new plantings. **DO NOT** exceed the applicable amounts as listed for the specific crop in the MAXIMUM ALLOWABLE MAXUNITECH CARFENTRAZONE-ETHYL 240EC HERBICIDE USE in Table 1. For optimum performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across. **Coverage is essential for good control.** Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a labeled burndown herbicides including glyphosate, glufosinate, paraquat, 2,4-D, or dicamba.

**Harvest Aid Desiccation Application**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide foliar to potatoes in the later stages of senescence for desiccation of potato foliage and vines. Maxunitech Carfentrazone-ethyl 240EC Herbicide will also desiccate late season susceptible broadleaf weeds to aid in tuber harvest. Adequate desiccation is achieved within 14 days after the initial treatment is applied. If the potato crop is in the active vegetative growth stage when desiccation is initiated, two applications may be required to provide desiccation of leaf and stem tissue. Dense potato canopy, large plant size and environmental conditions not conducive to product absorption or activity will reduce initial application efficacy and increase the need for a second application. If a second application is necessary, apply at 7 to 14 days after the first application. **Thorough coverage of the potato plant to be desiccated is essential.** Use a sufficient volume of water to obtain thorough coverage of the potato leaves and vines.

**Ground Application**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide in at least 20 gallons of water per acre. Vary the spray volume and spray pressure as indicated by the density of the potato canopy and vines to assure thorough spray coverage. Increase the spray volume and pressure if the potato canopy is dense or under cool, cloudy or dry conditions. Increased spray volumes will enhance performance.

**Aerial Application**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide with aerial equipment using 5 to 10 gallons of water per acre, using higher volumes when potato canopies and vines are dense. Adjust the nozzles to provide a uniform pattern and a droplet size of 350 to 450 microns.

**Adjuvant Requirements**

A nonionic surfactant (NIS), methylated seed oil (MSO), crop oil concentrate (COC) or other suitable surfactant mixture is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient, or a methylated seed oil, or crop oil concentrate (COC)(petroleum or seed oil) at 1 to 2 v/v (1 to 2 gallons per 100 gallons of spray solution. The use of a high quality sprayable liquid nitrogen fertilizer at 2

to 4 % v/v (2 to 4 gallons per 100 gallons spray solution) or ammonium sulfate (AMS) at the rate of 2 to 4 pounds per acre in addition to the nonionic surfactant methylated seed oil or crop oil is allowed.

Adjuvant rates should increase as spray volumes exceed 20 gallons per acre.

**Tank Mix**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide as a tank mix or as a sequential application with other potato desiccants. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**RESTRICTIONS:**

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per application for Fallow Systems and Preplant Burndown Applications.

**DO NOT** apply more than 5.8 fl oz (0.09 lb. ai) per acre per application for Harvest Aid Applications.

**DO NOT** apply more than 11.6 fl oz (0.181 lb. ai) per acre per year as a desiccant.

**DO NOT** apply more than 11.6 fl oz (0.181 lb. ai) per acre per year for all applications.

**DO NOT** make more than 3 applications per year.

**DO NOT** make applications less than 7 days apart.

**DO NOT** apply within 7 days of harvest.

**DO NOT** apply when conditions favor drift or wind is above 10 mph.

**FALLOW SYSTEMS**

Methods and Timing	Target Weeds	Application Rates
Postemergence Weed Control	Refer to table 3	Up to 2.0 fl oz (0.031 lb. ai) per acre.

**Directions for Use:**

Apply Maxunitech Carfentrazone-ethyl 240EC Herbicide by ground or air alone or with other herbicides in the fallow period prior to planting or the emergence of any crop listed on this label to control or suppress weeds. For optimum performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across. **Coverage is essential for good weed control.**

Maxunitech Carfentrazone-ethyl 240EC Herbicide may be utilized in Fallow Cropping Systems for chemical weed control to aid in moisture conservation between cropping periods.

**Adjuvant Requirements**

A nonionic surfactant, crop oil concentrate or methylated seed oil is required. Use a nonionic surfactant (NIS) at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient or a petroleum or oil seed based crop oil concentrate (COC) at 1.0 to 2 % v/v (1.0 to 2.0 gallons per 100 gallons of spray solution) or a methylated seed oil (MSO). A high quality sprayable liquid nitrogen fertilizer at 2 to 4 % v/v (2 to 4 gallons per 100 gallons) or ammonium sulfate at 2 to 4 pounds per acre in addition to the selected NIS, MSO or COC is allowed.

Optimum broad-spectrum control of annual and perennial weeds requires a tank mix with a broad-spectrum burndown herbicide including glyphosate, glufosinate or paraquat. Refer to Table 3 for proper use rate for weed spectrum. For specific mixing instructions, refer to the Mixing and Loading Instructions under the PRODUCT INFORMATION section.

For crop planting information following fallow treatments, refer to the preplant burnout for planting interval instructions.

**Tank Mix**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**RESTRICTIONS:**

**DO NOT** apply more than 2.0 fl oz (0.031 lb. ai) per acre per application.

**DO NOT** apply more than 7.9 fl oz (0.124 lb. ai) per acre per year.

**DO NOT** make more than 5 applications per year at reduced rates.

**DO NOT** make applications less than 14 days apart.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage or disposal.

### Pesticide storage

Not for use or storage in or around the home.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. **DO NOT** put formulated or dilute material into food or drink containers. **DO NOT** contaminate other pesticides, fertilizers, water, food, or feed by inappropriate storage or disposal.

To confine spill: Dike surrounding area, sweep up spillage. Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in an approved manner. Place damaged container in a larger holding container. Identify contents per required hazardous waste labeling regulations.

### Pesticide Disposal

Waste resulting from the use of this product may be disposed of at an approved waste disposal facility.

### Container Handling

**[NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS): DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.]

**[NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS): DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.]

**[REFILLABLE CONTAINER:** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.]

## LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES and LIMITATIONS OF LIABILITY. CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Maxunitech North America LLC. All such risks shall be assumed by the user or buyer. **DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Maxunitech North America LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Maxunitech North America LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Maxunitech North America LLC disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Maxunitech North America LLC's election, the replacement of product.

[EPA approval date]

**{[LANGUAGE ON LABEL AFFIXED TO CONTAINER]}**

CARFENTRAZONE-ETHYL	GROUP	14	HERBICIDE
---------------------	-------	----	-----------

**Maxunitech Carfentrazone-ethyl 240EC Herbicide**

<b>ACTIVE INGREDIENT:</b>	<b>By Wt.</b>
Carfentrazone-ethyl .....	23.93%
<b>OTHER INGREDIENTS:</b> .....	76.07%
<b>TOTAL:</b> .....	100.00%

This product contains 2.0 lbs. active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

FIRST AID	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li><b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li><b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12 PM Pacific Standard Time. In the event of a medical emergency, call your poison control center at 1-800-222-1222.	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS**

Carfentrazone-ethyl is very toxic to algae and moderately toxic to fish. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the high-water mark, except as specified on this label. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

**For ground water:**

Residues of this chemical have properties and characteristics associated with chemicals detected in ground water. Residues of this chemical may leach into ground water if the chemical is used in areas where soils are permeable, particularly where the water table is shallow.

**For Surface Water:**

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of carfentrazone-ethyl residues from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

**STORAGE AND DISPOSAL**

**DO NOT** contaminate water, food or feed by storage or disposal.

**Pesticide storage**

Not for use or storage in or around the home.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. **DO NOT** put formulated or dilute material into food or drink containers. **DO NOT** contaminate other pesticides, fertilizers, water, food, or feed by inappropriate storage or disposal.

To confine spill: Dike surrounding area, sweep up spillage. Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in an approved manner. Place damaged container in a larger holding container. Identify contents per required hazardous waste labeling regulations.

**Pesticide Disposal**

Waste resulting from the use of this product may be disposed of at an approved waste disposal facility.

**Container Handling**

**[NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS): DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.]

**[NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS): DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.]

**[REFILLABLE CONTAINER:** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.]

See label booklet for additional Precautionary Statements and Directions for Use.

**EPA Reg. No.: 95009-XX**

**EPA Est. No.:**

**Manufactured for:**

Maxunitech North America LLC  
853 Valparaiso Ave.  
Menlo Park, CA 94025

**Net Contents:**