



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505T)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

95009-14

Date of Issuance:

9/19/25

NOTICE OF PESTICIDE:

☒ Registration  
☐ Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Maxunitech Prothio 480 SC

Name and Address of Registrant (include ZIP Code):

Maxunitech North America, Inc.  
11601 Shadow Creek Pkwy, Suite 111-573  
Pearland, TX 77584

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

*Continues page 2*

Signature of Approving Official:

Stephanie Suarez, Ph.D., Acting Product Manager 22  
Fungicide Branch, Registration Division (7505T)

Date:

9/19/25

2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 95009-14."
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 06/25/2023

If you have any questions, please contact James Orrock by phone at 202-566-2862 or by email at [orrock.james@epa.gov](mailto:orrock.james@epa.gov).

Enclosure

{Note to reviewer: [Text] in brackets denotes optional text. {Text} in braces denotes where in the final label text will appear and notes to reviewer.}

{BOOKLET FRONT PANEL}



PROTHIOCONAZOLE	GROUP	<b>3</b>	FUNGICIDE
-----------------	-------	----------	-----------

## Maxunitech Prothio 480 SC

FOR CONTROL OF SPECIFIED DISEASES ON LISTED CROPS

Active Ingredient:	By Wt
Prothioconazole*	41.48%
Other Ingredients	58.52%
Total:	100.00%

\*2-[2-(1-Chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]-1,2-dihydro-3H-1,2,4-triazole-3-thione

Maxunitech Prothio 480 SC contains 4.0 pounds prothioconazole per gallon.

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

[See inside label booklet for additional] [Precautionary Statements][,] [and] [Directions for Use] [including] [Storage and Disposal] [instructions][.]

EPA Reg. No.: 95009-XX

EPA Est. No.:

**Manufactured for:**

Maxunitech North America, Inc.

11601 Shadow Creek Pkwy, Suite 111-573

Pearland, TX 77584

**ACCEPTED**

**09/19/2025**

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 95009-14

[Lot][Batch]No:

**Net Contents:**

## {LANGUAGE INSIDE BOOKLET}

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li></ul>
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
IF INHALED:	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>HOTLINE NUMBER</b> <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12 PM Pacific Standard Time. In the event of a medical emergency, call your poison control center at 1-800-222-1222.</p>	

**For Chemical Emergency:**  
**Spill, Leak, Fire, Exposure, or Accident,**  
**Call CHEMTREC Day or Night**  
**Within USA and Canada: 1-800-424-9300**

### **PRECAUTIONARY STATEMENTS** **HAZARDS TO HUMANS & DOMESTIC ANIMALS** **CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

#### **PERSONAL PROTECTIVE EQUIPMENT (PPE):**

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material including barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils, shoes and socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### **ENGINEERING CONTROL STATEMENTS**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS	
Users should:	<ul style="list-style-type: none"><li>• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.</li><li>• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li><li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li></ul>

## ENVIRONMENTAL HAZARDS

This product is toxic to estuarine/marine invertebrates, and freshwater/estuaries/marine aquatic plants. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Degradates of prothioconazole are known to leach through soil into groundwater under certain conditions as a result of label use. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### Surface Water Advisory:

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of prothioconazole and degradates from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

## PHYSICAL OR CHEMICAL HAZARDS

**DO NOT** mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### RESTRICTIONS:

- **DO NOT** apply prothioconazole with mechanically pressurized handgun equipment.
- **DO NOT** apply prothioconazole with mechanically pressurized handgun equipment to orchards or vineyards (Bushberry subgroup 13-07B, Low growing berry subgroup, except strawberry subgroup 13-07H), field crops (Cucurbit Vegetables (Crop Group 9), Corn, sweet, Garbanzos (including chickpeas); Lentils) or nursery pine and conifer seedlings (Shortleaf Loblolly, Slash, Longleaf and other pines, other conifers, other hardwoods).

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Crop-specific REIs are listed in the Directions for Use section associated with the crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil or water is: coveralls, chemical resistant gloves made of waterproof material including barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, polyvinyl chloride (PVC)  $\geq$  14 mils, or Viton  $\geq$  14 mils, shoes plus socks.

## PRODUCT INFORMATION

Maxunitech Prothio 480 SC is a broad-spectrum systemic fungicide for the control of Ascomycetes, Basidiomycetes and Deuteromycetes diseases in barley, buckwheat, bushberry subgroup, low growing berry subgroup (except strawberry), corn, cotton, cucurbit vegetables, dry shelled pea and bean crop subgroup, millet, oats, peanuts, rapeseed subgroup 20A (including canola and *Brassica carinata*), rice, rye, soybean, sugar beets, triticale, wheat; conifer and hardwood nursery seeds and seedlings. Under conditions conducive to extended infection periods or high disease pressure, another registered fungicide may be needed once this product's maximum application rates have been reached. Under these conditions use another fungicide registered for the crop/disease. Equipment must be properly calibrated before use.

**RESISTANCE MANAGEMENT** For resistance management, Maxunitech Prothio 480 SC is a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to Maxunitech Prothio 480 SC and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides/bactericides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance take one or more of the following steps:

- Rotate the use of Maxunitech Prothio 480 SC or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact a Maxunitech North America, Inc. retailer or representative.

**SPRAY EQUIPMENT/VOLUMES:** Maxunitech Prothio 480 SC may be applied by either ground, aerial and/or chemigation application equipment. Refer to the USE DIRECTIONS FOR SPECIFIC CROPS section of this label for approved applications for each crop.

Apply in a minimum of 10 gallons of spray solution per acre by ground sprayer. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment unless stated differently elsewhere in this label. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

**MIXING PROCEDURES:** It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Prepare no more spray mixture than is necessary for the immediate operation. Thoroughly clean spray equipment before using this product. Maintain maximum agitation throughout the spray operation. **DO NOT** let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to the previously treated area or dispose of the rinsate according to local regulations.

**Maxunitech Prothio 480 SC Alone:** Add ½ of the required amount of water to the mix tank. With the agitator running, add the product to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the product has completely and uniformly dispersed into the mix water. Maintain agitation until

all of the mixture has been applied.

**Maxunitech Prothio 480 SC with Tank-Mix Partners:** Add ½ of the required amount of water to the mix tank. Start the agitator running before adding any of the tank-mix partners. In general, add tank-mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables), liquid flowables, liquids, and emulsifiable concentrates. Always allow each tank-mix partner to become fully and uniformly dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

When using Maxunitech Prothio 480 SC in tank mixtures, add all products in water-soluble packaging to the tank before any other tank-mix partner. Allow the water-soluble packaging to completely disperse before adding any other tank-mix partner to the tank.

If using Maxunitech Prothio 480 SC in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and restrictions; which appear on the tank-mix product label. No label dosage rate may be exceeded, and the most restrictive label precautions and limitations must be followed. This product must not be mixed with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

Maxunitech Prothio 480 SC is compatible with most insecticide, fungicide, herbicide and foliar nutrient products. However, test the physical compatibility of Maxunitech Prothio 480 SC with tank-mix partners before use. To determine the physical compatibility of Maxunitech Prothio 480 SC with other products, use a jar test, as described below.

Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water-dispersible granular products first, then liquids, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank. For further information contact Maxunitech North America, Inc.

The crop safety of all potential tank mixes including additives and other pesticides on all crops has not been tested. Before applying any tank mixture not specified on this label, confirm the safety to the target crop. To test for crop safety, apply Maxunitech Prothio 480 SC to the target crop in a small area and in accordance with label instructions for the target crop.

### Soilborne/Seedling Disease Control

Maxunitech Prothio 480 SC can provide control of certain soilborne/seedling diseases when applied as an in-furrow application at time of planting or as a banded application applied over the row at time of planting and up to row closure on certain crops. Refer to the USE DIRECTIONS FOR SPECIFIC CROPS section of the label to determine which crops contain specifications for in-furrow and/or banded applications and for the optimal timings, rates and band widths of these applications.

### In-Furrow Application

- Apply Maxunitech Prothio 480 SC as an in-furrow application in 2.5 to 20 gallons of water at planting.
- Mount the spray nozzle such that the spray is directed into the furrow just before the seeds are covered.

### Banded Application

- Apply Maxunitech Prothio 480 SC as a directed spray to the soil, using single or multiple nozzles which can be adjusted to provide uniform coverage of the lower stems and the soil surface surrounding the plants.

Rate per 1000 Row Feet	Rate of Maxunitech Prothio 480 SC for In-Furrow and Banded Applications (fl oz/A)								
	15" rows	20" rows	22" rows	30" rows	32" rows	34" rows	36" rows	38" rows	40" rows
0.075	2.61								



Rate per 1000 Row Feet	Rate of Maxunitech Prothio 480 SC for In-Furrow and Banded Applications (fl oz/A)								
fl oz product	15" rows	20" rows	22" rows	30" rows	32" rows	34" rows	36" rows	38" rows	40" rows
0.100	3.48	2.61							
0.125	4.36	3.27	2.97						
0.150	5.23	3.92	3.56	2.61					
0.175	6.10	4.57	4.16	3.05	2.86	2.69			
0.200	6.97	5.23	4.75	3.48	3.27	3.07	2.90	2.75	2.61
0.225		5.88	5.35	3.92	3.68	3.46	3.27	3.10	2.94
0.250		6.53	5.94	4.36	4.08	3.84	3.63	3.44	3.27
0.275			6.53	4.79	4.49	4.23	3.99	3.78	3.59
0.300				5.23	4.90	4.61	4.36	4.13	3.92
0.325				5.66	5.31	5.00	4.72	4.47	4.25
0.350				6.10	5.72	5.38	5.08	4.81	4.57
0.375				6.53	6.13	5.77	5.45	5.16	4.90
0.400				6.97	6.53	6.15	5.81	5.50	5.23
0.425					6.94	6.53	6.17	5.85	5.55
0.450						6.92	6.53	6.19	5.88
0.475							6.90	6.53	6.21
0.500								6.88	6.53
0.525									6.86
0.543									7.10

**AERIAL APPLICATION:** Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. **DO NOT** apply directly to humans or animals.

**CHEMIGATION APPLICATION:** Apply Maxunitech Prothio 480 SC through irrigation equipment only to crops for which chemigation is specified on this label.

Maxunitech Prothio 480 SC alone or in combination with other pesticides, which are registered for application through irrigation systems, may be applied through irrigation systems. Apply this product only through center pivot, solid set, drip, linear, or moving wheel irrigation systems. **DO NOT** apply this product through any other type of irrigation system. Illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

#### Operating Instructions:

1. The system must contain a functional check-valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump),



effectively designed, and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

#### **Center Pivot Irrigation Equipment:**

Notes: (1) Use only with drive systems, which provide uniform water distribution. (2) **DO NOT** use end guns when chemigating Maxunitech Prothio 480 SC through center pivot systems because of non-uniform application.

Determine the size of the area to be treated. Determine the time required to apply 1/8-1/2 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as advised by the equipment manufacturer. When applying Maxunitech Prothio 480 SC through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80-95% of the manufacturer's rated capacity. Using water, determine the injection pump output when operated at normal line pressure. Determine the amount of Maxunitech Prothio 480 SC required to treat the area covered by the irrigation system. Add the required amount of Maxunitech Prothio 480 SC L and sufficient water to meet the injection time requirements to the solution tank. Make sure the system is fully charged with water before starting injection of the Maxunitech Prothio 480 SC solution. Time the injection to last at least as long as it takes to bring the system to full pressure. Maintain constant solution tank agitation during the injection period. Continue to operate the system until the Maxunitech Prothio 480 SC solution has cleared the sprinkler head.

#### **Solid Set and Moving Wheel Irrigation Equipment:**

When applying Maxunitech Prothio 480 SC through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Determine the amount of Maxunitech Prothio 480 SC required to treat the area covered by the irrigation system. Add the required amount of Maxunitech Prothio 480 SC into the same quantity of water used to calibrate the injection period. Operate the system at the same pressure and time interval established during the calibration. Stop injection equipment after treatment is completed. Continue to operate the system until the Maxunitech Prothio 480 SC solution has cleared the last sprinkler head.

#### **Using Water from Public Water Systems:**

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

**Adjuvants:** It is advisable to use Maxunitech Prothio 480 SC with a registered non-ionic surfactant at the lowest specified labeled rate for most crops. Refer to the USE DIRECTIONS FOR SPECIFIC CROPS for specific uses where a surfactant is not specified.

#### **SPRAY DRIFT ADVISORIES**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.  
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

## **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### **Controlling Droplet Size – Ground Boom**

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure specified for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **Controlling Droplet Size – Aircraft**

- Adjust Nozzles – Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles must be oriented parallel with the airflow in flight.

## **BOOM HEIGHT – Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

## **RELEASE HEIGHT – Aircraft**

Higher release heights increase the potential for spray drift.

## **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

## **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

## **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators used to be familiar with local wind patterns and terrain that could affect spray drift.

### **Handheld Technology Applications:**

- Take precautions to minimize spray drift.

## **ROTATIONAL RESTRICTIONS**

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. For crops not listed on this label, **DO NOT** plant back within 30 days of last application.

## **USE DIRECTIONS FOR SPECIFIC CROPS**

Maxunitech Prothio 480 SC provides control or suppression of many important diseases of barley, buckwheat, bushberry subgroup, low growing berry subgroup (except strawberry), corn, cotton, cucurbit vegetables, dry shelled pea and bean crop subgroup, rice, millet, oats, peanuts, rapeseed subgroup 20A (including canola and

*Brassica carinata*), rye, soybean, sugar beets, triticale, wheat; conifer and hardwood nursery seeds and seedlings. When reference is made to disease suppression, suppression can mean either erratic control from good to fair or consistent control at a level below that obtained with the best commercial disease control products.

BARLEY	
Disease Controlled	Rate of Maxunitech Prothio 480 SC
Fusarium Head Blight ( <i>Fusarium</i> spp.) (Suppression Only)	5.0-5.7 fl oz/A (0.156-0.178 lb ai/A)
<b>APPLICATION TIMING:</b> The optimal time to apply Maxunitech Prothio 480 SC is as a preventative foliar spray when barley heads on the main stem are fully emerged (~ Feekes Growth Stages 10.5). Spray equipment must be set to provide good coverage to barley heads. For thorough coverage of barley head using ground application equipment, it is advisable to use forward and backward mounted nozzles or nozzles that have a two-directional spray. Operate nozzles within the spray pressure directions suggested by the manufacturer.	
<b>Leaf and Stem Diseases:</b> Net Blotch ( <i>Pyrenophora teres</i> ) Powdery Mildew ( <i>Blumeria graminis</i> f. sp. <i>Hordei</i> ) Rusts ( <i>Puccinia</i> spp.) Scald ( <i>Rhynchosporium secalis</i> ) Spot Blotch ( <i>Cochliobolus sativus</i> )	2.8-4.3 fl oz/A (0.088-0.134 lb ai/A)
<b>APPLICATION TIMING:</b> Apply Maxunitech Prothio 480 SC as a preventative foliar spray when the earliest disease symptoms appear on the leaves or stems. Observe barley fields closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.	
Maxunitech Prothio 480 SC may be applied by either ground, aerial, or chemigation application equipment. For aerial applications made prior to heading (prior to Feekes Growth Stage 10.5), apply a minimum of 2 gpa spray solution. For aerial applications made at the heading growth stage or later, apply in a minimum of 5 gpa spray solution. Chemigation use is allowed only for applications made prior to heading.	
For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Maxunitech Prothio 480 SC.	
<b>USE RESTRICTIONS:</b> Apply up to two (2) applications of Maxunitech Prothio 480 SC per year. Repeat applications using a 14-day spray interval if conditions remain favorable for continued or increasing disease development. A maximum of 9.37 fl oz (0.293 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. <b>DO NOT</b> apply within 32 days of harvest.	

BUSHBERRY SUBGROUP (SUBGROUP 13-07B)	
Aronia berry; blueberry (highbush and lowbush); Chilean guava; highbush cranberry; currant (black, buffalo, and red); elderberry; European barberry; gooseberry; edible honeysuckle; huckleberry; jostaberry; juneberry (Saskatoon berry); lingonberry; native currant; salal; sea buckthorn; and cultivars, varieties, and/or hybrids of these.	
Disease Controlled	Rate of Maxunitech Prothio 480 SC
Septoria Leaf Spot ( <i>Septoria</i> spp.) Monilinia Blight ( <i>Monilinia vaccinii-corymbosi</i> ) Valdensinia Leaf Spot ( <i>Valdensinia heterodoxa</i> ) Leaf Rust ( <i>Thekopsora minima</i> ) Anthracnose ( <i>Colletotrichum gloeosporioides</i> ) Botrytis Blight ( <i>Botrytis cinerea</i> ) Phomopsis Canker and Twig Blight <sup>[1]</sup> ( <i>Phomopsis vaccinii</i> ) Alternaria Fruit Rot <sup>[1]</sup> ( <i>Alternaria</i> spp.) White Pine Blister Rust <sup>[1]</sup> ( <i>Cronartium ribicola</i> )	5.7 fl oz/A (0.178 lb ai/A)
<b>APPLICATION TIMING:</b> Apply Maxunitech Prothio 480 SC at the first sign of disease.	
Maxunitech Prothio 480 SC may be applied by either ground or chemigation application equipment.	
<b>USE RESTRICTIONS:</b> Apply up to two (2) applications of Maxunitech Prothio 480 SC per year. Repeat applications as needed using a 7- to 10-day spray interval if conditions remain favorable for continued or increasing disease development. A maximum of 11.4 fl oz (0.356 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. <b>DO NOT</b> apply within 7 days of harvest.	
[ <sup>1</sup> Not registered for use by California.]	

<b>LOW GROWING BERRY SUBGROUP, EXCEPT STRAWBERRY (SUBGROUP 13-07H) <sup>[1]</sup></b>	
Bearberry; bilberry; cloudberry; cranberry; muntries; partridgeberry; and cultivars, varieties, and/or hybrids of these	
<b>Disease Controlled</b>	<b>Rate of Maxunitech Prothio 480 SC</b>
<b>Fruit Rot:</b> <i>Coleophoma empetri</i> <i>Glomerella cingulata</i> <i>Phyllosticta vaccinii</i> <i>Physalospora vaccinii</i> <i>Allantophomopsis lycopodina</i> <i>Allantophomopsis cytispora</i> <i>Fusicoccum putrefaciens</i> <i>Penicillium</i> spp. <i>Phomopsis vaccinii</i> <i>Colletotrichum acutatum</i> <i>Botrytis</i> spp. <i>Monilinia</i> spp. Valdensinia Leaf Spot ( <i>Valdensinia heterodoxa</i> )	5.0 fl oz/A (0.156 lb ai/A)
<b>APPLICATION TIMING:</b> For best control of fruit rots begin applications of Maxunitech Prothio 480 SC at early bloom. Make a second application of Maxunitech Prothio 480 SC or another approved fungicide 7-10 days later.	
Maxunitech Prothio 480 SC may be applied by either ground or chemigation application equipment.	
<b>USE RESTRICTIONS:</b> Apply up to two (2) applications of Maxunitech Prothio 480 SC per year. Repeat applications as needed using a 7- to 10-day spray interval if conditions remain favorable for continued or increasing disease development. A maximum of 10.0 fl oz (0.313 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. <b>DO NOT</b> apply within 45 days of harvest.	
[ <sup>1</sup> Not registered for use by California.]	

<b>BUCKWHEAT, MILLET (PEARL), MILLET (PROSO), OATS, RYE</b>	
<b>Disease Controlled</b>	<b>Rate of Maxunitech Prothio 480 SC</b>
Rusts ( <i>Puccinia</i> spp.) Glume Blotch ( <i>Stagonospora nodorum</i> ) Head Blight or Scab ( <i>Fusarium graminearum</i> ) – Suppression Powdery Mildew ( <i>Erysiphe graminis</i> ) Scald ( <i>Rhynchosporium secalis</i> ) Speckled Blotch ( <i>Septoria avenae</i> ; <i>Septoria tritici</i> ) Spot Blotch ( <i>Bipolaris sorokiniana</i> ) Tan Spot or Yellow Leaf Spot ( <i>Pyrenophora tritici-repentis</i> )	5.0-5.7 fl oz/A (0.156-0.178 lb ai/A)
<b>APPLICATION TIMING:</b> Apply Maxunitech Prothio 480 SC as a preventative foliar spray when the earliest disease symptoms appear on the leaves or stems. Observe fields closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.	
Maxunitech Prothio 480 SC may be applied by either ground or aerial spray equipment.	
<b>USE RESTRICTIONS:</b> Apply only one (1) application of Maxunitech Prothio 480 SC per year. A maximum of 5.7 fl oz (0.178 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. <b>DO NOT</b> apply within 30 days of harvest.	

<b>CORN (FIELD CORN, FIELD CORN GROWN FOR SEED AND POPCORN)</b>	
<b>Disease Controlled</b>	<b>Rate of Maxunitech Prothio 480 SC</b>
Anthracnose Leaf Blight ( <i>Colletotrichum graminicola</i> ) Eye Spot ( <i>Aureobasidium zeae</i> ) Gray Leaf Spot ( <i>Cercospora zeae-maydis</i> ) Northern Corn Leaf Blight ( <i>Setosphaeria turcica</i> ) <sup>1</sup> Northern Corn Leaf Spot ( <i>Cochliobolus carbonum</i> ) <sup>1</sup> Rust ( <i>Puccinia</i> spp.) Southern Corn Leaf Blight ( <i>Cochliobolus heterostrophus</i> ) <sup>1</sup>	5.7 fl oz/A (0.178 lb ai/A)
<b>APPLICATION TIMING:</b> Apply Maxunitech Prothio 480 SC at the first sign of disease. Repeat applications as needed on a 7- to 14-day interval if favorable conditions for disease development persist. <b>DO NOT</b> use adjuvants in sprays made between V8 (8 leaf collar) and VT (lowest branch of the tassel is visible, but the silks have not yet emerged).	
<b>SUPPRESSION ONLY:</b> Fusarium <sup>[2]</sup> Ear Rot ( <i>Fusarium</i> spp.) Gibberella <sup>[2]</sup> Ear Rot ( <i>Gibberella</i> spp.) Aspergillus <sup>[2]</sup> Ear Rot ( <i>Aspergillus</i> spp.)	5.7 fl oz/A (0.178 lb ai/A)
<b>APPLICATION TIMING:</b> For optimum suppression, apply Maxunitech Prothio 480 SC from the R1 (initial silk emergence) to the R2 (brown silk) corn growth stages. Maxunitech Prothio 480 SC will reduce both disease symptoms and levels of mycotoxin in the grain.	
<b>Soilborne/Seedling Diseases:</b> Rhizoctonia Root <sup>[2]</sup> and Stalk Rot <sup>[2]</sup> ( <i>Rhizoctonia solani</i> )	2.6-4.0 fl oz/A (0.081-0.125 lb ai/A)
<b>APPLICATION TIMING:</b> In furrow application for soilborne/seedling disease control, see use directions and rate chart for various row spacing under the SOILBORNE/SEEDLING DISEASE CONTROL section.	
Maxunitech Prothio 480 SC may be applied by either ground, aerial, or chemigation application equipment. For aerial applications apply Maxunitech Prothio 480 SC using a minimum of 3 gpa spray solution. An adjuvant may be used to improve spray coverage. Refer to the adjuvant product label for specific use directions.	
Application of Maxunitech Prothio 480 SC is not advised at times when corn is under severe environmental stress conditions.	
<b>USE RESTRICTIONS:</b> <b>DO NOT</b> exceed 22.8 fl oz (0.713 lb prothioconazole) total from all uses, including soil and foliar applications, per acre of Maxunitech Prothio 480 SC per year. <b>DO NOT</b> apply within 14 days of harvest for grain and fodder. Forage may be harvested the same day of application.	
<sup>1</sup> The above diseases are also known as Helminthosporium leaf blights. <sup>[2]</sup> Not registered for use by California.	

<b>COTTON</b>	
<b>Disease Suppressed</b>	<b>Rate of Maxunitech Prothio 480 SC</b>
<b>In Furrow and Banded:</b> Damping Off ( <i>Rhizoctonia solani</i> ) Fusarium Wilt <sup>[1]</sup> ( <i>Fusarium</i> spp.)	5.7-7.1 fl oz/A (0.178-0.222 lb ai/A) (0.4-0.5 fl oz per 1000 row feet)
<b>APPLICATION DIRECTIONS:</b> For in-furrow applications: Apply 5.7-7.1 fl oz per acre (0.393-0.489 fl oz per 1000 row feet if on 36 inch row spacing) in the furrow at planting. For banded applications: Maxunitech Prothio 480 SC may also be applied in a 4- to 6-inch band over the row at or near emergence.	
<b>Foliar:</b> Target Spot ( <i>Corynespora cassicola</i> ) Rust ( <i>Puccinia</i> spp.)	5.0-5.7 fl oz/A (0.156-0.178 lb ai/A)
<b>APPLICATION DIRECTIONS:</b> Maxunitech Prothio 480 SC may be applied by ground, aerial, or chemigation application equipment.	
<b>USE RESTRICTIONS:</b> When used at 5.0-5.7 fl oz (0.156-0.178 lb ai) per acre, <b>DO NOT</b> apply more than three (3) applications of Maxunitech Prothio 480 SC per year including the in-furrow, banded and foliar applications. If any application is made at a rate above 5.7 fl oz (0.178 lb ai) per acre, <b>DO NOT</b> make more than two (2) total applications per year including in-furrow, banded, and foliar applications. Repeat applications as needed using a 14-day spray interval if conditions remain favorable for continued or increasing diseases development. A maximum of 17.1 fl oz (0.534 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year including all soil and foliar applications. <b>DO NOT</b> apply within 30 days of harvest.	
<sup>[1]</sup> Not registered for use by California.	

CHICKPEA	
Disease Controlled	Rate of Maxunitech Prothio 480 SC
Ascochyta Blight ( <i>Ascochyta</i> spp.)	5.0-5.7 fl oz/A (0.156-0.178 lb ai/A)
<b>APPLICATION TIMING:</b> Apply Maxunitech Prothio 480 SC at the first sign of disease. Use the higher use rate when conditions are favorable for severe disease pressure and/or when growing susceptible varieties. Maxunitech Prothio 480 SC may be applied by either ground, aerial or chemigation application equipment.	
<b>USE RESTRICTIONS:</b> Apply up to three (3) applications of Maxunitech Prothio 480 SC per year. Repeat applications as needed using a 10- to 14-day spray interval if conditions remain favorable for continued or increasing disease development. To optimize disease control, tank mix the lowest labeled rate of a spray surfactant with Maxunitech Prothio 480 SC. A maximum of 17.1 fl oz (0.534 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. Allow a minimum of 7 days from the last application until cutting or swathing the crop for harvest.	

CUCURBIT VEGETABLES (CROP GROUP 9)	
Chayote (fruit) ; Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; edible gourd (includes hyotan, cucuzza, hechima, Chinese okra); <i>Momordica</i> spp. (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon (includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); pumpkin; squash (summer and winter, includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini, butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon	
Disease Controlled	Rate of Maxunitech Prothio 480 SC
Fusarium Wilt Fusarium Blight ( <i>Fusarium oxysporum</i> ; <i>Fusarium</i> spp.) Gummy Stem Blight ( <i>Didymella</i> spp.) Southern Blight <sup>[1]</sup> ( <i>Sclerotium rolfii</i> ) Powdery Mildew ( <i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i> ) ( <i>Erysiphe cichoracearum</i> )	<b>Soil:</b> 5.7 fl oz/A (0.178 lb ai/A) <b>Foliar:</b> 5.7 fl oz/A (0.178 lb ai/A)
<b>APPLICATION DIRECTIONS:</b> Maxunitech Prothio 480 SC may be applied by either ground or chemigation application equipment (including drip irrigation). <b>DO NOT</b> use in water used for hand transplanting. Not for use in greenhouse/transplant house.	
<b>USE RESTRICTIONS:</b> Apply up to one (1) soil application and two (2) foliar applications of Maxunitech Prothio 480 SC per year. Repeat applications as needed using a 5- to 10-day spray interval if conditions remain favorable for continued or increasing disease development. A maximum of 17.1 fl oz (0.534 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. <b>DO NOT</b> apply within 7 days of harvest.	
<sup>[1]</sup> Not registered for use by California.	



<b>DRIED SHELLED PEAS AND BEANS SUBGROUP (EXCEPT SOYBEANS) (SUBGROUP 6C)</b>	
<i>Lupinus</i> spp. (grain, sweet, white and white sweet lupins); <i>Phaseolus</i> spp. (field, kidney, dry lima, navy, pinto and tepary beans); <i>Vigna</i> spp. (adzuki bean, blackeye pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea and urd bean); dry broad bean; guar; lablab bean; <i>Pisum</i> spp. (pea (including field pea) and pigeon pea)	
<b>Disease Controlled</b>	<b>Rate of Maxunitech Prothio 480 SC</b>
<b>In Furrow:</b> Rhizoctonia rots <sup>[1]</sup> ( <i>Rhizoctonia</i> spp.)	2.6-5.0 fl oz/A (0.081-0.156 lb ai/A)
<b>APPLICATION DIRECTIONS:</b> Apply up to 5.0 fl oz (0.156 lb ai) per acre (0.288 fl oz per 1000 row feet if on 30 inch row spacing; 0.192 fl oz per 1000 row feet if on a 20 inch row spacing) in the furrow at planting. See use directions and rate chart for various row spacing under the SOILBORNE/SEEDLING DISEASE CONTROL section.	
<b>Foliar:</b> Ascochyta Blight ( <i>Ascochyta pinodes</i> ) Rust ( <i>Uromyces appendiculatus</i> ) White Mold ( <i>Sclerotinia sclerotiorum</i> )	5.7 fl oz/A (0.178 lb ai/A)
<b>APPLICATION DIRECTIONS:</b> For rust control, apply Maxunitech Prothio 480 SC at the first sign of disease. For white mold control, apply Maxunitech Prothio 480 SC at 25% flower.	
Maxunitech Prothio 480 SC may be applied by either ground, aerial or chemigation application equipment. For ground applications, apply in a minimum of 20 gpa.	
<b>USE RESTRICTIONS:</b> Apply up to three (3) applications of Maxunitech Prothio 480 SC per year. Repeat applications as needed using a 5- to 14-day spray interval if conditions remain favorable for continued or increasing disease development. To optimize disease control, tank mix the lowest labeled rate of a spray surfactant with Maxunitech Prothio 480 SC. A maximum of 17.1 fl oz (0.534 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. Allow a minimum of 7 days from the last application until cutting or swathing the crop for harvest.	
[ <sup>1</sup> Not registered for use by California.]	

<b>LENTILS</b>	
<b>Disease Controlled</b>	<b>Rate of Maxunitech Prothio 480 SC</b>
Ascochyta Blight ( <i>Ascochyta</i> spp.)	4.3-5.7 fl oz/A (0.134-0.178 lb ai/A)
<b>APPLICATION TIMING:</b> Apply Maxunitech Prothio 480 SC at early flower or the first sign of disease. Use the higher use rate when conditions are favorable for severe disease pressure and/or when growing less disease resistant varieties.	
Maxunitech Prothio 480 SC may be applied by either ground, aerial or chemigation application equipment.	
<b>USE RESTRICTIONS:</b> Apply up to three (3) applications of Maxunitech Prothio 480 SC per year. Repeat applications as needed using a 10- to 14-day spray interval if conditions remain favorable for continued or increasing disease development. To optimize disease control, tank mix the lowest labeled rate of a spray surfactant with Maxunitech Prothio 480 SC. A maximum of 17.1 fl oz (0.534 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. Allow a minimum of 7 days from the last application until cutting or swathing the crop for harvest.	

<b>RAPESEED SUBGROUP 20A (INCLUDING CANOLA AND BRASSICA CARINATA<sup>[1]</sup>)</b>	
Borage; <i>Brassica carinata</i> ; crambe; cuphea; echium; flax; gold of pleasure; hare's ear mustard; lesquerella; lunaria; meadowfoam; milkweed; mustard seed; oil radish; poppy seed; rapeseed; sesame; sweet rocket; cultivars, varieties, and/or hybrids of these	
<b>Disease Controlled</b>	<b>Rate of Maxunitech Prothio 480 SC</b>
Sclerotinia Stem Rot White Mold ( <i>Sclerotinia sclerotiorum</i> )	4.3-5.7 fl oz/A (0.134-0.178 lb ai/A)
<b>APPLICATION TIMING:</b> Apply Maxunitech Prothio 480 SC when the crop is in the 20-50% bloom stage. Best protection will be achieved when the fungicide is applied prior to petals beginning to fall, and will allow for the maximum number of petals to be protected. Utilize the higher rate for fields with a history of heavy disease pressure or for dense crop stands. Good spray coverage of the plants is essential. Maxunitech Prothio 480 SC may be applied until the 50% bloom stage.	
Maxunitech Prothio 480 SC may be applied by either ground, aerial or chemigation application equipment.	
The lowest labeled rate of a non-ionic surfactant may be tank-mixed with Maxunitech Prothio 480 SC.	
<b>USE RESTRICTIONS:</b> Apply up to two (2) applications of Maxunitech Prothio 480 SC per year. A second application may be made after 14 days if conditions remain favorable for continued or increasing disease development. A maximum of 11.4 fl oz (0.356 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. <b>DO NOT</b> apply within 36 days of harvest.	
[ <sup>1</sup> Not registered for use by California.]	



PEANUT <sup>[1]</sup>	
Disease Suppressed	Rate of Maxunitech Prothio 480 SC
<b>In Furrow and Banded:</b> Sclerotium Rot, White Mold, Southern Blight, Southern Stem Rot ( <i>Sclerotium rolfsii</i> ) Rhizoctonia Limb Rot ( <i>Rhizoctonia solani</i> ) Early Leaf Spot ( <i>Cercospora arachidicola</i> ) Late Leaf Spot ( <i>Cercosporidium personatum</i> ) Cylindrocladium Black Rot (CBR) ( <i>Cylindrocladium crotalariae</i> ) (Suppression with in furrow only)	5.7 fl oz/A (0.178 lb ai/A) (0.4 fl oz per 1000 row feet)
<b>APPLICATION TIMING:</b> Apply 5.7 fl oz (0.178 lb ai) per acre (0.4 fl oz per 1000 row feet if on 36 inch row spacing) in the furrow at planting. Maxunitech Prothio 480 SC may also be applied in a 4- to 6- inch band over the row at or near emergence. Maxunitech North America, Inc. advises a minimum application volume of 20 gpa.	
<b>Soil-Borne:</b> Sclerotium Rot( <i>Sclerotium rolfsii</i> ), (White Mold, Southern Blight, Southern Stem Rot) Rhizoctonia Limb Rot, Peg Rot, Pod Rot ( <i>Rhizoctonia solani</i> ) Cylindrocladium Black Rot (CBR) ( <i>Cylindrocladium crotalariae</i> ) (Suppression only)	5.7 fl oz/A (0.178 lb ai/A)
<b>APPLICATION TIMING:</b> For optimum control of the specified soil-borne diseases, four (4) consecutive applications of Maxunitech Prothio 480 SC must be made at 14-day intervals. In a typical 7-spray application program beginning 30-40 days after planting or as advised by the local Extension Service, apply Maxunitech Prothio 480 SC for sprays 3, 4, 5, and 6. Make application of fungicides with a different mode of action prior to and following applications of Maxunitech Prothio 480 SC to discourage development of resistant strains of fungi. Use Maxunitech Prothio 480 SC in conjunction with cultural practices that are known to reduce the severity of soil-borne diseases, including proper crop rotation practices.  For control of soil-borne diseases when using a <b>Leaf Spot Advisory Program</b> schedule, apply Maxunitech Prothio 480 SC in the first advisory spray in July and continue Maxunitech Prothio 480 SC applications at 14-day intervals.  Maxunitech Prothio 480 SC must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by <i>Sclerotium rolfsii</i> and <i>Rhizoctonia solani</i> . Drought conditions will decrease the effectiveness of Maxunitech Prothio 480 SC against the root and pod rots.	
<b>Foliar:</b> Early Leaf Spot ( <i>Cercospora arachidicola</i> ) Late Leaf Spot ( <i>Cercosporidium personatum</i> ) Leaf Rust ( <i>Puccinia arachidis</i> ) Web Blotch ( <i>Phoma arachidicola</i> ) Leaf Scorch and Pepper Spot ( <i>Leptosphaerulina crassiasca</i> )	5.0-5.7 fl oz/A (0.156-0.178 lb ai/A)
<b>APPLICATION TIMING:</b> Apply the specified rate in a preventative spray schedule. Apply up to four (4) sprays using a 14-day interval. Use the higher use rate when conditions are favorable for severe disease pressure and/or when growing less disease resistant varieties.	
Maxunitech Prothio 480 SC may be applied by either ground, aerial, or chemigation application equipment.  <b>USE RESTRICTIONS:</b> Apply up to four (4) applications of Maxunitech Prothio 480 SC per year, including the in-furrow and banded applications. When planting varieties with good to excellent levels of resistance to foliar diseases, the application interval may be extended up to 21 days in the absence of soil borne diseases. A maximum of 22.8 fl oz (0.713 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. Maxunitech Prothio 480 SC may be applied up to 14 days before harvest. <b>DO NOT</b> feed hay or threshings or allow livestock to graze in treated areas.	
<sup>[1]</sup> Not registered for use by California.	

RICE <sup>[1]</sup>	
Disease Controlled	Rate of Maxunitech Prothio 480 SC
<b>Sheath/Stem Diseases:</b> Sheath Blight ( <i>Rhizoctonia solani</i> ) <b>Foliar Diseases:</b> Brown Spot ( <i>Cochliobolus miyabeanus</i> ) Narrow Brown Leafspot ( <i>Cercospora oryzae</i> ) Leaf Smut ( <i>Entyloma oryzae</i> ) False Smut ( <i>Ustilaginoidea virens</i> )	4.5 fl oz/A (0.141 lb ai/A)
<b>APPLICATION TIMING:</b> Apply Maxunitech Prothio 480 SC at initial sign of disease. Exact timing for rice disease control is dependent on rice growth stage, rice variety, the type of disease to be controlled and disease severity. Applications typically will occur from panicle differentiation to late boot. Consult with your local extension personnel or Maxunitech North America, Inc. representative to determine if treatment is needed.	
Maxunitech Prothio 480 SC may be applied by either ground or aerial spray equipment.	
<b>USE RESTRICTIONS:</b> Apply only one (1) application of Maxunitech Prothio 480 SC per year. A maximum of 4.5 fl oz (0.141 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. DO NOT apply within 40 days of harvest. <b>DO NOT</b> apply Maxunitech Prothio 480 SC later than 70% panicle emergence from the boot.	
<sup>[1]</sup> Not registered for use by California.	

SOYBEAN	
Disease Controlled	Rate of Maxunitech Prothio 480 SC
Asian Soybean Rust ( <i>Phakopsora pachyrhizi</i> ) Frog Eye Leaf Spot ( <i>Cercospora sojina</i> ) Powdery Mildew ( <i>Microsphaera diffusa</i> ) Brown Spot <sup>[1]</sup> ( <i>Septoria glycines</i> )	2.5-3.0 fl oz/A (0.078-0.094 lb ai/A)
Alternaria Leaf Spot <sup>[1]</sup> ( <i>Alternaria</i> spp.) Anthracnose <sup>[1]</sup> ( <i>Colletotrichum truncatum</i> ) Asian Soybean Rust <sup>[1]</sup> ( <i>Phakopsora pachyrhizi</i> ) Brown Spot <sup>[1]</sup> ( <i>Septoria glycines</i> ) Cercospora Blight <sup>[1]</sup> ( <i>Cercospora kikuchii</i> ) Frog Eye Leaf Spot <sup>[1]</sup> ( <i>Cercospora sojina</i> ) Pod & Stem Blight <sup>[1]</sup> ( <i>Diaporthe phaseolorum</i> ) Powdery Mildew <sup>[1]</sup> ( <i>Microsphaera diffusa</i> ) Rhizoctonia Aerial Blight <sup>[1]</sup> ( <i>Rhizoctonia solani</i> ) Sclerotinia Stem Rot AKA White Mold ( <i>Sclerotinia sclerotiorum</i> ) (Suppression only)	3.0-5.0 fl oz/A (0.094-0.156 lb ai/A)
<b>APPLICATION TIMING:</b> Apply Maxunitech Prothio 480 SC as a broadcast, preventative foliar spray or at first visible symptoms of the disease. Repeat applications on a 10- to 21-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rate and shorter spray intervals are advised when disease pressure is severe.	
<b>Sclerotinia Stem Rot (Suppression Only):</b> Apply Maxunitech Prothio 480 SC as a broadcast foliar spray at R1 (beginning bloom) when conditions are favorable for disease development. A sequential treatment of Maxunitech Prothio 480 SC or other registered fungicide (Prothioconazole+Trifloxystrobin) may be made at R3-R4 (beginning to full pod). Maxunitech Prothio 480 SC may be applied by ground or air. Apply in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft equipment.	
<b>In Furrow:</b> Rhizoctonia Rots <sup>[1]</sup> ( <i>Rhizoctonia</i> spp.)	2.6-5.0 fl oz/A (0.081-0.156 lb ai/A)
<b>APPLICATION TIMING:</b> Apply up to 5.0 fl oz (0.156 lb ai) per acre (0.288 fl oz per 1000 row feet if on 30 inch row spacing; 0.192 fl oz per 1000 row feet if on a 20 inch row spacing) in the furrow at planting. See use directions and rate chart for various row spacing under the SOILBORNE/SEEDLING DISEASE CONTROL section.	
Maxunitech Prothio 480 SC may be applied by either ground, aerial, or chemigation application equipment. For aerial application, apply in a minimum spray volume of 2 gpa.	
<b>USE RESTRICTIONS: DO NOT</b> apply more than three (3) applications per year including all soil and foliar applications. A maximum of 12.9 fl oz (0.403 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. Applications may not be made within 21 days of harvest.	
<sup>[1]</sup> Not registered for use by California.	

SUGAR BEETS	
Disease Controlled	Rate of Maxunitech Prothio 480 SC
<b>In Furrow and Banded:</b> Rhizoctonia Stem Canker, Root Rot, Crown Rot ( <i>Rhizoctonia solani</i> )	5.7 fl oz/A (0.178 lb ai/A)
<b>APPLICATION TIMING:</b> For in-furrow use: Apply up to 5.7 fl oz (0.178 lb ai) per acre (0.4 fl oz per 1000 row feet if on 36 inch row spacing) in the furrow at planting. See use directions and rate chart for various row spacing under the SOILBORNE/SEEDLING DISEASE CONTROL section. For soil-borne disease control: Apply Maxunitech Prothio 480 SC in a seven-inch band at the 4-leaf to row closure growth stage.	
<b>Foliar:</b> Cercospora Leaf Spot ( <i>Cercospora beticola</i> ) Powdery Mildew ( <i>Erysiphe polygoni</i> )	5.0-5.7 fl oz/A (0.156-0.178 lb ai/A)
<b>APPLICATION TIMING:</b> Apply Maxunitech Prothio 480 SC at the first sign of disease. Use the higher use rate and shorter intervals when conditions are favorable for severe disease pressure and/or when growing less disease resistant varieties. Maxunitech Prothio 480 SC may be applied by either ground, aerial or chemigation application equipment.  To optimize disease control, the lowest labeled rate of a spray surfactant may be tank-mixed with Maxunitech Prothio 480 SC.  <b>USE RESTRICTIONS:</b> Apply up to three (3) applications of Maxunitech Prothio 480 SC per year including all soil and foliar applications. Repeat applications as needed using a 14- to 21-day spray interval depending on disease pressure. Use a 14-day spray interval under normal to heavy disease pressure and a 21-day spray interval under light disease pressure. A maximum of 17.1 fl oz (0.534 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. Allow a minimum of 7 days from the last application before harvesting. Maxunitech Prothio 480 SC is a Group 3 fungicide. To limit the potential for development of disease resistance alternate every application of Maxunitech Prothio 480 SC with a non-Group 3 fungicide.	

WHEAT (SPRING, DURUM AND WINTER), TRITICALE	
Disease Controlled	Rate of Maxunitech Prothio 480 SC
Fusarium Head Blight ( <i>Fusarium</i> spp.) (Suppression Only)	5.0-5.7 fl oz/A (0.156-0.178 lb ai/A)
<b>APPLICATION TIMING:</b> The optimal time to apply Maxunitech Prothio 480 SC is as a preventative foliar spray at early flower (Feekes Growth Stage 10.51). Spray equipment must be set to provide good coverage to wheat heads. For thorough coverage of the wheat head using ground application equipment, use forward and backward mounted nozzles or nozzles that have a two-directional spray. Operate nozzles within the spray pressure directions suggested by the manufacturer.	
<b>Leaf and Stem Diseases:</b> Powdery Mildew ( <i>Blumeria graminis</i> f. sp. <i>tritici</i> ) Rusts ( <i>Puccinia</i> spp.) Septoria Leaf and Glume Blotch ( <i>Septoria tritici</i> ) Stagonospora Blotch ( <i>Stagonospora nodorum</i> ) Tan Spot ( <i>Pyrenophora tritici-repentis</i> )	4.3-5.0 fl oz/A (0.134-0.156 lb ai/A)
<b>APPLICATION TIMING:</b> Apply Maxunitech Prothio 480 SC as a preventative foliar spray when the earliest disease symptoms appear on the leaves or stems. Observe wheat fields closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.  Maxunitech Prothio 480 SC may be applied by either ground, aerial or chemigation application equipment. For aerial application made prior to early flower (prior to Feekes Growth Stage 10.51), apply a minimum of 2 gpa spray solution. For aerial applications made at the early flower growth state or later, apply in a minimum of 5 gpa spray solution. Chemigation use is allowed only for applications made prior to early flower.  For optimal disease control, the lowest labeled rate of a spray surfactant may be tank-mixed with Maxunitech Prothio 480 SC.  <b>USE RESTRICTIONS:</b> Apply up to two (2) applications of Maxunitech Prothio 480 SC per year. Repeat applications using a 14-day spray interval if conditions remain favorable for continued or increasing disease development. A maximum of 9.37 fl oz (0.293 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per year. <b>DO NOT</b> apply within 30 days of harvest.	

NURSERY SEEDLINGS	
Shortleaf, loblolly, slash, longleaf and other pines and other conifers and hardwoods	
Disease Controlled	Rate of Maxunitech Prothio 480 SC
Fusiform Rust ( <i>Cronartium quercum</i> f. sp. <i>Fusiforme</i> ) Pitch Canker ( <i>Fusarium</i> spp.) Rhizoconia Foliar Blight ( <i>Rhizoconia</i> spp.)	5.0 fl oz/A (0.156 lb ai/A)
<b>APPLICATION TIMING:</b> Foliar disease control: Apply Maxunitech Prothio 480 SC preventatively or at the first sign of disease using ground equipment only. Repeat applications as needed using a 14- to 21-day spray interval depending on your region. Consult your local extension agent on locally advised spray intervals. Use shorter intervals when conditions are favorable for severe disease pressure and/or when growing less disease resistant varieties. To optimize disease control, the lowest labeled rate of a spray surfactant may be tank-mixed with Maxunitech Prothio 480 SC.	
The crop safety and mix compatibility on all tree species and in tank-mixes with other products (spray surfactants, fertilizers, insecticides, etc.) has not been confirmed. Maxunitech North America, Inc. advises small scale testing with your planned use pattern. The user assumes all risks with the use of this product on trees.	
<b>USE RESTRICTIONS: DO NOT use in forest planting or established woodlands.</b> A maximum of 25 fl oz (0.781 lb prothioconazole) of Maxunitech Prothio 480 SC may be applied per acre per crop year.	

NURSERY SEEDS	
Shortleaf, loblolly, slash, longleaf and other pines and other conifers and hardwoods	
Disease Controlled	Rate of Maxunitech Prothio 480 SC
Fusiform Rust ( <i>Cronartium quercum</i> f. sp. <i>Fusiforme</i> ) Pitch Canker ( <i>Fusarium</i> spp.)	10 fl oz per 50 lb seed
<b>APPLICATION DIRECTIONS:</b> Apply specified dosage to seed in a commercial treater or other suitable tumbler apparatus. Allow to mix for at least 10 minutes. Thoroughly air dry before sowing.	
The crop safety and mix compatibility on all tree species and in tank-mixes with other products (spray surfactants, fertilizers, insecticides, etc.) has not been confirmed. Maxunitech North America, Inc. advises small scale testing with your planned use pattern. The user assumes all risks with the use of this product on trees.	
<b>USE RESTRICTIONS: DO NOT use in forest planting or established woodlands. DO NOT use treated seed for food or feed purposes.</b> Seed that has been treated with this product that is then packaged or bagged for future use must contain the following labeling found in the SEED TAG LABELING section of this label.	

### SEED TAG LABELING

The Federal Seed Act requires that the container of seed treated with Maxunitech Prothio 480 SC must be labeled with the following statements:

- This seed has been treated with Maxunitech Prothio 480 SC, a fungicide which contains 41.48% Prothioconazole.
- **DO NOT** Use for Food, Feed, or Oil Purposes.

In addition, the following statements are required on the container of seed treated with Maxunitech Prothio 480 SC:

- Treated areas may be replanted with any crop specified on this label as soon as practical after last application. For crops not listed on this label, **DO NOT** plant back within 30 days of last application.
- **Store treated seeds away from food and feedstuffs.**
- **DO NOT allow children, pets, or livestock to have access to treated seed.**
- **When opening this bag or loading/pouring the treated seed/seed-pieces, wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves.**
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting (such as in row ends).
- Dispose of all excess treated seed. Leftover treated seed may be buried away from water sources in accordance with local requirements.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash waters.
- Treated at manufacturer's recommended rate.
- Treated seed must be adequately covered with soil at planting. Plant at a minimum depth of ½ inch.

- **DO NOT** plant treated seed by broadcasting to the soil surface. Ensure that all planted seeds are thoroughly incorporated by the planter during planting, additional incorporation may be required to thoroughly cover exposed seeds.
- Dispose of seed packaging or containers in accordance with local requirements. **DO NOT** use empty seed bags for any other purpose.
- To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's recommendations. Surplus seed or empty seed containers must be stored or disposed according to local federal regulations.
- After the seeds have been planted, **DO NOT** enter or allow worker entry into the treated areas during the restricted-entry interval (REI) of 12 hours. Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with the treated seeds in the soil or with the planting media.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage, disposal or cleaning of equipment.

### PESTICIDE STORAGE

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. **DO NOT** walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

### PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed on site or at an approved waste disposal facility.

### CONTAINER HANDLING:

**[Nonrefillable Plastic Containers 5 Gallons or Less:** Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

**[Nonrefillable Plastic Containers Larger Than 5 Gallons:** Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

**[Refillable Plastic Containers Larger Than 5 Gallons:** Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. When this container is empty, replace the cap and seal all openings that have been made during usage and return the container to the point of purchase, or to an alternate location designated by the manufacturer at the time of purchase of this product. If not returned, clean container the empty container and offer for recycling, if available. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the re-filler. To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or re-circulate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times. If the container cannot be refilled, offer for

recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

**[Nonrefillable Plastic and Metal Containers (e.g., Intermediate Bulk Containers [IBC]) (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down):** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into formulation equipment and before final disposal using the following pressure rinsing procedure: Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure spray duration and/or spray volume. If the manufacturer's instructions are not available pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain pour or pump rinsate into formulation equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.]

#### **LIMITATION OF WARRANTY AND LIABILITY**

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES and LIMITATIONS OF LIABILITY. CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Maxunitech North America, Inc. All such risks shall be assumed by the user or buyer. **DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Maxunitech North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Maxunitech North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Maxunitech North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Maxunitech North America, Inc.'s election, the replacement of product.

[EPA approval date]



{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

PROTHIOCONAZOLE GROUP 3 FUNGICIDE



Maxunitech Prothio 480 SC

ACTIVE INGREDIENT:	By Wt.
Prothioconazole.....	41.48%
OTHER INGREDIENTS:.....	58.52%
TOTAL:.....	100.00%

KEEP OUT OF REACH OF CHILDREN  
CAUTION

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li></ul>
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
IF INHALED:	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12 PM Pacific Standard Time. In the event of a medical emergency, call your poison control center at 1-800-222-1222.	

**For Chemical Emergency:**  
**Spill, Leak, Fire, Exposure, or Accident,**  
**Call CHEMTREC Day or Night**  
**Within USA and Canada: 1-800-424-9300**

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to estuarine/marine invertebrates, and freshwater/estuaries/marine aquatic plants. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water

when disposing of equipment washwater or rinsate.

Degradates of prothioconazole are known to leach through soil into ground water under certain conditions as a result of label use. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of prothioconazole and degrades from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. **DO NOT** walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

**[Nonrefillable Plastic Containers 5 Gallons or Less:**  
Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

**[Nonrefillable Plastic Containers Larger Than 5 Gallons:**  
Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution,



for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

**[Refillable Plastic Containers Larger Than 5 Gallons:** Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. When this container is empty, replace the cap and seal all openings that have been made during usage and return the container to the point of purchase, or to an alternate location designated by the manufacturer at the time of purchase of this product. If not returned, clean container the empty container and offer for recycling, if available. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the re-filler. To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or re-circulate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times. If the container cannot be refilled, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

**[Nonrefillable Plastic and Metal Containers (e.g., Intermediate Bulk Containers [IBC]) (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down):** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into formulation equipment and before final disposal using the following pressure

rinsing procedure: Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure spray duration and/or spray volume. If the manufacturer's instructions are not available pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain pour or pump rinsate into formulation equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.]

[See] [label] [booklet] [for] [additional] [Precautionary Statements] [and] [Directions for Use].

**EPA Reg. No.: 95009-XX**

**Net Contents:**

**EPA Est. No.:**

**Manufactured for:**

Maxunitech North America, Inc.

11601 Shadow Creek Pkwy, Suite 111-573

Pearland, TX 77584