



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

January 25, 2024

Oliver Pelz
Telone Regulatory Leader
Salt Lake Holding LLC
2211 H. H. Dow Way
Midland, MI 48674

Subject: Approval of Label Amendment; Only Indicated Changes Reviewed – Removal of reference to supplemental labeling for two California counties
Product Name: TELONE II
EPA Registration Number: 95290-1
Application Date: 8/14/2023
Case Number: 488992

Dear Oliver Pelz:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. However, EPA reviewed only the label changes highlighted, marked, or otherwise indicated on the submitted label. Any other changes to the previously approved label that were not clearly highlighted, marked, or otherwise indicated in your submission were not reviewed and may form the basis of regulatory and/or enforcement action if later discovered by the Agency. Further, submission of a label amendment application with unidentified changes may be considered a knowing submission of false information to the Agency. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

The label submitted with the application has been stamped "Accepted Only Indicated Revisions Reviewed" and is enclosed for your records.

This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 C.F.R. § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 C.F.R. § 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website contains any false or misleading statement, design, or graphic, the product may be misbranded and unlawful to sell or distribute under FIFRA Sections 2(q)(1)(A) and 12(a)(1)(E). 40 C.F.R. § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on the product label, claims made as part of the product's sale or distribution may not substantially differ from those claims approved through the registration process under FIFRA Section 12(a)(1)(B). Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the product will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Stephanie Suarez at 202-566-2918 or at Suarez.Stephanie@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to be 'Stephanie', with a long horizontal flourish extending to the right.

Stephanie Suarez, Ph.D.
Acting Product Manager 21
Fungicide Branch
Registration Division (7505T)
Office of Pesticide Programs

(Base label):

RESTRICTED USE PESTICIDE

Due to high acute inhalation toxicity and carcinogenicity.
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

TELONE™ II

Soil Fumigant

A multi-purpose liquid fumigant for preplant treatment of soil to control plant parasitic nematodes, symphylans and to help manage certain soil borne diseases in cropland

Not for use in greenhouses or other enclosed areas.

Not for formulation or manufacturing use. Do not formulate this product into other products.

Active Ingredient:	(by weight)
1,3-dichloropropene.....	97.5%
Other Ingredients	2.5%
Total	100.0%

1 gallon of TELONE™ II weighs 10.15 lb at 70°F. Contains 9.85 lb of 1,3-dichloropropene per gallon.

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See back panel for First Aid statements.

Precautionary Statements

Hazards to Humans and Domestic Animals

Hazardous Liquid and Vapor

- Do not swallow any of this product. May be fatal if swallowed.
- Do not get in eyes. Causes substantial, but temporary eye injury.
- Do not get on skin. May be fatal if absorbed through the skin. Causes skin irritation and, if confined, skin burns. May cause allergic skin reaction.
- Do not breathe vapor. May be fatal if inhaled. May cause lung, liver, and kidney damage and respiratory system irritation upon prolonged contact.
- The use of this product may be hazardous to your health. This product contains 1,3-dichloropropene, which has been determined to cause tumors in laboratory animals. Risks can be reduced by exactly following directions for use, precautionary statements, by wearing the personal protective equipment specified in this labeling.

Personal Protective Equipment (PPE)

ACCEPTED
ONLY INDICATED
REVISIONS REVIEWED

01/25/2024

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

95290-1

No label revisions other than those indicated were reported to the Agency.

Chemical-Resistant Materials: Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category H on an EPA chemical resistance category selection chart. PPE constructed of saranex, neoprene, and chlorinated polyethylene provide short-term contact or splash protection against liquid in this product. Longer-term protection is provided by PPE constructed of viton, Teflon, and EVAL barrier laminates (for example, responder suits manufactured by Life-guard or silvershield gloves manufactured by North). Where chemical-resistant materials are required, leather, canvas, or cotton materials offer no protection from this product and must not be worn as the sole article of protection when contact with this product is possible. Where coveralls are required, they must be loose-fitting and constructed of woven fabrics (e.g., tight knit cotton or cotton/polyester), non-woven fabrics (e.g., tyvek or sontara), or fabrics containing microporous Teflon.

1. Handlers Performing Tasks with Liquid Contact Potential

Tasks with liquid contact potential are tasks performed outdoors or in a well-ventilated area. They include:

- Equipment calibration or adjustment
- Equipment clean-up and repair
- Product sampling
- Any activity less than 6 feet from an unshielded pressurized hose containing this product
- Rinsate disposal
- Fumigant transfer
- Clean-up of small spills
- Preparing containers for aeration
- Any other task not otherwise listed in (2), (3), or (4) below

Handlers performing tasks with liquid contact potential must wear at minimum:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate (EVAL) or viton
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron
- A face shield or safety glasses with brow and temple shields (do not wear chemical goggles)
- A half-face respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G). See further respirator requirements in the User Safety Requirements section on this label.

2a. Handlers Performing Tasks with No Liquid Contact Potential – Broadcast, In-Bed Applications, or Applications at the Time of Bedding Except as in 2b.

Tasks with no liquid contact potential are tasks performed outdoors or in a well-ventilated area. These tasks include:

- Tractor driving
- Soil sealing
- Field activities on the day of application that do not disrupt the soil at the depth of liquid injection

Handlers performing tasks with no liquid contact potential must wear at minimum:

- Loose fitting or well ventilated long-sleeved shirt and long pants
- Shoes and socks
- A face shield or safety glasses with brow and temple shields (do not wear chemical goggles)
- A half-face respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G).
- A respirator is not required (not applicable in California) if the occupants are within an enclosed cab that is in conformance with one of the following: 1) ANSI/ASAE S525-1.1 MAY98 sections 7.1.5, 7.1.7, 7.2.3, and 9, or 2) the requirements listed in the Worker Protection Standard (WPS) for

agricultural pesticides -- 40 CFR 170.240(d)(5). The cab must be equipped with a vapor-adsorptive filter containing a minimum of 1000 grams activated charcoal. The filter must be changed after no more than 50 hours of application time. See further respirator requirements in the User Safety Requirements section on this label.

- In addition, the PPE specified in (1) for activities with direct liquid contact potential must be immediately available and must be worn if the handler is to perform any direct-contact activity with a potential for liquid contact

2b. Handlers Performing Tasks with No Liquid Contact Potential - Pre-Bed, Row Product Applications (e.g., Yetter Rig) (Not Applicable in California)

Tasks with no liquid contact potential are tasks performed outdoors or in a well-ventilated area. These tasks include:

- Tractor driving
- Soil sealing
- Field activities on the day of application that do not disrupt the soil at the depth of liquid injection

Handlers performing tasks with no liquid contact potential must wear at minimum:

- Loose fitting or well ventilated long-sleeved shirt and long pants
- Shoes and socks
- A face shield or safety glasses with brow and temple shields (do not wear chemical goggles)
- In addition, the PPE specified in (1) for activities with direct liquid contact potential must be immediately available and must be worn if the handler is to perform any direct-contact activity with a potential for liquid contact

3. Handlers in Treated Area 1 to 5 Days After Application

Only the following handler tasks may be performed in the treated area within 5 days after the application is complete:

- Assessing/adjusting the soil seal
- Assessing pest control, application technique, or application efficacy
- Sampling air or soil for this product
- Removing tarp or plastic film

All other tasks are prohibited until the 5-day period has expired.

Handlers in treated area 1 to 5 days after application must wear at minimum:

- Loose fitting or well ventilated long-sleeved shirt and long pants
- Shoes and socks
- A face shield or safety glasses with brow and temple shields (do not wear chemical goggles)
- A half-face respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G).
- A respirator is not required if the occupants are within an enclosed cab (not applicable in California) that is in conformance with one of the following: 1) ANSI/SAE S525-1.1 MAY98 sections 7.1.5, 7.1.7, 7.2.3, and 9, or 2) the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides -- 40 CFR 170.240(d)(5). The cab must be equipped with a vapor-adsorptive filter containing a minimum of 1000 grams activated charcoal. The filter must be changed after no more than 50 hours of application time. See further respirator requirements in the User Safety Requirements section on this label.
- In addition, the PPE specified in (1) for activities with direct liquid contact potential must be immediately available and must be worn if the handler is to perform any direct-contact activity with a potential for liquid contact

4. Handlers Exposed to High Concentrations

Handlers exposed to high airborne concentrations of this product, such as cleanup following large spills and exposure to this product in poorly ventilated areas, must wear at minimum:

- Chemical-resistant suit
- Chemical-resistant gloves, such as barrier laminate (EVAL) or viton
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear
- Supplied-air respirator with MSHA/NIOSH approval number prefix TC-19C **or** self-contained breathing apparatus (SCBA) with MSHA/NIOSH approval number prefix TC-13F. See further respirator requirements in the User Safety Requirements section on this label.

Note: In-tank cleaning of bulk tanks must be performed only by persons who have been specifically trained for this activity. Refer to OSHA 29 CFR Part 1910.146 and the Storage and Handling Guide.

Engineering Controls Requirements

Mechanical Transfer System: Personal protective equipment specified for Direct Contact Activities must be worn by the operator of the mechanical transfer system. The operator of the mechanical transfer system must follow instructions on proper operation of the system found in the “TELONE™ Soil Fumigants - A Guide to Application” manual. Contact your distributor for TELONE™ II for more information or for these materials.

End-Row Spillage Control: The dispensing system must shut off the feed stream when chisels are raised out of the ground. Do not stop or park near any area where dribble from chisel tips has fallen. The applicator must follow instructions on proper operation and maintenance of the system found in the “TELONE™ Soil Fumigants - A Guide to Application” manual. Contact your distributor for TELONE™ II for more information or for these materials.

- A flow shutoff device must be placed as close as is technically feasible to the fluid discharge point. This can be a ball, poppet, or diaphragm check valve, or full flow shutoff device such as an electric or pneumatically actuated valve.
- Service any system immediately if continuous drip occurs.
- If mechanical check valves and orifices are used, place the check valve above the orifice. Also, isolate the check valve from upstream pressure by installing a main line shut off or bypass valve prior to the manifold.
- Pipe diameter from check valve to injection point must not exceed 1/4 inches ID National Pipe Standard (NPS). Preferably, use the smallest diameter pipe or tubing possible which achieves the required flow rate.
- Alternate end-row spillage devices or methods such as, but not limited to, micro-bore restricted flow tubing or line purge systems may be used if they provide equal or superior control versus check valves.

With all bulk and non-bulk containers, TELONE™ II must be transferred through connecting hoses, pipes, and/or couplings sufficiently tight to prevent workers or other persons from coming in contact with liquid TELONE™ II.

- All hoses, piping, and tanks used in connection with TELONE™ II shall be of the type appropriate for use under the pressure and vacuum conditions to be encountered.
- External sight gauges shall be equipped with valves so that pipes to sight gauge can be shut off in case of breakage or leakage.
- The mechanical transfer system must be adequate to make necessary measurements of the pesticide being used.
- Shut-off devices must be installed on the exit end of all hoses and at all disconnect points to prevent leakage of TELONE™ product when the transfer is stopped and hose is removed or disconnected. A dry coupler that will **minimize pesticide leakage** must be installed at the disconnect point.
- The pressure in hoses used to move TELONE™ II beyond a pump **must not exceed** the manufacturer's maximum pressure specification.

User Safety Requirements

- 1. Respirator Requirements:** When a respirator is required for use with this product, the following criteria must be met:
 - a. Cartridges or canisters must be replaced daily or when odor or irritation from this product becomes apparent, whichever is sooner.
 - b. Respirators must be fit-tested and fit-checked using a program that conforms to OSHA's requirements (described in 29 CFR Part 1910.134).
 - c. Respirator users must be trained using a program that conforms to OSHA's requirements (described in 29 CFR Part 1910.134).
 - d. Respirator users must be examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn.
- 2. Dispose of Contaminated Clothing:** Discard clothing and other absorbent materials that have been drenched or heavily contaminated with liquid from this product. Do not reuse them.
- 3. Clean and Maintain PPE:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Wash PPE after each day's use.
- 4. Contact with Mouth:** Never siphon this product by mouth or use mouth to blow out clogged lines, nozzles, etc.
- 5. Heat Illness Avoidance:** Use measures to avoid or minimize heat illness while using this product. These measures include gradual adjustment to heat and respirator stress, fans for cooling, cooling vests, frequent breaks to cool down, frequent intake of drinking water, and maintaining weight from day to day.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. If breathing is difficult, give oxygen. Call a poison control center or doctor for further treatment advice.

If on skin or clothing: Immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Call a poison control center or doctor for treatment advice. If water is not immediately available, remove excess chemical from skin with adsorbent material such as towel or dry soil, then proceed at once to a location where water is available and thoroughly wash contaminated skin with plenty of water. Call a poison control center or doctor for treatment advice.

If in eyes: Immediately flush eyes with plenty of water. Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

Note to physician: Because rapid absorption may occur through lungs if product is aspirated and cause systemic effects, the decision to induce vomiting or not should be made by a physician. If lavage is

performed, endotracheal and/or esophageal control is suggested. Danger from lung aspiration must be weighed against toxicity when considering emptying the stomach.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsate. See Storage and Disposal section. In case of spills properly dispose of contaminated materials.

Groundwater advisory: 1,3-dichloropropene is known to move through soil and under certain conditions has the potential to reach groundwater as a result of agricultural use. Application in areas where soils are permeable and groundwater is near the surface could result in groundwater contamination. Do not apply within 100 feet of any well used for potable water. Do not apply this product within 100 feet from the edge of karst topographical features. Karst topography is identified from landscape features that result from the dissolving activity of water in carbonate rock formations (limestone, dolomite and marble). Surface features that are associated with karst topography include sinkholes, caverns, springs, and sinking or disappearing streams. In North Dakota, South Dakota, Wisconsin, Minnesota, New York, Maine, New Hampshire, Vermont, Massachusetts, Utah, and Montana: Where groundwater aquifers exist at a depth of 50 feet or less from the surface, do not apply this product where soils are Hydrologic Group A.

Reporting Ecological Incidents: To report ecological incidents, including mortality, injury, or harm to plants and animals, call 989-636-4400.

Physical or Chemical Hazards

Combustible. Do not use or store near heat or open flame.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

(Storage and Disposal for rigid containers 5 gal or less)

Storage and Disposal

Do not contaminate water, food and feed by storage and disposal.

Pesticide Storage: Store in tightly-closed original container away from dwellings. Prolonged exposure of the container to direct sunlight must be avoided. Do not allow contamination of seeds, plants, fertilizers, or other pesticide chemicals.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide and rinsates is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

Because TELONE™ II Soil Fumigant is corrosive under certain conditions, flush all application equipment with fuel oil, kerosene or a similar type of petroleum solvent immediately after use. Fill pumps and meters with new motor oil or a 50% motor oil/fuel oil mixture before storing. **Do not use water.** Dispose of rinsate by applicable Federal, state and local regulations. Never introduce rinsate or unused TELONE™ II into surface or underground water supplies.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect

rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

(Storage and Disposal for refillable rigid containers larger than 5 gal)

Storage and Disposal

Do not contaminate water, food and feed by storage and disposal.

Pesticide Storage: Store in tightly-closed original container away from dwellings. Prolonged exposure of the container to direct sunlight must be avoided. Do not allow contamination of seeds, plants, fertilizers, or other pesticide chemicals.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide and rinsates is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

Because TELONE™ II is corrosive under certain conditions, flush all application equipment with fuel oil, kerosene or a similar type of petroleum solvent immediately after use. Fill pumps and meters with new motor oil or a 50% motor oil/fuel oil mixture before storing. **Do not use water.** Dispose of rinsate by applicable Federal, state and local regulations. Never introduce rinsate or unused TELONE™ II into surface or underground water supplies.

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)

Storage and Disposal

Do not contaminate water, food and feed by storage and disposal.

Pesticide Storage: Store in tightly-closed original container away from dwellings. Prolonged exposure of the container to direct sunlight must be avoided. Do not allow contamination of seeds, plants, fertilizers, or other pesticide chemicals.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide and rinsates is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

Because TELONE™ II is corrosive under certain conditions, flush all application equipment with fuel oil, kerosene or a similar type of petroleum solvent immediately after use. Fill pumps and meters with new motor oil or a 50% motor oil/fuel oil mixture before storing. **Do not use water.** Dispose of rinsate by applicable Federal, state and local regulations. Never introduce rinsate or unused TELONE™ II into surface or underground water supplies.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-424-9300 .

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No.95290-1

EPA Est. _____

**Produced for
Salt Lake Holding LLC
2211 H. H. Dow Way
Midland, MI 48674**

Net Contents _____

(Cover, shipping container):

RESTRICTED USE PESTICIDE

Due to high acute inhalation toxicity and carcinogenicity.
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

TELONE™ II

Soil Fumigant

A multi-purpose liquid fumigant for preplant treatment of soil to control plant parasitic nematodes, symphylans and to help manage certain soil borne diseases in cropland

Not for use in greenhouses or other enclosed areas.

Not for formulation or manufacturing use. Do not formulate this product into other products.

Active Ingredient:	(by weight)
1,3-dichloropropene.....	97.5%
Other Ingredients	2.5%
Total	100.0%

1 gallon of TELONE™ II weighs 10.15 lb at 70°F. Contains 9.85 lb of 1,3-dichloropropene per gallon.

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-424-9300 .

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No.95290-1

EPA Est. _____

**Produced for
Salt Lake Holding LLC**

**2211 H. H. Dow Way
Midland, MI 48674**

Net Contents _____

(Page 1 through end):

Table of Contents	Page
Precautionary Statements	-
Hazards to Humans and Domestic Animals	-
Personal Protective Equipment (PPE)	-
Engineering Controls Requirements	-
User Safety Requirements	-
User Safety Recommendations	-
First Aid	-
Environmental Hazards	-
Physical or Chemical Hazards	-
Directions for Use	-
Agricultural Use Requirements	-
Storage and Disposal	-
General Information	-
General Use Precautions	-
Application Directions	-
Application Timing	-
Soil Conditions	-
Soil Moisture	-
Soil Preparation	-
Placement of Fumigant	-
Application Methods and Equipment	-
Sealing the Soil After Application	-
Soil Fumigation Interval	-
Uses	-
Control of Nematodes	-
Control of Plant Diseases	-
Control of Soil Insects	-
Terms and Conditions of Use	-
Warranty Disclaimer	-
Inherent Risks of Use	-
Limitation of Remedies	-

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

Hazardous Liquid and Vapor

- Do not swallow any of this product. May be fatal if swallowed.
- Do not get in eyes. Causes substantial, but temporary eye injury.
- Do not get on skin. May be fatal if absorbed through the skin. Causes skin irritation and, if confined, skin burns. May cause allergic skin reaction.
- Do not breathe vapor. May be fatal if inhaled. May cause lung, liver, and kidney damage and respiratory system irritation upon prolonged contact.
- The use of this product may be hazardous to your health. This product contains 1,3-dichloropropene, which has been determined to cause tumors in laboratory animals. Risks can be reduced by exactly following directions for use, precautionary statements, by wearing the personal protective equipment specified in this labeling.

Personal Protective Equipment (PPE)

Chemical-Resistant Materials: Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category H on an EPA chemical resistance category selection chart. PPE constructed of daraneX, neoprene, and chlorinated polyethylene provide short-term contact or splash protection against liquid in this product. Longer-term protection is provided by PPE constructed of viton, Teflon, and EVAL barrier laminates (for example, responder suits manufactured by Life-guard or silvershield gloves manufactured by North). Where chemical-resistant materials are required, leather, canvas, or cotton materials offer no protection from this product and must not be worn as the sole article of protection when contact with this product is possible. Where coveralls are required, they must be loose-fitting and constructed of woven fabrics (e.g., tight knit cotton or cotton/polyester), non-woven fabrics (e.g., tyvek or sontara), or fabrics containing microporous Teflon.

1. Handlers Performing Tasks with Liquid Contact Potential

Tasks with liquid contact potential are tasks performed outdoors or in a well-ventilated area. They include:

- Equipment calibration or adjustment
- Equipment clean-up and repair
- Product sampling
- Any activity less than 6 feet from an unshielded pressurized hose containing this product
- Rinsate disposal
- Fumigant transfer
- Clean-up of small spills
- Preparing containers for aeration
- Any other task not otherwise listed in (2), (3), or (4) below

Handlers performing tasks with liquid contact potential must wear at minimum:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate (EVAL) or viton
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron
- A face shield or safety glasses with brow and temple shields (do not wear chemical goggles)
- A half-face respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides

(MSHA/NIOSH approval number prefix TC-14G). See further respirator requirements in the User Safety Requirements section on this label.

2a. Handlers Performing Tasks with No Liquid Contact Potential – Broadcast, In-Bed Applications, or Applications at the Time of Bedding Except as in 2b.

Tasks with no liquid contact potential are tasks performed outdoors or in a well-ventilated area.

These tasks include:

- Tractor driving
- Soil sealing
- Field activities on the day of application that do not disrupt the soil at the depth of liquid injection

Handlers performing tasks with no liquid contact potential must wear at minimum:

- Loose fitting or well ventilated long-sleeved shirt and long pants
- Shoes and socks
- A face shield or safety glasses with brow and temple shields (do not wear chemical goggles)
- A half-face respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G).
- A respirator is not required (not applicable in California) if the occupants are within an enclosed cab that is in conformance with one of the following: 1) ANSI/SAE S525-1.1 MAY98 sections 7.1.5, 7.1.7, 7.2.3, and 9, or 2) the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides -- 40 CFR 170.240(d)(5). The cab must be equipped with a vapor-adsorptive filter containing a minimum of 1000 grams activated charcoal. The filter must be changed after no more than 50 hours of application time. See further respirator requirements in the User Safety Requirements section on this label.
- In addition, the PPE specified in (1) for activities with direct liquid contact potential must be immediately available and must be worn if the handler is to perform any direct-contact activity with a potential for liquid contact

2b. Handlers Performing Tasks with No Liquid Contact Potential - Pre-Bed, Row Product Applications (e.g., Yetter Rig) (Not Applicable in California)

Tasks with no liquid contact potential are tasks performed outdoors or in a well-ventilated area.

These tasks include:

- Tractor driving
- Soil sealing
- Field activities on the day of application that do not disrupt the soil at the depth of liquid injection

Handlers performing tasks with no liquid contact potential must wear at minimum:

- Loose fitting or well ventilated long-sleeved shirt and long pants
- Shoes and socks
- A face shield or safety glasses with brow and temple shields (do not wear chemical goggles)
- In addition, the PPE specified in (1) for activities with direct liquid contact potential must be immediately available and must be worn if the handler is to perform any direct-contact activity with a potential for liquid contact

3. Handlers in Treated Area 1 to 5 Days After Application

Only the following handler tasks may be performed in the treated area within 5 days after the application is complete:

- Assessing/adjusting the soil seal
- Assessing pest control, application technique, or application efficacy
- Sampling air or soil for this product
- Removing tarp or plastic film

All other tasks are prohibited until the 5-day period has expired.

Handlers in treated area 1 to 5 days after application must wear at minimum:

- Loose fitting or well ventilated long-sleeved shirt and long pants
- Shoes and socks
- A face shield or safety glasses with brow and temple shields (do not wear chemical goggles)
- A half-face respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G).
- A respirator is not required if the occupants are within an enclosed cab (not applicable in California) that is in conformance with one of the following: 1) ANSI/ASAE S525-1.1 MAY98 sections 7.1.5, 7.1.7, 7.2.3, and 9, or 2) the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides -- 40 CFR 170.240(d)(5). The cab must be equipped with a vapor-adsorptive filter containing a minimum of 1000 grams activated charcoal. The filter must be changed after no more than 50 hours of application time. See further respirator requirements in the User Safety Requirements section on this label.
- In addition, the PPE specified in (1) for activities with direct liquid contact potential must be immediately available and must be worn if the handler is to perform any direct-contact activity with a potential for liquid contact

4. Handlers Exposed to High Concentrations

Handlers exposed to high airborne concentrations of this product, such as cleanup following large spills and exposure to this product in poorly ventilated areas, must wear at minimum:

- Chemical-resistant suit
- Chemical-resistant gloves, such as barrier laminate (EVAL) or viton
- Chemical -resistant footwear plus socks
- Chemical-resistant headgear
- Supplied-air respirator with MSHA/NIOSH approval number prefix TC-19C **or** self-contained breathing apparatus (SCBA) with MSHA/NIOSH approval number prefix TC-13F. See further respirator requirements in the User Safety Requirements section on this label.

Note: In-tank cleaning of bulk tanks must be performed only by persons who have been specifically trained for this activity. Refer to OSHA 29 CFR Part 1910.146 and the Storage and Handling Guide.

Engineering Controls Requirements

Mechanical Transfer System: Personal protective equipment specified for Direct Contact Activities must be worn by the operator of the mechanical transfer system. The operator of the mechanical transfer system must follow instructions on proper operation of the system found in the “TELONE™ Soil Fumigants - A Guide to Application” manual. Contact your distributor for TELONE™ II for more information or for these materials.

End-Row Spillage Control: The dispensing system must shut off the feed stream when chisels are raised out of the ground. Do not stop or park near any area where dribble from chisel tips has fallen. The applicator must follow instructions on proper operation and maintenance of the system found in the “TELONE™ Soil Fumigants - A Guide to Application” manual. Contact your distributor for TELONE™ II for more information or for these materials.

- A flow shutoff device must be placed as close as is technically feasible to the fluid discharge point. This can be a ball, poppet, or diaphragm check valve, or full flow shutoff device such as an electric or pneumatically actuated valve.
- Service any system immediately if continuous drip occurs.
- If mechanical check valves and orifices are used, place the check valve above the orifice. Also, isolate the check valve from upstream pressure by installing a main line shut off or bypass valve prior to the manifold.

- Pipe diameter from check valve to injection point must not exceed 1/4 inches ID National Pipe Standard (NPS). Preferably, use the smallest diameter pipe or tubing possible which achieves the required flow rate.
- Alternate end-row spillage devices or methods such as, but not limited to, micro-bore restricted flow tubing or line purge systems may be used if they provide equal or superior control versus check valves.

With all bulk and non-bulk containers, TELONE™ II must be transferred through connecting hoses, pipes, and/or couplings sufficiently tight to prevent workers or other persons from coming in contact with liquid TELONE™ II.

- All hoses, piping, and tanks used in connection with TELONE™ II shall be of the type appropriate for use under the pressure and vacuum conditions to be encountered.
- External sight gauges shall be equipped with valves so that pipes to sight gauge can be shut off in case of breakage or leakage.
- The mechanical transfer system must be adequate to make necessary measurements of the pesticide being used.
- Shut-off devices must be installed on the exit end of all hoses and at all disconnect points to prevent leakage of TELONE™ product when the transfer is stopped and hose is removed or disconnected. A dry coupler that will **minimize pesticide leakage** must be installed at the disconnect point.
- The pressure in hoses used to move TELONE™ II beyond a pump **must not exceed** the manufacturer's maximum pressure specification.

User Safety Requirements

- 1. Respirator Requirements:** When a respirator is required for use with this product, the following criteria must be met:
 - a. Cartridges or canisters must be replaced daily or when odor or irritation from this product becomes apparent, whichever is sooner.
 - b. Respirators must be fit-tested and fit-checked using a program that conforms to OSHA's requirements (described in 29 CFR Part 1910.134).
 - c. Respirator users must be trained using a program that conforms to OSHA's requirements (described in 29 CFR Part 1910.134).
 - d. Respirator users must be examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn.
- 2. Dispose of Contaminated Clothing:** Discard clothing and other absorbent materials that have been drenched or heavily contaminated with liquid from this product. Do not reuse them.
- 3. Clean and Maintain PPE:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Wash PPE after each day's use.
- 4. Contact with Mouth:** Never siphon this product by mouth or use mouth to blow out clogged lines, nozzles, etc.
- 5. Heat Illness Avoidance:** Use measures to avoid or minimize heat illness while using this product. These measures include gradual adjustment to heat and respirator stress, fans for cooling, cooling vests, frequent breaks to cool down, frequent intake of drinking water, and maintaining weight from day to day.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. If breathing is difficult, give oxygen. Call a poison control center or doctor for further treatment advice.

If on skin or clothing: Immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Call a poison control center or doctor for treatment advice. If water is not immediately available, remove excess chemical from skin with adsorbent material such as towel or dry soil, then proceed at once to a location where water is available and thoroughly wash contaminated skin with plenty of water. Call a poison control center or doctor for treatment advice.

If in eyes: Immediately flush eyes with plenty of water. Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

Note to physician: Because rapid absorption may occur through lungs if product is aspirated and cause systemic effects, the decision to induce vomiting or not should be made by a physician. If lavage is performed, endotracheal and/or esophageal control is suggested. Danger from lung aspiration must be weighed against toxicity when considering emptying the stomach.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsate. See Storage and Disposal section. In case of spills properly dispose of contaminated materials.

Groundwater advisory: 1,3-dichloropropene is known to move through soil and under certain conditions has the potential to reach groundwater as a result of agricultural use. Application in areas where soils are permeable and groundwater is near the surface could result in groundwater contamination. Do not apply within 100 feet of any well used for potable water. Do not apply this product within 100 feet from the edge of karst topographical features. Karst topography is identified from landscape features that result from the dissolving activity of water in carbonate rock formations (limestone, dolomite and marble). Surface features that are associated with karst topography include sinkholes, caverns, springs, and sinking or disappearing streams. In North Dakota, South Dakota, Wisconsin, Minnesota, New York, Maine, New Hampshire, Vermont, Massachusetts, Utah, and Montana: Where groundwater aquifers exist at a depth of 50 feet or less from the surface, do not apply this product where soils are Hydrologic Group A.

Reporting Ecological Incidents: To report ecological incidents, including mortality, injury, or harm to plants and animals, call 989-636-4400.

Physical or Chemical Hazards

Combustible. Do not use or store near heat or open flame.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

ENDANGERED SPECIES PROTECTION REQUIREMENTS

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species and certain threatened species, under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the

Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/espp/>, call 1-844-447-3813, or email ESPP@epa.gov. You must use the Bulletin valid for the month in which you will apply the product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry intervals, and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Entry Restriction:

Entry (including early entry that would otherwise be permitted under the WPS) by any person -- other than a correctly trained and equipped handler who is performing a handling task permitted on this labeling -- is **prohibited** from the start of application until 5 days after application. **In addition**, if tarps are used for the application, non-handler entry is prohibited while tarps are being removed.

Notification:

Notify workers of the application by warning them orally and by posting fumigant warning signs at entrances to treated areas. The sign must bear the skull and crossbones symbol and state: (1) "**DANGER/PELIGRO**," (2) Areas under fumigation, **DO NOT ENTER/NO ENTRE**," (3) the date and time of fumigation, (4) "TELONE™ II Soil Fumigant in use," and (5) name, address, and telephone number of the applicator." Post the fumigant warning sign instead of the WPS sign for this application, but follow all WPS requirements pertaining to location, legibility, size and timing of posting and removal.

PPE for Reentry During the Entry-Restricted Period:

PPE for entry that is permitted by this labeling is listed in the "Hazards to Humans and Domestic Animals" section of this labeling.

Storage and Disposal

Do not contaminate water, food and feed by storage and disposal.

Pesticide Storage: Store in tightly-closed original container away from dwellings. Prolonged exposure of the container to direct sunlight must be avoided. Do not allow contamination of seeds, plants, fertilizers, or other pesticide chemicals.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide and rinsates is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

Because TELONE™ II Soil Fumigant is corrosive under certain conditions, flush all application equipment with fuel oil, kerosene or a similar type of petroleum solvent immediately after use. Fill pumps and meters with new motor oil or a 50% motor oil/fuel oil mixture before storing. **Do not use water.** Dispose of rinsate by applicable Federal, state and local regulations. Never introduce rinsate or unused TELONE™ II into surface or underground water supplies.

Nonrefillable containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate

into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers 5 gallons or larger:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers 5 gallons or larger:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

General Information

TELONE™ II Soil Fumigant is a multi-purpose liquid fumigant for preplant treatment of cropland soil. TELONE™ II can be used as part of a nematode management program involving crop rotation, planting of resistant varieties, sanitation, and other cultural practices designed to reduce nematode infestations.

TELONE™ II may be applied as a preplant soil treatment as part of a management program to aid in reducing the damaging effects of certain soil pests; **plant parasitic** nematodes [citrus, burrowing, cyst formers (golden, sugarbeet, soybean, carrot and wheat), dagger, lance, pin, needle, reniform, ring, root knot, root lesion, spiral, sting and stubby root]; **symphylans** (garden centipedes); and **wireworms**.

TELONE™ II can also be used to suppress sugar beet Rhizomania disease, *Fusarium* wilt of cotton and *Verticillium* wilt of mint, and aid in the control of bacterial canker of peaches.

Soil sampling for the type and number of pests present is recommended before fumigation. In fields where pre-treatment soil samples indicate the presence of high population levels of nematodes, a successful fumigation cannot be expected to eradicate entire populations. Therefore, post-treatment (mid-season and/or preharvest) sampling is recommended to determine the need for additional pest management practices.

Supplemental labels are available for certain crops in selected geographies. Refer to these supplemental labels for specific use directions. Consult a company representative for additional information.

Consult State Agricultural Experiment Station or Extension Service specialists for information on other practices such as post-harvest destruction of crop residues, weed control or other cultural practices, and use of nematode resistant crop varieties that may aid in reducing crop losses from soil borne pests.

General Use Precautions

Soil fumigation using TELONE™ II Soil Fumigant should be conducted only according to directions and conditions of use described in this labeling.

Recontamination Prevention: TELONE™ II will help manage certain soil borne pests that are present in the soil treatment zone at time of fumigation. It will not control pests that are introduced into soil after fumigation. To avoid reinfestation of treated soil do not use irrigation water, transplants, seed pieces, or equipment that could carry soil borne pests from infested land. Avoid contamination from moving infested soil onto treated beds through cultivation, movement of soil from below the treated zone, dumping contaminated soil in treated fields and soil contamination from equipment or crop remains. Clean equipment carefully before entering treated fields. Cultural practices, which provide post-harvest destruction of crop residues and weeds prior to fumigation and practices which prevent weed infestation following fumigation and prior to planting, will help prevent recontamination.

Do not use containers, pumps or other transfer equipment made of aluminum, magnesium or their alloys, as under certain conditions TELONE™ II may be severely corrosive to such metals.

Equipment Clean-Up: Because TELONE™ II is corrosive under certain conditions, flush all application equipment with fuel oil, kerosene or a similar type of petroleum solvent immediately after use. Fill pumps and meters with new motor oil or a 50% motor oil/fuel oil mixture before storing. **Do not use water.** Dispose of rinsate by incorporation into field just treated or by other approved means. Never introduce rinsate or unused TELONE™ II into surface or underground water supplies.

Chemigation: Do not apply TELONE™ II through any type of irrigation system.

Fertility Interactions: Fumigation may temporarily raise the level of ammonia nitrogen and soluble salts in the soil. This is most likely to occur when heavy rates of fertilizer and fumigant are applied to soils that are either cold, wet, acidic, or high in organic matter. To avoid injury to certain crops including red beets, carrots, corn, radishes, cole crops, legumes (beans), lettuce, onions, and sugarbeets, fertilize as indicated by soil tests made after fumigation. To avoid ammonia injury or nitrate starvation (or both) to crops grown on high organic soils, fertilizers containing ammonium salts are not recommended.

When using high rates of TELONE™ II as required by certain state nursery regulations, liming of highly acid soils before fumigation may stimulate nitrification and reduce the possibility of ammonia toxicity. Certain nursery crops such as citrus seedlings, *Cornus* sp., *Crataegus* sp., spruce, and vegetable crops such as cauliflower have shown evidence of phosphorus deficiency following fumigation. To avoid this possible effect, additional phosphate fertilizer (foliar applied) is recommended where experience indicates a deficiency may occur.

Use Restrictions for Certain Florida Counties: For application of this product in Brevard, Charlotte, Citrus, Collier, DeSoto, Glades, Hardee, Hendry, Hernando, Highlands, Hillsborough, Indian River, Lake, Lee, Manatee, Martin, Monroe, Okeechobee, Orange, Osceola, Palm Beach, Pasco, Pinellas, Polk, Sarasota, Seminole, St. Lucie, Sumter, and Volusia counties, applicators must have labeling for FIFRA Section 24(c) Special Local Need (SLN) FL-200004 in their possession and comply with stated requirements. Use of TELONE™ II is prohibited in Broward and Dade counties.

Use Restrictions for Certain New York Counties: This product is prohibited from sale, use or distribution in Nassau and Suffolk counties.

Application Directions

Application Timing

TELONE™ II Soil Fumigant can be applied at any time of the year when soil conditions permit. Conditions that allow rapid diffusion of the fumigant as a gas through the soil normally give best results. Because TELONE™ II does not provide residual control of soil pests, it should be used as a preplant application before planting each crop. The following soil temperature and moisture conditions should exist at time of treatment. Failure to meet these conditions may result in unsatisfactory product performance:

Soil Conditions

Soil temperature at the depth of application must be between 40°F and 80°F. In areas where the soil temperature in the spring may not reach 40°F in time to allow application of TELONE™ II prior to planting, late summer or early fall treatment is recommended.

Soil Moisture

It is critical to manage soil moisture properly before fumigation. Plan fumigation for seasons, crop rotations, or irrigation schedules which leave moisture in the soil. For application depths greater than 18 inches, the soil should be moist within a 16-inch radius upwards from the point of injection as determined by the feel method (see below). For all other applications, the soil must be moist from 2 inches below the soil surface to at least 12 inches deep as determined by the feel method (see below). The amount of moisture needed in this zone will vary according to soil type. The surface soil generally dries very rapidly and should not be considered in this determination. If there is insufficient moisture at the 2 to 6 inch depth, the soil moisture must be adjusted. If irrigation is not available and there is adequate soil moisture below 6 inches, it may be brought to the surface by disking or plowing before or during the injection. To conserve existing soil moisture, pretreatment or treatment tillage practices should be done as close to the time of application as possible. For fields with more than one soil texture, soil moisture content in the lightest textured (most sandy) areas must comply with this soil moisture requirement. Whenever possible, the field should be divided into areas of similar soil texture and the soil moisture of each area should be adjusted as needed. Coarser textured soils can be fumigated under conditions of higher soil moisture than finer textured soils; however, if the soil moisture is too high, fumigant movement will be retarded and effectiveness of the treatment will be reduced. Previous and/or local experience with the soil to be treated or the crop to be planted can often serve as a guide to conditions that will be acceptable. If you do not know how to determine the soil moisture content of the area to be treated, consult your local extension service or soil conservation service specialist or pest control advisor (ag consultant) for assistance.

In general, no irrigation should immediately precede subsoiling or fumigation; however, when irrigation is available and surface soil moisture conditions are not likely to provide an adequate seal against fumigant loss, a very light sprinkler irrigation to wet the top 1 to 2 inches of soil is recommended before and/or immediately after fumigation.

The following descriptions will aid in determining acceptable soil moisture conditions by the “feel method.” For coarse soils (sand and loamy sand), there must be enough moisture to allow formation of a weak ball when compressed in the hand. Due to soil texture, this ball is easily broken with little disturbance. In loamy, moderately coarse, or medium textured soils (coarse sandy loam, sandy loam, and fine sandy loam), a soil sample with the proper moisture content can be formed into a ball which holds together with moderate disturbance, but does not stick between the thumb and forefinger. Fine textured soils (clay loam, silty clay loam, sandy clay, silty clay, sandy clay loam and clay), should be pliable and not crumbly, but should not form a ribbon when compressed between the thumb and forefinger.

Soil Preparation

The soil should be free of clods. Large clods can prevent effective soil sealing and reduce effectiveness of TELONE™ II. Plant residues should be thoroughly incorporated into the soil prior to treatment to avoid interfering with application. Non-decomposed plant material may harbor pests that will not be controlled

by fumigation. Little or no crop residue should be present on the soil surface. Crop residue that is present should lie flat to permit the soil to be sealed effectively. Compacted soil layers within the desired treatment zone should be fractured before or during application of the fumigant. Deviation from the above conditions may result in unsatisfactory results.

Placement of Fumigant

TELONE™ II may be applied as either a broadcast (overall) or row treatment. It must be placed at least 12 inches below the final soil surface. When soil conditions allow, placement at a minimum of 14 inches below the final soil surface is recommended. Deeper placement is required when fumigating soil to be planted to deep-rooted plants, such as perennial fruit and nut crops, or to control deeply distributed pests. For row application, the fumigant must be placed at least 12 inches from the nearest soil/air interface (e.g., furrow or bed top).

Application Methods and Equipment

Broadcast Application: Use chisel (shank) or coulter (e.g., Yetter 30-inch Avenger), offset wing shank, Nobel (sweep) plow, or plow-sole application equipment. For best results when using chisel equipment, use ripper-type, forward-swept shanks. Nobel plow equipment is particularly useful for fall fumigation when the soil still contains some non-decomposed standing plant material. Subsoiling may be necessary before application as described under Soil Preparation. Choose application equipment that allows the deepest application and best soil seal under existing conditions.

The fumigant outlet spacing varies with the type of application equipment used.

With chisel and coulter equipment, a fumigant shank spacing of 12 to 24 inches is recommended. Do not exceed the maximum shank and outlet spacing of 24 inches. The outlet spacing for this equipment may be up to 1 1/2 times the application depth but generally should be equal to the application depth and should not exceed the soil-shattering capability of the chisels.

With plow-sole equipment, a 12-inch outlet spacing is recommended. Do not exceed an outlet spacing of 18 inches.

With Nobel (sweep) plow equipment, use an outlet spacing of 9 to 12 inches along the sweeps. Application should be made to a depth of at least 15 inches.

Broadcast application can be made in the same direction or at an angle to the direction of row planting. Refer to Table 1 for broadcast treatment rates for various crops.

Row Application (for row spacing greater than 24 inches): Use chisel equipment to treat a band of soil where the crop is to be planted, i.e., the plant row. When multiple chisels per plant row are used, space the chisels (fumigant outlets) no more than 12 inches apart. Regardless of the number or spacing of chisels used, the fumigant must be placed at least 12 inches from the nearest soil/air interface (e.g., furrow or bed top).

With certain deeper rooted crops such as potatoes and sugarbeets, higher rates may be necessary to ensure adequate treatment of the zone of soil where primary root growth occurs.

To prevent seed germination problems caused by improper seed-to-soil contact or improper planting depth regardless of application method, do not place the seed directly over the furrow left by the applicator chisel(s)/coulter(s). When 1 chisel is used per plant row, place the seed about 4 inches to one side of the chisel furrow. When 2 chisels are used per plant row, plant the seed offset from the chisel trace.

Sealing the Soil After Application

For broadcast treatment (flat fumigation), immediately after chisel application of TELONE™ II, the soil must be "sealed" to prevent fumigant loss and ensure that an effective concentration of fumigant is maintained within the soil for a period of several days. To create an effective seal it is important that the

shank traces be disrupted and the soil surface compacted. Disruption of shank traces can be accomplished with equipment that will uniformly mix the soil to a depth of 3 to 4 inches to eliminate chisel or plow traces which can allow direct escape of the fumigant. A tandem disc or similar equipment may be used for this purpose. To maximize soil sealing, steps should also be taken to compact the soil surface to further retard the rate of fumigant loss by following with a ring roller or cultipacker in combination with the aforementioned tillage equipment. Compaction of the soil surface alone does not effectively disrupt chisel or plow traces. When using coulters (e.g., Yetter 30-inch Avenger) applications, additional sealing may not be necessary when soil moisture conditions are optimal and a beaver tail is used.

For row treatment, forming the beds at the time of application should be accomplished in a manner that places the fumigant at least 12 inches from the nearest soil/air interface (e.g., furrow or bed top). The closest soil/air interface could be the furrow for multiple knife applications or the top of the bed for single knife applications. It is recommended that additional soil sealing be accomplished by going over the bed with a bed shaper, press sealer, rolling cultivator, ring roller, or rolling basket.

Sealing can also be improved by applying non-perforated plastic film, such as polyethylene, over the entire area or in strips. Use of a film to seal the soil surface does not eliminate the need to eliminate chisel traces prior to application of the plastic film. When using coulters (e.g., Yetter prebedder) applications, a beaver tail may be used for sealing.

Proper soil conditions at the time of application (see Soil Preparation section) are important to ensure proper placement of fumigant (see Placement of Fumigant section) and obtaining adequate sealing. Prior tillage should be adequate to eliminate clods and thoroughly mix crop residues into the soil.

Soil Fumigation Interval

Leave the soil undisturbed and unplanted for at least 7 days after application of the fumigant. A longer undisturbed fumigation interval is required if the soil becomes cold or wet, and for deep-rooted tree, shrub and vine planting sites.

Following completion of the fumigation interval, to prevent phytotoxicity, allow the fumigant to dissipate completely before planting the crop. Dissipation is usually complete when TELONE™ II can no longer be detected at the application depth. Under optimum soil conditions for dissipation, a period of 1 week for each 10 gallons per treated acre is generally required for complete dissipation. If virtually impermeable films (VIF) are used a longer dissipation period may be needed. Rapidly germinating seed (i.e., lettuce or radish) and/or seed or transplants to be grown may be used as a bioassay to determine if TELONE™ II is present in the soil at concentrations sufficient to cause plant injury.

To hasten dissipation especially if heavy rains or low temperatures occur during the treatment period, till the soil to the depth of fumigant application. Use a knife-like chisel without turning the soil to reduce the possibility of recontaminating the treated soil. Dissipation is usually complete when the odor of TELONE™ II is no longer evident at the application depth. Seed may be used as a bioassay to determine if TELONE™ II is present in the soil at concentrations sufficient to cause plant injury. Do not plant if the odor of TELONE™ II is present within the zone of fumigation.

Buffer Zone: An application of TELONE™ II shall not be made within 100 feet of an occupied structure, such as a school, hospital, business or residence. No person shall be present at this structure at any time during the seven consecutive day period following application. **This buffer zone does not apply to use on soils that will not experience an additional 1,3-D treatment for at least three years. For example, on soils to be planted with fruit trees, nut and nursery crops, perennial vines, hops, mint or pineapple. Note:** TELONE™ II shall not be applied to soils more frequently than once each year.

Uses

Control of Nematodes

TELONE™ II Soil Fumigant is recommended for control of nematodes and symphylans, and suppression of wireworms in soils to be planted to vegetable crops, field crops, fruit and nut crops, and nursery crops.

Table 1. Broadcast Application Rates and Use Information for Control of Nematodes and Symphylans†, Suppression of Wireworms†, and to Help Manage Certain Soil Borne Diseases in Soils Planted to Crops Listed

Crops (listed but not limited to)	Soil Type	Broadcast Application Rates ¹ (Gallons/Acre)
Vegetable Crops ²	Mineral ³	9 - 12 ⁴
	Muck or Peat ⁵	25
Field Crops	Mineral	9 - 12 ⁴
	Muck or Peat	18
Fruit and Nut Crops ^{6,7,8,9,10}	Mineral	27 - 35
Nursery Crops ¹¹	Mineral	42 - 55

†**Note:** For control of symphylans (garden centipedes) or suppression of wireworms consult the Soil Insects section below for more specific directions and application rates.

¹Rates given may be concentrated in the row, but in no case should the amount applied per acre exceed the maximum broadcast application rates [gallons per acre (gpa)] given in the above table.

²Potatoes: Before fumigation, soil sampling for the type and number of pests present is recommended and can help to determine the need for additional treatment with a contact nematicide. Preharvest tuber sampling for nematodes also is recommended. If the nematode population is high enough to damage the crop, potatoes can be harvested early. Do not store potatoes with a detectable nematode infestation.

Row treatment is not recommended for potatoes in irrigated areas of western and northwestern states.

In Colorado, Idaho, Nevada, Oregon, Utah and Washington, and in Modoc and Siskiyou counties of California refer to supplemental labeling for TELONE™ II entitled: "For the Control of Nematodes and the Suppression of Wireworms in Soils to be Planted to Potatoes, Onions, or Carrots" for directions for use.

³Mineral soil includes sand, sandy loam, silt, and clay loam. Use the higher rates for finer textured (heavier) soils.

⁴For cyst-forming nematodes increase dosage to 18 gpa.

⁵Greater than 20% organic matter content.

⁶Pineapple: Application may be made at the time of planting. For best results, seal the soil with polyethylene film, which acts as a gas permeability barrier.

⁷Tree Planting Sites in the Western U.S.: Use 24 fl oz (1.5 pints) of TELONE™ II by applying the fumigant at a single point in the center of each planting site at a depth of 5 feet below the original soil surface, or into at least 3 points per planting site, at a depth of 3 feet below the original soil surface. The recommended procedure is to prepare the site by backhoeing to break up restrictive soil layers that may retard fumigant movement. The backhoe site should be dug in the approximate dimensions of 10 x 10 x 10 feet. The hole should then be backfilled and the fumigant applied using a closed-system application tube. For sites where no restrictive soil layers are present, the fumigant can be applied to a depth of 5 feet using an injection auger. If backhoe procedure is not used, product performance may be reduced. To prevent phytotoxicity, assure that the chemical has dissipated completely before planting. Dissipation is slower in cold, wet soils. Prepare and treat planting sites in the fall and plant in the spring. In other areas of the U.S., the above may be followed. Regardless of method, ensure thorough fumigation of the desired treatment area. Do not place in groundwater.

⁸For shallow-rooted plants grown only year, use 15 to 27 gpa.

⁹Citrus Fruits: For burrowing nematode control, inject TELONE™ II on 18-inch centers at least 12 inches deep. For buffers within existing groves or for tree planting sites within existing groves, do not apply within 5 feet of living trees. Keep the field free of plants susceptible to burrowing nematodes for 2 years before replanting to citrus.

¹⁰Stone Fruits: Within existing groves or for tree planting sites within existing groves, do not apply within 5 feet of living trees.

¹¹When used according to state nursery regulations, TELONE™ II may be used in the production of certified nursery stock.

Control of Plant Diseases

Bacterial Canker of Peaches: To aid in the control of this disease apply TELONE™ II as a preplant broadcast treatment to light (sandy) soils at the rate of 35 gpa preferably in the fall when the soil is warm (55 to 80°F at injection depth) and moist. Inject the fumigant at least 18 inches deep with chisels mounted on 12- to 18-inch centers.

Fusarium Wilt of Cotton: The effects of this disease can be suppressed by controlling the root knot nematodes associated with this disease/nematode complex. Use TELONE™ II as a row treatment at the rate of 12 gpa.

Sugar Beet *Rhizomania* Disease: Use TELONE™ II to suppress the effects of this disease by preplant broadcast application at the rate of 10 to 18 gpa broadcast equivalent. Use the higher rates for heavier (finer textured) soils and/or for higher levels of disease infestation. TELONE™ II is believed to reduce the activity of *Polymyxa beta*, which has been identified as the vector of the *Rhizomania* disease virus.

Verticillium Wilt of Mint: To aid in the control of this disease, apply TELONE™ II as a broadcast treatment at 25 to 30 gpa in the spring, or preferably in the fall.

Control of Soil Insects

Symphylans (Garden Centipedes): Use TELONE™ II for treatment of soil to be planted to crops where these pests have been shown to be a problem. Apply the fumigant only as a broadcast treatment at the rate of 18 to 35 gpa. Applications made during late summer or early fall when the soil is warm are recommended.

Wireworms: Use TELONE™ II for treatment of soil to be planted to crops where these pests have been shown to be a problem. Apply the fumigant as a broadcast treatment at 20 gpa by injection at least 14 inches below the final soil surface.

Supplemental labels are available for certain crops in selected geographies. Refer to these supplemental labels for specific use directions. Consult a company representative for additional information.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Salt Lake Holding LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Salt Lake Holding LLC MAKES NO OTHER EXPRESS OR

IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Salt Lake Holding LLC or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Salt Lake Holding LLC's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Salt Lake Holding LLC shall not be liable for losses or damages resulting from handling or use of this product unless Salt Lake Holding LLC is promptly notified of such loss or damage in writing. In no case shall Salt Lake Holding LLC be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Salt Lake Holding LLC or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™ Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow
EPA accepted: [MM/DD/YY 12/02/22](#)

01/25/2024

Under the Federal Insecticide, Fungicide and
Rodenticide Act as amended, for the pesticide
registered under EPA Reg. No.

95290-1

No label revisions other than those indicated were
reported to the Agency.H1A / TELONE™ II / Sec 3 Supp Potato-Onion-Carrot Amend with Edits / 08-14-23

Supplemental Labeling



Salt Lake Holding LLC

~~Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA~~
~~Salt Lake Holding LLC 2211 H. H. Dow Way Midland, MI 48674 USA~~

Restricted Use Pesticide

Due to high acute inhalation toxicity and carcinogenicity. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

TELONE™ II~~Telone* II~~ Soil Fumigant

(EPA Reg. No. 95290-1 ~~62749-32~~)

**For the Control of Nematodes and the Suppression of Wireworms in Soils
to be Planted to Potatoes, Onions or Carrots**

**(For Distribution and Use Only in Colorado, Idaho, Nevada, Oregon, Utah, and Washington
~~and in Modoc and Siskiyou Counties of California~~)**

This supplemental label expires on January 25, 2027 and must not be used or distributed after this date.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for TELONE™ II ~~Telone* II~~ Soil Fumigant before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of TELONE™ II ~~Telone* II~~ according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use

For control of Columbia root knot nematode (*Meloidogyne chitwoodi*) in soils to be planted to potatoes, apply TELONE™ II ~~Telone* II~~ Soil Fumigant as described below at the rate of 20 gallons per acre in mineral soil and 25 gallons per acre in muck soil. **In soils to be planted to onions**, apply TELONE™ II ~~Telone* II~~ as described below at the rates of 18 and 25 gallons per acre in mineral and muck soils, respectively. **In soils to be planted to carrots**, apply TELONE™ II ~~Telone* II~~ at the rate of 18 to 20 gallons per acre. For best results, apply the fumigant consistently at least 18 inches below the final soil surface.

For control of northern root knot nematode (*Meloidogyne hapla*) in soils to be planted to either potatoes, onions, or carrots, apply TELONE™ II ~~Telone* II~~ as described below at the rate of 15 to 18 gallons per acre in mineral soil. **In soils to be planted to potatoes or onions**, use 25 gallons per acre in muck soil. For best results, apply the fumigant consistently at least 18 inches below the final soil surface. For high populations, use the higher recommended application rate.

For control of stubby-root nematode (*Paratrichodorus* spp.) in soils to be planted to potatoes, apply ~~TELONE™ II~~ as described below at the rate of 20 to 25 gallons per acre in mineral and muck soils. **In soils to be planted to onions,** apply ~~TELONE™ II~~ as described below at the rates of 18 and 25 gallons per acre for mineral and muck soils, respectively. For best results, apply the fumigant consistently at least 18 inches below the final soil surface.

For suppression of wireworms (*Limonius* spp.) in soils to be planted to potatoes, apply ~~TELONE™ II~~ as described below at the rate of 20 gallons per acre in mineral soil and 25 gallons per acre in muck soil. **In soils to be planted to onions,** apply ~~TELONE™ II~~ as described below at the rates of 18 and 25 gallons per acre in mineral and muck soils, respectively. **In soils to be planted to carrots,** apply ~~TELONE™ II~~ as described below at the rate of 18 to 20 gallons per acre. For best results, apply the fumigant consistently at least 18 inches below the final soil surface in the fall.

Application Methods and Equipment: Apply ~~TELONE™ II~~ as a broadcast treatment using either chisel (shank), Nobel (sweep) or modified ParaTill application equipment according to the following recommendations: Except in those conditions described in the next paragraph for Nobel plow equipment, use either chisel equipment with ripper-type shanks or ParaTill equipment modified so that outlet spacing is evenly distributed under the tool bar. With chisel and ParaTill equipment, a shank spacing of 12 to 24 inches is recommended. Do not exceed a shank spacing of 24 inches. Outlet depth should be at least 18 inches below the final soil surface.

Nobel plow equipment may be used only when either shallow soils (those less than 18 inches deep) or soils containing excessive live root material such as alfalfa or corn stubble prevents the use of shank application. Nobel plow outlet spacing should not exceed 12 inches and application should be made to a depth of at least 15 inches. Fumigant penetration may be limited if a plow pan exists below the depth of the Nobel blade.

Do not use plow-sole application.

Soil Sealing: Immediately after fumigation application, use a disc, paddle-wheel, or similar device to uniformly mix the top 4 to 6 inches of soil to effectively eliminate chisel traces. Then follow immediately with a ring roller or cultipacker to seal the soil surface. Little or no crop residue should be exposed at the surface following the sealing operation. Any remaining crop residue should lie flat following sealing.

Soil Fumigation Interval: Leave the soil undisturbed for 7 to 14 days after application of the fumigant. A longer undisturbed fumigation interval is required if the soil becomes cold or wet.

After the fumigation interval, to prevent phytotoxicity, allow the fumigant to dissipate completely before planting the crop. As a guide, under optimum soil conditions for dissipation, 1 week for each 10 gallons per treated acre is recommended. To hasten dissipation, after the proper fumigation interval, till the soil no deeper than the depth of the fumigant application. During this process be careful not to bring in untreated soil which could contribute to a reinfestation of pests. Use a knife-like chisel without turning the soil to reduce the possibility of recontaminating the treated soil. Dissipation is usually complete when ~~TELONE™ II~~ can no longer be detected at the application depth. Seed may be used as a bioassay to determine if ~~TELONE™ II~~ is present in the soil at concentrations sufficient to cause plant injury. Do not plant if ~~TELONE™ II~~ is detected.

Special Precautions:

- Use of ~~TELONE™ II~~ does not guarantee pest-free crop at harvest.
- Use of ~~TELONE™ II~~ according to these use directions will control only the nematode populations present within the fumigated zone at the time of fumigation. The fumigated zone can vary depending upon a number of factors such as fumigant rate, application methods used, depth of fumigant application, soil moisture, soil type, soil temperature and soil tilth (including soil compaction and soil porosity). ~~TELONE™ II~~ will not control or prevent reinfestation subsequent to the treatment. Subsequent pest populations may infest the fumigated zone from irrigation water, equipment or other sources of contamination or may invade the fumigated zone from surrounding untreated soil such as from beneath the fumigated zone or from unfumigated pockets within the fumigated zone.

- Before fumigation, soil sampling following university soil sampling guidelines for the type and number of pests present is recommended. Fumigation cannot be expected to eradicate the entire pest population. Therefore, post-treatment/preplant soil sampling, again following university soil sampling guidelines, is recommended to determine the need for additional pest management practices.
- Preharvest soil sampling, following university soil sampling guidelines, and preharvest tuber sampling is recommended to detect developing nematode populations or early tuber infection. For best timing and sampling methods, consult your local extension service, pest control advisor, or a company your local Dow AgroSciences representative for assistance. If the nematode population in the soil is high enough that the crop may be damaged, or if any nematode damage is detected in the tubers, the potatoes should be harvested and marketed immediately. **Do not store nematode-infested tubers.**
- Do not plow the ground in the spring in such a way that inverts the soil prior to a spring fumigation. Such tillage operations should be conducted in the fall to allow winter-kill of residual nematode populations in the top 1 to 2 inches of the soil profile.
- Following a fall soil fumigation and undisturbed soil interval as described above, a cover crop, such as wheat or grass, can be planted to reduce the potential for soil erosion.
- Refer to the product label affixed to the container for TELONE™ II for recommended soil conditions and additional use precautions for optimum fumigant performance. The user should understand that there is a range of recommended soil conditions under which TELONE™ II may be applied in compliance with the product label affixed to the container for TELONE™ II. Within that range of recommended soil conditions, product performance can be expected to improve as the soil conditions move toward optimum. Use of TELONE™ II under soil conditions outside the recommended range can be expected to yield less than satisfactory performance.

™ Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow. *Trademark of Dow AgroSciences LLC

01-Supp-B-03D06-018-034

EPA accepted 06-19-03MM-DD-YY

Replaces prior Supplemental label accepted by EPA on 06-19-03

Replaces D06-018-029

H1A / TELONE™ II / Sec 3 Supp Potato-Onion-Carrot Amend with Edits / 08-14-23
H1A / Telone II / Sec 3 Supp Potato-Onion-Carrot FPL / 06-23-03
file: D06-018-031 Telone II Sec 3 CO-ID NV-OR UT-WA-CA Potato-Onion-Carrot 23June03f.doc

TELONE™ Telone* II

EPA Reg. No. 95290-1 62719-32

Registration Notes:

Source Label Text~~Final printed labeling~~ based on EPA-accepted ~~text~~copy dated June 19, 2003.

1. Removal of text “and in Modoc and Siskiyou counties of California” from the Distribution and Use Only section.

2. Added 3-year expiration date of Supplemental Label.

3. Organizational (company) changes related to the October 11, 2019 Transfer of Pesticide Registrations and Data From Company Number 62719 to Company Number 95290.

a. Company name changed to Salt Lake Holding LLC.

b. EPA Registration No. changed to 95290-1.

c. Trademark shown as TELONE™ II.

d. Trademark statement changed to represent Dow or its affiliates.

*Trademark of Dow AgroSciences LLC™ Trademark of The Dow Chemical Company (“Dow”) or an affiliated company of Dow