



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

95670-1

Date of Issuance:

5/20/21

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

BGT

Name and Address of Registrant (include ZIP Code):

CIECH Sarzyna S.A.
c/o Wagner Regulatory Associates, Inc.
P.O.Box 640
Hockessin, DE 19707

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Emily Schmid

Emily Schmid, Product Manager 25
Herbicide Branch, Registration Division (7505P)

Date:

5/20/21

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 95670-1.”
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 10/6/2020

If you have any questions, please contact Lydia Crawford by phone at 703-347-0622, or via email at Crawford.Lydia@epa.gov.

Enclosure

GLYPHOSATE	GROUP	9	HERBICIDE
------------	-------	----------	-----------

[MASTER LABEL]

BGT

ACTIVE INGREDIENT:	% BY WT.
*Glyphosate (n-(phosphonomethyl) glycine), in the form of its isopropylamine salt	20.02%
OTHER INGREDIENTS:	79.98%
TOTAL:	100.00%

*Contains 243 grams per liter or 2 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt (equal to approximately 1.5 pounds acid equivalent (ae) per U.S. gallon of the acid, glyphosate).

KEEP OUT OF REACH OF CHILDREN
CAUTION - PRECAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail)

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> ● Hold eye open and rinse slowly and gently with water for 15-20 minutes. ● Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ● Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> ● Take off contaminated clothing. ● Rinse skin immediately with plenty of water for 15-20 minutes. ● Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> ● Move person to fresh air. ● If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. ● Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> ● Call a poison control center or doctor immediately for treatment advice. ● Have person sip a glass of water if able to swallow. ● Do not induce vomiting unless told to do so by a poison control center or doctor. ● Do not give anything by mouth to an unconscious person.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222.</p>	

[Optional referral statements when booklets and container labels are used:

See label booklet for [complete] [additional] [First Aid,] [Precautionary Statements,] [Directions For Use,] and [Storage and Disposal.]


EPA Reg. No. 95670-XX

EPA Est. No. XXXXX-XX-XXX

Net Contents: _____ [Gals/L]

Manufactured For [By]:
CIECH SARZYNA S.A.
[Chemików 1
37-310 Nowa Sarzyna, Poland]

[P.O. Box 640
Hockessin, DE 19707-0640]



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before re-use.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers. Do not use with or store near any oxidizing agents or chemical reaction may occur.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas, which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette, or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area at the time of application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated area during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

PRODUCT INFORMATION

BGT mixes readily with water to be applied as a foliar spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water in accordance with label instructions. Shake the container if its size permits before pouring to the tank of sprayer when product was stored for the long time.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visible effects of control. Visible effects are a gradual wilting and yellowing of the plant, which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Unless otherwise specified on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the **WEEDS CONTROLLED** section of this label.

Un-emerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at later growth stages – at best when perennials are approaching maturity.

Always use the higher listed rate of this product per acre within the specified range when (1) weed growth is heavy or dense, or (2) weeds are growing in an undisturbed (non-cultivated) area.

BGT contains a built-in adjuvant system that effectively overcomes water hardness, biotic and abiotic stress of weeds, however do not treat weeds under extremely poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Reduced control may result when applications are made to annual and perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the specified stage for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required. To ensure optimum effect do not irrigate sooner than 6 hours after application.

This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the precautionary statements and all other information appearing on the labels of all herbicides used.

To the extent consistent with applicable law, buyers and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly specified in this labeling. Mixing this product with herbicides or other materials not specified on this label may result in reduced performance.

For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

This product may be applied during fallow intervals preceding planting, prior to planting or transplanting, at-planting, or preemergence to annual and perennial crops listed on this label, except where specifically limited. For any crop not listed on this label, application must be made a minimum of 30 days prior to planting.

Use of this product in any manner not consistent with this label may result in injury to persons, animals, or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

RESISTANCE MANAGEMENT

BGT contains the active ingredient glyphosate, classified in Group 9 – Glycine (inhibitor of 5-enolpyruvyl-shikimate-3-phosphate synthase [EPSPS]) chemical classes.

Herbicide resistance is defined as the inherited ability of a plant to survive and reproduce following exposure to a dose of herbicide normally lethal to the wild type. In a plant, resistance may be naturally occurring or induced by such techniques as genetic engineering or selection of variants produced by tissue culture or mutagenesis. Any weed population may contain or develop plants that are naturally resistant to **BGT** and other Group 9 herbicides. Weed species with acquired resistance to Group 9 herbicides may eventually dominate the weed population if Group 9 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by **BGT** or other Group 9 herbicides.

To delay herbicide resistance, consider the below best practices for resistance management:

- Plant into weed-free fields and keep fields as weed-free as possible.
- To the extent possible, use a diversified approach toward weed management. Whenever possible incorporate multiple weed-control practices such as mechanical cultivation, biological management practices, and crop rotation.
- Fields with difficult to control weeds must be rotated to crops that allow the use of herbicides with alternative mechanisms of action or different management practices.

- To the extent possible do not allow weed escapes to produce seeds, roots or tubers. Manage weed seeds at harvest and post-harvest to prevent a buildup of the weed seed-bank.
- Prevent field-to-field and within-field movement of weed seed or vegetative propagules. Thoroughly clean plant residues from equipment before leaving fields.
- Prevent an influx of weeds into the field by managing field borders.
- Identify weeds present in the field through scouting and field history and understand their biology. The weed-control program must consider all of the weeds present.
- Difficult to control weeds may require sequential applications of herbicides with differing mechanisms of action.
- Apply this herbicide at the correct timing and rate needed to control the most difficult weed in the field.
- Use a broad-spectrum soil-applied herbicide with a mechanism of action that differs from this product as a foundation in a weed-control program. Do not use more than two applications of this or any other herbicide with the same mechanism of action within a single growing season unless mixed with an herbicide with another mechanism of action with an overlapping spectrum for the difficult-to-control weeds.
- If resistance is suspected, treat weed escapes with an herbicide with a different MOA or use non-chemical methods to remove escapes.
- Monitor treated weed populations for loss of field efficacy.
- Scout field(s) before and after application.
- Report lack of performance to your local representative.

Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species.

Contact your local sales representative, extension agent, or certified crop advisors to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of action for each target weed.

Glyphosate Resistance

Some naturally occurring weed biotypes resistant to glyphosate may exist through normal genetic variability in any weed population. The repeated use of herbicides with the same mode of action is known to lead, under certain conditions, to a selection of resistant weeds. Certain agronomic practices reduce the likelihood that resistant weed populations will develop and integrated strategies are known to manage such problem weeds.

Glyphosate is the active ingredient in **BGT**, so glyphosate-resistance management is critical. When applying **BGT** to broadleaf weeds that are suspected or known to be resistant to glyphosate, tank mix with atrazine or dicamba to provide an additional mode of action.

BGT will not provide control of emerged grasses that are resistant to glyphosate. For control of glyphosate-resistant grass weeds, a weed control program that includes a pre-emergence grass herbicide will reduce the dependence on glyphosate.

Use cultural and mechanical control practices, including crop rotation or tillage, as appropriate.

When a naturally occurring resistant biotype(s) is present, tank mix or apply sequentially with an appropriate herbicide with a different mode of action to achieve control.

The Best Weed Management practice includes the diversification of glyphosate-dependent weed control programs with alternative mode of action herbicides or cultural practices.

1. In Roundup Ready (RR™) corn systems do not use more than two applications of a glyphosate-based herbicide over a two-year period. Diversify with alternative mode of action herbicides and/or cultural practices.
2. Use alternative (non-glyphosate) burndown and/or residual herbicides for glyphosate tolerant crops likely to require more than one application of glyphosate.
3. To help manage glyphosate tolerant-resistant volunteers, rotate glyphosate tolerant crops with conventional or non-glyphosate tolerant crops.
4. Use full labeled rates of glyphosate and tank mix partners. Minimize weed escapes.
5. Monitor treated weed populations for any loss of field efficacy.
6. Contact your local Cooperative Extension specialist, certified crop advisor, and/or local representative for herbicide resistance management and/or integrated weed management practices for specific crops and resistant weed biotypes.

MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS

Spray solutions of this product must be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL

(EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. DO NOT APPLY WHEN WIND OR OTHER CONDITIONS FAVOR DRIFT. HAND-HELD APPLICATIONS MUST BE PROPERLY DIRECTED TO AVOID SPRAYING DESIRABLE PLANTS.

NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

Eliminate any risk of siphoning the contents of the tank back into the carrier source while mixing. Use approved anti-back-siphoning devices where required by State or local regulations.

Clean sprayer parts immediately after use of this product by thoroughly flushing with water.

Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of water. Add the specified amount of this product (see the **WEEDS CONTROLLED** section of this label) near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by State or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

Tank Mixtures

This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control, a broader weed control spectrum or an alternate mode of action. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Under certain growth stages and/or under other circumstances, some tank mixtures have the potential to cause crop injury. Prior to use, read all labels for products to be used in the tank mixture to determine the potential for crop injury.

To the extent consistent with applicable law, buyers and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are specified in this labeling.

USE PRECAUTIONS

- Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers may result in reduced weed control or crop injury.
- Prior to using a product in tank mixture, it is advised that the user has knowledge of how the tank mixture will perform on a use site area from a crop safety and efficacy standpoint by performing a field bioassay on a small section of use area prior to making application to entire area. Do not use tank mixture if reduced performance and/or crop safety is unacceptable to user.
- Always predetermine the compatibility of tank mixtures with this product with water carrier by mixing small proportional quantities in advance. Do not use if tank mixture results show the mixture is not compatible.

Mixing order is as follows:

1. Water
2. Agitation
3. **BGT**
4. Tank mix partner

Note: **BGT** formulation is loaded with adjuvant system to manage water hardness as well as to improve droplets retention, absorption and translocation of active substance into weed tissues. **BGT** as such does not require ammonium sulphate (AMS) or any other surfactant or adjuvant.

Never tank mix without constant and complete agitation.

Mix labeled tank mixtures of **BGT** with water as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it slowly through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one-part water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
6. Continue filling the spray tank with water and add the required amount of this product near the end of the filling process.
7. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control

additive, water soluble liquid

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. Keep by-pass line on or near bottom of the tank to minimize foaming. Screen size in nozzle or line strainers must be no finer than 50-mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles. Clean sprayer and parts immediately after using this product by thoroughly flushing with water.

Additives

BGT formulation is loaded with adjuvant system to manage water hardness as well as to improve droplets retention, absorption and translocation of active substance into weed tissues. **BGT** as such does not require ammonium sulphate (AMS) or any other surfactant or adjuvant.

- **Colorants or Dyes:** Agriculturally approved colorants or marking dyes may be added to this product. Use colorants or dyes according to the manufacturer's directions.
- **Drift Control:** When a drift control additive is used, read and carefully observe the precautionary statements and all other information appearing on the additive label.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator and the grower. The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase nozzle pressure.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended.

AERIAL SPRAY DRIFT MANAGEMENT

Use the following information when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops:

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.

The following drift management requirements must also be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications or to public health uses:

1. The distance of the outermost nozzles on the boom must not exceed 0.75 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they must be observed.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the **Wind, Temperature and Humidity**, and **Temperature Inversions** sections of this label).

Controlling Droplet Size

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Use the lower spray pressures as specified for the nozzle. Higher pressure reduces droplet size and does not improve canopy protection. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles:** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation:** Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

- **Application Height:** Applications must not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application must be avoided below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications must not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide must only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion.

APPLICATION INFORMATION

USE PRECAUTION:

- Avoid herbicide contact with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees because severe injury or destruction may result.

USE RESTRICTION:

- Do not apply this product plus 2,4-D, dicamba, or sulfometuron-methyl tank mixtures by air in California.

FOR AERIAL APPLICATIONS IN CALIFORNIA ONLY

This label must be in the possession of the user at the time of the herbicide application.

See **PRODUCT INFORMATION** and **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** sections of this label for essential product performance information.

See the **CROPPING SYSTEMS** section of this label for specific instructions on the use of this product.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE, GREEN STEMS, OR FRUIT OF DESIRABLE CROPS, PLANTS, TREES, OR OTHER DESIRABLE VEGETATION SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

Aerial applications of this product are allowed in the following situations:

1. In fallow and reduced tillage systems prior to the emergence or transplanting of labeled crops.
2. Prior to harvest in cotton, soybeans, wheat and glyphosate tolerant canola, corn, and cotton.

Do not plant subsequent crops other than those listed in this label for 30 days following application.

When applied as directed, under the conditions described, **BGT** controls annual and perennial weeds listed in this label.

DO NOT EXCEED MAXIMUM RATE OF 1 QT. PER ACRE (0.5 lb. a.i./A) OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR WITH

THE FOLLOWING EXCEPTIONS:

DO NOT EXCEED A MAXIMUM RATE OF 2 QTS. PER ACRE (1 lb. a.i./A) OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR IN FALLOW AND REDUCED TILLAGE SYSTEMS, AND PRIOR TO HARVEST IN GLYPHOSATE TOLERANT COTTON.

**FOR AERIAL APPLICATION IN FRESNO COUNTY CALIFORNIA ONLY
FROM FEBRUARY 15th THROUGH MARCH 31st ONLY.**

NOTE: For aerial application outside these dates, refer to FOR AERIAL APPLICATION IN CALIFORNIA ONLY section.

This label must be in the possession of the user at the time of the herbicide application.

See **PRODUCT INFORMATION** and **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** sections of this label for essential product performance information.

See the **CROPPING SYSTEMS** section of this label for specific instructions on the use of this product.

Applicable Area

The following information only applies to the area contained inside the following boundaries within Fresno County California only:

- **North:** Fresno County line
- **South:** Fresno County line
- **East:** State Highway 99
- **West:** Fresno County line

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of **BGT**.

Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor, and aerial applicator.

Written Instructions

A written instruction **MUST** be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to application. This written instruction **MUST** state the proximity of the surrounding crops, and that conditions of each manufacturer's applicable product label(s) and this label have been satisfied.

Aerial Applicator Training and Equipment

Aerial application of **BGT** is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight, and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to ensure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commission approved fly-ins constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

Application at Night - Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

FOR AERIAL APPLICATION IN ARKANSAS ONLY

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL GLYPHOSATE TOLERANT CROPS), DESIRABLE PLANTS AND TREES, AS SEVERE INJURY OR DESTRUCTION MAY RESULT.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

See **PRODUCT INFORMATION** and **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** sections of the container label for this product for essential product performance information.

AERIAL APPLICATIONS

Use the specified rate of this product in 3 to 15 gals. of water per acre.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended.

Apply only when the wind speed is less than or equal to 10 mph at the application site.

Do not apply by air if drift can occur to sensitive non-target crops or plants that are within 100 ft. of the application site.

If wind up to 5 mph is blowing toward desirable vegetation or crops, do not apply within 500 ft. upwind of the desirable vegetation or crops.

Winds blowing from 5 to 10 mph toward desirable vegetation or crops will likely require buffer zones in excess of 500 ft.

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron ranges are specified.

Applications should typically be made with the nozzle release point at 8 to 15 ft. above the top of the target plants unless a greater height is required for aircraft safety.

The boom length must not exceed 75% of the wingspan or rotor blade diameter. In many cases, reducing this distance to 65% of the length of the wingspan or rotor will improve drift control without affecting the swath width.

Do not make any type of application into temperature inversions. Do not apply into still air where there is a temperature inversion layer low enough for fine spray particles to become suspended and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 mph.

Nozzles must always discharge backward parallel with the air stream and never discharge downwards more than 45 degrees on fixed wing aircraft or forward of the prevailing air flow on rotary winged aircraft. Avoid the use of nozzles with wide-angle discharge.

APPLICATION EQUIPMENT AND TECHNIQUES

DO NOT apply this product through any type of irrigation system.

This product may be applied with the following application equipment:

- **Aerial** - Fixed wing and helicopter
- **Broadcast Spray – Ground** - Boom or boomless systems, pull-type sprayers, floaters, pick-up sprayers, spray coupes and other broadcast equipment.
- **Controlled Droplet Applicator (CDA)** - Hand-held or boom-mounted applicators that produce a spray consisting of a narrow range of droplet sizes.
- **Hand-Held and High-Volume Spray Equipment** - Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, handwands, mist blowers*, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.
*This product is not registered in California or Arizona for use in mist blowers.
- **Selective Equipment** - Recirculating sprayers, shielded sprayers, and wiper applicators.
- **Injections Systems** - Ground or aerial injections systems.

SPRAY SOLUTIONS MUST BE APPLIED IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT THAT IS CAPABLE OF DELIVERING VOLUMES DESIRED.

Aerial Equipment

Use the specified rates of this herbicide in 3 to 15 gals. of water per acre unless otherwise specified on this label. See the **WEEDS CONTROLLED** section of this label for specific rates by weed species. Unless otherwise specified, do not exceed 1 qt. per acre. Aerial applications of this product may be made as listed in the individual use area sections of this label. Refer to the individual use areas sections for specified volumes and application rates.

Do not apply to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters.

AVOID DRIFT - DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops:

1. Do not apply within 100 ft. of all desirable vegetation or crop(s).
2. If wind up to 5 mph is blowing toward desirable vegetation or crop(s), do not apply within 500 ft. of the desirable vegetation or crop(s).
3. Winds blowing from 5 - 10 mph toward desirable vegetation or crop(s) may require buffer zones in excess of 500 ft.
4. Do not apply when winds are in excess of 10 mph or when inversion conditions exist.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets.

Do not angle nozzles forward into the airstream and do not increase nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the precautionary statements and all other information appearing on the additive label.

Ensure Uniform Application - To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

Ground Broadcast Equipment

For control of annual or perennial weeds listed on this label using broadcast equipment, use the specified rates of this product in 8 – 20 gals. of water per acre as a broadcast spray unless otherwise specified on this label. See the **WEEDS CONTROLLED** section of this label for specific rates. As density of weeds increases, spray volume must be increased within the specified range to ensure complete coverage. Carefully select proper nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

Controlled Droplet Application (CDA)

The rate of this product applied per acre by vehicle-mounted CDA equipment must not be less than the amount specified in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 3 - 15 gals. of water per acre.

For the control of labeled annual weeds with hand-held CDA units, apply a 20% solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 mph (1 qt. per acre). For the control of labeled perennial weeds, apply a 20 - 40% solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mph (2 - 4 qts. per acre).

Controlled droplet application equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

Hand-Held and High-Volume Equipment

Use coarse sprays only.

Mix this product in clean water and apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage must be uniform and complete.

USE RESTRICTIONS:

- Do not spray to the point of runoff.
- Do not use any additional surfactants and adjuvants.

For control of annual weeds listed on this label, apply a 0.5% solution of this product weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. Allow 3 or more days before tillage or mowing.

For annual weeds over 6 inches tall, unless otherwise specified, use a 1% solution. For best results, use a 2% solution on harder-to-control perennials.

When using application methods that result in less than complete coverage, use a 5% solution for annual and perennial weeds and a 5 - 10% solution for woody brush and trees.

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the below table.

Spray Solution

Desired Volume	Amount of BGT					
	0.5%	1%	1.5%	2%	5%	10%
1 Gal.	0.66 oz.	1.33 oz.	2 oz.	2.66 oz.	6.5 oz.	13 oz.
25 Gals.	1 pt.	1 qt.	1.5 qts.	2 qts.	5 qts.	10 qts.
100 Gals.	2 qts.	1 gal.	1.5 gals.	2 gals.	5 gals.	10 gals.
2 tablespoons = 1 fluid ounce						

For use in knapsack sprayers, it is suggested that the specified amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution.

Selective Equipment

This product may be applied through a recirculating spray system, a shielded applicator, hooded sprayers, wiper applicators, or sponge

bars after dilution and thorough mixing with water to listed weeds growing in any non-crop site specified on this label.

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

A shielded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide.

A wiper applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution.

USE PRECAUTIONS:

- AVOID CONTACT WITH DESIRABLE VEGETATION.
- Contact of the herbicide solution with the desirable vegetation may result in damage or destruction.
- Applicators used above desired vegetation must be adjusted so that the lowest spray stream or wiper contact point is at least 2" above the desirable vegetation. Droplets, mist, foam, or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction.
- Applications made above the crops must be made when the weeds are a minimum of 6" above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded Applicators/Hooded Sprayers

When applied as directed under conditions described for shielded applicators and hooded sprayers, **BGT** will control those weeds listed in the **WEEDS CONTROLLED** section of this label.

Use the following equation to convert from a broadcast rate per acre to a band rate per acre:

$$\frac{\text{Band Width in Inches}}{\text{Row Width in Inches}} \times \text{Herbicide Broadcast RATE / Acre} = \text{Herbicide Band RATE / Acre}$$

$$\frac{\text{Band Width in Inches}}{\text{Row Width in Inches}} \times \text{Herbicide Broadcast VOLUME of Solution / Acre} = \text{Band VOLUME of Solution / Acre}$$

Use nozzles that provide uniform coverage within the treated area. Keep shields on shielded sprayers adjusted to protect desirable vegetation. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT WITH DESIRABLE VEGETATION.**

A hooded sprayer is a type of shielded sprayer where the spray pattern is fully enclosed including the top, sides, front and back, thereby shielding the crop from the spray solution. To protect desirable vegetation, adjust the shields on these sprayers. When applications are made to crops grown on raised beds, make sure the hood is capable of completely enclosing the spray pattern. If necessary, extend the front and rear flaps of the hooded applicator downward to reach the ground in deep furrows. **EXTREME CARE MUST BE TAKEN TO AVOID CONTACT OF THIS HERBICIDE WITH DESIRABLE VEGETATION.**

For specific rates of application and instructions for control of various annual weeds and perennial weeds, see the **WEEDS CONTROLLED** section of this label.

Wiper Applicators

Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed.

Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. When more of the weed is exposed to the herbicide solution, better results may be obtained. Weeds must be a minimum of 6" above the desirable vegetation. To ensure adequate contact with weeds, adjust the height of the applicator. Weeds not contacted by the solution will not be affected. Poor contact may occur when weeds are growing in dense clumps, in severe weed infestations or when weed height varies dramatically. In these instances, repeat applications may be necessary. Operate this equipment at ground speeds no greater than 5 mph. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if 2 applications are made in opposite directions.

Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using this product by thoroughly flushing with water.

USE RESTRICTIONS:

- Do not use wiper equipment when weeds are wet.

- Do not add surfactant to the herbicide solution.

WEEDS CONTROLLED

This herbicide will provide control of the listed weeds below for the labeled uses.

ANNUAL WEEDS

- Apply to actively growing grass and broadleaf weeds.
- Allow at least 3 days after treatment before tillage.
- For maximum agronomic benefit, apply when weeds are 6" or less in height.
- To prevent seed production, applications must be made prior to seedbed formation.
- This product does not provide residual control; therefore, delay application until maximum weed emergence. Repeat treatments may be necessary to control later germinating weeds.

PERENNIAL WEEDS

NOTE: If weeds have been mowed or tilled, do not treat until plants have resumed active growth. Best results are obtained when perennial weeds are treated after they reach the reproductive stage of growth (seed head initiation in grasses and bud formation in broadleaves). For non-flowering plants, best results are obtained when the plants reach a mature stage of growth. In many situations, treatments are required prior to these growth stages. Under these conditions, use the higher application rate within the labeled range.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

Additives

BGT formulation is loaded with adjuvant system to manage water hardness as well as to improve droplets retention, absorption and translocation of active substance into weed tissues. **BGT** as such does not require ammonium sulphate (AMS) or any other surfactant or adjuvant. Refer to the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label.

When applied as specified under the conditions described, **BGT** will control the listed weeds below.

Common Name	Use Rate [except in California] (fl. oz. product/A)	[Use Rate California (fl. oz. product/A)]	Common Name	Use Rate [except in California] (fl. oz. product/A)	[Use Rate California (fl. oz. product/A)]
Alfalfa	32-45	[32-64]	Junglerice	32-45	[32-64]
American marshpennywort	32-45	[32-64]	Kenya white clover	32-45	[32-64]
Annual Ryegrass	32-45	[32-64]	Mexican lovegrass	32-45	[32-64]
Bahiagrass	32-45	[32-64]	Mexican primrose-willow	32-45	[32-64]
Barnyard grass	32-45	[32-64]	Prostrate spurge	32-45	[32-64]
Bermuda grass	32-45	[32-64]	Puncturevine	32-45	[32-64]
Bistrly oxtongue	32-45	[32-64]	Purple Nutsedge	32-45	[32-64]
California Burclover	32-45	[32-64]	Purslane	32-45	[32-64]
Carelessweed	32-45	[32-64]	Redroot pigweed	32-45	[32-64]
Carpetweed	32-45	[32-64]	<i>Sida pseudocordifolia</i>	32-45	[32-64]
Cocklebur	32-45	[32-64]	Smutgrass	32-45	[32-64]
Common Lambsquarters	32-45	[32-64]	Sowthistle	32-45	[32-64]
Crabgrass	32-45	[32-64]	Spiny amaranth	32-45	[32-64]
Creeping buttercup	32-45	[32-64]	Tall fescue	32-45	[32-64]
Doveweed	32-45	[32-64]	Tomatillo group	32-45	[32-64]
Downy brome	32-45	[32-64]	Tufted lovegrass	32-45	[32-64]
Fall panicum	32-45	[32-64]	Walnut - Sucker Control	32-45	[32-64]
Field Bindweed	32-45	[32-64]	Weeping lovegrass	32-45	[32-64]
Flatsedge	32-45	[32-64]	White Clover	32-45	[32-64]
Florida pusley	32-45	[32-64]	Wild carrot	32-45	[32-64]
Hairy nightshade	32-45	[32-64]	Witchgrass	32-45	[32-64]
Healall	32-45	[32-64]	Yellow Nutsedge	32-45	[32-64]
Horehound	32-45	[32-64]			
Horsenettle	32-45	[32-64]			

Rate Conversion Chart

Quarts	Pints	Fluid Ounces	Pounds Active Ingredient*
1	2	32	0.5 lbs. a.i.
1.03	2.1	33	0.516 lbs. a.i.
1.41	2.8	45	0.703 lbs. a.i.
1.50	3.0	48	0.8 lbs. a.i.
2	4	64	1 lbs. a.i.
4	8	128	2 lbs. a.i.

*Glyphosate, In the form of its isopropylamine salt

CROP-SPECIFIC USE DIRECTIONS

CORN[*]

Hooded Sprayers - BGT may be used through hooded sprayers for weed control between the rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used.

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set-up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

When applying to corn that is grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

USE RESTRICTIONS:

- The spray hoods must be operated on the ground or skimming across the ground.
- Do not apply more than 1 qt. (0.5 lbs. a.i.) of this product per acre per application.
- Do not apply more than 16.2 qts. (8.1 lbs ai) of this product per acre per year.
- Do not apply more than 3 qts. (1.5 lbs. a.i./A) of this product per acre per year for hooded sprayer applications.
- Corn must be at least 12" tall, measured without extending leaves.
- Leave at least an 8" untreated strip over the drill row. For example, if the crop row width is 38", the maximum width of the spray hood must be 30".
- Maximum tractor speed: 5 mph
- Maximum wind speed: 10 mph
- Use low-drift nozzles.
- Do not graze or feed corn forage or fodder following application of this product through hooded sprayers.

USE PRECAUTIONS:

- Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.
- Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

For specific rates of application and instructions for control of various annual and perennial weeds, see the **WEEDS CONTROLLED** section of this label.

Fallow and Reduced Tillage Systems

[For aerial application in California, refer to the **FOR AERIAL APPLICATION IN CALIFORNIA ONLY** and **FOR AERIAL APPLICATION IN FRESNO COUNTY CALIFORNIA ONLY** sections of this label.]

Use **BGT** in fallow and reduced tillage systems for control of annual weeds prior to emergence of crops listed in this label. Refer to the **WEEDS CONTROLLED** section of this label for specific rates and instructions. This product may be applied using ground or aerial spray equipment. See the **APPLICATION EQUIPMENT AND TECHNIQUES** section of this label for instructions.

[*Not registered for sale or use in California.]

POST-HARVEST GRAIN SORGHUM, SORGHUM REGROWTH CONTROL[*]

BGT may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 1 qt. of this product per acre for control, or 1.5 pts. of this product per acre for suppression.

[*Not registered for sale or use in California.]

PASTURES[*]

Apply **BGT** prior to planting forage grasses and legumes.

Pasture or Hay Crop Renovation - When applied as a broadcast spray, this product controls the annual and perennial weeds listed in this label prior to planting forage grasses or legumes.

USE RESTRICTIONS

- Remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.
- Do not apply more than 21.6 qts. (10.8 lbs ai) of this product per acre per year.

Spot Treatment - When applied as a spot treatment as directed, this product controls annual and perennial weeds listed in this label which are growing in pastures, forage grasses and forage legumes composed of bahiagrass, Bermuda grass, bluegrass, brome, fescue, orchardgrass, ryegrass, timothy, wheatgrass, alfalfa, or clover.

Wiper Application - When applied as directed, this product controls or suppresses the weeds listed under Wiper Applicators in the Selective Equipment section of this label.

USE RESTRICTIONS FOR SPOT TREATMENT AND WIPER APPLICATION

- Apply in areas where the movement of domestic livestock can be controlled.
- No more than one-tenth of any acre must be treated at one time.
- Further applications may be made in the same area at 30-day intervals.
- Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

[*Not registered for sale or use in California.]

GLYPHOSATE TOLERANT CROPS[*]

[*Not registered for sale or use in California]

USE RESTRICTION

- Do not apply more than 16.2 qts. (8.1 lbs ai) of this product per acre per year.

USE PRECAUTION:

- Glyphosate tolerant crops contain a gene that provides tolerance to glyphosate, the active ingredient in this product. This product will cause severe crop injury or destruction and yield loss if applied to crops that are not glyphosate tolerant. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants that do not contain a glyphosate tolerant gene, since severe plant injury or destruction will result.

Information on glyphosate tolerant crops may be obtained from your seed supplier.

The instructions in the sections that follow, include all applications of this product that may be made onto specified glyphosate tolerant crops during the complete cropping season. **DO NOT** combine these instructions with other instructions for crops within this label that do not contain a glyphosate tolerant gene.

USE INSTRUCTIONS

See the **WEEDS CONTROLLED** section of this label for rate specifications for specific weeds. When applied as directed, **BGT** will control these annual and perennial grasses and broadleaf weeds. Observe the maximum application rates and crop stage timings specified for individual glyphosate tolerant crops in the sections that follow.

For Ground Broadcast Applications: Apply this product in 5 - 20 gals. of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat spray nozzles. Check for even distribution of spray droplets.

For Aerial Applications: All labeled treatment may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this label, particularly, in the **Aerial Equipment** section. Apply this product in 3 - 15 gals. of water per acre. See the **APPLICATION EQUIPMENT AND TECHNIQUES** section of this label for procedures on avoiding spray drift that may cause injury to any vegetation not intended for treatment. Use of appropriate buffer zones will help prevent injury to adjacent vegetation.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN A GLYPHOSATE TOLERANT GENE.

See the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** and **APPLICATION EQUIPMENT AND TECHNIQUES** sections of this label for additional directions and restrictions on the application of this product.

TANK MIXTURES

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers may result in reduced weed control or crop injury and are NOT specified for post-emergence (in-crop) applications of this product over-the-top of glyphosate tolerant crops, unless otherwise noted in this product label. Some tank mixture products have the potential to cause crop injury under certain conditions, at certain crop growth stage and/or under other circumstances. Read all labels for products used in the tank mixture prior to use to determine the potential for crop injury. Always predetermine the compatibility of tank mix products together in the carrier by mixing small proportional quantities in advance. A tank mixture of this product with other herbicides may cause incompatibility, antagonism, or a reduction in product efficacy. Ciech Sarzyna S.A. has not tested all tank mix product formulations for compatibility or performance. See the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label for more information on tank mixtures.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product. Follow the cleaning procedures specified on the label of the product(s) previously used. **THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.**

GLYPHOSATE TOLERANT CORN [*]

[*Not registered for sale or use in California.]

Types of Applications: Pre-plant, At-planting, Pre-emergence, Post-emergence (In-crop), Spot treatment, Pre-harvest, Post-harvest
Refer to the following table for maximum application rates of this product with glyphosate tolerant corn hybrids.

Maximum Application Rates/Acre	
Combined total per year for all applications	16.2 qts (8.1 lbs. a.i.)
Total of all pre-plant, at-planting, pre-emergence applications	5 qts. (2.5 lbs. a.i.)
Total of all in-crop applications from emergence through 48" corn	96 fl. oz. (1.5 lbs. a.i.) (48 fl. oz. per acre per application/0.8 lb. a.i.)
Maximum pre-harvest application rate after maximum kernel fill is complete and the crop is physiologically mature (black layer formation) until 7 days before harvest*	33 fl. oz. (0.5 lb. a.i.)
*See RESTRICTIONS within the Pre-Harvest section.	

For specific rates of application and instructions for control of annual and perennial weeds see the **WEEDS CONTROLLED** section of this label.

USE PRECAUTION:

- See the **GLYPHOSATE TOLERANT CROPS** section of this label for general precautionary instructions for use in glyphosate tolerant crops.

USE RESTRICTIONS:

- The use of the in-crop (over-the-top) rates described in these instructions on corn hybrids that are *other than* glyphosate tolerant, may cause crop injury and reduced yields.
- The maximum combined total amount of this product that may be applied is 16.2 qts per acre per year.

PRE-PLANT, AT-PLANTING, PRE-EMERGENCE

This product may be applied alone or in a tank mixture before, during or after planting.

Tank Mixtures: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Ensure that the specific product being used is labeled for application prior to emergence of corn. Apply these tank mixtures in 10 – 20 gals. of water, per acre. See the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label for more information on tank mixtures.

USE RESTRICTION:

- Maximum quantity of this product that may be applied for all pre-plant, at-planting, and pre-emergence applications combined is 5 qts. per acre per year.

NOTE: For maximum weed control, apply a post-emergence (in-crop) application of this product following the use of a labeled pre-emergence residual product.

POST-EMERGENCE (IN-CROP)

This product may be applied alone or in tank mixtures over-the-top of glyphosate tolerant corn from emergence through the V8 stage (8 leaves with collars), or until corn height reaches 30" (free standing), whichever comes first. Drop nozzles are specified for optimum spray coverage and weed control when corn height is 24" - 30". For corn heights 30" - 48" (free standing), apply this product **only** using ground application equipped with drop nozzles aligned to avoid spraying into the whorls of the corn plants. Single in-crop applications of this product up to 48" corn must not exceed 48 fl. oz. per acre. Sequential in-crop application of this product from emergence through 48" in height must not exceed 96 fl. oz. per acre per growing season.

When applied as directed, this product will control annual grasses and broadleaf weeds listed on this label. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of this product. A post-emergence application of 32 to 45 fl. oz. of this product per acre can be made before weeds exceed 4" in height, or, generally, before they become competitive with the crop. If new flushes of weeds occur, a sequential application of this product at 32 to 45 fl. oz. per acre can be made before weeds exceed 4" in height.

Tank Mixtures: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Ensure that the specific product being used in the tank mixture is registered for application post-emergence (in-crop) to corn. See the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label for more information on tank mixtures.

USE RESTRICTIONS:

- Allow a minimum of 10 days between in-crop applications of this product.
- Pre-Harvest Interval (PHI): 50 days between application of this product and harvest of corn forage or grain.

PRE-HARVEST

This product may be applied for annual and perennial weed control prior to harvest. Make applications at 35% grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed).

USE RESTRICTIONS:

- Pre-Harvest Interval (PHI): 7 days between application and harvest or feeding of corn stover or grain.
- A pre-harvest application may only be made if the combined total of previously applied over-the-top or drop nozzle applications does not exceed 66 fl. oz. of this product per acre.
- Pre-Harvest Interval (PHI): 7 days between treatment and harvest or feeding of treated vegetation.

POST-HARVEST

This product may be applied for weed control after crop harvest. The higher listed rates may be required for control of large weeds. Tank mixtures with 2,4-D or dicamba may be used.

GLYPHOSATE TOLERANT COTTON[*]

[*Not registered for sale or use in California.]

Types of Applications: Pre-plant, At-planting, Pre-emergence, Post-emergence (In-crop), Selective equipment (In-crop), Pre-harvest

For specific rates of application and instructions for control of annual and perennial weeds see the WEEDS CONTROLLED section of this label.

Maximum Application Rates/Acre	
Combined total per year for all applications	16.2 qts/ (8.1 lbs. ai.)
Total of all pre-plant, at-planting, pre-emergence applications	5 qts. (2.5 lbs. a.i.)
Total of all In-crop applications from ground crackling to lay-by	4 qts. (2 lbs. a.i.)
Maximum Pre-harvest application rate	2 qts. (1 lbs. a.i.)
Combined total of all In-crop applications from emergence through harvest	6 qts. (3 lbs. a.i.)

USE PRECAUTION:

- See the **GLYPHOSATE TOLERANT CROPS** section of this label for precautionary instructions for use in glyphosate tolerant crops.

USE RESTRICTION:

- Pre-Harvest Interval (PHI): 7 days between application and harvest. Allow a minimum of 7 days between application and harvest.
- The combined total application of this product from cotton emergence through harvest must not exceed 6 qts per acre.
- DO NOT ADD ADDITIONAL SURFACTANT OR ADDITIVES CONTAINING SURFACTANT TO THIS PRODUCT (OTHER THAN THOSE CONTAINED IN ANY TANK MIX PRODUCT) FOR OVER-THE-TOP APPLICATIONS TO GLYPHOSATE TOLERANT COTTON.

PRE-PLANT, AT-PLANTING, PRE-EMERGENCE

This product may be applied before, during or after planting glyphosate tolerant cotton.

Tank Mixtures: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Ensure that the specific product being used in the tank mixture is registered for application prior to the emergence of cotton. See the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label for more information on tank mixtures.

USE RESTRICTION:

- Maximum quantity of this product that may be applied for all pre-plant, at-planting and pre-emergence applications combined is 5 qts. per acre per year.

POST-EMERGENCE (IN-CROP)

This product may be applied over-the-top of glyphosate tolerant cotton (in-crop).

The product may be applied over-the-top of glyphosate tolerant cotton at rates up to 33 fl. oz. per acre per application from ground cracking until the 4-leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter).

USE RESTRICTIONS:

- NO MORE THAN 2 OVER-THE-TOP BROADCAST APPLICATIONS MAY BE MADE FROM CROP EMERGENCE THROUGH THE 4-LEAF (NODE) STAGE OF DEVELOPMENT.
- SEQUENTIAL OVER-THE-TOP OR POST-DIRECTED APPLICATIONS OF THIS PRODUCT IN-CROP MUST BE AT LEAST 10 DAYS APART AND COTTON MUST HAVE AT LEAST 2 NODES OF INCREMENTAL GROWTH BETWEEN APPLICATIONS.

USE PRECAUTIONS:

- Over-the-top applications made after the 4-leaf (node) stage of development may result in boll loss, delayed maturity and/or yield loss.
- Some tank mixtures may cause leaf yellowing and/or leaf crinkling when applied post-emergence (in-crop). Refer to specific tank mix partner product label for information.
- Dual MAGNUM applied over-the-top of glyphosate tolerant cotton may cause leaf injury in the form of necrotic spotting.

Tank Mixtures:

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Ensure that the specific product being used in the tank mixture is registered for application post-emergence (in-crop) to cotton. See the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label for more information on tank mixtures.

Salvage Treatment:

This treatment may be used after the 4-leaf stage of development and must only be used where weeds threaten to cause the loss of the crop.

Apply 33 fl. oz. per acre either as an over-the-top application or as a post-directed treatment sprayed higher on the cotton plants and over the weeds.

USE PRECAUTION:

- Salvage treatments will result in significant boll loss, delayed maturity and/or yield loss.

USE RESTRICTIONS:

- No more than one salvage treatment must be used per growing year.
- Maximum quantity of this product that may be applied for all in-crop applications from ground-cracking to lay-by combined is 3.75 qts. per acre per year.
- DO NOT ADD ADDITIONAL SURFACTANT OR ADDITIVES CONTAINING SURFACTANT TO THIS PRODUCT (OTHER THAN THOSE CONTAINED IN ANY TANK MIX PRODUCT) FOR OVER-THE-TOP APPLICATIONS TO GLYPHOSATE TOLERANT COTTON.

Selective Equipment (In-Crop)

This product may be applied using precision post-directed or hooded sprayers at rates up to 33 fl. oz. per acre per application through

lay-by. At this stage, use post-directed equipment that directs the spray to the base of the cotton plants. Avoid contact of the herbicide spray with leaves of the cotton plant to the maximum extent possible. To minimize spray contact, maintain a low spray pressure (less than 30 lbs. per square inch) and place nozzles in a low position directing a horizontal spray pattern under the leaves of the cotton plant and onto the weeds in the row. For best results, make applications while weeds are small (less than 3" in height). See additional use instructions in the **APPLICATION EQUIPMENT AND TECHNIQUES** section of this label.

Tank Mixtures: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Ensure that the specific product being used in the tank mixture is registered for application post-emergence (in-crop) to cotton. See the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label for more information on tank mixtures.

USE PRECAUTIONS:

- Some tank mixtures may cause leaf yellowing and/or leaf crinkling when applied post-emergence (in-crop). Refer to specific tank mix partner product label for information.
- Staple may cause leaf yellowing and/or leaf crinkling when applied post-emergence (in-crop).

USE RESTRICTIONS:

- NO MORE THAN 2 APPLICATIONS OF THIS PRODUCT MUST BE MADE FROM THE 5-LEAF STAGE THROUGH LAY-BY.
- SEQUENTIAL OVER-THE-TOP OR POST-DIRECTED APPLICATIONS OF THIS PRODUCT IN-CROP MUST BE AT LEAST 10 DAYS APART AND COTTON MUST HAVE AT LEAST 2 NODES OF INCREMENTAL GROWTH BETWEEN APPLICATIONS.

GLYPHOSATE TOLERANT SOYBEANS[*]

[*Not registered for sale or use in California.]

Types of Applications: Pre-plant, At-planting, Pre-emergence, Post-emergence (In-crop), Pre-harvest, Post-harvest

For specific rates of application and instructions for control of annual and perennial weeds see the **WEEDS CONTROLLED** section of this label.

Maximum Application Rates/Acre	
Combined total per year for all applications	16.2 qts (8.1 lbs. a.i.)
Total of all pre-plant, at-planting, pre-emergence applications	5 qts. (2.5 lbs. a.i.)
Total of all in-crop applications from cracking through flowering (R2 stage soybeans)	96 fl. oz. (1.5 lbs. a.i.)
Maximum pre-harvest application rate	33 fl. oz. (0.5 lbs. a.i.)

USE PRECAUTION:

- See the **GLYPHOSATE TOLERANT CROPS** section of this label for general precautionary instructions for use in glyphosate tolerant crops.

USE RESTRICTIONS:

- The maximum combined total quantity of this product for all applications in a year is 8 qts. per acre.

PRE-PLANT, AT-PLANTING, PRE-EMERGENCE

This product may be applied before, during, or after planting glyphosate tolerant soybeans.

Tank Mixtures: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Ensure that the specific product being used in the tank mixture is registered for application prior to emergence of soybeans. See the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label for more information on tank mixtures.

USE RESTRICTION:

- Maximum quantity of this product that may be applied for all pre-plant, at-planting and pre-emergence applications combined is 5 qts. per acre per year.

POST-EMERGENCE (IN-CROP)

This product may be used to control listed annual grasses and broadleaf weeds in glyphosate tolerant soybeans. Applications of this product can be made in glyphosate tolerant soybeans from emergence (cracking) through flowering (R2 stage soybeans). R2 stage soybeans ends when a pod 5 millimeters (3/16 inch) long appears at one of the four uppermost nodes on the main stem with a fully developed leaf (R3 stage). See the **WEEDS CONTROLLED** section of this label for rate instructions for specific weeds. In general, an initial application of 33 fl. oz. per acre on 2" - 8" tall weeds is specified. Weeds will generally be 2" - 8" tall, 2 - 5 weeks after planting.

If the initial application is delayed and weeds are larger, apply the higher listed rate of this product. This product may be applied to up to 2 qts. per acre as a single, in-crop application for control of annual weeds and where dense weed populations exist.

A 33 - 66 fl. oz. per acre rate (single or multiple applications) of this product will control or suppress listed perennial weeds. For best results, allow perennial weed species to achieve at least 6" of growth before spraying with this product.

USE PRECAUTIONS:

- Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this product may be necessary to control late flushes of weeds.
- IN THE SOUTHERN STATES, A SEQUENTIAL APPLICATION OF THIS PRODUCT WILL BE REQUIRED TO CONTROL NEW FLUSHES OF WEEDS IN THE GLYPHOSATE TOLERANT SOYBEAN CROP.

Tank Mixtures: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Ensure that the specific product being used in the tank mixture is registered for application post-emergence (in-crop) to soybeans. See the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label for more information on tank mixtures.

USE PRECAUTION:

- In some cases, tank mix products will cause visual soybean injury.

USE RESTRICTIONS:

- The combined total application from crop emergence through harvest must not exceed 96 fl. oz. per acre (1.5 lb. a.i.).
- The maximum rate for any single in-crop application is 66 fl. oz. per acre (1.03 lbs. a.i.).
- The maximum combined total of this product that can be applied during flowering (R2 stage soybeans) is 66 fl. oz. per acre (1.03 lbs. a.i.).

PRE-HARVEST

This product may be applied to glyphosate tolerant soybeans for weed control prior to harvest.

USE PRECAUTION:

- Care must be taken to avoid excessive seed shatter loss due to ground application equipment.

USE RESTRICTION:

- Pre-Harvest Interval (PHI): 14 days between final application and harvest of soybean grain or feeding of soybean grain, forage or hay.

POST-HARVEST

This product may be applied for weed control after harvest of glyphosate tolerant soybeans. The higher listed rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

CONSERVATION TILLAGE, MINIMUM TILLAGE, AND NO-TILL SYSTEMS

PRE-HARVEST APPLICATIONS COTTON[*], SOYBEANS[*], SORGHUM (GRAIN) [*] AND WHEAT[*]

When applied as directed under the conditions described, **BGT** controls annual and perennial weeds listed on this label prior to the harvest of cotton, soybeans, grain sorghum (milo)[*], and wheat.

For specific rates and application instructions for control of various annual and perennial weeds, see the **WEEDS CONTROLLED** section of this label.

This product may be applied by both ground and aerial application equipment.

See the **APPLICATION EQUIPMENT AND TECHNIQUES** section of this label for instructions for ground and aerial applications.

USE RESTRICTIONS:

- DO NOT APPLY MORE THAN 1 QT. (0.5 lbs a.i.) PER ACRE OF THIS PRODUCT BY AIR.
- Do not apply to crops grown for seed. Reduction in germination or vigor may occur.

[*Not registered for sale or use in California.]

PRE-HARVEST APPLICATIONS COTTON[*]

Broadcast Applications - This product may be applied using either aerial or ground spray equipment. For ground applications with broadcast equipment, apply this product in 10 - 20 gals. of water per acre. For aerial applications, apply this product in 3 - 10 gals. of

water per acre.

For specific rates of application and instructions for control of annual and perennial weeds see the **WEEDS CONTROLLED** section of this label.

This product provides weed control and cotton regrowth inhibition when applied prior to the harvest of cotton. Apply 1 - 2 qts. of this product in 3 - 10 gals. of water per acre for cotton regrowth inhibition. Do not apply more than 2 qts. (1 lb a.i.) of this product per acre for pre-harvest applications.

Make application after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential.

USE RESTRICTIONS:

- THE USE OF ADDITIVES FOR PRE-HARVEST APPLICATION TO COTTON IS PROHIBITED.
- Preharvest Interval (PHI): 7 days between application and harvest of cotton.
- Do not feed or graze treated cotton forage or hay following pre-harvest applications.

[*Not registered for sale or use in California.]

PRE-HARVEST APPLICATIONS SOYBEANS[*]

Apply after all pods have set and lost all green color. For specific rates of application and instructions for control of annual and perennial weeds see the **WEEDS CONTROLLED** section of this label.

USE PRECAUTIONS:

- Care must be taken to avoid excessive seed shatter loss due to ground application equipment.

USE RESTRICTIONS:

- Pre-Harvest Interval (PHI): 7 days between application and harvest of soybeans.
- Do not graze or harvest treated crop for livestock feed within 25 days of last pre-harvest application.
- DO NOT APPLY MORE THAN 6 QTS. (3 lbs. a.i.) PER ACRE OF THIS PRODUCT FOR PRE-HARVEST APPLICATIONS.

[*Not registered for sale or use in California.]

PRE-HARVEST APPLICATIONS GRAIN SORGHUM (MILO)[*]

Make applications at 30% grain moisture or less and at least 7 days prior to harvest.

Apply up to 2 qts. of this product per acre.

USE RESTRICTIONS:

- Pre-Harvest Interval (PHI): 7 days between application and harvest.
- Do not feed or graze treated sorghum forage or hay following pre-harvest applications.
- DO NOT APPLY MORE THAN 2 QTS. (1 lb a.i.) PER ACRE OF THIS PRODUCT FOR PRE-HARVEST APPLICATIONS.

[*Not registered for sale or use in California.]

PRE-HARVEST APPLICATIONS WHEAT[*]

Apply after hard dough stage of grain (30% or less grain moisture). For specific rates of application and instructions for control of annual and perennial weeds see the **WEEDS CONTROLLED** section of this label.

USE RESTRICTIONS:

- Pre-Harvest Interval (PHI): 7 days prior to harvest.
- DO NOT APPLY MORE THAN 1 QT. (0.5 lbs. a.i.) PER ACRE OF THIS PRODUCT FOR PRE-HARVEST APPLICATIONS TO WHEAT.

[*Not registered for sale or use in California.]

TREE AND VINE CROPS: Apples[*], Grape[*], Lemon[*], Orange[*], Peach[*], Plum[*], Walnut[*]

BGT is specified for weed control in established groves, vineyards, and orchards, or for site preparation prior to transplanting crops listed in this section. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, orchard guns or with wiper applicator equipment, except as directed in this section. See the **APPLICATION EQUIPMENT AND TECHNIQUES** section of this label for specific information on use of equipment.

When applying this product, refer to the **WEEDS CONTROLLED** section of this label and to specific instructions in this section for rates to be used.

NOTE: Repeat treatments may be necessary to control weeds originating from underground parts of untreated weeds or from seeds. This product does not provide residual weed control. For subsequent weed control, use repeated applications of this product.

USE RESTRICTIONS:

- Do not apply more than 10.6 qts. (5.3 lbs. ai.) of this product per acre per year.

USE PRECAUTIONS:

- EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT OR OTHER PARTS OF TREES OR VINES.
- CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.
- AVOID PAINTING CUT STUMPS WITH THIS PRODUCT, AS INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES.
- Reduced control may result when applications are made to annual or perennial weeds that have been mowed grazed or cut and have not been allowed to regrow to the specified stage for treatment.

Tank Mixtures: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. See the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label for more information on tank mixtures.

Middle Management (For Annual and Perennial Listed Weeds in Middle Between Rows of Listed Tree and Vine Crops)

For lemon or orange crops, treat uniformly between trees.

Apply the specified rates of this product, either alone or in mixtures by spray volume in 3 - 10 gals. of water per acre. Apply when weeds are actively growing and less than 6" in height or diameter. If weeds are under drought stress, irrigate prior to application. Reduced control may occur if weeds have been mowed prior to application. Up to 48 fl. oz. per acre of this product may be used to control weeds, which have been mowed, are stressed or are growing in dense populations.

[*Not registered for sale or use in California.]

TREE CROPS[*]

For specific rates of application and instructions for control of annual and perennial weeds see the **WEEDS CONTROLLED** section of this label.

CITRUS[*]

Lemon and Orange

Use Restriction:

- Pre-Harvest Interval (PHI): 1 day between last application and harvest.

WALNUTS[*]

Use Restriction:

- Pre-Harvest Interval (PHI): 3 days between last application and harvest.

POME FRUIT[*]

Apple

Use Restriction:

- Pre-Harvest Interval (PHI): 1 day between last application and harvest.

STONE FRUIT[*]

Peaches and Plums/Prunes.

Use Restriction:

- Pre-Harvest Interval (PHI): 17 days between last application and harvest.

For Peaches grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina and Tennessee only - Apply with a shielded boom sprayer or shielded wiper applicator, which prevents any contact with the foliage or bark of trees. Apply no later than 90 days after first bloom. Applications made after this time may result in severe damage. Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees that have been planted in the orchard for 2 or more years. EXTREME CARE MUST BE TAKEN TO ENSURE NO PART OF THE PEACH TREE IS CONTACTED.

Any application equipment listed in this section may be used in peaches and plums/prunes growing in Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah and Washington, except for peaches grown in states specified in the following paragraph. In all other states, use wiper equipment only.

[*Not registered for sale or use in California.]

GRAPES[*]

For specific rates of application and instructions for control of annual and perennial weeds see the **WEEDS CONTROLLED** section of this label.

- Any variety of table, wine, or raisin grapes may be treated with any equipment listed in this section.
- Applications must not be made when green shoots, canes, or foliage are in the spray zone.
- In the Northeast and Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury.

Use Restriction:

- Pre-Harvest Interval (PHI): 14 days between last application and harvest.

[*Not registered for sale or use in California.]

NON-CROP USES[*]

See **PRODUCT INFORMATION** and **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** sections of this label for essential product performance information and the following Non-Crop sections for specific uses.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE TURFGRASSES, TREES, SHRUBS OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds. Where repeat applications are necessary, do not exceed 10.6 qts. of this product per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate. This product does not provide residual weed control. For subsequent weed control, follow a label-approved herbicide program.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. See the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label for more information on tank mixtures.

[*Not registered for sale or use in California.]

INDUSTRIAL, RECREATIONAL, AND PUBLIC AREAS[*]

When applied as directed for **NON-CROP USES**, under conditions described, **BGT** controls annual and perennial weeds listed on this label growing in areas including airports, ditch banks, dry ditches, dry canals, fencerows, golf courses, highways, industrial plant sites, lumberyards, parking areas, parks, petroleum tank farms and pumping installations, pipelines, power and telephone rights-of-way, railroads, roadsides, schools, storage areas, and utility substations.

For specific rates of application and instructions for control of annual and perennial weeds see the **WEEDS CONTROLLED** section of this label.

This product may be applied with recirculating sprayers, shielded applicators, or wiper applicators in any non-crop site specified on this label. See the **Selective Equipment** part of **APPLICATION EQUIPMENT AND TECHNIQUES** section of this label for information on proper use and calibration of this equipment.

[*Not registered for sale or use in California.]

FARMSTEAD WEED CONTROL[*]

When applied as directed for **NON-CROP USES** under conditions described, **BGT** controls undesirable vegetation listed on this label around farmstead building foundations, along and in fences, shelterbelts and for nonselective farmstead weed control.

For specific rates of application and instructions for control of annual and perennial weeds see the **WEEDS CONTROLLED** section of this label.

Where broadleaf weed control or suppression is desired, tank mix this product with the appropriate labeled broadleaf weed herbicide. See the **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** section of this label for more information on tank mixtures.

Farm Ditches

This product will suppress perennial grasses along farm ditches.

Apply treatments in 10 - 20 gals. of spray solution per acre to actively growing perennial grass covers. For best spray distribution and coverage, use flat fan nozzles.

[*Not registered for sale or use in California.]

CONSERVATION RESERVE PROGRAM (CRP ACRES)[*]

BGT can be used to control undesirable vegetation when rotating out of CRP acres or to suppress competitive growth and seed production of undesirable vegetation in CRP acres.

For specific rates of application and instructions for control of annual and perennial weeds see the **WEEDS CONTROLLED** section of this label.

[*Not registered for sale or use in California.]

ORNAMENTALS, TREE NURSERIES, AND CHRISTMAS TREES[*]

[*Not registered for sale or use in California.]

When applied as instructed for the conditions described for **NON-CROP USES**, **BGT** controls undesirable vegetation listed on this label prior to planting, within and around greenhouses and shadehouses, and as a post-directed spray around established ornamentals and Christmas trees.

For specific rates of application and instructions for control of various annual and perennial weeds, see the **WEEDS CONTROLLED** section of this label.

USE RESTRICTION:

- DO NOT USE THIS PRODUCT AS AN OVER-THE-TOP BROADCAST SPRAY IN ORNAMENTALS AND CHRISTMAS TREES.
- Where repeat applications are necessary, do not exceed 10.6 qts (5.3 lbs. a.i.) of this product per acre per year.

USE PRECAUTIONS:

- Precautions must be taken to protect non-target plants during site preparation applications.
- Protect desirable plants from the spray solution by using shields or coverings made of cardboard or other impermeable material.

Site Preparation

Following pre-plant applications of this product, any ornamental, nursery species, or Christmas tree species may be planted.

Greenhouse/Shadehouse Use

This product may be used to control weeds listed on this label that are growing inside greenhouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

Post-Directed Spray

Use a post-directed spray around established woody ornamental species, nursery species, or Christmas trees including those listed below. Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of established ornamental species.

Common Name	Scientific Name	Common Name	Scientific Name
Arborvitae	<i>Thuja</i> spp.	Lilac	<i>Syringa</i> spp.
Azalea	<i>Rhododendron</i> spp.	Magnolia	<i>Magnolia</i> spp.
Boxwood	<i>Buxus</i> spp.	Maple	<i>Acer</i> spp.
Crabapple	<i>Malus</i> spp.	Oak	<i>Quercus</i> spp.
Douglas Fir	<i>Pseudotsuga</i> spp.	Privet	<i>Ligustrum</i> spp.
Euonymus	<i>Euonymus</i> spp.	Pine	<i>Pinus</i> spp.
Fir	<i>Abies</i> spp.	Spruce	<i>Picea</i> spp.
Jojoba	<i>Simmondsia chinensis</i>	Yew	<i>Taxus</i> spp.
Hollies	<i>Ilex</i> spp.		

CUT STUMP TREATMENTS[*]

[*Not registered for sale or use in California.]

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with **BGT**. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. Apply a 50 - 100% solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results,

application must be made during periods of active growth and full leaf expansion.

When used according to directions for cut stump application, this product will Control, Partially Control, or Suppress many types of woody brush and tree species, some of which are listed below:

Partial List - Species Controlled or Suppressed - Cut Stump Application			
Common Name	Scientific Name	Common Name	Scientific Name
Alder	<i>Alnus</i> spp.	Saltcedar	<i>Tamarix</i> spp.
Eucalyptus	<i>Eucalyptus</i> spp.	Sweetgum	<i>Liquidambar styraciflua</i>
Madrone	<i>Arbutus menziesii</i>	Tanoak	<i>Lithocarpus densiflorus</i>
Oak	<i>Quercus</i> spp.	Willow	<i>Salix</i> spp.
Reed, Giant	<i>Arundo donax</i>		

WOODY BRUSH AND TREES[*]

[*Not registered for sale or use in California.]

USE RESTRICTION:

- If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has reached the specified stages of growth.
- Allow 7 or more days after application before tillage, mowing or removal.

USE PRECAUTIONS

- Some Autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if Fall treatments are made following a frost.

Apply **BGT** when plants are actively growing and, unless otherwise directed, after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in the late Summer or Fall after fruit formation.

In arid areas, best results are obtained when application is made in the Spring to early Summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with Fall treatments.

Repeat treatments may be necessary to control plants regenerating from underground parts or seed.

When applied as specified under the conditions described, **BGT** Controls or Partially Controls the following woody brush plants and trees (see additional notes, by species, below this listing):

See **PRODUCT INFORMATION** and **MIXING, ADDITIVES, AND APPLICATION INSTRUCTIONS** sections of this label for labeled uses and specific application instructions.

Apply **BGT** as follows to control or partially control the following woody brush and trees:

Species	Application Directions
Alder / Dewberry / Honeysuckle / Post Oak / Raspberry	For control, apply 3 - 4 qts. per acre of this product as a broadcast spray or as a 1 - 1.5% solution with hand-held equipment.
Aspen, Quaking / Cherry: Bitter, Black, Pin / Hawthorn / Oak, Southern Red / Sweetgum / Trumpet Creeper	For control, apply 2 - 3 qts. of this product per acre as a broadcast spray or as a 1 - 1.5% solution with hand-held equipment.
Birch / Elderberry / Hazel / Salmonberry / Thimbleberry	For control, apply 2 qts. per acre of this product as a broadcast spray or as a 1% solution with hand-held equipment.
Blackberry	For control, apply 3 - 4 qts. per acre of this product as a broadcast spray, or 1 - 1.5% solution with hand-held equipment. Make application after plants have reached full leaf maturity. Best results are obtained when applications are made in the late Summer or Fall. After berries have set or dropped in late Fall, blackberry can be controlled by applying a 0.75% solution of this product by total spray volume with hand-held equipment. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3 - 4 qts. of this product in 10 - 40 gals. of water per acre.
Broom: French, Scotch	For control, apply a 1.5 - 2% solution with hand-held equipment.
Buckwheat, California / Hasardia / Monkey Flower / Tobacco, Tree	For partial control of these species, apply a 1 - 2% solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.
Catclaw	For partial control, apply a 1 - 1.5% solution with hand-held equipment.
Coyote Brush	For control, apply a 1.5 - 2% solution with hand-held equipment when at least 50% of the new leaves are fully developed.
Eucalyptus	For control of eucalyptus resprouts, apply a 2% solution with hand-held equipment when resprouts are 6 - 12 ft. tall. Ensure complete coverage. Apply when plants are growing actively. Avoid application

	to drought stressed plants.
Kudzu	For control, apply 4 qts. of this product per acre as a broadcast spray or as a 2% solution with hand-held equipment. Repeat applications will be required to maintain control.
Madrone Resprouts	For suppression or partial control, apply a 2% solution of this product to resprouts less than 3 - 6 ft. tall. Best results are obtained with Spring/early Summer treatments.
Maple, Red	For control, apply as a 1 - 1.5% solution with hand-held equipment when at least 50% of the new leaves are fully developed. For partial control, apply 2 - 4 qts. of this product per acre as a broadcast spray.
Maple, Sugar / Oak, Northern Pin / Oak, Red	For control, apply as a 1 - 1.5% solution with hand-held equipment when at least 50% of the new leaves are fully developed.
Poison Ivy / Poison Oak	For control, apply 4 - 5 qts. of this product per acre as a broadcast spray or as a 2% solution with hand-held equipment. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.
Rose, Multiflora	For control, apply 2 qts. of this product per acre as a broadcast spray or as a 1% solution with hand-held equipment. Treatments must be made prior to leaf deterioration by leaf-feeding insects.
Sage, Black / Sagebrush, California / Chamise / Tallowtree, Chinese	For control of these species, apply a 1% solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.
Tanoak Resprouts	For suppression or partial control, apply a 2% solution of this product to resprouts less than 3 - 6 ft. tall. Best results are obtained with Fall applications.
Willow	For control, apply 3 qts. of this product per acre as a broadcast spray, or as a 1% solution with hand-held equipment.
Other Woody Brush and Trees listed on this label	For partial control, apply 2 - 5 qts. of this product per acre as a broadcast spray or as a 1 - 2% solution with hand-held equipment.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Open dumping is prohibited. Store in original container only. Keep containers closed when not in use. Separate pesticides during storage to prevent cross-contamination of other pesticides, fertilizers, food, and feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed must be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed.

[CONTAINER HANDLING:

Nonrefillable container (equal to or less than 5 gallons): Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.]

[Nonrefillable Container (greater than 5 gallons to 270 gallons): Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.]

[Refillable Container (275 gallons or larger): Refill this container with glyphosate only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Dispose of in a sanitary landfill or by other procedures approved by State and local authorities.]

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control

of Ciech Sarzyna S.A. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Ciech Sarzyna S.A. and Seller harmless for any claims relating to such factors.

Ciech Sarzyna S.A. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Ciech Sarzyna S.A., and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CIECH SARZYNA S.A. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Ciech Sarzyna S.A. nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CIECH SARZYNA S.A. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CIECH SARZYNA S.A. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Ciech Sarzyna S.A. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Ciech Sarzyna S.A.

[All trademarks are the property of their respective owners.]