



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

October 6, 2026

Nancy Huddy
Agent for Agrotecnologías Naturales S.L.
SciReg, Inc.
12733 Director's Loop
Woodbridge, VA 22192

Subject: Pesticide Registration Improvement Act (PRIA) Labeling Amendment – To Add the First Residential Use for the Active Ingredient and Additional Crops
Product Name: Tricoten WP
EPA Registration Number: 96029-2
EPA Receipt Date: 01/11/2025
Action Case Number: 00642663

Dear Nancy Huddy:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable under FIFRA section 3(c)(5).

You must submit and/or cite all data required for registration or registration review of your product when the U.S. Environmental Protection Agency requires all registrants of similar products to submit such data.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. Pursuant to 40 CFR 156.10(a)(6), you must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by EPA. See FIFRA section 2(p)(2). If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process, FIFRA section 12(a)(1)(B). Therefore, should EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to EPA's Office of Enforcement and Compliance Assurance.

Page 2 of 2

EPA Reg. No. 96029-2

Action Case No. 00642663

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Jennifer Odom-Douglas via email at odomdouglas.jennifer@epa.gov.

Sincerely,



Digitally signed by
JEANNINE KAUSCH
Date: 2026.10.06
16:40:19 -04'00'

Jeannine Kausch, Senior Advisor
Immediate Office
Biopesticides and Pollution
Prevention Division
Office of Pesticide Programs

Enclosure (1): Stamped Labeling Dated 10/06/2026

TRICOTEN WP

For control of fungal diseases on crops caused by: *Alternaria* spp., *Botrytis cinerea*, *Cephalosporium maydis*, *Didymella bryoniae*, Esca and wood diseases (*Phaeomoniella chlamydospora*, *Togninia minima*, *Fomitiporia mediterranea*, *Botryosphaeria* spp.), *Fusarium* spp., *Gibberella fujikuroi*, *Leptosphaeria lindquistii*, *Macrophomina phaseolina*, *Magnaporthe oryzae*, *Mycosphaerella* spp., *Phytophthora* spp., *Pythium* spp., *Rhizoctonia* spp., *Sclerotinia* spp., *Sclerotium* spp., and *Verticillium* spp.

ACTIVE INGREDIENT:

Trichoderma atroviride strain AT10 100%

Product contains a minimum concentration of 1×10^8 CFU/gram

KEEP OUT OF REACH OF CHILDREN

CAUTION

{Optional referral statements when booklets and container labels are used, or when First Aid statement is not on front panel:} [See [side][back] panel for [additional][complete] Precautionary Statements [,First Aid] [and Directions for Use].] [See booklet for [additional] Precautionary Statements, Directions for Use, and Storage and Disposal.] [See booklet for complete Directions for Use.] [See booklet for additional Precautionary Statements and Directions for Use.]


FIRST AID	
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER: Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For non-emergency information concerning this product, call the National Pesticide Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PT (NPIC Web site: www.npic.orst.edu). For emergencies, call the poison control center 1-800-222-1222, 24 hours a day, 7 days a week.	

EPA Reg. No. 96029-2

EPA Est. No.

Batch Code:

Net Weight:



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear
- Mixer/loaders and applicators must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air-purifying respirator with an HE filter. (Repeated exposures to high concentrations of microbial proteins can cause allergic sensitization.)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water) is:

- Protective eyewear
- Coveralls
- Chemical resistant gloves (made of any waterproof material)
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until [sprays have dried] [or] [seeds have dried and been packaged].

Product Information

TRICOTEN WP is a preventative and curative biological fungicide for control of plant diseases. It is composed of the active ingredient, *Trichoderma atroviride* strain AT10. For control of fungal diseases on crops caused by: *Alternaria* spp., *Botrytis cinerea*, *Cephalosporium maydis*, *Didymella bryoniae*, Esca and wood diseases (*Phaeomoniella chlamydospora*, *Togninia minima*, *Fomitiporia mediterranea*, *Botryosphaeria* spp.), *Fusarium* spp., *Gibberella fujikuroi*, *Leptosphaeria lindquistii*, *Macrophomina phaseolina*, *Magnaporthe oryzae*, *Mycosphaerella* spp. *Phytophthora* spp., *Pythium* spp., *Rhizoctonia* spp., *Sclerotinia* spp., *Sclerotium* spp., and *Verticillium* spp.

Use Information

TRICOTEN WP must not be tank mixed with chemicals that contain the following active ingredients: benomyl, imazalil, propiconazole, tebuconazole, and triflumizole. Fungal treatments can affect or inhibit the development of *Trichoderma* fungi. When tank mixing, follow the most restrictive of the labeling limitations and precautions of all products used in mixtures. Consult the technical service for more information. The cultivation should be subject to regular fertilization, not only to the application of the product. TRICOTEN WP contains live spores of a microbe that must be used prior to disease onset. TRICOTEN WP becomes active in soil or on plants when temperatures are above 50°F and is not effective while temperatures remain cold. If sterilization or fumigation of the soil is employed, TRICOTEN WP must be applied afterwards. Apply TRICOTEN WP as a foliar spray (including via sprinkler irrigation) or via drip irrigation according to the application instructions below.

Application Instructions

All Chemigation

Apply this product only through sprinkler, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move, or drip (trickle) irrigation system(s). Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

It is recommended to mix TRICOTEN WP in a pesticide supply tank with the specified dose of water. Agitation is recommended to homogenize the solution and facilitate the complete dilution of the product. TRICOTEN WP has to be diluted in the adequate volume for its application. For further information about the volume of water, check the Field Application table below.

Following the timing indicated in the Field Application table below, apply the complete specified dose of TRICOTEN WP. The dilution process is recommended to be carried out gradually to avoid flocculation of the product.

The product should be applied during the second third of the irrigation water application, after the soil has been adequately wetted. The final third of the irrigation water should then be applied to incorporate the product into the soil and move it into the treated root zone.

Drip (Trickle) Chemigation

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional inter-locking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Chemigation through systems connected to public water systems

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Sprinkler Chemigation

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Field Applications

Crop	Application Method	Application Rate	Application Instructions
Fruiting Vegetables, including: Eggplant, Sweet and Hot Peppers, Tomatillos, Tomatoes	Drip irrigation	0.5 to 1 lb. in 500 gal of water per acre. Do not apply more than 3 lb. per acre per crop season.	Begin applications at germination and continue until harvest. For the first application, apply 1 lb. per acre. For the remaining four applications, apply 0.5 lb. per acre. Apply at 10-day intervals. Do not make more than 5 applications per crop cycle.
Leafy Vegetables (including Brassica Vegetables): Lettuce (head, leaf, romaine), Spinach, Amaranth (Chinese spinach)			
Brassica (Cole) Leafy Vegetables: Broccoli (including Chinese), Brussels sprout, Cabbage, Cauliflower, Chinese cabbage, (bok choy), Choi sum, Collard, Kohlrabi, Chinese Mustard Cabbage			
Cucurbit Vegetables, including: Cucumbers, Melons (i.e., Chinese Waxgourd, Citron Melon, Muskmelons, or Watermelon), Gourds, Pumpkins, Squash			
Berries and Small Fruits, including: Blackberries, Blueberries, Currants, Elderberries, Gooseberries, Huckleberries, Loganberries, Raspberries, Strawberries, Grapes			
Root and Tuber Vegetables, including: Beets, Sugar Beets, Carrots, Celeriac, Chicory,			

Crop	Application Method	Application Rate	Application Instructions
Horseradish, Parsnip, Radish, Rutabaga, Salsify, Turnips, Potatoes, Sweet Potatoes, Yams, Jerusalem Artichoke, Cassava, Ginger, Ginseng			
Bulb Vegetable (Allium spp.): Garlic, Leek, Onion (dry, green)			
Legume Vegetables: Beans, snap, Phaseolus, chickpea, lentil, pea, soybean, Chinese longbean, Mung bean including all hybrids and varieties of these			
Pome Fruit: Apple, Pear, Crabapple, Loquat, Mayhaw, Oriental Pear, Quince			
Stone Fruit: Bearing and Non-Bearing Apricot, Sweet and Tart Cherry, Nectarine, Peach, Plum, Plumcot, Prune, and all hybrids and cultivars of these			
Citrus Fruits: Calamondin, Citrus Citron, Citrus Hybrids (i.e., chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Orange, Pummelo, Satsuma Mandarin, and all cultivars and hybrids of these			
Olive			
Pomegranate			

Crop	Application Method	Application Rate	Application Instructions
Tree Nuts: Almonds, Beechnut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory, Hazelnuts, Macadamia, Pecans, Pistachios, Walnuts			
Tropical and Subtropical Fruit: Acerola, Atemoya, Avocado, Banana, Canistel, Cherimoya, Custard Apple, Guava, Illama, Kiwifruit, Longan, Loquat, Lychee, Mango, Mangosteen, Papaya, Passionfruit, Pawpaw, Persimmon, Pineapple, Rambutan, Black Sapote, Mamey Sapote, Sapodilla, Soursop, Starfruit, Star Apple, Sugar Apple, Tamarind			
Small Fruit Vine Climbing: Table, Wine, Raisin Grapes, Kiwi			
Hops			
Hemp			
Muscadines			
Stalk, Stem and Leaf Petiole Vegetable: Asparagus, Celery, Kale			
Herbs and Spices: Angelica, Basil, Catnip, Chervil, Catnip, Chamomile, Chive, Cilantro-Leaf, Coriander-Leaf, Curry, Dillweed, Fennel, Hyssop, Lavender, Lemongrass, Marigold, Marjoram, Nasturtium, Parsley, Rosemary, Sage, Savory-			

Crop	Application Method	Application Rate	Application Instructions
Summer, Savory- Winter, Sweet Bay, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood Mint, Peppermint, Spearment			
Flowers, Bedding Plants, and Ornamentals: Carnation, Chrysanthemums, Cyclamen, Geranium, Gerbera, Hollyhock, Lily, Rose, Pansy, Petunia, Poinsettia, Primrose, Snapdragon			
Ornamental Trees Non-Flowering Deciduous and Non- Deciduous Landscape, Ornamental, Forest, and Nursery Trees: Ash, Elm, Linden, Maple, And Sycamore			
Evergreen/Conifers and Christmas Trees: Cedar, Fir, Pine, Spruce, Yew, Arborvitae			
Flowering and Non- Bearing Fruit Trees: Crabapple, Hawthorn, Cherry, Plum, Pear			
Tropical Foliage and Flowering Plants: Fern, English ivy, Pothos, Poinsettia			
Nursery, Shrubs, and Woody Ornamentals: Azalea, Holly, Pittosporum, Rose			

Crop	Application Method	Application Rate	Application Instructions
Turf Grass: Turf, golf courses-all uses, sports fields, sod and seed production	Drip irrigation	0.5 to 1 lb. in 500 gal of water per acre. Do not apply more than 3 lb. per acre per crop season.	Begin applications at turfgrass seeding or establishment. Make the first application at 1 lb. per acre, followed by up to four applications at 0.5 lb. per acre at 10-day intervals. Do not make more than 5 applications per year.
Grapevine	Foliar spray (ground)	Gray mold (<i>Botrytis cinerea</i>) 0.5 to 1 lb. per 30 to 80 gals of water per acre. Do not apply more than 3 lb. per acre per crop season.	Begin applications at fruit set and continue until harvest. For the first application, apply 1 lb. per acre. For the remaining four applications, apply 0.5 lb. per acre. Apply at 10-day intervals. Do not make more than 5 applications per crop cycle.
		Esca and wood diseases (<i>Phaeomoniella chlamydospora</i>, <i>Togninia minima</i>, <i>Fomititiporia mediterranea</i>, <i>Botryosphaeria</i> spp.) 1 lb. per 20 to 40 gal. of water per acre Do not apply more than 2 lb. per acre per crop season.	Make first application after winter pruning and second application after spring thinning. Make applications at least 90 days apart. Do not make more than 2 applications per crop cycle.

Greenhouse Applications

Crop	Application Method	Application Rate	Application Instructions
Fruiting Vegetables, including: Eggplant, Sweet and Hot Peppers, Tomatillos, Tomatoes Cucurbit Vegetables, including: Cucumbers, Melons (i.e., Chinese Waxgourd, Citron Melon, Muskmelons, or Watermelon), Gourds, Pumpkins, Squash Berries and Small Fruits, including: Blackberries, Blueberries, Currants, Elderberries, Gooseberries, Huckleberries, Loganberries, Raspberries, Strawberries, Grapes Leafy Vegetables (including Brassica Vegetables): Lettuce (head, leaf, romaine), Spinach, Amaranth (Chinese spinach) Brassica (Cole) Leafy Vegetables: Broccoli (including Chinese), Brussels sprout, Cabbage, Cauliflower, Chinese cabbage, (bok choy), Choi sum, Collard, Kohlrabi, Chinese Mustard Cabbage	Drip irrigation	0.5 to 1 lb. in 500 gal. of water per acre. Do not apply more than 3 lb. per acre per crop season.	Begin applications at germination and continue until harvest. For the first application, apply 1 lb. per acre. For the remaining four applications, apply 0.5 lb. per acre. Apply at 10-day intervals. Do not make more than 5 applications per crop cycle.

Herbs and Spices: Angelica, Basil, Catnip, Chervil, Catnip, Chamomile, Chive, Cilantro-Leaf, Coriander-Leaf, Curry, Dillweed, Fennel, Hyssop, Lavender, Lemongrass, Marigold, Marjoram, Nasturtium, Parsley, Rosemary, Sage, Savory- Summer, Savory- Winter, Sweet Bay, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood Mint, Peppermint, Spearmint			
Flowers, Bedding Plants, and Ornamentals: Carnation, Chrysanthemums, Cyclamen, Geranium, Gerbera, Hollyhock, Lily, Rose, Pansy, Petunia, Poinsettia, Primrose, Snapdragon			
Ornamental Trees Non-Flowering Deciduous and Non- Deciduous Landscape, Ornamental, Forest, and Nursery Trees: Ash, Elm, Linden, Maple, And Sycamore			
Evergreen/Conifers and Christmas Trees: Cedar, Fir, Pine, Spruce, Yew, Arborvitae			
Flowering and Non- Bearing Fruit Trees: Crabapple, Hawthorn, Cherry, Plum, Pear			
Tropical Foliage and Flowering Plants:			

Fern, English ivy, Pothos, Poinsettia			
Nursery, Shrubs, and Woody Ornamentals: Azalea, Holly, Pittosporum, Rose			

Seed Treatment Applications

Crop	Application Method	Application Rate	Application Instructions
Cereal seeds (winter and spring wheat, triticale, barley, oats, and rye)	Seed treatment (coating)	0.5 lb. in 2 gal. of water per ton of seed	Apply TRICOTEN WP as a slurry, a coating, in a pellet, or during seed priming using standard slurry or mist-type seed treatment application equipment. To assure uniform application, add half the required amount of TRICOTEN WP to half the seed in the hopper, mix with a wooden paddle, and then add the remaining seed and TRICOTEN WP. Dilute with sufficient amount of water to achieve uniform application.
Grass, Forage, Fodder, and Hay (including pasture grasses and grasses grown for hay or silage such as Bermuda grass, bluegrass, brome grass, and fescue)			
Sunflower			
Rice			

Corn (all types, including field corn, popcorn, sweet corn, and corn produced for seed)	Seed treatment (coating)	1 lb. in 2 gal. of water per ton of seed	Apply TRICOTEN WP as a slurry, a coating, in a pellet, or during seed priming using standard slurry or mist-type seed treatment application equipment. To assure uniform application, add half the required amount of TRICOTEN WP to half the seed in the hopper, mix with a wooden paddle, and then add the remaining seed and TRICOTEN WP. Dilute with sufficient amount of water to achieve uniform application.
Alfalfa (for use in forage and seed crops, including birdsfoot trefoil)			
Clover (for use in forage and seed crops)			
Soybean			
Cotton			
Peanut			
Tobacco			
Canola			

For seed treated on the farm, do not use for food or feed purposes or process for oil. Additionally, treat only those seeds needed for immediate use, minimizing the interval between treatment and planting.

All seed treated commercially with this product must be adequately dyed with a suitable color to prevent accidental use as food for humans or feed for animals. Use an EPA-approved dye or colorant that imparts an unnatural color to the seed. Any dye or colorant added to treated seed must be cleared for use pursuant to 40 CFR 153.155.

Seed Labeling

Seed treated with TRICOTEN WP that is then packaged for future use must contain the following labeling on the outside of the seed package.

- This seed has been treated with *Trichoderma atroviride* strain AT10.
- Do not use this treated seed for food, feed or oil production.
- Wear long-sleeved shirt, long pants, shoes, socks and gloves when handling treated seed.

- Store treated seed away from food and feed.
- Do not allow access by children, pets or livestock.
- Do not reuse empty seed containers.
- Store or dispose of surplus treated seed or empty seed containers according to local requirements.
- Do not contaminate bodies of water when disposing of excess treated seed or wash waters from planting equipment.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store this product in its original container. Store product away from direct sunlight between $4 \pm 2^{\circ}\text{C}$.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Warranty

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. Seller warrants that this product conforms to the description on the label and is reasonably fit for the purposes stated on the label when used and stored in accordance with directions under normal conditions of use. To the extent permitted by applicable law, this warranty does not extend to use of this product under conditions not reasonably foreseeable by the Seller, and Buyer and User assume the risk of any such use. To the extent permitted by applicable law, Seller disclaims all other warranties express or implied, including any warranty of fitness or merchantability. To the extent permitted by applicable law, Seller shall not be liable for consequential, special or indirect damages resulting from use or handling of this product and Seller's sole liability and Buyer's and User's exclusive remedy shall be limited to refund of the purchase price. This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified.

Agrotecnologías Naturales S.L.
Ctra.T-214, s/n Km 4,125
43762 Riera de Gaià La
Tarragona
Spain