



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

June 26, 2025

SENT BY EMAIL

Dr. Shiv Sharma
shiv@cropprotectdirect.com
JABCO, LLC

Subject: Labeling Notification per Pesticide Registration Notice (PRN) 98-10 - add ABN
Product Name: CROP PROTECT DIRECT AZOXYPROP SE FUNGICIDE
Admin Number: 101458-9
EPA Receipt Date: 05/20/2025
Action Case Number: 00657783

Dear Dr. Shiv Sharma:

The U.S. Environmental Protection Agency is in receipt of your application for notification under Pesticide Registration Notice 98-10 for the above referenced product. The EPA has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The labeling submitted with this application has been stamped "Notification" and will be placed in our records.

The alternate brand name: CROP PROTECT DIRECT AZOXYPROP SE has been added to the registration. Our records have been updated accordingly.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

If you have questions, please contact Elisha Graham via email at graham.elisha@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Elisha Graham". The script is elegant and cursive, with the first letters of both first and last names being capitalized and prominent.

Elisha Graham, Risk Manager
FB, RD
Office of Pesticide Programs

[Note to reviewer: [Text] in brackets denotes optional text].

[Note to reviewer: {Text} in braces denotes where in the final label text will appear.]

{BOOKLET FRONT PANEL}

Azoxystrobin	GROUP	11	FUNGICIDE
Propiconazole	GROUP	3	FUNGICIDE

Crop Protect Direct AzoxyProp SE

ABN: Crop protect Direct AzoxyProp SE Fungicide

BROAD-SPECTRUM FUNGICIDE FOR CONTROL OF PLANT DISEASES

ACTIVE INGREDIENTS:	% By Weight
Azoxystrobin	13.5%
Propiconazole	11.7%
OTHER INGREDIENTS	74.8%
TOTAL	100.0%

Contains 1.18 lb. a.i. azoxystrobin and 1.02 lb. a.i. propiconazole per gallon. Contains petroleum distillate.
Crop Protect Direct AzoxyProp SE is a suspo-emulsion formulation.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

[See] [inside] [label] [booklet] [for] [First Aid][,] [additional] [Precautionary Statements][,] [and]
[Directions for Use] [including] [Storage and Disposal Instructions][.]

EPA Reg. No.: 101458-9

EPA Est. No.:

Manufactured for:

JABCO, LLC
550 W. Pioneer Blvd
Suite 140
Mesquite, NV 89027

Net Contents:

ACCEPTED

Jun 26, 2025

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No.

101458-9

{LANGUAGE INSIDE BOOKLET}

FIRST AID	
IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	• Immediately call a poison control center or doctor. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give <i>any</i> liquid to the person. • DO NOT give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For 24-Hour Medical Emergency Assistance (Human or Animal), call: 1-800-222-1222. For Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), call CHEMTREC: 1-800-424-9300. For general information on this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu .	
NOTE TO PHYSICIAN: Contains petroleum distillate. Vomiting may cause aspiration pneumonia.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing vapor or spray mist. Avoid contact with eyes, skin or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as Barrier Laminate, Butyl Rubber $\geq$ 14 mils, Nitrile Rubber $\geq$ 14 mils, and Viton $\geq$ 14 mils
- Shoes plus socks
- Protective eyewear (goggles, face shield, or safety glasses)

USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)]. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Azoxystrobin can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination.

Azoxystrobin and propiconazole are toxic to freshwater and estuarine/marine fish; and azoxystrobin is toxic to aquatic invertebrates. Propiconazole is toxic to shrimp. **DO NOT** apply directly to water except as specified on this label. For terrestrial uses, **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Notify state and/or Federal authorities and JABCO, LLC immediately if you observe any adverse environmental effects due to use of this product.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT mix or allow coming in contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE USE DIRECTIONS AND RESTRICTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL AND/OR ILLEGAL RESIDUES.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers,

and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves such as Barrier Laminate, Butyl Rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils, and Viton ≥ 14 mils
- Shoes plus socks
- Protective eyewear

PRODUCT INFORMATION

Crop Protect Direct AzoxyProp SE is a broad-spectrum, preventative fungicide with systemic properties for the control of many important plant diseases. **Crop Protect Direct AzoxyProp SE** may be applied as a foliar spray in alternating spray programs or in tank mixes with other crop protection products. All applications must be made according to the use directions that follow.

USE RESTRICTIONS

- **DO NOT** spray **Crop Protect Direct AzoxyProp SE** where spray drift may reach apple trees.
- **DO NOT** spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your state extension agent for spray drift prevention guidelines in your area.
- **DO NOT** use spray equipment which has been previously used to apply **Crop Protect Direct AzoxyProp SE** to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.
- **DO NOT** use in nurseries, greenhouses, or landscape plantings.
- Human flagging is prohibited.

USE PRECAUTIONS

- **Crop Protect Direct AzoxyProp SE** is extremely phytotoxic to certain apple varieties.
- AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).
- AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

PRODUCT USE INSTRUCTIONS

Application

Thorough coverage is necessary to provide good disease control. Make up no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

Adjuvants

For some uses on this label, a spreading/penetrating type adjuvant including a non-ionic surfactant, crop oil concentrate, or blend may be added at the manufacturer's specified rates. Adjuvants that contain some form of silicone can contribute to phytotoxicity. When an adjuvant is used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification program is advised.

Crop Tolerance/Phytotoxicity

Crop Protect Direct AzoxyProp SE demonstrates some phytotoxic effects when mixed with products that are formulated as ECs. These effects are enhanced if applications are made under

cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone can contribute to phytotoxicity. Under certain environmental conditions, tank mixes of **Crop Protect Direct AzoxyProp SE** plus herbicides and/or fertilizers may cause crop injury in barley, triticale, and wheat.

Efficacy

Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of **Crop Protect Direct AzoxyProp SE** has been used. If resistant isolates to Group 3 or Group 11 fungicides are present, efficacy can be reduced. The higher rates in the rate range and/or shorterspray intervals may be required under conditions of heavy infection pressure, highly susceptible varieties, or when environmental conditions conducive to disease exist.

Integrated Pest Management: Crop Protect Direct AzoxyProp SE must be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development must be followed. The **SPECIFIC USE DIRECTIONS** section in this label identifies specific IPM advisories for each crop. Consult your local agricultural authorities for additional IPM strategies established for your area. **Crop Protect Direct AzoxyProp SE** may be used in State Agricultural Extension advisory (disease forecasting) programs which specify application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

Azoxystrobin	GROUP	11	FUNGICIDE
Propiconazole	GROUP	3	FUNGICIDE

For resistance management, please note that **Crop Protect Direct AzoxyProp SE** contains both a Group 3 (propiconazole) and Group 11 (azoxystrobin) fungicide. Any fungal population may contain individuals naturally resistant to **Crop Protect Direct AzoxyProp SE** and other Group 3 or Group 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **Crop Protect Direct AzoxyProp SE** or other Group 3 or Group 11 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact your JABCO, LLC retailer

or representative. You can also contact your pesticide distributor or university extension specialist to report resistance.

ROTATIONAL CROPS

Rotational Crops	Planting Time From Last Crop Protect Direct AzoxyProp SE Application
Bulb crops Carrots Celery (and other leaf petiole crops - subgroup 4B) Cereals (wheat, barley, triticale) Corn (field, seed, popcorn, and sweet) Grasses grown for seed Mint Oats Peanuts Rice Rye Sorghum Soybeans Strawberries Sugar beets Wild rice	0 Days
Buckwheat Millet	12 Months
Alfalfa (if propiconazole rate does not exceed 0.22 lb. a.i./acre/year)	75 Days
All Other Crops Intended for Food and Feed	105 Days

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 feet above the ground or crop canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select nozzles that deliver Medium to coarse spray droplets in accordance with ASABE Standard S-572.1.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- **DO NOT** apply during temperature inversions.

Groundboom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

Airblast Applications:

- Sprays must be directed into the canopy.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer rows.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size—Groundboom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size—Aircraft

- Adjust Nozzles – Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

MIXING AND APPLICATION METHODS

Crop Protect Direct Azoxystrobin SE may be applied with all types of spray equipment commonly used for making ground and aerial applications. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control.

Spray Equipment

Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Nozzles must be the same size and uniformly spaced across the boom.
- Calibrate sprayer before use.
- It is suggested that screens be used to protect the pump and to prevent nozzles from clogging.
- Screens placed on suction side of pump must be 16-mesh or coarser.
- **DO NOT** place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's specifications.

Pump

- Use a pump with capacity to:
 1. Maintain 35-40 psi at nozzles.
 2. Provide sufficient agitation in tank to keep mixture in suspension. Use a jet agitator or liquid sparge tube for agitation.

For more information on spray equipment and calibration, consult sprayer manufacturer's and state specifications. For specific local directions and spray schedules, consult the current state agricultural specifications.

Mixing Instructions

- **Crop Protect Direct AzoxyProp SE** is a suspo-emulsion (SE) formulation.
- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

Crop Protect Direct AzoxyProp SE Alone (no tank mix)

- Add 1/2 - 2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add **Crop Protect Direct AzoxyProp SE** to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after **Crop Protect Direct AzoxyProp SE** has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

Crop Protect Direct AzoxyProp SE + Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Crop Protect Direct AzoxyProp SE is usually compatible with all tank-mix partners listed on this label. **DO NOT** combine **Crop Protect Direct AzoxyProp SE** in the spray tank with pesticides, surfactants, or fertilizers unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective, and non-injurious to the crop under your conditions.

of use. To determine the physical compatibility of **Crop Protect Direct AzoxyProp SE** with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water dispersible granular products first, then liquid flowables (which includes suspo-emulsions), followed by emulsifiable concentrates and additives/adjuvants last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Mixing in the Spray Tank

- Add 1/2 to 2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above in the “**Crop Protect Direct AzoxyProp SE + Tank Mixtures**” section.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and the **Crop Protect Direct AzoxyProp SE** to the spray tank.
- Allow **Crop Protect Direct AzoxyProp SE** to completely disperse.
- Spray the mixture with the agitator running.
- Observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix product label.
- Do not exceed dosage rate, and the most restrictive label directions and limitations must be followed.
- This product must not be mixed with any product which prohibits such mixing.

Application Instructions

Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur.

DO NOT apply in a manner that will result in exposure to humans or animals.

Ground Application

- For field crops (non-trees), apply in a minimum of 10 gallons of water per acre unless specified otherwise.
- For tree crops, apply in a minimum of 50 gallons of water per acre unless specified otherwise.
- Thorough coverage is necessary to provide good disease control.

Aerial Application

- Use only on crops where aerial applications are indicated.
- For field crops (non-trees), apply in a minimum spray volume of 2 gallons per acre unless specified otherwise.
- For ULV applications (corn), apply in a minimum spray volume of 1 gallon per acre. For ULV applications, thorough coverage is necessary to provide good results. Please refer to the “**Application**” instructions section for details regarding best practices to achieve good coverage. ULV applications are not approved in California.
- For tree crops, apply in a minimum of 10 gallons of water per acre unless specified otherwise.
- Thorough coverage is necessary to provide good disease control.
- **Crop Protect Direct AzoxyProp SE** is extremely phytotoxic to certain apple varieties.
- AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).
- **DO NOT** spray **Crop Protect Direct AzoxyProp SE** where spray drift may reach apple trees.

Application Through Irrigation Systems (Chemigation)

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems.
- **DO NOT** apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.125 - 0.25 inches per acre of water. Excessive water may reduce efficacy.
- If you have questions about calibration, you must contact State Extension Service specialists, equipment manufacturers, or other experts.
- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, must shut the system down and make necessary adjustments if the need arises.

Spray Preparation

Chemical tank and injector system must be thoroughly cleaned. Flush system with clean water.

Operating Instructions

1. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, including a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.
8. **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
9. **DO NOT** apply when wind speed favors drift beyond the area intended.

Center Pivot Irrigation Equipment

Notes: (1) Use only with drive systems which provide uniform water distribution. (2) **DO NOT** use end guns when chemigating **Crop Protect Direct AzoxyProp SE** through center pivot systems because of non-uniform application.

- Determine the size of the area to be treated.

- Determine the time required to apply 0.125 - 0.25 inches per acre of water over the entire area to be treated when the system and injection equipment are operated at normal pressures as specified by the equipment manufacturer. When applying **Crop Protect Direct AzoxyProp SE** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80-95% of the manufacturer's rated capacity.
- Using water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of **Crop Protect Direct AzoxyProp SE** required to treat the area covered by the irrigation system.
- Add the required amount of **Crop Protect Direct AzoxyProp SE** and sufficient water to meet the injection time requirements to the solution tank.
- Make sure the system is fully charged with water before starting injection of the **Crop Protect Direct AzoxyProp SE** solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Continue to operate the system until the **Crop Protect Direct AzoxyProp SE** solution has cleared the sprinkler head.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20- to 30-minute interval. When applying **Crop Protect Direct AzoxyProp SE** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of **Crop Protect Direct AzoxyProp SE** required to treat the area covered by the irrigation system.
- Add the required amount of **Crop Protect Direct AzoxyProp SE** into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the **Crop Protect Direct AzoxyProp SE** solution has cleared the last sprinkler head.

Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), back-flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no

water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

6. Systems must use a metering pump, including a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

Crop Protect Direct Azoxystrobin SE Rate Conversion Table

Fl. Oz. Product/A	Lb. A.I. Azoxystrobin	Lb. A.I. Propiconazole
7	0.056	0.06
10.5	0.10	0.08
14.0	0.13	0.11
15.75	0.15	0.125
17.5	0.16	0.14
21	0.19	0.17
26	0.24	0.21
27	0.25	0.22
28	0.26	0.22

SPECIFIC DIRECTIONS FOR USE

ALMONDS

Almond diseases are more effectively controlled by ground application, using sufficient water volume to provide thorough and uniform coverage. **Crop Protect Direct Azoxystrobin SE** may be applied by ground or by air (minimum of 15 gals./A). Aerial application may be used if necessary but disease control may be reduced. **Crop Protect Direct Azoxystrobin SE** may be applied by air only at growth stages prior to and including 5 weeks after petal fall.

USE RESTRICTIONS:

- **DO NOT** apply more than 112 fl. oz./A (1.04 lb. a.i. azoxystrobin + 0.88 lb. a.i. propiconazole) of **Crop Protect Direct Azoxystrobin SE** per year.
- **DO NOT** apply more than 26 fl. oz./A (0.24 lb. a.i. azoxystrobin + 0.21 lb. a.i. propiconazole) of **Crop Protect Direct Azoxystrobin SE** per application.
- **DO NOT** apply more than 0.9 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 1.5 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 4 applications at the highest rate of 26 fl. oz./A (0.24 lb. a.i. azoxystrobin + 0.21 lb. a.i. propiconazole) or 8 applications at the lowest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct Azoxystrobin SE** per year.
- **DO NOT** apply within 60 days of harvest (60-day PHI). **DO NOT** graze livestock in treated areas or cut treated covercrop for feed.
- **Minimum Retreatment Interval:** 7 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Brown Rot Blossom Blight (<i>Monilinia</i> spp.)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) to 26 (0.24 lb. a.i. azoxystrobin + 0.21 lb. a.i. propiconazole)	Apply Crop Protect Direct Azoxystrobin SE at early bloom stage. If disease pressure is low, a second application of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) may be made as needed through petal fall.

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
		Under conditions of high disease pressure and/or very susceptible varieties, applications may be needed at 50-75% bloom and petal fall. Crop Protect Direct AzoxyProp SE may be used on only 2 blossom blight applications. Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.
Alternaria Leaf Spot (<i>A. alternata</i>) Anthracnose (<i>Colletotrichum acutatum</i>) Leaf Blight (<i>Seimatosporium lichenicola</i>) Leaf Rust (<i>Tranzschelia discolor</i>) Scab (<i>Cladosporium carpophilum</i>) Shothole (<i>Wilsonomyces carpophilus</i>)	17.5 (0.16 lb. a.i. azoxystrobin + 0.14 lb. a.i. propiconazole) to 26 (0.24 lb. a.i. azoxystrobin + 0.21 lb. a.i. propiconazole)	Apply Crop Protect Direct AzoxyProp SE beginning at bud break on a 7- to 14-day interval. Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

BANANAS, PLANTAINS

Crop Protect Direct AzoxyProp SE may be applied by ground (minimum of 15 gals./A) or aerial application (minimum of 5 gals./A).

USE RESTRICTIONS:

- **DO NOT** apply more than 84 fl. oz. (0.76 lb. a.i. azoxystrobin + 0.68 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year (this includes any pre-harvest sprays).
- **DO NOT** apply more than 10.5 fl. oz./A (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.67 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 1.08 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** make more than 8 applications of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply **Crop Protect Direct AzoxyProp SE** within 100 yards of non-bagged bananas.
- **DO NOT** apply **Crop Protect Direct AzoxyProp SE** on bananas unless they are protected by polyethylene bags.
- **DO NOT** apply **Crop Protect Direct AzoxyProp SE** on plantains if the fruit present are not protected with polyethylene bags.
- **DO NOT** feed whole bananas and plantains to animals.
- Minimum Retreatment Interval: 21 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Black Sigatoka (<i>Mycosphaerella fijiensis</i>) Yellow Sigatoka	10.5 (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole)	Apply Crop Protect Direct AzoxyProp SE before disease symptoms appear at the onset of the rainy season.

<i>(Mycosphaerella musicola)</i>		Apply 10.5 fl. oz. /A (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) of Crop Protect Direct AzoxyProp SE/A in 10-20 gallons of water. Apply no more than 2 consecutive applications on a 21- to 25-day schedule before rotating to another labeled product with a different mode of action for at least 2 sprays. If possible, it is advised to have at least 2 consecutive months 'triazole free' during the period of lower disease pressure.
----------------------------------	--	---

BEANS, DRY and SUCCULENT

Cicer arietinum (chickpea, garbanzo bean); *Lupinus* spp. (including sweet lupine, white sweet lupine, white lupine, and grain lupine); *Phaseolus* spp. (including kidney bean, lima bean, mung bean, navy bean, pinto bean, snap bean, and wax bean); *Vicia faba* (broad bean, fava bean); *Vigna* spp. (including asparagus bean, black-eyed pea, and cowpea)

USE RESTRICTIONS:

- **DO NOT** apply more than 42 fl. oz./A (0.39 lb. a.i. azoxystrobin + 0.33 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.34 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 1.5 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 3 applications of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 7 days of harvest (7-day PHI) for succulent beans.
- **DO NOT** apply within 14 days of harvest (14-day PHI) for dry beans.
- Not for use on cowpea cultivars intended for livestock feeding only.
- **Minimum Retreatment Interval:** 7 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Alternaria Blight (<i>Alternaria</i> spp.) Alternaria Leaf Spot (<i>Alternaria alternata</i>) Anthracnose (<i>Colletotrichum lindemuthianum</i>) Ascochyta Blight (<i>Mycosphaerella pinodes</i>) Ascochyta Leaf and Pod Spot (<i>Ascochyta</i> spp.) Ascochyta Leaf Spot (<i>Ascochyta phaseolorum</i>) Bean Rust (<i>Uromyces appendiculatus</i>) Rust (<i>Phakopsora</i> spp.) Southern Blight (<i>Sclerotium rolfsii</i>) Web Blight (<i>Rhizoctonia solani</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground or air (minimum of 15 gals./A). Apply when conditions are conducive for disease. Up to three applications may be made on a 7-14 day interval. NOTE: On certain bean varieties azoxystrobin application may cause crinkled and/or greener leaves. Yields of beans displaying these characteristics have not been reduced.

BERRIES, BUSHBERRY SUBGROUP 13-07B

Aronia berry; Blueberry, Highbush; Blueberry, Lowbush; Buffalo currant; Chilean guava; Cranberry, Highbush; Currant, black; Currant, red; Elderberry; European barberry; Gooseberry; Honeysuckle, Edible; Huckleberry; Jostaberry; Juneberry (Saskatoon berry); Lingonberry; Native currant; Salal; Sea buckthorn and cultivars and/or hybrids of these

USE RESTRICTIONS:

- **DO NOT** apply more than 63 fl. oz./A (0.57 lb. a.i. azoxystrobin + 0.51 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.84 lb. a.i./A of a propiconazole-containing products per year.
- **DO NOT** apply more than 0.75 lb. a.i./A azoxystrobin-containing products per year on bushberries.
- **DO NOT** make more than 3 applications of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 30 days of harvest (30-day PHI).
- **Minimum Retreatment Interval:** 7 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Botryosphaeria Canker (<i>Botryosphaeria</i> spp.) Leaf Spot and Stem Canker (<i>Septoria albopunctata</i>) Leaf Spot (<i>Septoria</i> spp.) Mummy berry (<i>Monilinia vaccinii-corymbosi</i>) Phomopsis Twig Blight, Fruit Rot, and Stem Canker (<i>P. vaccinii</i>) Powdery Mildew (<i>Microsphaera vaccini</i>) Rust (<i>Pucciniastrum vaccinii</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) to 21 (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground or by air (minimum of 15 gals./A). For Mummyberry, make the first application of Crop Protect Direct AzoxyProp SE beginning at green tip and repeat in 7 to 10 days. If conditions are favorable for disease development, additional application may need to be made at pink bud and repeating every 7 to 10 days through petal fall. For other diseases listed, Crop Protect Direct AzoxyProp SE must be applied prior to disease development and continue throughout the season on a 7- to 14- day interval. Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

BERRIES, CANEBERRY SUBGROUP 13-07A

Blackberry, Bingleberry, Boysenberry, Dewberry, Loganberry, Lowberry, Marionberry, Olallieberry, Raspberry, red and black, Wild Raspberry, Youngberry including cultivars and/or hybrids of these.

USE RESTRICTIONS:

- **DO NOT** apply more than 63 fl. oz./A (0.57 lb. a.i. azoxystrobin + 0.51 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 21 fl. oz./A (0.167 lb. a.i. propiconazole + 0.194 lb. a.i. azoxystrobin) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.84 lb. a.i./A of a propiconazole-containing products per year.
- **DO NOT** apply more than 1.5 lb. a.i. /A of an azoxystrobin-containing products per year on caneberries.
- **DO NOT** apply more than 3 applications of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 30 days of harvest (30-day PHI).
- Minimum Retreatment Interval: 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Anthrachnose (<i>Sphaceloma necator</i> , <i>Elsinoe veneta</i>) Botryosphaeria Canker (<i>B. dothidea</i>) Leaf and Cane Spot (<i>Septoria rubi</i>) Leaf Spot (<i>Septoria</i> spp.) Powdery Mildew (<i>Sphaerotheca</i> <i>macularis</i>) Rosette or Double Blossom of Blackberries (<i>Cercospora rubi</i>) Rust (<i>Phragmidium violaceum</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) to 21 (0.167 lb. a.i. propiconazole + 0.194 lb. a.i. azoxystrobin)	Crop Protect Direct AzoxyProp SE may be applied by ground or by air (minimum of 15 gals./A). Crop Protect Direct AzoxyProp SE applications must begin prior to disease development and continue throughout the season on a 14-day interval. Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

BULB VEGETABLES Subgroup 3-07A and 3-07B

Chive, fresh leaves; chive, Chinese, fresh leaves; daylily, bulb; elegans hosta; fritillaria, bulb; fritillaria, leaves; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; kurrat; lady's leek; leek; leek, wild; lily, bulb; onion, Beltsville bunching; onion, bulb; onion, Chinese, bulb; onion, fresh; onion, green; onion, macrostem; onion, pearl; onion, potato, bulb; onion, tree, tops; onion, Welsh, tops; shallot, bulb; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these.

USE RESTRICTIONS:

- **DO NOT** apply more than 56 fl. oz./A (0.52 lb. a.i. azoxystrobin + 0.44 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.45 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 1.5 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 2 applications at the highest rate of 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) or 4 applications at the lowest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 14 days of harvest (14-day PHI) on dry bulb onions.
- **Crop Protect Direct AzoxyProp SE** may be applied the day of harvest (0-day PHI) for green onion types.
- Minimum Retreatment Interval: **7 days**

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Cladosporium Leaf Blotch (<i>C. allii</i>) Purple Blotch (<i>Alternaria porri</i>) Rust (<i>Puccinia allii</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) to 21 (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground (15 gals./A minimum) or aerial application (minimum of 5 gals./A). Begin applications when conditions favor disease development and continue on a 7- to 10-day interval.
Botrytis Leaf Blight (<i>B. squamosa</i>) Downy Mildew (<i>Peronospora destructor</i>) White Rot (<i>Sclerotium cepivorum</i>)	17.5 (0.16 lb. a.i. azoxystrobin + 0.14 lb. a.i. propiconazole) to 21 (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole)	Use the higher rate and shorter interval when disease conditions are severe. NOTE: Mixing with products formulated as an EC may result in phytotoxicity. Make only one application of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

CARROTS

USE RESTRICTIONS:

- **DO NOT** apply more than 56 fl. oz./A (0.52 lb. a.i. azoxystrobin + 0.44 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.45 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 2.0 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 4 applications of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 14 days of harvest (14-day PHI).
- **Minimum Retreatment Interval:** 7 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Alternaria Leaf Blight (<i>Alternaria dauci</i>) Early Blight (<i>Cercospora carotae</i>) Powdery Mildew (<i>Erysiphe polygoni</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground (15 gals./A minimum) or aerial application (minimum of 5 gals./A). Apply Crop Protect Direct AzoxyProp SE when conditions favor disease development. Continue applications on a 7- to 10-day interval, using the shorter interval when disease conditions are severe. Make only one application of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

CELERY

USE RESTRICTIONS:

- **DO NOT** apply more than 56 fl. oz./A (0.52 lb. a.i. azoxystrobin + 0.44 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.45 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 1.5 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 4 applications of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 14 days of harvest (14-day PHI).
- **Minimum Retreatment Interval:** 7 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Early Blight (<i>Cercospora apii</i>) Late Blight (<i>Septoria apiicola</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground, air (5 gals./A minimum), or chemigation. Apply Crop Protect Direct AzoxyProp SE on a 7- to 10-day schedule in alternation with propiconazole-containing products or another product with a different mode of action than Group 11 fungicides. Make only one application of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

CEREALS - WHEAT

See next section for Other Cereals.

Crop Protect Direct AzoxyProp SE is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage.

USE RESTRICTIONS:

- **DO NOT** apply more than 28 fl. oz./A (0.26 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.22 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 0.40 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** apply more than 2 applications at the highest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) or 4 applications at the lowest rate of 7 fl. oz./A (0.06 lb. a.i. azoxystrobin + 0.056 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 7 days of harvest (7-day PHI) for forage and hay.
- **DO NOT** apply after Feekes 10.54.
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Early Season Suppression of: Glume Blotch (<i>Stagonospora nodorum</i>) Leaf Blight (<i>Septoria tritici</i>) Powdery Mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Tan Spot (<i>Pyrenophora tritici-repentis</i>)	7 (0.06 lb. a.i. azoxystrobin + 0.056 lb. a.i. propiconazole) to 14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground, air, or chemigation. Apply Crop Protect Direct AzoxyProp SE in the spring for suppression of early season diseases. Follow up with a second application (see below) for full season control. Under certain environmental conditions, tank mixes of Crop Protect Direct AzoxyProp SE plus herbicides and/or fertilizers may cause crop injury.

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Control of Leaf Diseases: Glume Blotch (<i>Stagonospora nodorum</i>) Helminthosporium Leaf Blight (<i>Drechsleratritici-repentis</i>) Leaf Blight (<i>Septoria tritici</i>) Powdery Mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Rust (<i>Puccinia</i> spp.) Spot Blotch (<i>Bipolaris sorokiniana</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>)	10.5 (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) to 14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground, air, or chemigation. Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when Crop Protect Direct AzoxyProp SE is applied when the flag leaf is 50% to fully emerged. Applications must not be made closer than a 14-day interval. Crop Protect Direct AzoxyProp SE can be applied through full head emergence (Feekes growth stage 10.54). DO NOT apply after this stage to avoid possible illegal residues.
Foot Rot/Eyespot (<i>Tapesia</i> spp.)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Apply full rate of Crop Protect Direct AzoxyProp SE plus half the rate specified on other EPA-registered fungicides containing thiophanate-methyl. Apply at tillering but before elongation has occurred.

CEREALS - BARLEY, OATS, RYE, TRITICALE

Crop Protect Direct AzoxyProp SE is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. An adjuvant may be added at specified rates to improve canopy coverage and penetration while reducing evaporation and drift.

USE RESTRICTIONS:

- **DO NOT** apply more than 28 fl. oz./A (0.26 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.22 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 0.40 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** apply more than 2 applications at the highest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) or 4 applications at the lowest rate of 7 fl. oz./A (0.06 lb. a.i. azoxystrobin + 0.056 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 7 days of harvest (7-day PHI) for forage and hay.
- **DO NOT** apply after Feekes 10.54.
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Early Season Suppression of: Glume Blotch (<i>Stagonospora nodorum</i>) Leaf Blight (<i>Septoria tritici</i>) Powdery Mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Tan Spot (<i>Pyrenophora tritici-repentis</i>)	7 (0.06 lb. a.i. azoxystrobin + 0.056 lb. a.i. propiconazole) to 14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground, air, or chemigation. Apply Crop Protect Direct AzoxyProp SE in the spring for suppression of early season diseases. Follow up with a second application (see below) for full season control. Under certain environmental conditions, tank mixes of Crop Protect Direct AzoxyProp SE plus herbicides and/or fertilizers may cause crop injury.
Control of Leaf Diseases: Barley Scald (<i>Rhynchosporium secalis</i>) Barley Stripe (<i>Pyrenophora graminea</i>) Glume Blotch (<i>Stagonospora nodorum</i>) Helminthosporium Leaf Blight (<i>Drechslera tritici-repentis</i>) Kernel Blight (<i>Alternaria</i> spp.) Leaf Blight (<i>Septoria tritici</i>) Net Blotch (<i>Pyrenophora teres</i>) Powdery Mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Rust (<i>Puccinia</i> spp.) Spot Blotch (<i>Bipolaris sorokiniana</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>)	10.5 (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) to 14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground, air, or chemigation. Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when Crop Protect Direct AzoxyProp SE is applied when the flag leaf is 50% to fully emerged. Applications must not be made closer together than a 14-day interval.
Foot Rot/Eyespot (<i>Tapesia</i> spp.)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Apply full rate of Crop Protect Direct AzoxyProp SE plus half the rate specified on other EPA-registered fungicides containing thiophanate- methyl. Apply at tillering but before elongation has occurred.

CORN, FIELD, and POP
(Includes Seed Production)

For best results, sufficient coverage is very important. For ULV aerial applications, **DO NOT** use less than 1.0 GPA. Use a higher water volume for aerial application if equipment and/or conditions will not provide good coverage. **Crop Protect Direct AzoxyProp SE** may be applied by ground, air (ULV), or chemigation.

USE RESTRICTIONS:

- **DO NOT** apply more than 38.5 fl. oz./A (0.36 lb. a.i. azoxystrobin + 0.30 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 28 fl. oz./A ((0.26 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) for field corn harvested for forage.
- **DO NOT** apply more than 0.45 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 2.0 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** apply more than 2 applications at the highest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) or 3 applications at the lowest rate of 10.5 fl. oz./A (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 30 days of harvest (30-day PHI) for forage, grain, or stover.
- ULV applications are not approved in California.
- **Minimum Retreatment Interval:** 7 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Anthracnose Leaf Blight (<i>Colletotrichum graminicola</i>) Eye Spot (<i>Aureobasidium zeae</i>) Gray Leaf Spot (<i>Cercospora zeae-maydis</i>) Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>) Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>) Physoderma Brown Spot (<i>Physoderma maydis</i>) Rusts (<i>Puccinia</i> spp.) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>) also known as Helminthosporium Leaf Blights (<i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i>) Suppression of: Diplodia Ear Rot (<i>D. maydis</i>)	10.5 (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole)	Early application (V4-V8): An early application (V4-V8) of Crop Protect Direct AzoxyProp SE may be applied for early season disease control. If mixing with herbicides other than solo glyphosate products, consult your local JABCO, LLC representative.

CORN, FIELD, and POP (cont.)
(Includes Seed Production)

For best results, sufficient coverage is very important. For ULV aerial applications, **DO NOT** use less than 1.0 GPA. Use a higher water volume for aerial application if equipment and/or conditions will not provide good coverage. **Crop Protect Direct AzoxyProp SE** may be applied by ground, air (ULV), or chemigation.

USE RESTRICTIONS:

- **DO NOT** apply more than 38.5 fl. oz./A (0.36 lb. a.i. azoxystrobin + 0.30 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 28 fl. oz./A ((0.26 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) for field corn harvested for forage.
- **DO NOT** apply more than 0.45 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 2.0 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** apply more than 2 applications at the highest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) or 3 applications at the lowest rate of 10.5 fl. oz./A (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 30 days of harvest (30-day PHI) for forage, grain, or stover.
- ULV applications are not approved in California.
- **Minimum Retreatment Interval:** 7 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Anthracnose Leaf Blight <i>(Colletotrichum graminicola)</i> Eye Spot <i>(Aureobasidium zeae)</i> Gray Leaf Spot <i>(Cercospora zeae-maydis)</i> Northern Corn Leaf Blight <i>(Setosphaeria turcica)</i> Northern Corn Leaf Spot <i>(Cochliobolus carbonum)</i> Physoderma Brown Spot <i>(Physoderma maydis)</i> Rusts <i>(Puccinia spp.)</i> Southern Corn Leaf Blight <i>(Cochliobolus heterostrophus)</i> also known as Helminthosporium Leaf Blights <i>(H. maydis, H. turcicum, H. carbonum)</i> Suppression of: Diplodia Ear Rot <i>(D. maydis)</i>	10.5 (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) to 14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Later season applications: For gray leaf spot, rusts, anthracnose, and eye spot, apply Crop Protect Direct AzoxyProp SE when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule. For leaf blights, apply Crop Protect Direct AzoxyProp SE when disease first appears. Continue on a 7- to 14-day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the high rate. DO NOT use adjuvants or other additives after the V8 growth stage and prior to the VT growth stage, as use during these development times may impose stress on the plant that could inhibit proper kernel development. VT is defined as when the last branch of the tassel is completely visible, but silks have not yet emerged from the ear shoot. Apply no more than 2 applications of Crop Protect Direct AzoxyProp SE or

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
		any other Group 11 fungicide per year. Use of an adjuvant including COC may provide additional disease control.

CORN, SWEET

Sweet Corn (Includes Seed Production)

For best results, sufficient coverage is very important. Use of a crop oil concentrate is advised for aerial applications to reduce evaporation and enhance canopy penetration and coverage. Consult your aerial applicator for advised concentration of crop oil concentrate. **DO NOT** use less than 1.0 GPA for the ULV applications. Use higher water volumes for aerial applications if equipment and/or conditions will not provide good coverage. **Crop Protect Direct AzoxyProp SE** may be applied by ground, air (ULV), or chemigation.

USE RESTRICTIONS:

- **DO NOT** apply more than 56 fl. oz./A (0.52 lb. a.i. azoxystrobin + 0.44 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.45 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 2.0 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** apply more than 4 applications at the highest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) or 5 applications at the lowest rate of 10.5 fl. oz./A (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply to sweet corn within 14 days of harvest (14-day PHI) for ears or forage.
- ULV applications are not approved in California.
- **Minimum Retreatment Interval:** 7 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Anthracnose Leaf Blight (<i>Colletotrichumgraminicola</i>) Eye Spot (<i>Aureobasidium zeae</i>) Gray Leaf Spot (<i>Cercospora zeae-maydis</i>)	10.5 (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) to 14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Apply Crop Protect Direct AzoxyProp SE when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule.
Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>) Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>) Rusts (<i>Puccinia spp.</i>) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>)		For leaf blights apply Crop Protect Direct AzoxyProp SE when disease first appears. Continue on a 7- to 14-day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the high rate. Make no more than one application before alternating with propiconazole-containing products or to a non-Group 11 fungicide.

CRANBERRIES

USE RESTRICTIONS:

Use is limited to Oregon, Washington, and Wisconsin only.

- **DO NOT** apply more than 63 fl. oz./A (0.57 lb. a.i. azoxystrobin + 0.51 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.67 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 1.5 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 3 applications of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 45 days of harvest (45-day PHI).
- **DO NOT** allow release of irrigation or flood water to non-target aquatic habitat for at least 14 days after the last application.
- **DO NOT** use cranberry fields used for aquaculture of fish and crustaceans.
- **DO NOT** apply when weather conditions favor drift from treated areas to non-target aquatic habitat. Applicators must use care in making applications near non-target aquatic habitats.
- **DO NOT** apply to flooded crop.
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Cottonball (<i>Monilinia oxycocci</i>) Fruit Rots (<i>Physalospora vaccinii</i>) (<i>Glomerella cingulata</i>) (<i>Coleophoma empetri</i>) Lophodermium Twig Blight (<i>Lophodermium</i> spp.)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) to 21 (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground (minimum of 10 gals./A) or aerial application (minimum of 20 gals./A). Make the first application at leaf bud break and repeat in 14 days. Additional applications must be made at early bloom. Under severe pressure, use the higher rate for control. Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

FILBERTS

USE RESTRICTIONS:

- **DO NOT** apply more than 112 fl. oz./A (1.04 lb. a.i. azoxystrobin + 0.88 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.9 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 1.2 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 5 applications at the highest rate of 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) or 8 applications at the lowest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 60 days of harvest (60-day PHI).
- **DO NOT** graze livestock in treated areas or cut treated cover crop for feed.
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Eastern Filbert Blight (<i>Anisogramma anomala</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) to 21 (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground or aerial application (minimum of 15 gals./A). Begin applications when green leaf tissue becomes visible and continue on a 2- to 3-week interval. Under severe disease conditions, use the higher rate and shorter interval. NOTE: On certain varieties, Crop Protect Direct AzoxyProp SE applications may cause smaller and/or greener leaves. Yields of filberts displaying these characteristics have not been reduced due to Crop Protect Direct AzoxyProp SE treatments. Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

GRASSES (Grown For Seed)

Crop Protect Direct AzoxyProp SE is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important.

USE RESTRICTIONS:

Use is limited to Idaho, Minnesota, Nebraska, Oregon, and Washington only.

- **DO NOT** apply more than 86 fl. oz./A (0.80 lb. a.i. azoxystrobin + 0.68 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 26 fl. oz./A (0.24 lb. a.i. azoxystrobin + 0.21 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.9 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 0.8 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** make more than 3 applications at the highest rate of 26 fl. oz./A (0.24 lb. a.i. azoxystrobin + 0.21 lb. a.i. propiconazole) or 6 applications at the lowest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 20 days of harvest (20-day PHI) of seed.
- **DO NOT** feed hay cut within 20 days of the last application.
- **DO NOT** graze treated areas within 140 days of the last application.
- **DO NOT** apply to bermudagrass grown for seed.
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Ergot Stem Diseases Powdery Mildew (<i>Erysiphe graminis</i>) Rusts (<i>Puccinia</i> spp.) Selenophoma Stem Eyespot (<i>Selenophoma</i> spp.)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) to 26 (0.24 lb. a.i. azoxystrobin + 0.21 lb. a.i. propiconazole)	Apply Crop Protect Direct AzoxyProp SE in a minimum of 20 gals. of water per acre for ground or in a minimum of 10 gals. of water per acre for aerial. Crop Protect Direct AzoxyProp SE may be applied by ground, air or chemigation. Apply Crop Protect Direct AzoxyProp SE when powdery mildew infections, <i>Selenophoma</i> infections, and/or rust pustules are noticeable and increasing in number in late spring or early summer. To maximize control of severe rust pressure, apply 26 fl. oz./A (except bluegrass apply 14 fl. oz./A) and make applications at 14-day intervals until the seed is mature. For bluegrass, it is important to begin application early in the growing season. Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

MINT

Peppermint tops, Spearmint tops

USE RESTRICTIONS:

- **DO NOT** apply more than 42 fl. oz./A (0.39 lb. a.i. azoxystrobin + 0.33 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.22 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 0.75 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 3 applications at the highest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) or 4 applications at the lowest rate of 10.5 fl. oz./A (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 7 days of harvest (7-day PHI).
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Powdery Mildew (<i>Erysiphe</i> spp.) Rust (<i>Puccinia menthae</i>)	10.5 (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) to 14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground (minimum of 20 gals./A) or chemigation. Begin applications when the plants are 2 - 4 inches high or when conditions become favorable for disease development. Make a second application 14 days after the first application.

PEANUTS

[Not for use in California]

When applying **Crop Protect Direct AzoxyProp SE** via irrigation or as a directed ground application, additional methods must be employed for leaf spot control. **Crop Protect Direct AzoxyProp SE** may be applied by ground, air, or chemigation.

USE RESTRICTIONS:

- **DO NOT** apply more than 56 fl. oz./A (0.52 lb. a.i. azoxystrobin + 0.44 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 28 fl. oz./A (0.26 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.45 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 0.80 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** make more than 2 applications at the highest rate of 28 fl. oz./A (0.26 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) or 4 applications at the lowest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 14 days of harvest (14-day PHI) when using a maximum rate of 14 fl. oz./A.
- **DO NOT** apply within 21 days of harvest (21-day PHI) when using rates above 14 fl. oz./A and
- **DO NOT** feed hay from treated fields to livestock if using rates higher than 14 fl. oz./A.
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Early Leaf Spot <i>(Cercospora arachidicola)</i> Late Leaf Spot <i>(Cercosporidium personatum)</i> Rust <i>(Puccinia arachidis)</i> Web Blotch <i>(Phoma arachidicola)</i>	10.5 (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) to 14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Apply Crop Protect Direct AzoxyProp SE beginning 35 to 40 days after planting or at the first appearance of disease. Continue applications on a 14-day schedule. Under heavy disease pressure use higher specified application rates. Crop Protect Direct AzoxyProp SE also may be used in State Agricultural Extension advisory (disease forecasting) programs which specify application timing based on environmental factors favorable for disease development. Make no more than two sequential applications of a Group 11 fungicide before alternating to another product with a different mode of action than Group 11 fungicides.
Soil-Borne Diseases - mid-late season Rhizoctonia Peg and Pod Rot <i>(R. solani)</i> Stem Rot/White Mold/ Southern Blight <i>(Sclerotium rolfsii)</i> Suppression Only: Cylindrocladium Black Rot <i>(C. crotalariae)</i> Pythium Pod Rot <i>(P. myriotylum)</i>	21 (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) to 28 (0.26 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole)	Apply Crop Protect Direct AzoxyProp SE at approximately 60 and 90 days after planting as a foliar application. This application regime may be applied earlier in the season if environmental conditions favor disease development. This application will provide protection against soil-borne diseases and will also provide control of the foliar diseases listed for a 10- to 14-day period after each spray. Under heavy pressure and/or heavy rainfall or irrigation, use 28 fl. oz. of Crop Protect Direct AzoxyProp SE per acre. Under lighter pressure and dry conditions (non-irrigated, low rainfall), use 21 - 28 fl. oz. of Crop Protect Direct AzoxyProp SE per acre.

PEANUTS (cont.)
[Not for use in California]

When applying **Crop Protect Direct AzoxyProp SE** via irrigation or as a directed ground application, additional methods must be employed for leaf spot control. **Crop Protect Direct AzoxyProp SE** may be applied by ground, air, or chemigation.

USE RESTRICTIONS:

- **DO NOT** apply more than 56 fl. oz./A (0.52 lb. a.i. azoxystrobin + 0.44 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 28 fl. oz./A (0.26 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.45 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 0.80 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** make more than 2 applications at the highest rate of 28 fl. oz./A (0.26 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) or 4 applications at the lowest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 14 days of harvest (14-day PHI) when using a maximum rate of 14 fl. oz./A.
- **DO NOT** apply within 21 days of harvest (21-day PHI) when using rates above 14 fl. oz./A and
- **DO NOT** feed hay from treated fields to livestock if using rates higher than 14 fl. oz./A.
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Soil-Borne Diseases - mid-late season Rhizoctonia Peg and Pod Rot (<i>R. solani</i>) Stem Rot/White Mold/Southern Blight (<i>Sclerotium rolfsii</i>) Suppression Only: Cylindrocladium Black Rot (<i>C. crotalariae</i>) Pythium Pod Rot (<i>P. myriotylum</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) to 28 (0.26 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) plus azoxystrobin in tank mix	Tank-mix option: Apply 14 fl. oz./A of Crop Protect Direct AzoxyProp SE in a tank mix with azoxystrobin-containing products or other fungicides for control of soil-borne diseases. A minimum of 0.15 lb. a.i./A azoxystrobin must be in the tank mix (see Crop Protect Direct AzoxyProp SE rate conversion table below). DO NOT exceed 0.4 lb. of azoxystrobin/A/application. Apply Crop Protect Direct AzoxyProp SE plus azoxystrobin at approximately 60 and 90 days after planting as a foliar application. This application regime may be applied earlier in the season if environmental conditions favor disease development. This application will provide protection against soil-borne diseases and will also provide control of the foliar diseases listed for a 10- to 14-day period after each spray. Under heavy pressure and/or heavy rainfall or irrigation, there must be 0.30 - 0.4 lb. a.i. of azoxystrobin in the tank. Under lighter pressure and dry conditions (non-irrigated, low rainfall), 0.2 - 0.4 lb. a.i. of azoxystrobin can be used.

PECANS

USE RESTRICTIONS:

- **DO NOT** apply more than 105 fl. oz./A (0.95 lb. a.i. azoxystrobin + 0.85 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.9 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 1.2 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** make more than 5 applications at the highest rate of 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) or 7 applications at the lowest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply after shuck split or within 45 days of harvest (45-day PHI), whichever is first.
- **DO NOT** graze livestock in treated areas or cut treated cover crops for feed.
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Anthracnose (<i>Glomerella cingulata</i>) Downy Spot (<i>Mycosphaerella caryigena</i>) Liver Spot (<i>Gnomonia caryae</i> pv <i>pecanae</i>) Pecan Scab (<i>Cladosporium caryigenum</i>) Powdery Mildew (<i>Microsphaera penicillata</i>) Vein Spot (<i>Gnomonia nerviseda</i>) Zonate Leaf Spot (<i>Cristulariella moricola</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) to 21 (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole)	Ground applications must be applied in sufficient water to provide for full coverage. Crop Protect Direct AzoxyProp SE maybe applied by ground or air (minimum of 20 gals./A). Pecan scab: Apply 14 - 21 fl. oz./A Crop Protect Direct AzoxyProp SE on a 14-day schedule during bud break and pre-pollination sprays. Apply 20 - 21 fl. oz./A during nutformation and cover sprays. Use higher rates whendisease pressure is heavier. DO NOT apply after shuck split. Other foliar diseases: Crop Protect Direct AzoxyProp SE may be applied for control of mid to late season foliar diseases at 14 - 20.5 fl. oz./A with other pecan products labeled for these diseases. Observe all directions, precautions, and limitations for the other products. Use of an adjuvant including COC may provide additional disease control. Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

PISTACHIOS

USE RESTRICTIONS:

- **DO NOT** apply more than 105 fl. oz./A (0.95 lb. a.i. azoxystrobin + 0.85 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.9 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 1.5 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 5 applications at the highest rate of 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) or 6 applications at the lowest rate of 17.5 fl. oz./A (0.16 lb. a.i. azoxystrobin + 0.14 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 60 days of harvest (60-day PHI).
- **DO NOT** graze livestock in treated areas or cut treated cover crop for feed.
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Alternaria Late Blight (<i>A. alternata</i>) Botryosphaeria Panicle and Shoot Blight (<i>B. dothidea</i>) Septoria Leaf Spot (<i>S. pistaciarum</i>)	17.5 (0.16 lb. a.i. azoxystrobin + 0.14 lb. a.i. propiconazole) to 21 (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground or aerial application (minimum of 15 gals./A). Begin applications when green leaf tissue becomes visible and continue on a 14- to 21-day interval. Under severe disease conditions, use the higher rate and the shorter interval. Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

RICE

Including Wild Rice

USE RESTRICTIONS:

- **DO NOT** apply more than 42 fl. oz./A (0.39 lb. a.i. azoxystrobin + 0.33 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 27 fl. oz./A (0.25 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.34 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 0.70 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** make more than 1 application at the highest rate of 27 fl. oz./A (0.25 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) or 2 applications at the lowest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 35 days of harvest (35-day PHI).
- **DO NOT** release floodwater within 14 days of an application.
- **DO NOT** apply to stubble or ratoon crop rice.
- **DO NOT** use in rice fields where commercial farming of crayfish will be practiced.
- **DO NOT** drain water from treated rice fields into ponds used for commercial fish farming.
- **DO NOT** use water drained from treated fields to irrigate other crops.
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Aggregate Sheath Spot <i>(Rhizoctoniaoryzae-sativa)</i> Black Sheath Rot <i>(Gaeumannomyces graminis)</i> Brown Leaf Spot <i>(Helminthosporium oryzae)</i> Kernel Smut <i>(Tilletia barclayana)</i> Leaf Blast <i>(Pyricularia grisea)</i> Leaf Smut <i>(Entyloma oryzae)</i> Narrow Brown Leaf Spot <i>(Cercospora oryzae)</i> Panicle Blast <i>(P. grisea)</i> Sheath Blight <i>(Rhizoctonia solani)</i> Sheath Spot <i>(Rhizoctonia oryzae)</i> Stem Rot <i>(Sclerotium oryzae)</i> For Disease Suppression of: False Smut <i>(Ustilaginoidea virens)</i>	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) to 27 (0.25 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole)	Timing of Crop Protect Direct AzoxyProp SE application will depend on disease severity, disease complex and rice variety/growth stage. Consult local extension experts for local economic thresholds established for various rice varieties and diseases. For aerial application, volumes must be 5-10 GPA. An adjuvant may be added at specified rates to improve canopy coverage and penetration while reducing evaporation and drift. Leaf blast: Crop Protect Direct AzoxyProp SE must be applied for preventive control. Apply 21 - 27 fl. oz./A. Panicle blast: Apply Crop Protect Direct AzoxyProp SE at 10% head emergence with an additional application of an azoxystrobin-containing product at 90% emergence. Refer to the azoxystrobin-containing product label for rates and timing. All other leaf/stem diseases: Apply 15.75 - 27 fl. oz./A at initial sign of disease. Apply higher rates when disease pressure is heavy and/or when environmental conditions are highly favorable for disease development. A second application may be made 14 days later.

RICE (cont.)
Including Wild Rice

USE RESTRICTIONS:

- **DO NOT** apply more than 42 fl. oz./A (0.39 lb. a.i. azoxystrobin + 0.33 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 27 fl. oz./A (0.25 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.34 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 0.70 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** make more than 1 application at the highest rate of 27 fl. oz./A (0.25 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) or 2 applications at the lowest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 35 days of harvest (35-day PHI).
- **DO NOT** release floodwater within 14 days of an application.
- **DO NOT** apply to stubble or ratoon crop rice.
- **DO NOT** use in rice fields where commercial farming of crayfish will be practiced.
- **DO NOT** drain water from treated rice fields into ponds used for commercial fish farming.
- **DO NOT** use water drained from treated fields to irrigate other crops.
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Aggregate Sheath Spot (<i>Rhizoctoniaoryzae-sativa</i>) Black Sheath Rot (<i>Gaeumannomyces graminis</i>) Brown Leaf Spot (<i>Helminthosporium oryzae</i>) Kernel Smut (<i>Tilletia barclayana</i>) Leaf Blast (<i>Pyricularia grisea</i>) Leaf Smut (<i>Entyloma oryzae</i>) Narrow Brown Leaf Spot (<i>Cercospora oryzae</i>) Panicle Blast (<i>P. grisea</i>) Sheath Blight (<i>Rhizoctonia solani</i>) Sheath Spot (<i>Rhizoctonia oryzae</i>) Stem Rot (<i>Sclerotium oryzae</i>) For Disease Suppression of: False Smut (<i>Ustilaginoidea virens</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) to 27 (0.25 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole)	Tank mix option: Apply 15.75 - 20.5 fl. oz./A of Crop Protect Direct AzoxyProp SE in a tank mix with azoxystrobin-containing products or other fungicides for control of rice diseases. A minimum of 0.15 lb. a.i./A azoxystrobin must be in the tank mix (see Crop Protect Direct AzoxyProp SE rate conversion table below). DO NOT exceed 0.3 lb. of azoxystrobin/A/per application to rice or 0.25 lb. of azoxystrobin/A/application to wild rice. The lower rate of 14 fl. oz./A may only be used for hybrids or varieties with at least moderate resistance to sheath blight. Apply from late boot to boot split for control of diseases (except leaf blast and false smut) of rice (including wild rice). When applying prior to late boot or after boot split growth stages, use the higher rates listed above. Make no more than 2 applications of a Group 11 (QoI) fungicide per year.

SORGHUM

USE RESTRICTIONS:

- **DO NOT** apply more than 56 fl. oz./A (0.52 lb. a.i. azoxystrobin + 0.44 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 28 oz. (0.26 lb. a.i. azoxystrobin + 0.22 lb. a.i. propiconazole) on sorghum harvested for forage.
- **DO NOT** apply more than 0.45 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 0.75 lb./A of azoxystrobin-containing products to sorghum grown for grain and/or stover per year.
- **DO NOT** apply more than 0.5 lb./A of azoxystrobin-containing products to sorghum grown for forage per year.
- **DO NOT** make more than 4 applications at the highest rate of 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) or 5 applications at the lowest rate of 10.5 fl. oz./A (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 30 days of harvest (30-day PHI) for forage.
- **DO NOT** apply within 21 days of harvest (21-day PHI) for grain or stover.
- **DO NOT** graze livestock or cut for green chop or silage within 30 days of application.
- **Minimum Retreatment Interval:** 5 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Anthracnose (<i>Colletotrichum graminicola</i>) Ergot (<i>Claviceps sorghi</i>) Gray Leaf Spot (<i>Cercospora sorghi</i>) Ladder Leaf Spot (<i>Cercospora fusimaculans</i>) Leaf Blight (<i>Exserohilum turcicum</i>) Zonate Leaf Spot (<i>Gloeocercospora sorghi</i>)	10.5 (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) to 14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground or aerial application. For ergot control, make the first application at or just prior to flowering. Repeat on a 5- to 7-day interval. For other diseases, apply at first sign of disease. Apply on a 14-day interval.

SOYBEANS

Crop Protect Direct AzoxyProp SE is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. **DO NOT** use less than 2.0 GPA. Use a higher water volume for aerial application if equipment and/or conditions will not provide for good coverage.

USE RESTRICTIONS:

- **DO NOT** apply more than 42 fl. oz./A (0.39 lb. a.i. azoxystrobin + 0.33 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.34 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 1.5 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 2 applications at the highest rate of 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) or 4 applications at the lowest rate of 10.5 fl. oz./A (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- Apply up to Stage R6.
- **Minimum Retreatment Interval:** 14 days
- **PHI:** 0 day

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Aerial Web Blight (<i>Rhizoctonia solani</i>) Alternaria Leaf Spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum truncatum</i>) Brown Spot (<i>Septoria glycines</i>) Cercospora Blight and Leaf Spot (<i>C. kikuchii</i>) Frogeye Leaf Spot (<i>Cercospora sojina</i>) Pod and Stem Blight (<i>Diaporthe</i> spp.) Soybean Rust (<i>Phakopsora pachyrhizi</i>)	10.5 (0.10 lb. a.i. azoxystrobin + 0.08 lb. a.i. propiconazole) to 21 (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground, air, or chemigation. Foliar diseases (except rust): Apply 14 - 21 fl.oz./A at growth stage R3 (early pod set) when pods are 1/8 - 1/4 inch long) and 14-21 days later at growth stage R5 (pod fill). Crop Protect Direct AzoxyProp SE maybe applied earlier if conditions are conducive for disease. Soybean rust: Apply 14 - 21 fl. oz./A at first indication that disease is in the area. For best control, preventive applications work best. Repeat on a 14- to 21-day interval. Use higher rate and shorter interval when diseases are present in the field and incidence is less than 2% (2 plants in 100 are infected). If incidence is greater than this or if disease is in mid-canopy, control will not be acceptable. Scouting for the disease and/or being aware of the proximity of the disease via monitoring systems will aid in the proper timing to maximize the effectiveness of the fungicide applications. On certain varieties, Crop Protect Direct AzoxyProp SE applications may cause crinkled, smaller and/or greener leaves. Yields of beans displaying these characteristics have not been reduced due to Crop Protect Direct AzoxyProp SE treatments.

STONE FRUITS Crop Group 12-12

Apricot, Cherry (sweet), Cherry (tart), Nectarine, Peach, Plum, Plumcot, Prune, Including all cultivars and hybrids of these

Stone fruit diseases are more effectively controlled by ground application, using sufficient water volume to provide thorough and uniform coverage. Aerial application (minimum of 15 gals./A) may be used if necessary but disease control may be reduced.

Applications of **Crop Protect Direct AzoxyProp SE** during bloom to Stanley plums have occasionally caused fruit to be less oval in shape and smaller in size at harvest. To avoid this, **DO NOT** apply **Crop Protect Direct AzoxyProp SE** to Stanley plums earlier than 21 days prior to harvest.

USE RESTRICTIONS:

- **DO NOT** apply more than 70 fl. oz./A (0.65 lb. a.i. azoxystrobin + 0.55 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.56 lb. a.i./A propiconazole-containing products per year.
- **DO NOT** apply more than 1.5 lb. a.i./A azoxystrobin-containing products per year.
- **DO NOT** make more than 5 applications of **Crop Protect Direct AzoxyProp SE** per year.
- **Crop Protect Direct AzoxyProp SE** may be applied the day of harvest (0-day PHI).
- **Minimum Retreatment Interval:** 7 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Alternaria Spot and Fruit Rot (<i>A. alternata</i>) Anthracnose (<i>Colletotrichum prunicola</i>) Brown Rot Blossom Blight (<i>Monilinia</i> spp.) Brown Rot on Fruit (<i>Monilinia</i> spp.) Cherry Leaf Spot (<i>Blumeriella jaapii</i>) Powdery Mildew (<i>Podosphaera clandestina</i> , <i>Sphaerotheca pannosa</i>) Rust (<i>Tranzschelia discolor</i>) Scab (<i>Cladosporium carpophilum</i>) Shothole (<i>Wilsonomyces carpophilus</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	For brown rot blossom blight, apply Crop Protect Direct AzoxyProp SE at early bloom stage. If disease pressure is low, a second application of 14 fl. oz./A may be made as needed through petal fall. Under conditions of high disease pressure and/or very susceptible varieties, applications may be needed at 50 - 75% bloom and petal fall. For brown rot on fruit, apply as needed, a maximum of 2 sprays of Crop Protect Direct AzoxyProp SE, during the preharvest period up to the day of harvest. Make the two applications no closer than 10 days apart. For powdery mildew, rust, and cherry leaf spot, follow the blossom blight schedule. Make up to 2 additional applications on a 10- to 14-day interval from the end of petal fall to harvest. For scab, begin applications at petal fall and continue on a 7- to 14-day interval. For other diseases, begin applications at onset of disease and continue on a 10- to 14-day interval. Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

STRAWBERRIES AND LOW GROWING BERRY SUBGROUP 13-07G (EXCEPT CRANBERRY)

Bearberry, Bilberry, Cloudberry, Muntries, Partridgeberry, Including all cultivars and/or hybrids of these

USE RESTRICTIONS:

- **DO NOT** apply more than 56 fl. oz. /A (0.52 lb. a.i. azoxystrobin + 0.44 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.45 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 1.0 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 4 applications of **Crop Protect Direct AzoxyProp SE** per year.
- **Crop Protect Direct AzoxyProp SE** may be applied the day of harvest (0-day PHI).
- **Minimum Retreatment Interval:** 10 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Anthracnose (<i>Colletotrichum</i> spp.) Leaf Rust (<i>Phragmidium potentillae</i>) Leaf Spot (<i>Cercospora fragariae</i>) Powdery Mildew (<i>Sphaerotheca maculans</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground (20 gals./A minimum) or aerial application (15 gals./A minimum). Begin applications prior to disease development. Repeat on a 10- to 14-day interval. Make no more than 4 applications per year of Crop Protect Direct AzoxyProp SE or other QoI containing product. Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

SUGAR BEETS

[Not for use in California]

For best results, sufficient water volume must be used to provide thorough coverage. A minimum of 15 gals./A for ground applications is specified. For aerial applications, a minimum of 5 gals./A of water is specified. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

USE RESTRICTIONS:

- **DO NOT** apply more than 42 fl. oz./A (0.39 lb. a.i. azoxystrobin + 0.33 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 14 fl. oz./A (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.34 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 2.0 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 3 applications of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 21 days of harvest (21-day PHI).
- **Minimum Retreatment Interval:** 10 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Cercospora Leaf Spot (<i>C. beticola</i>) Powdery Mildew (<i>Erysiphe polygoni</i>) Rhizoctonia Crown Rot (<i>R. solani</i>)	14 (0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be applied by ground, chemigation, or aerial application. Begin applications preventively or on a forecast system. For powdery mildew, apply at first sign of disease. Apply Crop Protect Direct AzoxyProp SE on a 10- to 21-day schedule. Make only one Crop Protect Direct AzoxyProp SE spray then alternate to a non-triazole fungicide (non-Group 3) that is registered on sugar beets for these diseases. If disease pressure is high, use the highest rate and shortest interval. For Rhizoctonia crown rot, apply 14 oz. in a 7-inch band over the row at the 4- to 8-leaf stage.

SUGARCANE

USE RESTRICTIONS:

- **DO NOT** apply more than 84 fl. oz./A (0.76 lb. a.i. azoxystrobin + 0.68 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 21 fl. oz./A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.67 lb. a.i. of propiconazole-containing products/A/year.
- **DO NOT** apply more than 0.80 lb. a.i. of azoxystrobin-containing products/A/year.
- **DO NOT** make more than 4 applications at the highest rate of 21 fl. oz/A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) or 5 applications at the lowest rate of 16 fl. oz./A (0.15 lb. a.i. azoxystrobin + 0.13 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply within 30 days of harvest (30-day PHI).
- **Minimum Retreatment Interval:** 14 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Brown Rust (<i>Puccinia melanocephala</i>) Orange Rust (<i>Puccinia kuehnii</i>)	16 (0.15 lb. a.i. azoxystrobin + 0.13 lb. a.i. propiconazole) to 21 (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole)	Crop Protect Direct AzoxyProp SE may be made by ground, air, or chemigation. Begin applications prior to rust development and continue throughout the season every 14-28 days following resistance management guidelines. Scout fields and begin applications at the earliest sign of rust.

TREE NUTS Crop Group 14-12

Almond (see specific directions), Beechnut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (see specific directions), Hickory, Macadamia, Pecan (see specific directions), Pistachios (see specific directions), Walnut

USE RESTRICTIONS:

- **DO NOT** apply more than 84 fl. oz./A (0.76 lb. a.i. azoxystrobin + 0.68 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per year.
- **DO NOT** apply more than 21 fl. oz/A (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole) of **Crop Protect Direct AzoxyProp SE** per application.
- **DO NOT** apply more than 0.9 lb. a.i./A of propiconazole-containing products per year.
- **DO NOT** apply more than 1.2 lb. a.i./A of azoxystrobin-containing products per year.
- **DO NOT** make more than 4 applications of **Crop Protect Direct AzoxyProp SE** or other QoI containing products per year.
- **DO NOT** apply within 60 days of harvest (60-day PHI) except for pecan (see specific use directions).
- **DO NOT** graze livestock in treated areas or cut treated cover crop for feed.
- **Minimum Retreatment Interval:** 7 days

Target Diseases	Use Rate Fl. Oz. Product/A	Application Instructions
Foliar Diseases	14 0.13 lb. a.i. azoxystrobin + 0.11 lb. a.i. propiconazole) to 21 (0.19 lb. a.i. azoxystrobin + 0.17 lb. a.i. propiconazole)	For best control of tree nut diseases, ground applications are advised. Crop Protect Direct AzoxyProp SE may be applied by ground or aerial application (15 gals./A minimum). Apply Crop Protect Direct AzoxyProp SE at first sign of disease. Repeat on a 7- to 14-day interval. Make no more than 2 sequential applications of a Group 11 fungicide prior to alternating with another product with a different mode of action than Group 11 fungicides.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Store in original container only. Store in a cool, dry and well-ventilated place. Protect from excessive heat. Keep container closed when not in use. **DO NOT** store near food or feed.

Pesticide Disposal

Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling:

[NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[REFILLABLE CONTAINER: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or reconditioning if appropriate.]

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of JABCO, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold JABCO, LLC and Seller harmless for any claims relating to such factors.

JABCO, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or JABCO, LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, JABCO, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither JABCO, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF JABCO, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF JABCO, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

JABCO, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of JABCO, LLC.

[All trademarks are the property of their respective owners.]

[20240912]