Case No. 495186



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505T) 1200 Pennsylvania Ave., N.W.

Washington, D.C. 20460

Torm	Ωf	leer	ıan	co.

EPA Reg. Number:

101563-409

Conditional

Name of Pesticide Product:

E2Y45 0.067%G INSECTICIDE ON FERTILIZER P

Date of Issuance:

5/29/25

NOTICE OF PESTICIDE:

X Registration

___ Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Environmental Science U.S., LLC 5000 CentreGreen Way, Suite 400 Cary, NC 27513, PA 19104

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Melissa Bridges, Product Manager 7 Invertebrate-Vertebrate Branch III Registration Division (7505T)

Date:

5/29/25

Page 2 of 2 EPA Reg. No. 101563-409 Case No. 495186

- 2. You are required to comply with the data requirements described in the generic data call-in (GDCI) identified below:
 - a. Chlorantraniliprole GDCI-090100-1895

You must comply with all of the data requirements within the established deadlines. If you have questions about the GDCI listed above, you may contact the Chemical Review Manager in the Pesticide Re-Evaluation Division: http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1

- 3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 101563-409."
- 4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 05/09/2025
- Alternate CSF 1 dated 05/09/2025
- Alternate CSF 2 dated 05/09/2025

If you have any questions, please contact Cassandra Luna at Luna. Cassandra@epa.gov.

CHLORANTRANILIPROLE GROUP 28 INSECTICIDE

E2Y45

0.067%G INSECTICIDE ON FERTILIZER P

Alternative Brand Name:

- E2Y45 0.067%G INSECTICIDE ON FERTILIZER
- DURENTIS® 0.067%G INSECTICIDE ON FERTILIZER

For foliar and systemic control of insect pests on turfgrass and ornamental plants on golf courses, sports and recreation turf, lawns and landscaped areas on and around residential, commercial and institutional properties, commercial sod farms, and interior plantscapes.

Use sites include residential, institutional, public, commercial, and industrial buildings, golf courses, parks, recreational areas and athletic fields, and sod farms.

EPA Reg. No. 101563-

EPA Est. XXX

Active Ingredients:	By Wt.
Chlorantraniliprole*	0.067%
Other Ingredients:	99.933%
Total:	100.000%

^{*}Contains 0.00067 lb of chlorantraniliprole per pound of product.

Chlorantraniliprole (CASRN 500008-45-7) belongs to the anthranilic diamide chemical class.

{Required Text if the product's Guaranteed Analysis is greater than 0-0-0}

Guaranteed Analysis XX-X-XX

Sulfate Of Potash, Iron Sucrate

Chlorine (CI) Max X.XX%

KEEP OUT OF REACH OF CHILDREN

See [other panels] [[inside]booklet] [back][panel] for additional precautionary information.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

PRODUCED FOR

Environmental Science U.S., LLC 5000 CentreGreen Way, Suite 400 Cary, NC 27513, PA 19104

Net Contents:

ACCEPTED

05/29/2025

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 404 F00 400

101563-409

5/1/2025

MEDICAL HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the Environmental Science U.S., LLC Emergency Response Telephone No. 1-800-424-9300 for emergency medical treatment information.

For Technical Support or information regarding the use of this product, call 1-800-331-2867

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

When used as directed, this product does not present a hazard to humans or domestic animals.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- · Shoes plus socks.

After the product has been loaded into properly calibrated application equipment, shirt, pants, socks, and shoes are sufficient Personal Protection Equipment (PPE). Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handing this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to aquatic invertebrates, oysters, and shrimp. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to use sites.

<u>Surface water advisory:</u> This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of chlorantraniliprole from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

<u>Groundwater advisory</u>: This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read the following "Directions for Use" carefully before applying this product.

STOP. READ AND UNDERSTAND THE ENTIRE LABEL BEFORE USE.

E2Y45 0.067%G insecticide on fertilizer P must be used only in accordance with the directions on this label, in separate EPA-approved labeling or exemptions under FIFRA (Supplemental Labels, Special Local Need Registrations, FIFRA Section 18 exemptions, FIFRA 2(ee) Bulletins), or as otherwise permitted by FIFRA. Always read the entire label, including the Limitation of Warranty and Liability.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN TURF INJURY AND POOR INSECT CONTROL.

5/1/2025 Page 2 of 9

RESTRICTIONS

- DO NOT apply this product with aerial application equipment.
- DO NOT apply this product through any type of irrigation system.
- DO NOT use for commercial grass seed production.
- DO NOT use in commercial ornamental nurseries and greenhouses.
- DO NOT apply more than 750 lbs E2Y45 0.067%G insecticide on fertilizer P (equivalent to 0.5 lb of active ingredient) per acre per calendar year.
- DO NOT apply more than 300 lbs E2Y45 0.067%G insecticide on fertilizer P (equivalent to 0.2 lb of active
 ingredient) in a single application on sod farms.
- Wait a minimum of 7 days to retreat.
- DO NOT formulate this product into other end-use products without written permission from Environmental Science U.S., LLC.

New York State restrictions:

- Not registered for sale, sale into, distribution and/or use in Nassau, Suffolk, Kings, Queens Counties of New York State unless permitted under FIFRA Section 24fc) Special Local Need Registration.
- Buffer Zones
 - Golf course greens and tee boxes:

 DO NOT apply within 25 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
 - Golf course fairways:
 DO NOT apply within 50 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
 - DO NOT apply within 50 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch)
 All other Sites:

DO NOT apply within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers (as applicable), and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Long-sleeved shirt and long pants
- Shoes plus socks.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Professional applications to golf courses, residential, industrial and commercial lawns and sports fields are not within the scope of the Worker Protection Standard.

Do not allow other workers to enter the treated area while application is in progress.

RESISTANCE MANAGEMENT

5/1/2025 Page 3 of 9

For resistance management, E2Y45 0.067%G insecticide on fertilizer P contains a Group 28 insecticide. Any insect population may contain individuals naturally resistant to E2Y45 0.067%G insecticide on fertilizer P and other Group 28 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same area. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of E2Y45 0.067%G insecticide on fertilizer P or other Group 28 insecticide within a growing season, or among growing seasons with different insecticide groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - o Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical
 information related to pesticide use, record keeping, and which considers cultural, biological and other
 chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or pest control advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.

For further information or to report suspected resistance, contact Environmental Science U.S., LLC at 800-331-2867. You can also contact your pesticide distributor or university extension specialist to report resistance.

INTEGRATED PEST MANAGEMENT

Environmental Science U.S., LLC supports the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an IPM program, which can include biological, cultural, and genetic practices, aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, rotation of insecticides with different modes-of-action and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants, or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

General Information

E2Y45 0.067%G insecticide on fertilizer P contains 0.00067 lb of chlorantraniliprole per pound of product.

E2Y45 0.067%G insecticide on fertilizer P is a member of the anthranilic diamide class of insecticides with a mode of action acting on insect ryanodine receptors. E2Y45 0.067%G insecticide on fertilizer P is effective through ingestion of treated plant material. After exposure to E2Y45 0.067%G insecticide on fertilizer P affected insects will rapidly stop feeding, become paralyzed, and typically die within 1 - 3 days. Time applications to the most susceptible insect pest stage, typically at egg lay, egg hatch and/or newly hatched larvae before populations reach damaging levels. When pest populations are high, use the highest listed application rate for that pest.

Table 1: List of Controlled Pests

List of Controlled Pests		
Annual Bluegrass Weevil (Hyperodes) adults	White Grubs including	
Armyworms	Asian Garden Beetle	
Billbugs	Black Turfgrass Ataenius (Adult)	
Chinch Bugs*	Japanese Beetle	
Cutworms	European Masked Chafer	

5/1/2025 Page 4 of 9

European Crane Fly (larvae)	Northern Masked Chafer	
Spittlebugs (two-lined)	 Southern Masked Chafer 	
Sod Webworm	Oriental Beetle	
	Green June Beetle	
	May/ June Beetle	
*Suppression only		

TURFGRASS

Established Turf

Use sites include residential, institutional, public, commercial, and industrial buildings, golf courses, parks, recreational areas and athletic fields, and sod farms. E2Y45 0.067%G insecticide on fertilizer P may be used on seeded, sodded, or sprigged turfgrasses.

Uniform application is necessary to provide good pest control. Apply as a broadcast application.

Restrictions and Other Use Precautions:

- DO NOT use for commercial grass seed production.
- DO NOT apply this product through any type of irrigation system.
- Do not apply more than once every 7 days.
- DO NOT apply more than 750 lbs product (equivalent to 0.5 lbs chlorantraniliprole) per acre per calendar year in broadcast application to turfgrass.
- DO NOT apply more than 300 lbs product (equivalent to 0.2 lbs chlorantraniliprole) per acre in a single application to sod farms.

USE RATE Conversion			
lbs product per 1000 sq ft	Ibs Chlorantraniliprole per	lbs product per acre	
	acre		
3.44	0.1	150	
5.17	0.15	225	
6.88	0.2	300	
17.2	0.5	750	

	Single Application Use Rate: E2Y45 0.067%G insecticide on fertilizer P	
Target Pest	lbs per 1000 sq ft	lbs per acre
White Grubs (including Aphodius spp., Asiatic garden beetle, black turfgrass ataenius, European chafer, green June beetle, Japanese beetle, May/June beetles (Phyllophaga spp.), northern masked chafer, oriental beetle and southern masked chafer) Billbugs European Crane Fly Turf caterpillars (including armyworms, cutworms and sod webworms)	3.44 - 6.88	150 - 300
Annual bluegrass weevil	5.17 - 6.88	225 - 300
Chinch bugs (suppression only)	3.44 - 6.88	150 - 300

5/1/2025 Page 5 of 9

ORNAMENTAL PLANTS

E2Y45 0.067%G insecticide on fertilizer P may be used to control the pests of landscape ornamentals and interior plantscapes.

For maximum efficiency, apply at rates up to 500 lbs product (0.33 lbs chlorantraniliprole) per acre in a single application to control any of the pests listed below.

Restrictions and Other Use Precautions:

• DO NOT apply more than 750 lbs product (0.5 lbs chlorantraniliprole) per acre per calendar year in broadcast application to landscape ornamentals.

USE RATE Conversion			
lbs product per 1000 sq ft	Ibs Chlorantraniliprole per Ibs product per		
	acre		
3.44	0.1	150	
6.88	0.2	300	
11.3	0.33	500	

	Single Application Use Rate: E2Y45 0.067%G insecticide on fertilizer P	
Target Pest	lbs per 1000 sq ft	lbs per acre
White Grubs (Asiatic garden beetle, European chafer, green June beetle, Japanese beetle, May/June beetles (Phyllophaga spp.), northern masked chafer, oriental beetle and southern masked chafer)	3.44 - 6.88	150 - 300

5/1/2025 Page 6 of 9

CLEANING APPLICATION EQUIPMENT

Ensure following best practices for application equipment post use. Thoroughly clean all application equipment to mitigate the risk of hardened deposits. Use clean water to rinse all associated application equipment and adhere to necessary safety precautions. Do not clean equipment near water sources, including wells. Dispose of waste rinse water in accordance with local regulations.

SPREADER EQUIPMENT SETTINGS

Broadcast application should be performed with a calibrated commercial broadcast spreader to evenly distribute the granules. [Approximate settings for several models are provided in the table below.]

Insert optional spreader setting table here.

{Note to Reviewer: The exact setting will depend on the fertilizer prill size and type of fertilizer in the bag. For this reason, a representative table cannot be provided here.}

[Optional metals information]

[Information regarding the contents and levels of metals in this product is available on the internet at <u>Trace Metals - Metal Content Analysis Information</u>]

[Information regarding the contents and levels of metals in this product is available on the internet at <u>Fertilizer Database</u> | <u>Washington State Department of Agriculture</u>]

[Information regarding the contents and levels of metals in this product is available on the internet at CDFA Fertilizer Product Database]

[Information regarding the contents and levels of metals in this product is available on the internet at <u>Search Fertilizer</u> <u>Registered Products</u>]

5/1/2025 Page 7 of 9

STORAGE AND DISPOSAL

Do not contaminate water, other pesticide, fertilizer, food, or feed in storage or disposal. Not for use or storage in or around the home.

PESTICIDE STORAGE: Keep out of reach of children and pets. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, close tightly.

In case of spill, avid contact, isolate area and keep out animals and unprotected persons. Confine spills. You may contact the Environmental Science U.S., LLC Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Environmental Science U.S., LLC Emergency Response Telephone No. is 1-800-424-9300.

To confine spill: Cover to prevent disposal. Place damaged package in a holding container. Identify contents.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: [Nonrefillable container.] Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling, if available, or dispose of empty bag in a sanitary landfill.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of Environmental Science U.S., LLC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Environmental Science U.S., LLC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ENVIRONMENTAL SCIENCE U.S., LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or Environmental Science U.S., LLC, and buyer assumes the risk of any such use.

To the extent consistent with applicable law, Environmental Science U.S., LLC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ENVIRONMENTAL SCIENCE U.S., LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ENVIRONMENTAL SCIENCE U.S., LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

Owned or Licensed U.S. Patents for this product can be found at https://patent.envu.com

Envu and the Envu logo are trademarks and DURENTIS® are registered trademarks owned by Environmental Science U.S., LLC or one of its affiliates.

5/1/2025 Page 8 of 9

[Optional State Specific Language]

{Note to Reviewer: These statements are typically added right after the Fertilizer composition section. Typically states have not raised questions concerning where these statements are on the labels, as long as the information is present. Only states where the product will be registered will be included on the commercial label.}

FLORIDA APPLICATIONS: For use on athletic turf in FL, we recommend you follow SL191, "Recommendations for N, P, K & Mg for Golf Course and Athletic Field Fertilization Base on Mehlich I Extractant", available on the internet at http://edis.ifas.ufl.edu/SS404.

FLORIDA APPLICATIONS: For commercial application to urban turf or lawns in FL, we recommend you follow the Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries, available on the internet at http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP Book final.pdf

Optional State Specific Language for Virginia

In Virginia this lawn/turf fertilizer contains phosphorous and is intended only for nonagricultural use on (1) turf during its first growing season, (2) on turf areas being repaired or renovated, and (3) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not intended for the routine maintenance of turf.

Optional State Specific Language for Maryland

This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article. Annotated Code of Maryland performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.

5/1/2025 Page 9 of 9