



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs

Registration Division (7505T)

1200 Pennsylvania Ave., N.W.

Washington, D.C. 20460

EPA Reg. Number:

103220-7

Date of Issuance:

11/24/25

NOTICE OF PESTICIDE:

☒ Registration

☐ Reregistration

(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Dicamba DMA

Name and Address of Registrant (include ZIP Code):

Agripro USA Limited

815 Brazos St

Austin TX 78701

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Submit one copy of the final printed label for the record before you release the product for shipment.

*Continues page 2*

Signature of Approving Official:

*Emily Schmid*

Emily Schmid, Product Manager 25

Herbicide Branch, Registration Division (7505T)

Date:

11/24/25

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Please note that the alternate brand name, "**AgriPro Dicamba 4 DMA**" has been added to the product record.

The record for this product currently contains the following CSF:

- Basic CSF dated 05/16/2025

If you have any questions, please contact Jamie Millard at (202) 566-2726 or by email at [millard.jamie@epa.gov](mailto:millard.jamie@epa.gov).

Enclosure

[MASTER LABEL]

DICAMBA GROUP 4 HERBICIDE

# Dicamba DMA

## ABN: AgriPro Dicamba 4 DMA

For weed control in Corn, Cotton, Sorghum, Small Grains, Pasture, Hay, Rangeland, General Farmstead (Non-Cropland), Rights-of-Way, Public Utility and Industrial Areas, Fallow, Sugarcane, Asparagus, Turf and Grass Seed Crops.

<b>ACTIVE INGREDIENT:</b>	<b>By Wt. %</b>
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid) *	49.41%
<b>OTHER INGREDIENTS:</b>	50.59%
<b>TOTAL:</b>	100.00%

\*This product contains 41.0% 3,6-dichloro-o-anisic acid (Dicamba) or 4 pounds per gallon (480 g/L).

### KEEP OUT OF REACH OF CHILDREN

### CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• <b>DO NOT</b> induce vomiting unless told to do so by the poison control center or doctor.</li><li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For 24-Hour Medical Emergency Assistance (Human or Animal), call: <b>1-800-222-1222</b> .	

[Optional referral statements when booklets and container labels are used:]


[See label booklet for [complete] [First Aid,] [Precautionary Statements], [Directions For Use], and [Storage and Disposal].]

EPA Reg. No. 103220-7

EPA Est. No. \_\_\_\_\_

Net Content: \_\_\_\_\_ Gals. (\_\_\_\_ L)

Manufactured For:  
AgriPro USA Limited  
815 Brazos St  
Austin TX 78701



### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION/PRECAUCIÓN**

Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**All mixers, loaders and applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of Barrier Laminate, Butyl Rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils, Neoprene Rubber ≥ 14 mils, Natural Rubber ≥ 14 mils, Polyethylene, Polyvinyl Chloride (PVC) ≥ 14 mils, and Viton ≥ 14 mils (except for applicators using ground boom equipment, pilots, and flaggers)
- Shoes plus socks

See **ENGINEERING CONTROLS** for additional requirements and exceptions.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROLS**

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-f)], the handler PPE requirements may be reduced or modified as specified in the WPS. Pilots must use cockpits in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-f)].

**USER SAFETY RECOMMENDATIONS****Users should:**

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Apply this product only as directed on label.

This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Ground And Surface Water Protection**

**Point source contamination:** To prevent point source contamination, do not mix or load this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers and natural or impounded lakes and reservoirs. Do not apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas as described below.

Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or moved across the pad. The pad must be self-contained to prevent surface water flow over or from the pad. The pad capacity must be maintained at 110% that of the largest pesticide container or application equipment used on the pad and have sufficient capacity to contain all product spills, equipment or container leaks, equipment washwaters and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding well-head setbacks and operational containment.

Care must be taken when using this product to prevent: a) back siphoning into wells, b) spills or c) improper disposal of excess pesticide, spray mixtures or rinsates. Check valves or anti siphoning devices must be used on all mixing equipment.

**Movement by surface runoff or through soil:** Do not apply under conditions which favor runoff. Do not apply to impervious substrates such as paved or highly compacted surfaces in areas with high potential for groundwater contamination. Groundwater contamination may occur in areas where soils are permeable or coarse and groundwater is near the surface. Do not apply to soils classified as *Sand* with less than 3% organic matter and where groundwater depth is shallow. To minimize the possibility of ground water contamination, carefully follow specified application rates as affected by soil type in the **CROPS** section of this label.

**Movement by water erosion of treated soil:** Do not apply or incorporate this product through any type of irrigation equipment nor

by flood or furrow irrigation. Ensure treated areas have received at least 0.5 inch of rainfall (or irrigation) before using tailwater for subsequent irrigation of other fields.

**Physical and Chemical Hazards:** Do not mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Before applying this product, read all directions and precautions appearing on this label. Failure to follow all directions, precautions and restrictions may result in unsatisfactory weed control, crop injury, or illegal residues.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS.

**Do not enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of 24 hours.**

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls worn over short-sleeved shirt and short pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant headgear for overhead exposure
- Protective eyewear

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter treated areas until sprays have dried.

### RESISTANCE MANAGEMENT

DICAMBA	GROUP	4	HERBICIDE
---------	-------	---	-----------

For resistance management, this product is a Group 4 mode of action herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 mode of action herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or

integrated weed-management recommendations for specific crops and weed biotypes.

- For further information or to report suspected resistance, contact Agripro USA Limited representatives.

### PRODUCT INFORMATION

The following directions apply to all uses of this product. Additional precautions and restrictions will be found in each specific use section.

#### Use Restrictions

- Do not treat irrigation ditches or water used for crop irrigation or domestic uses.
- Do not apply this product through any type of irrigation system.
- Do not apply more than 1 lb. ae (32 oz. **Agripro Dicamba DMA**) per acre per application, and no more than 2 applications per year.

#### Mixing And Application

UNLESS OTHERWISE SPECIFIED UNDER THE INDIVIDUAL USE HEADING OF THIS LABEL, THE FOLLOWING DIRECTIONS APPLY TO ALL CROP AND NON-CROP USES OF THIS PRODUCT. REFER TO INDIVIDUAL USE SECTIONS FOR ADDITIONAL PRECAUTIONS, RESTRICTIONS, APPLICATION RATES AND TIMINGS.

This product is a water-soluble formulation that can be applied using water or sprayable fluid fertilizer as the carrier. If a fluid fertilizer is to be used, perform a compatibility test (see the **Compatibility Test** section of this label) prior to tank-mixing.

Use ground or aerial application equipment which will give good spray coverage of weed foliage. However, do not use aerial application equipment if spray particles can be carried by wind into areas where sensitive crops or plants are growing or when temperature inversions exist.

Apply 3 to 50 gallons of a diluted spray per treated acre when using ground application equipment or 1 to 10 gallons of a diluted spray per treated acre (2 to 20 gallons of diluted spray per acre for pre-harvest uses) in a water-based carrier when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays.

Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

To avoid uneven spray coverage, this product should not be applied during periods of gusty wind or when wind is in excess of 15 mph.

Avoid disturbing (e.g., cultivating or mowing) treated areas for at least 7 days following application.

#### Best Stewardship Practices

This product provides effective broadleaf weed and brush control when properly applied. Best stewardship practices in all mixing, loading and application operations not only maximize weed control, but also protect ground and surface waters and minimize off-target movement.

#### Sensitive Crop Precautions

This product may cause injury to desirable trees and plants, particularly Beans, Cotton, Flowers, Fruit trees, Grapes, Ornamentals, Peas, Potatoes, Soybeans, Sunflowers, Tobacco, Tomatoes, and other broadleaf plants when contacting their roots, stems or foliage. These plants are most sensitive to this product during their development or growing stage. Follow the precautions listed below when using this product:

- Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing, or when temperature inversions exist. Leave an adequate buffer zone between area to be treated and sensitive plants. Coarse sprays are less likely to drift out of the target area than fine sprays.
- Use coarse sprays to avoid potential herbicide drift. Select nozzles which are designed to produce minimal amounts of fine spray particles. Examples of nozzles designed to produce coarse spray via ground applications are Delavan Raindrops, Spraying Systems XR flat fans or large capacity flood nozzles such as D10, TK10 or greater capacity tips. Keep the spray pressure at or below 20 PSI and the spray volume at or above 20 gallons per acre, unless otherwise required by the manufacturer of drift-reducing nozzles. Consult with your spray nozzle supplier concerning the choice of drift-reducing nozzles.
- Agriculturally approved drift-reducing additives may be used.
- To avoid injury to desirable plants, equipment used to apply this product should be thoroughly cleaned (see **Procedure For Cleaning Spray Equipment** section of this label) before reusing to apply any other chemicals.

#### Use Restrictions on Sensitive Crops

- Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of this product with the roots of desirable plants such as trees and shrubs.
- Do not spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of adjacent sensitive crops.
- Do not apply this product adjacent to sensitive crops when the temperature on the day of application is expected to exceed 85°F, as drift is more likely to occur.

All crop uses of this product are intended for a normal growing interval between planting and harvest. No crop rotation restrictions exist if normal harvest of treated crop has occurred. If this interval is shortened, such as in cover crop that will be plowed under, do not follow up with the planting of a sensitive crop.

Crops growing under stress conditions such as drought, poor fertility or foliar damage due to hail, wind or insects, can exhibit various injury symptoms that may be more pronounced if herbicides are applied.

Consult your local or State authorities for possible application restrictions and advice concerning these and other special local use situations. TANK-MIX DIRECTIONS ARE FOR USE ONLY IN STATES WHERE THE TANK-MIX PRODUCT AND APPLICATION SITE ARE REGISTERED.

### Band Treatments

This product may be applied as a band treatment. Use the formula below to determine the appropriate rate and volume per acre:

Band width in inches

Row width in inches      ×      Broadcast RATE per treated acre      =      Band RATE per treated acre

Band width in inches

Row width in inches      ×      Broadcast VOLUME per treated acre      =      Band VOLUME per treated acre

### Compatibility Test

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

Amount of Herbicide to Add to One Pint of Spray Carrier (Assuming Volume is 25 Gallons per Acre)		
Herbicide Formulations	Rate Per Acre	Level Teaspoons
Dry	1 lb.	1.5
Liquid	1 pt.	0.5

If herbicide(s) do not ball up or form flakes, sludge, gels, oily films or layers or other precipitates, then the tested spray mix is compatible. Usually, incompatibility in any of the above described forms will occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is needed. Re-run the above compatibility test with a suitable compatibility agent (0.25 teaspoon is equivalent to 2 pints per 100 gallons of fluid fertilizer).

### Procedure For Cleaning Spray Equipment

The steps listed below are suggested for thorough cleaning of spray equipment following applications of this product or tank-mixes of this product or tank-mixes of this product plus 2,4-D Amine.

1. Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
2. Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Remove the nozzles and screens and flush the system with two full tanks of water.

The steps listed below are suggested for thorough cleaning of spray equipment used to apply this product as a tank-mix with wettable powders (WP), emulsifiable concentrates (EC) or other types of water-dispersible formulations. This product tank-mixed with water-dispersible formulations requires the use of a water/detergent rinse.

5. Complete Step 1.
6. Fill tank with water while adding 2 pounds of detergent for every 40 gallons of water. Operate the pump to circulate the detergent solution through the sprayer system for 2 to 10 minutes and discharge a small amount of the solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
7. Flush the detergent solution out of the spray tank through the boom.
8. Repeat Step 1, and follow with Steps 2, 3 and 4.

### WEED LIST

This is a list of weeds which may be treated with **Agripro Dicamba DMA** in accordance with this label as specified under the rates and timing sections of the individual use headings. Proper usage of **Agripro Dicamba DMA** will give control or growth suppression of many annual, biennial, and perennial broadleaf weeds and many woody brush and vine species, including:

ANNUALS	
Amaranth, Spiny (Spiny Pigweed)	Nightshade, Black
Aster, Slender	Pennycress, Field (Fanweed, Frenchweed, Stinkweed)
Bedstraw	Pepperweed, Virginia (Peppergrass)
Beggarweed, Florida	Pigweed, Prostrate
Broomweed, Common	Pigweed, Redroot (Carelessweed)
Buckwheat, Wild	Pigweed, Rough
Buffalobur	Pigweed, Smooth
Burclover, California	Pigweed (Triazine-resistant)



Burcucumber Buttercup, Roughseed Carpetweed Catchfly, Nightflowering Chamomile, Corn Chickweed, Common Clovers (Annual) Cockle, Corn Cockle, Cow Cocklebur, Common Croton, Tropic Croton, Woolly Daisy, English Evening Primrose, Cutleaf Fleabane, Annual Goosefoot, Nettleleaf Henbit Jimsonweed Knotweed Kochia Ladysthumb Lambsquarters, Common Lambsquarters (Triazine-resistant) Lettuce, Prickly Mallow, Common Mallow, Venice Maretail (Horseweed) Mayweed Morningglory, Ivyleaf Morningglory, Tall Mustard, Tansy Mustard, Wild Mustard, Yellowtops	Pigweed, Tumble Poorjoe Puncturevine Purslane, Common Pusley, Florida Radish, Wild Ragweed, Common Ragweed, Giant (Buffaloweed) Ragweed, Lance-leaf Rubberweed, Bitter (Bitterweed) Sesbania, Hemp Shepherd's Purse Sicklepod Sida, Prickly (Teaweed) Smartweed, Green Smartweed, Pennsylvania Sneezeweed, Bitter Sowthistle, Annual Sowthistle, Spiny Spikeweed, Common Spurge, Prostrate Spurry, Corn Starbur, Bristly Sumpweed, Rough Sunflower, Common (Wild) Sunflower, Volunteer Thistle, Russian Velvetleaf Waterhemp Waterprimrose, Winged Wormwood, Annual
--	---

**BIENNIALS**

Burdock, Common Carrot, Wild (Queen Anne's Lace) Cockle, White Evening Primrose, Common Geranium, Carolina Gromwell Knapweed, Diffuse Knapweed, Spotted Mallow, Dwarf	Plantain, Bracted Ragwort, Tansy Starthistle, Yellow Sweetclover Teasel Thistle, Bull Thistle, Milk Thistle, Musk Thistle, Plumeless
---	--

**PERENNIALS**

Alfalfa* Artichoke, Jerusalem Aster, Spiny Aster, Whiteheath Bedstraw, Smooth Bindweed, Field Bindweed, Hedge Blueweed, Texas Bursage (Bur Ragweed) (Lakeweed) (Povertyweed)* Bursage, Woollyleaf (Lakeweed) Buttercup, Tall Campion, Bladder Chickweed, Field Chickweed (Mouseear, Canada) Chicory Clover, Hop* Dandelion, Common* Dock, Broadleaf (Bitterdock)* Dock, Curly* Dogbane, Hemp	Milkweed, Common Milkweed, Honeyvine Milkweed, Western Whorled Nettle, Stinging Nightshade, Silverleaf (White Horsenettle) Onion, Wild Plantain, Broadleaf* Plantain, Buckhorn Pokeweed Ragweed, Western Redvine Sericia Lespedeza Smartweed, Swamp Snakeweed, Broom Sorrel, Red (Sheep Sorrel)* Sowthistle Sowthistle, Perennial Spurge, Leafy Sundrop, Halfshrub (Evening Primrose) Thistle, Canada
---	--



Dogfennel (Cypressweed)* Fern, Bracken Garlic, Wild Goldenrod, Canada Goldenrod, Missouri Goldenweed, Common Hawkweed Henbane, Black Horsenettle, Carolina Ironweed Knapweed, Black Knapweed, Russian Milkweed, Climbing	Toadflex, Dalmation Tropical Soda Apple Trumpet creeper (Buckvine) Vetch Waterhemlock Waterprimrose, Creeping Woodsorrel, Creeping Common Yellow* Wormwood, Common Wormwood, Louisiana Yankee weed* Yarrow, Common
*These perennials may be controlled using this product at rates lower than those specified for other listed perennial weeds (see <b>Application Rates and Timings</b> sections).	

WOODY SPECIES	
Alder Ash Aspen Basswood Beech Birch Blackberry* Blackgum* Cedar* Cherry Chinquapin Cottonwood Creosotebush* Cucumbertree Dewberry* Dogwood* Elm Grape Hawthorn (Thornapple)* Hemlock Hickory Honeylocust Honeysuckle Hornbeam Huckleberry Huisache Ivy, Poison Kudzu	Locust, Black Maple Mesquite Oak Oak, Poison Olive, Russian Persimmon, Eastern Pine Plum, Sand (Wild Plum)* Poplar Rabbitbrush Redcedar, Eastern* Rose McCartney* Rose Multiflora* Sagebrush, Fringe Sassafras Serviceberry Spicebush Spruce Sumac Sweetgum* Sycamore Tarbush Willow Witchhazel Yaupon* Yucca*
*Growth suppression	

## CROPS

### CORN (FIELD, POP\*, SEED\*, SILAGE)

Observe all precautions and restrictions. Read and follow mixing, application and cleaning instructions on this label, as well as the following:

\*DO NOT apply this product to Popcorn or Seed corn without first verifying with your local Seed corn company (supplier) the selectivity of this product on your inbred line or variety of Popcorn. This precaution will help avoid potential injury of sensitive varieties.

Direct contact of this product with Corn seed must be avoided. If Corn seeds are less than 1.5 inches below the surface, delay application until Corn has emerged.

Applications of this product to corn during periods of rapid growth may result in temporary leaning. Corn will usually become erect within 3 to 7 days. Cultivation should be delayed until after corn is growing normally to avoid breakage.

Applications of this product to Corn during periods of rapid growth may result in temporary leaning. Corn will usually become erect within 3 to 7 days. Delay cultivation until after Corn is growing normally to avoid breakage.

Agriculturally approved surfactants or sprayable fertilizer (0.5 to 1 gallon per acre of 28%, 30%, 32% urea Ammonium nitrate or 2.5 pounds per acre spray-grade ammonium sulfate<sup>1</sup>) may be added to the spray mixture to improve post-emergence weed control, particularly in dry growing conditions.

<sup>1</sup> Not for use in California

Several synthetic Pyrethroid insecticides are labeled for tank-mix applications with this product. Refer to their label for specific directions.

#### Weeds Controlled

**Agripro Dicamba DMA** will control many annual broadleaf weeds or give growth suppression of many perennial broadleaf weeds commonly found in Corn (refer to the **WEED LIST** section).

For best performance, make application when weeds have emerged and are actively growing.

Pre-emergence control of Cocklebur, Jimsonweed and Velvetleaf may be reduced if conditions such as low temperature or lack of soil moisture cause delayed or deep germination of weeds.

#### Pre-Plant/Pre-Emergence In No-Tillage Corn (Preplant)

Applications of **Agripro Dicamba DMA** may be made before, during or after planting to emerged and actively growing broadleaf weeds. Apply this product at 1 pint per treated acre on medium- or fine-textured soils containing 2% or greater organic matter. Use 0.5 pint per treated acre on coarse-textured soils (*Sand, Sandy loam, and Loamy sand*) or medium- and fine-textured soils with less than 2% organic matter.

When planting into a legume sod (e.g., Alfalfa or Clover), apply this product after 4 to 6 inches of regrowth has occurred.

#### Pre-Emergence In Conventional Or Reduced Tillage Corn

**Agripro Dicamba DMA** may be applied after planting and prior to Corn emergence. Application at 1 pint per treated acre may be made to medium- or fine-textured soils which contain 2% or greater organic matter. Do not apply to coarse-textured soil (*Sand, Sandy loam, and Loamy sand*) until after crop emergence (see **Early Post-Emergence** uses below).

Pre-emergence application of this product does not require mechanical incorporation to become active. Perform a shallow mechanical incorporation if application is not followed by adequate rainfall or sprinkler irrigation. Avoid tillage equipment (e.g., drags, harrows) which concentrate treated soil over seed furrow.

#### Early Post-Emergence (All Tillage Systems)

**Spike Through 8 Inch Tall Corn:** **Agripro Dicamba DMA** at 1 pint per treated acre may be applied during the period from Corn emergence through the 5-leaf stage or 8 inches tall, whichever comes first. Reduce the rate to 0.5 pint per treated acre if Corn is growing on coarse-textured soils (*Sand, Sandy loam, and Loamy sand*). See late post-emergence application rates below if the 6<sup>th</sup> true-leaf is emerging from whorl, or Corn is greater than 8 inches tall.

#### Late Post-Emergence (All Tillage Systems)

**8 to 36 Inch Tall Corn:** Application of **Agripro Dicamba DMA** at 0.5 pint per treated acre may be made from 8 to 36 inch tall Corn or 15 days before tassel emergence, whichever comes first. For best performance, make applications when weeds are less than 3 inches tall.

Make directed spray application when:

- 1) Corn leaves prevent proper spray coverage;
- 2) sensitive crops are growing nearby;
- 3) tank-mixing with 2,4-D.

#### Use Restrictions:

- This product is not registered for use on sweet corn.
- Do not apply this product when Soybeans are growing nearby if any of these conditions exist:
  - Corn is more than 24 inches tall.
  - Soybeans are more than 10 inches tall.
  - Soybeans have begun to bloom.
- Up to 2 applications of this product may be made per year.
- Do not exceed a total of 1.5 pints (0.75 lb. ae) of this product per treated acre per crop year.
- Allow two weeks or more between applications of this product.
- Do not use adjuvants containing penetrants such as petroleum-based oils after crop emergence or crop injury may result.
- Corn may be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity.

#### Overlay (Sequential) Treatments

**Agripro Dicamba DMA** may be applied to ground previously treated with one or more of the following active ingredients:

Active Ingredient(s)
Acetochlor
Alachlor
Alachlor + Atrazine
Atrazine
Atrazine + Metolachlor
Atrazine + Dicamba
Butylate
Dicamba
Dimethenamid
Dimethenamid-p + Atrazine
EPTC
Flumetsulam
Glyphosate
Halosulfuron
S-Metolachlor
Paraquat
Pendimethalin
Propachlor
Simazine

Apply **Agripro Dicamba DMA** at 0.5 pint per treated acre to ground previously treated with full rates of Dicamba and Atrazine + Dicamba herbicides. **Use Restrictions:** Total Dicamba a.e. applied from all sources must not exceed 1.0 pound per acre. Allow at least 2 weeks between applications. No more than 2 total applications of Dicamba from any source may be made per acre per year.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

#### Tank-Mix Treatments For Corn

**Agripro Dicamba DMA** may be tank-mixed with one or more of, but not limited to, the following herbicides for control of grasses or additional broadleaf weeds in Corn. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

Active Ingredient
Acetochlor
Alachlor
Atrazine
Clopyralid
Dimethenamid
2,4-D
Glyphosate
Paraquat
Pendimethalin
Primisulfuron
Pyridate
Simazine

#### COTTON (EXCEPT CA)

**Pre-Plant Application:** Apply up to 8 fluid ounces of **Agripro Dicamba DMA** per acre to control emerged broadleaf weeds prior to planting cotton in conventional or conservation tillage systems.

For best performance, apply **Agripro Dicamba DMA** when weeds are in the 2- to 4-leaf stage and rosettes are less than 2 inches across.

#### Use Restrictions:

- Do not plant Cotton for at least 21 days after application and after allowing for a minimum accumulation of 1 inch of rainfall or overhead irrigation.
- Do not apply to Cotton west of the Rockies or to geographic areas with average annual rainfall of less than 25 inches.
- Do not apply more than 1 lb. ae (32 oz. Agripro Dicamba DMA) per acre per application.
- Do not apply more than twice per year.
- If applying a Spring pre-plant treatment following application of a Fall pre-plant (post-harvest) treatment, the combination of both treatments may not exceed 4 pints of this product (2 lbs. ae) per acre per year.

**Tank-Mix Treatments**

For control of grasses or additional broadleaf weeds, **Agripro Dicamba DMA** may be tank-mixed with glyphosate. Glyphosate, Paraquat and Prometryn.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

**TRITICALE (FALL AND SPRING SEEDED) (EXCEPT CA)****Early Season Applications**

Apply 2 to 4 fluid ounces of **Agripro Dicamba DMA** to Triticale.

Make early season applications to Fall seeded Triticale prior to jointing stage. Make early season applications to Spring seeded Triticale before crop reaches the 6-leaf stage.

**Tank-Mixes**

For best results, use **Agripro Dicamba DMA** in tank-mixture with Bromoxynil.

**Use Restrictions:**

- Do not apply more than 1 lb. ae (32 oz. Agripro Dicamba DMA) per acre per application.
- Do not apply more than twice per year.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

**SORGHUM (MILO)**

Observe all precautions and restrictions on this label including the reference to crops growing under stress. Read and follow mixing, application, and cleaning instructions on this label.

If Sorghum is grown for pasture or hay, refer to the **PASTURE, HAY, RANGELAND AND FARMSTEAD (NON-CROPLAND)** section of this label.

**Use Restrictions:**

- Do not graze or feed treated Sorghum forage or silage prior to mature Grain stage.
- Pre-harvest Interval (PHI) restrictions are as follows: Grain sorghum—30 days; Fodder—30 days; Forage—20 days.
- Do not apply this product to Sorghum grown for seed production.
- Make no more than one application per year.
- Applications of **Agripro Dicamba DMA** to Sorghum during periods of rapid growth may result in temporary leaning of plants or rolling of leaves. These effects are usually outgrown within 10 to 14 days.

**Weeds Controlled**

**Agripro Dicamba DMA**, when applied at the specified rate for Sorghum, will control many actively growing annual broadleaf weeds and will reduce competition from established perennial broadleaf weeds as well as control their seedlings (refer to the **WEED LIST** section).

**Rates and Timings**

**Agripro Dicamba DMA** may be applied to emerged and actively growing weeds at least 15 days prior to planting. Make post-emergence application of this product after Sorghum is in the spike stage (all Sorghum emerged) but before Sorghum is 15 inches tall. For best performance, apply when Sorghum is in the 3- to 5-leaf stage and weeds are small (less than 3 inches tall). Use drop pipes (drop nozzles) if Sorghum is taller than 8 inches. Keeping the spray off the Sorghum leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of weed foliage.

**Broadcast Rate Per Treated Acre:** 0.5 pint (0.25 lb. ae)

**Tank-Mix Treatments**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

**Agripro Dicamba DMA plus Atrazine**

For improved control of emerged, actively growing broadleaf weeds including triazine-resistant species and added suppression of perennial broadleaf weeds, tank-mix 0.5 pint of this product with the labeled rate of Atrazine per treated acre. For control of grasses (less than 1.5 inches tall), tank-mix 0.5 pint of this product with the labeled rate of Atrazine per treated acre. For best performance

and minimal crop injury, make application when Sorghum is 3 to 8 inches and when broadleaf weeds are small (less than 6 inches tall). Apply Atrazine before Sorghum is beyond 12 inches in height.

The Atrazine rate will depend upon soil texture and length of residual weed control desired. Follow all State and Federal restrictions pertaining to Atrazine applications.

#### Agripro Dicamba DMA plus Bromoxynil

For improved control of broadleaf weeds, tank-mix bromoxynil at the labeled rate per treated acre. Make application at 4-leaf to 15 inch tall Sorghum. Use drop nozzles to direct spray beneath Sorghum leaves when Sorghum is greater than 8 inches in height.

#### Overlay (Sequential) Treatments

Agripro Dicamba DMA may be applied to ground previously treated with one or more of the following active ingredients:

Active Ingredient	Maximum Rate Per Treated Acre (Lbs. a.i.)
Alachlor (Screen® treated seed)	See product label for rates related to this ingredient
Atrazine*	See product label for rates related to this ingredient
Metolachlor (Concep® treated seed)	See product label for rates related to this ingredient
Propachlor	See product label for rates related to this ingredient
*Maximum use rate for Atrazine is determined by soil type, tillage practices used, surface residue and State or local restrictions. Follow the more restrictive requirements when determining the maximum use rate for Atrazine.	

#### Pre-Harvest Uses (OK & TX Only)

Agripro Dicamba DMA may be applied for weed suppression any time after the Sorghum has reached the soft dough stage. An agriculturally approved surfactant may be used to improve performance. For aerial applications, use at least 2 gallons of water-based carrier per treated acre.

Delay harvest until 30 days after treatment.

**Broadcast Rate Per Treated Acre:** 0.5 pint (0.25 lb. a.e.)

#### SMALL GRAINS (BARLEY, OATS AND WHEAT) NOT UNDERSEED TO LEGUMES

Observe all precautions and restrictions. Read and follow mixing, application and cleaning instructions on this label.

#### Use Restrictions:

- Pre-harvest Interval (PHI) for Grain is 7 days.
- If Small grains are used for pasture hay, the following restrictions apply:
  - Animals cannot be removed from treated area for slaughter prior to 30 days after last application.
  - Treated areas may not be grazed by lactating dairy animals before 7 days after treatment.
  - Do not harvest hay from treated areas before 37 days after treatment.
- Do not apply more than 1 lb. ae (32 oz. Agripro Dicamba DMA) per acre per application.
- Do not apply more than twice per year.

**Note:** There is no waiting period between treatment and grazing for non-lactating dairy animals.

#### Weeds Controlled

Agripro Dicamba DMA or combinations with listed tank-mix partners, will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds, this product may be applied in a tank-mix with other herbicides. Refer to specific tank-mix options.

Alkanet*	Mustard, Tansy
Bedstraw, Catchweed*	Mustard, Treacle*
Bindweed, Field**	Mustard, Tumble (Jim Hill)*
Buckwheat, Tartary	Mustard, Wild*
Buckwheat, Wild	Nightshade, Black
Carpetweed*	Nightshade, Cutleaf*
Chamomile, Corn	Nightshade, Silverleaf (White Horsenettle)**
Chervil, Bur*	Pennycress, Field (Fanweed, Frenchweed, Stinkweed)
Chickweed, Common*	Pepperweed, Peppergrass*
Cockle, Corn	Pigweed, Redroot (Carelessweed)
Cockle, Cow	Pigweed, Rough
Cocklebur, Common	Pigweed, Tumble
Cornflower (Bachelor button)*	Pineappleweed*
Dandelion, Common**	Plantain, Broadleaf**
Dock, Curly**	Poppy, Red Horned*

Dragonhead, American* Evening Primrose, Cutleaf* Falseflax, Smallseeded* Fiddleneck (Tarweed)* Flixweed* Fumitory* Gromwell, Corn* Groundsel, Common* Hempnettle* Henbit Jacobs Ladder* Knawel (German Moss) Knotweed, Prostrate Kochia Ladysthumb Lambsquarters, Common Lettuce, Miners* Lettuce, Prickly Mallow, Common Mayweed, Chamomile (Dogfennel)* Mustard, Blue (Purple)*	Puncturevine* Purslane, Common* Radish, Wild* Ragweed, Common* Ragweed, Giant (Buffaloweed)* Rocket, London* Rocket, Yellow* Salsify (Goatsbeard)* Shepherd's Purse* Smartweed, Green Smartweed, Pennsylvania Sorrel, Red (Sheep Sorrel)* Sowthistle, Annual Starthistle, Yellow* Sunflower, Common (Wild) Thistle, Canada** Thistle, Russian Velvetleaf Vetch* Yarrow, Common**
*These weeds will be controlled with tank-mixtures of <b>Agripro Dicamba DMA</b> . Refer to tank-mix label for specific weeds controlled. **Tank-mixes of <b>Agripro Dicamba DMA</b> will provide suppression of established broadleaf weeds and control of their seedlings.	

### Rates and Timings

Application of **Agripro Dicamba DMA** may be made before, during or after planting Small grains. For best performance, apply when weeds are in the 2- to 3-leaf stage and rosettes are less than 2 inches across. Application of this product to Small grains during period of rapid growth may result in crop leaning. This condition is temporary and will not reduce crop yields.

Use this product at 2 to 4 fluid ounces per treated acre in Wheat, Fall seeded Barley, and Oats, and at 2 to 3 fluid ounces per treated acre in Spring seeded Barley. Use the higher level of listed rate ranges when treating difficult-to-control weeds such as Cow cockle, Kochia, Prickly lettuce, Prostrate knotweed, Russian thistle, and Wild buckwheat or when dense vegetative growth occurs.

This product used in a tank-mix with other herbicides offers best spectrum of weed control and herbicide tolerant or resistant weed management. Refer to specific crop for rate and application timing of this product.

For applications prior to the emergence of weeds or when Sulfonylurea-resistant weeds are present or suspected, use a minimum of 3 fluid ounces per treated acre of this product with a tank-mix herbicide. Non-sulfonylurea herbicides such as 2,4-D or MCPA tank-mixed with this product will offer more consistent control of Sulfonylurea-resistant weeds.

When tank-mixing with Sulfonylurea herbicides, such as metsulfuron-methyl, triasulfuron, tribenuron-methyl, chlorsulfuron + metsulfuron-methyl, chlorsulfuron and thifensulfuron-methyl + tribenuron-methyl, use an agriculturally approved surfactant of at least 80% active ingredient at the rate of 1 to 4 pints per 100 gallons of spray or not more than 0.25 to 0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank-mix and/or when treating more mature and difficult-to-control weeds or dense vegetative growth.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

### FALL AND SPRING SEEDED WHEAT

APPLY THIS PRODUCT TO FALL SEEDED WHEAT PRIOR TO THE JOINTING STAGE. APPLY THIS PRODUCT TO SPRING SEEDED WHEAT BEFORE WHEAT EXCEEDS THE 5-LEAF STAGE.

### Tank-Mix Treatments

**Agripro Dicamba DMA** may be tank-mixed with one or more of, but not limited to the following herbicides.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

### Broadcast Rate per Treated Acre

<sup>1</sup>Apply 2 to 4 fluid ounces of **Agripro Dicamba DMA** with:

Active Ingredient
Bromoxynil
Bromoxynil + MCPA
Chlorsulfuron
Chlorsulfuron + Metsulfuron-methyl



Clorpyralid
Clorpyralid + 2,4-D
2,4-D
Diuron <sup>2</sup>
Fenoxaprop-ethyl <sup>3</sup>
Fenoxaprop-ethyl + MCPA <sup>3</sup>
Fenoxaprop-ethyl + MCPA + 2,4-D <sup>3</sup>
MCPA
Metribuzin <sup>2</sup>
Metsulfuron-methyl
Triasulfuron
Thifensulfuron + Tribenuron-methyl
<sup>1</sup> Apply to early developing Wheat varieties such as TAM 107, Madison or Wakefield between early tillering and the jointing stage. Care should be taken in staging these varieties to be certain that the application occurs prior to the jointing stage.
<sup>2</sup> Tank-mixtures for Fall seeded Wheat only.
<sup>3</sup> Use <b>Agripro Dicamba DMA</b> only at 2 fluid ounces per acre with this product. Do not use if Wild oats is the target weed. Do not use as a tank-mixture with this product on Durum wheat.

### Special Use Tank-Mixes For Spring and Fall Seeded Wheat

Apply 3 to 4<sup>1</sup> fluid ounces of **Agripro Dicamba DMA** with:

<b>Active Ingredient</b>
Chlorsulfuron
Chlorsulfuron + 2,4-D <sup>3</sup>
Chlorsulfuron + Metsulfuron-methyl
(Chlorsulfuron + Metsulfuron-methyl ) + 2,4-D <sup>3</sup>
2,4-D or MCPA <sup>4</sup>
2,4-D or MCPA <sup>4</sup>
Glyphosate <sup>5</sup>
Metsulfuron-methyl
Metsulfuron-methyl + 2,4-D <sup>3</sup>
Tribenuron-methyl
Thifensulfuron + Tribenuron-methyl
(Thifensulfuron + Tribenuron-methyl) + 2,4-D <sup>3</sup>
(Thifensulfuron + Tribenuron-methyl) + 2,4-D <sup>3</sup>
Triasulfuron
Triasulfuron + 2,4-D <sup>3</sup>

<sup>1</sup>This product may be used at 6 fl. oz. on Fall seeded Wheat in Western Oregon as a Spring application only. In CO, KS, NM, OK and TX, up to 8 fl. oz. of this product may be applied on Fall seeded Wheat after it exceeds the 3-leaf stage for suppression of perennial weeds, such as Field bindweed. Applications may be made in the Fall following a frost but before a killing freeze. This product may be tank-mixed with 2,4-D Amine at 8 fl. oz. after Wheat begins to tiller. Periods of extended stress such as cold and wet weather may enhance the possibility of crop injury. For Fall applications only, do not use if the potential for crop injury is not acceptable.

<sup>2</sup>Do not use low rates of Sulfonylurea herbicides on more mature weeds and/or on dense vegetative growth.

<sup>3</sup>Use for improved control of Fiddleneck, Flixweed, Gromwell, Mayweed and Russian thistle.

<sup>4</sup>For use on Fall seeded Wheat only. Do not use unless potential crop injury will be acceptable.

<sup>5</sup>This product may be applied with Glyphosate as a pre-plant application to Small grains with no waiting period prior to planting. Add 0.5% by volume of an agriculturally approved non-ionic surfactant.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

### FALL SEEDED BARLEY

THIS PRODUCT MUST BE APPLIED TO FALL SEEDED BARLEY PRIOR TO THE JOINTING STAGE.

For Spring Barley varieties that are seeded during the Winter months or later, follow the rates and timings given for Spring seeded Barley.

### Tank-Mix Treatments

**Agripro Dicamba DMA** may be tank-mixed with one or more, but not limited to the following herbicides. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

### Broadcast Rate Per Treated Acre

Apply 2 to 4 fluid ounces of **Agripro Dicamba DMA** with:

<b>Active Ingredient</b>
Bromoxynil



Bromoxynil + MCPA
Chlorsulfuron
Chlorsulfuron + Metsulfuron-methyl
2,4-D
MCPA
Metribuzin
Metsulfuron-methyl
Tribenuron-methyl
Thifensulfuron + Tribenuron-methyl
Triasulfuron
*Do not use low rates of Sulfonylurea herbicides on more mature weeds and/or on dense vegetative growth.

**SPRING SEEDED BARLEY**

APPLY THIS PRODUCT BEFORE SPRING SEEDED BARLEY EXCEEDS THE 4-LEAF STAGE.

**Tank-Mix Treatments**

**Agripro Dicamba DMA** may be tank-mixed with one or more of, but not limited to the following herbicides. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

**Broadcast Rate Per Treated Acre**

Apply 2 to 3 fluid ounces of **Agripro Dicamba DMA** with:

Active Ingredient
Bromoxynil
Bromoxynil + MCPA
Chlorsulfuron
Chlorsulfuron + Metsulfuron-methyl
MCPA
Metribuzin
Metsulfuron-methyl
Triasulfuron
Tribenuron-methyl
Thifensulfuron + Tribenuron-methyl
*Do not use low rates of Sulfonylurea herbicides on more mature weeds and/or on dense vegetative growth.

**FALL AND SPRING SEEDED OATS**

APPLY THIS PRODUCT BEFORE SPRING SEEDED OATS EXCEED THE 5-LEAF STAGE. APPLY THIS PRODUCT TO FALL SEEDED OATS PRIOR TO THE JOINTING STAGE.

**Tank-Mix Treatments**

**Agripro Dicamba DMA** may be tank-mixed with one or more, but not limited to the following herbicides. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

**Broadcast Rate Per Treated Acre**

Apply 2 to 4 fluid ounces of **Agripro Dicamba DMA** with:

Active Ingredient
MCPA

**Pre-Harvest Applications For Barley And Wheat (Except CA)**

**Agripro Dicamba DMA** can be used to control weeds that may interfere with harvest of Fall and Spring seeded Barley and Wheat.

Apply 8 fluid ounces of this product per acre as a broadcast or spot treatment to annual broadleaf weeds when Barley or Wheat is in the hard dough stage and the green color is gone from the nodes (joints) of the stem. For best results, apply this product when weeds are actively growing but before weeds canopy.

A waiting interval of 7 days is required before harvest.

Do not use pre-harvest treated Barley or Wheat for seed unless a germination test is performed on the seed with an acceptable result of 95% or more germination.

For control of additional broadleaf weeds or grasses, this product may be tank-mixed with other herbicides, such as 2,4-D that are labeled for pre-harvest uses in Barley or Wheat. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-

mixture.

### SUGARCANE

Observe all precautions and restrictions. Read and follow mixing, application and cleaning instructions on this label.

Consult your local or State authorities for possible application restrictions, especially concerning aerial application and advice concerning special local use situations.

#### Use Restriction:

- Pre-harvest Interval (PHI) is 87 days.

#### Weeds Controlled

**Agripro Dicamba DMA**, when applied at specified rates, will control many annual, biennial, and perennial broadleaf weeds commonly found in Sugarcane (refer to the **WEED LIST** section).

#### Rates and Timings

Application of **Agripro Dicamba DMA** may be made any time after weeds have emerged and are actively growing but before the close-in stage of Sugarcane. Application rates and timings of this product are given below. Use the higher level of the listed rate ranges when treating dense vegetative growth.

Weed Stage and Type	Amount of Agripro Dicamba DMA	Broadcast Rate Per Treated Acre (lbs. a.e.)
Annual		
Small, actively growing	0.5 to 1 pt.	0.25 to 0.5
Established weed growth	1 to 1.5 pts.	0.5 to 0.75
Biennial	1 to 2 pts.	0.5 to 1
Perennial	2 pts.	1

When possible, direct the spray beneath the Sugarcane canopy in order to minimize the likelihood of crop injury. The use of directed sprays will also aid in maximizing spray coverage of weed foliage.

#### Use Restrictions:

- Do not exceed a total of 4 pints of this product (2 lbs. a.e.) per acre per year.
- Do not exceed a total of 2 applications of this product per acre per year.

Do not apply more than 1 lb. ae (32 oz. Agripro Dicamba DMA) per acre per application.

#### Tank-Mix Treatments

**Agripro Dicamba DMA** may be tank-mixed with one or more of, but not limited to, the following herbicides for control of grasses or additional broadleaf weeds. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

Active Ingredient
Ametryn
Asulam
Atrazine
2,4-D*
*Application of <b>Agripro Dicamba DMA</b> plus 2,4-D tank-mix at the higher listed rate range may result in crop injury.

### PASTURE, HAY, RANGELAND AND FARMSTEAD (NON-CROPLAND)

**Agripro Dicamba DMA** can be used on Pasture, Hay, Rangeland, General farmstead (Non-cropland [including fence rows and non-irrigation ditch banks]) for broadleaf weed and brush control. This product may also be applied to non-cropland areas for the control of broadleaf weeds in Noxious Weed Control Programs, districts or areas including broadcast or spot treatment of roadsides and highways, utilities, rail road and pipeline rights-of-way. Noxious weeds must be recognized at the State level but programs may be administered at State, County, or other levels.

Observe all precautions and restrictions. Read and follow mixing, application and cleaning instructions on this label.

Uses of this product described in this section also pertain to Small grains (such as Barley, Forage, Oats, Rye, Sorghum, Sudangrass or Wheat) grown for pasture use only.

Bentgrass, Buffalograss, Carpetgrass and St. Augustinegrass may be injured at rates exceeding 1 pint of this product (0.5 lb. a.e.) per treated acre. Usually, Colonial bentgrasses are more tolerant than creeping types. Velvetgrasses are most easily injured. Treatments will injure or kill Alfalfa, Clovers, Lespedeza, Wild winter peas, Vetch, and other Legumes.

#### Use Restrictions:

- Pre-harvest Interval (PHI): Grass forage - 0 days; Grass hay - 7 days.
- Do not remove animals from treated area for slaughter prior to 30 days after last application of this product.
- Do not apply more than 1 lb. ae (32 oz. Agripro Dicamba DMA) per acre per application.
- Do not apply more than twice per year.
- Newly seeded areas, including Small grains grown for pasture may be severely injured if rates of this product greater than 1 pint per acre are applied. Established grass crops growing under stress can exhibit various injury symptoms that may be more pronounced if herbicides are applied.

**Note:** There is no waiting period between treatment and grazing for non-lactating animals.

#### Timing Restrictions For Lactating Dairy Animals Following Treatment

Agripro Dicamba DMA (Rate/Treated Acre)	Days Before Grazing	Days Before Hay Harvest
Up to 1 pt. (0.5 lb. a.e.)	7	37
Up to 2 pts. (1 lb. a.e.)	21	51

#### Mixing and Application

**Agripro Dicamba DMA** can be applied using water, oil in water emulsions (including invert systems) or sprayable fluid fertilizer as a carrier. If compatibility is not known, perform a compatibility test (see **Compatibility Test** section) prior to tank-mixing.

To prepare oil-in-water emulsions, half-fill spray tank with water, then add the appropriate amount of emulsifier. With continuous agitation, slowly add the herbicide and then the oil (such as diesel or fuel oil) or a pre-mix of oil plus additional emulsifier to spray tank. Complete filling of spray tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers.

This product may be applied broadcast using either ground or aerial application equipment. When using ground equipment, apply 3 to 600 gallons of diluted spray per treated acre. Volume of spray applied will depend on the height, density and type of weeds or brush being treated and on the type of equipment being used. When using aerial equipment, apply 1 to 40 gallons of diluted spray per treated acre in a water-based carrier.

This product may be applied to individual clumps or small areas (Spot Treatment) of undesirable vegetation using handgun or similar types of application equipment. Apply diluted spray to allow complete wetting (up to runoff) of foliage and stems.

Herbicide adjuvants or other spray additives (emulsifiers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Spray additives must be agriculturally approved when used in pasture applications. If spray additives are used, read and follow all use directions and precautions on product label.

#### Weeds Controlled

**Agripro Dicamba DMA**, when applied at specified rates, will give control of many annual, biennial, and perennial broadleaf weeds, and many woody brush and vine species commonly found in pasture, hay, rangeland, and farmstead (non-cropland) areas (refer to the **WEED LIST** section). Noted (\*) perennial weeds may be controlled with lower rates of either this product or this product plus 2,4-D. See **Rates and Timings** below.

#### Rates and Timings

Application rates and timing of **Agripro Dicamba DMA** are given below. Use the higher level of the listed rates ranges when treating dense or tall vegetative growth.

Weed Stage and Type	Amount of Agripro Dicamba DMA	Broadcast Rate Per Treated Acre (Lbs. a.e.)*
Annuals Small, actively growing Established weed growth	0.5 to 1 pt. 1 to 1.5 pts.	0.25 to 0.5 0.5 to 0.75
Biennials <sup>1</sup> Rosette diameter Less than 3 inches 3 inches or more Bolting	0.5 to 1 pt. 1 to 2 pts. 2 pts.	0.25 to 0.5 0.5 to 1 1
Perennials Suppression or Top growth control Noted (*) perennials Other perennials	1 to 2 pts. 2 pts. 2 pts.	0.5 to 1 1 1
Woody Brush and Vines Top growth suppression Top growth control <sup>2</sup> Stems and Stem suppression	1 to 2 pts. 2 pts. 2 pts.	0.5 to 1 1 1

<sup>1</sup>For best performance, make application when biennial weeds are in the rosette stage.

<sup>2</sup>Species noted in **WEED LIST** will require tank-mixtures for adequate control.

\*Do not apply more than 1 lb. a.e./Ac. of this product per application. Retreatments may be made as needed. However, do not exceed a total of 4 pints of this product (2 lbs. a.e.) per treated acre per year.

**Tank-Mix Treatments**

**Agripro Dicamba DMA** may be tank-mixed with one or more of, but not limited to the following herbicides for control of grasses, additional broadleaf weeds and woody brush and vines.

Active Ingredient
2,4-D
Glyphosate
Metsulfuron-methyl
Paraquat
Picloram
Triclopyr

Due to the variations that may occur between formulated products and specific use ingredients (e.g., water supplies), if compatibility is not known, perform a compatibility test (see **Compatibility Test** section) prior to actual tank-mixing.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture

**Cut Surface Tree Treatments**

**Agripro Dicamba DMA** may be applied as a cut surface treatment to control unwanted trees and to prevent sprouts of cut trees. Use a mix of 1 part of this product with 1 to 3 parts water for application.

Use the lower dilution when treating difficult-to-control species.

**Frill or Girdle Treatments**

Make a continuous cut or series of overlapping cuts using an axe to girdle tree trunk. Spray or paint cut surface with this product/water mix.

**Stump Treatments**

Spray or paint freshly cut surface with the water mix. The area adjacent to the bark should be thoroughly wet. For more rapid foliar effects, 2,4-D may be added to this product/water mix.

**Dormant Applications For Control of Multiflora Rose**

**Agripro Dicamba DMA** can be applied when plants are dormant as an undiluted spot concentrate directly to the soil or as a Lo-Oil basal bark treatment using an oil-water emulsion solution.

**Spot Concentrate Application:** Apply **Agripro Dicamba DMA** directly to the soil as close as possible to the root crown but within 6 to 8 inches of the crown. On sloping terrain, apply to the uphill side of the crown. The use rate of this product is dependent on the canopy diameter of the Multiflora Rose. Examples: Use this product at 0.25, 1 or 2.25 fluid ounces, respectively, for 5, 10 or 15 feet canopy diameters.

**Use Restrictions:** Do not make application when snow or water prevents applying this product directly to the soil. Do not exceed a total of 4 pints of this product per acre per year.

**Lo-Oil Basal Application:** Apply **Agripro Dicamba DMA** to the basal stem region from the ground up to a height of 12 to 18 inches. Spray until runoff, with special emphasis on covering the root crown. For best results, apply when plants are dormant. Refer to the **Mixing and Application** section above for instructions on preparing oil-in-water emulsions. Example for making approximately 2 gallons of a LO-OIL spray solution mixture: Combine 1.5 gallons of water plus 1 ounce of emulsifier plus 1 pint of this product plus 2.5 pints of No. 2 diesel fuel. Adjust amounts of materials used proportionately to the amount of final spray solution desired.

**Use Restrictions:** Do not make application after budbreak or when plants are showing signs of active growth. Do not make application when snow or water prevents applying this product to the ground line. Do not exceed 8 gallons of spray solution mix applied per acre per year.

**CONSERVATION RESERVE PROGRAMS (CRP) ACRE**

**Agripro Dicamba DMA** can be used on both newly-seeded and established grasses grown in Conservation Reserve or Federal Set-Aside Programs.

Observe all precautions and restrictions. Read and follow mixing, application and cleaning instructions on this label.

**Use Restriction:**

- Treatment of this product will cause injury or may kill Alfalfa, Clovers, Lespedeza, Vetch, Wild winter peas, and other legumes.

- Agriculturally approved surfactants may be added to the spray mixture to improve post-emergence weed control, particularly in dry growing conditions.
- Do not use adjuvants containing penetrants such as petroleum-based oils after grass emergence on newly seeded grasses.

### Newly Seeded Areas

**Agripro Dicamba DMA** may be applied either pre-plant or post-emergence to newly seeded grasses or Small grains such as Barley, Oats, Rye, Sudangrass, Wheat or other Grain species grown as a cover crop. Post-emergence applications may be made after seedling grasses exceed the 3-leaf stage. Rates of this product greater than 1 pint per treated acre may severely injure newly seeded grasses.

**Pre-Plant Applications:** Injury to new seedlings may occur if intervals between application and grass planting is less than 45 days per pint of this product per treated acre West of the Mississippi River or 20 days per pint East of the Mississippi River.

### Established Grass Stands

Established grass stands are perennial grasses planted one or more seasons prior to treatment. Certain species, such as Bentgrass, Buffalograss, Carpetgrass, Smooth brome or St. Augustinegrass may be injured when treated with this product at rates exceeding 1 pint per treated acre.

### Weeds Controlled

This product when applied at specified rates will control many annual and biennial weeds and provide control and suppression of many perennial weeds (see **WEED LIST** section).

### Rates and Timings

Application rates and timing of treatment of this product are given below. Use the higher rate of the rate range when vegetation is either dense or tall or when weeds are growing under stressed conditions such as drought or cool temperature.

Weed Type* and Stage	Broadcast Rate Per Treated Acre	
	Amount of This Product (Pts.)	Equivalent Pounds of Dicamba
Annual Small, actively growing Established weed growth	0.25 to 1 1	0.125 to 0.5 0.5
Biennials** Rosette diameter a) Less than 3 inches b) 3 inches or more c) Bolting biennial	0.5 to 1 1 to 2 2	0.25 to 0.5 0.5 to 1 1
Perennials** Suppression/Control	2	1
*For best results, treat biennial weeds with this product when they are in the rosette stage of growth. Retreatment may be made as needed. However, do not exceed a total of 4 pts. (2 lbs. a.e.) per treated acre per year. **Biennial and perennial weeds will require follow-up (sequential) treatments for seedling control and escapes.		

### TANK-MIX TREATMENTS

To control grasses and additional broadleaf weeds, this product may be tank-mixed with other herbicides registered for use in Conservation Reserve Programs (CRP) such as 2,4-D, Glyphosate, Metsulfuron, Paraquat, and others.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

### ASPARAGUS (CA, OR, WA Only)

Observe all precautions and restrictions. Read and follow mixing, application and cleaning instructions on this label.

If spray contacts emerged spears, crooking (twisting) of some spears may result. If such crooking occurs, discard affected spears.

### Use Restrictions:

- Do not harvest prior to 24 hours after treatment.
- Do not use in the Coachella Valley of California.
- Maximum 2 applications per year.
- Do not exceed a total of 1 pint of this product (0.5 lb. a.e.) per treated acre per crop year.

### RATES AND TIMINGS

Apply this product to emerged and actively growing weeds in 40 to 60 gallons of diluted spray per treated acre immediately after cutting the field, but at least 24 hours before the next cutting.

Weeds	Rate Per Acre
Mustard, Black Pigweed, Redroot (Carelessweed)	0.5 to 1 pt. (0.25 to 0.5 lb. a.e.)

Sowthistle, Annual Thistle, Canada* Thistle, Russian	
Bindweed, Field* Chickweed, Common Goosefoot, Nettleleaf Radish, Wild Thistle, Milk	1 pt. (0.5 lb. a.e.)
*This product may be applied in a tank-mixture with either 2,4-D or Glyphosate (e.g., Imitator, Roundup) for improved control of Canada thistle and Field bindweed. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.	

### TURF AND LAWNS — FOR USE IN FARMSTEAD (NON-CROPLAND) AND SOD FARMS

Observe all precautions and restrictions. Read and follow mixing, application and cleaning instructions on this label.

To avoid injury to newly seeded grasses, delay application of this product until after the second mowing. Furthermore, application rates in excess of 1 pint (0.5 lb. a.e.) per treated acre may cause noticeable stunting or discoloration of sensitive grass species such as Bentgrass, Buffalograss, Carpetgrass and St. Augustinegrass.

#### Use Restriction:

- Do not make repeat applications in these areas for 30 days and until previous applications of this product have been activated in the soil by rain or irrigation.
- In areas where roots of sensitive plants extend, do not apply in excess of 0.25 pint of this product (0.125 lb. a.e.) per treated acre on coarse-textured (*Sandy* type) soils or in excess of 0.5 pint (0.25 lb. a.e.) per treated acre on fine-textured (*Clay* type) soils.

#### Weeds Controlled

This product when applied at specified rates will give control of many annual, biennial and noted (\*) perennial broadleaf weeds commonly found in Turf. This product will also give growth suppression of many other listed perennial broadleaf weeds and woody brush and vine species (refer to the **WEED LIST** section).

#### Mixing and Application

Apply 30 to 200 gallons of diluted spray per treated acre (6 pts. to 4.25 gals. per 1,000 sq. ft.) depending on density or height of weeds treated and on the type of equipment used.

#### Rates and Timings

Use the higher level of listed rate ranges when treating dense vegetative growth.

Weed Stage and Type	This Product		
	Pints Per Treated Acre	Pounds A.E. Per Treated Acre	Teaspoons Per 1,000 sq. ft.
Annuals Small, actively growing Established weed growth	0.5 to 1 1 to 1.5	0.25 to 0.5 0.5 to 0.75	1 to 2.25 2.25 to 3.25
Biennials, Rosette diameter Less than 3 inches 3 inches or more	0.5 to 1 1 to 2	0.25 to 0.5 0.5 to 1	1 to 2.25 2.25 to 4.5
Perennials and Woody brush and Vines	1 to 2	0.5 to 1	2.25 to 4.5

For best performance, apply when weeds have emerged and are actively growing.

#### Use Restrictions:

- Do not exceed a total of 2 pints of this product (1 lb. a.e.) per treated acre per application.
- Do not exceed a total of 2 applications of this product per acre year.

#### Tank-Mix Treatments

Tank-mix treatments of this product may be made with 2,4-D, MCPA, MCPP or Bromoxynil for control of additional weeds listed on the tank-mix product.

Use the higher level of the listed rate ranges when treating established weeds. Repeat treatments may be made as needed. However, do not exceed 2 pints of this product (1 lb. a.e.) per treated acre during the year.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.



**GRASS SEED CROPS — GRASSES GROWN FOR SEED**  
**(Such as Bermudagrass, Bluegrass, Fescue and Ryegrass)**

Observe all precautions and restrictions. Read and follow mixing, application and cleaning instructions on this label.

**Use Restrictions:**

- Do not use on Bentgrass unless possible crop injury can be tolerated.
- Refer to the **PASTURE, HAY, RANGELAND AND FARMSTEAD (NON-CROPLAND)** section for grazing and feeding restrictions. The pre-harvest interval (PHI) for Grass forage is 0 days. The PHI for Grass hay is 7 days.

**WEEDS CONTROLLED**

This product will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds plus additional weeds, this product may be applied in tank-mixture with other herbicides.

Alfalfa*	Hemlock, Poison
Bedstraw, Catchweed	Knapweed, Russian*
Bindweed, Field	Knawel
Buttercup, Corn	Knotweed, Prostrate
Buttercup, Creeping	Kochia
Buttercup, Western Field	Ladysthumb
Catchfly, Nightflowering	Lambsquarter, Common
Chamomile, Corn	Lettuce, Prickly
Chickweed, Common	Mayweed (Dogfennel)
Chickweed, Mouseear	Ragwort, Tansy
Clover	Sorrel, Red (Sheep Sorrel)
Cockle, White	Sowthistle, Annual
Dock, Broadleaf	Starwort, Little
Dock, Curly	Thistle, Canada*
*Top growth only.	

**RATES AND TIMINGS**

Apply 0.5 to 1 pint of this product per treated acre on seedling grass after the crop reaches the 3- to 5-leaf stage. Apply up to 2 pints of this product on well-established perennial grass. Apply before grass seed crop begins to joint. For best performance, make applications when weeds are in the 2- to 4-leaf stage and rosettes are less than 2 inches across. Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth.

**TANK-MIX TREATMENTS**

For control of grasses or additional broadleaf weeds, this product may be tank-mixed with broadleaf herbicides registered for use in grass seed production.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

**Broadcast Rate Per Treated Acre**

Apply 0.5 to 2 pints of this product with:

Active Ingredient
Bromoxynil
Clopyralid
Clopyralid + 2,4-D
2,4-D
Diuron
MCPA

**ANNUAL GRASS CONTROL**

For suppression of annual grass weeds such as:

Brome, Downy Cheatgrass)	Brome, Ripgut	Fescue, Rattail	Windgrass
--------------------------	---------------	-----------------	-----------

Apply up to 2 pints of this product per treated acre in the Fall or late Summer after harvest and burning of established grass seed crops. Apply immediately following the first irrigation when the soil is moist and before weeds have more than 2 leaves.

**PRE-PLANT DIRECTIONS (POST-HARVEST/FALLOW/CROP STUBBLE/SET-ASIDE) FOR BROADLEAF WEED CONTROL BEFORE CORN, SORGHUM, SOYBEANS AND WHEAT**

Observe all precautions and restrictions. Read and follow mixing, application and cleaning instructions on this label.



**WEEDS CONTROLLED**

This product may be applied alone or in tank-mix combinations with other herbicides registered for this use.

This product can be applied either post-harvest in the Fall, Spring or Summer, during the fallow period or to crop stubble/set-aside acres.

This product when applied at the specified rates will control many annual broadleaf weeds (see the **Weeds Controlled** under **SMALL GRAINS (BARLEY, OATS AND WHEAT) NOT UNDERSEEDED TO LEGUMES** section). In addition, this product will control or suppress the following biennial and perennial broadleaf weeds:

Alfalfa*	Nightshade, Silver
Artichoke, Jerusalem	Redvine
Bindweed, Field	Smartweed, Swamp
Bindweed, Hedge	Sowthistle, Perennial*
Blueweed, Texas	Spurge, Leafy
Bursage (Bur ragweed, Povertyweed, Lakeweed)*	Thistle, Bull
Dandelion, Common*	Thistle, Canada**
Dock, Curly*	Thistle, Milk
Dogbane, Hemp	Thistle, Musk
Garlic, Wild**	Thistle, Plumeless
Horsenettle, Carolina	Thistle, Scotch
Knapweed, Diffuse	Trumpet creeper (Buckvine)
Knapweed, Spotted	

\*These perennials may be controlled using this product at rates lower than those specified for other listed perennial weeds. See **RATES AND TIMINGS** in this section.

\*\*See the **SPECIAL TANK-MIX TREATMENTS** under this section for specific control program for these weeds.

**RATES AND TIMINGS**

Apply this product as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (post-harvest) and before a killing frost or in the fallow cropland or crop stubble the following Spring or Summer.

Agriculturally approved spray additives, such as surfactants or oils may be used to enhance spray coverage and the herbicide's penetration of weed foliage. See **CROPPING RESTRICTIONS** section for specified interval between application and planting to prevent crop injury.

For best performance, make application when annual weeds are less than 6 inches tall, when biennial weeds are in the rosette stage and to perennial weed regrowth in late Summer or Fall following a mowing or tillage treatment. Most effective control of upright perennial broadleaf weeds, such as Canada thistle and Jerusalem artichoke occurs if application is made when the majority of weeds, such as Field bindweed and Hedge bindweed are in or beyond the full bloom stage. Avoid disturbing treated areas following application. Treatments may not kill weeds which develop from seed or underground plant parts, such as rhizomes or bulblets after the effective period for this product.

For seedling control, a follow-up program or other cultural practices could be instituted. For Small grain in-crop uses of this product, see the **Rates and Timings** under **SMALL GRAINS (BARLEY, OATS AND WHEAT) NOT UNDERSEEDED TO LEGUMES** section for details.

Weed Type	Amount of This Product per Treated Acre (Pts.)
Annuals	0.5 to 1
Biennials	1 to 2
Perennials	
Suppression of perennials	1 to 2
Noted (*) perennials	2
Other perennials	2

**Use Restrictions:**

- Do not exceed a total of 4 pints of this product (2 lbs. a.e.) per treated acre per year. Do not apply more than 1 lb. ae (32 oz. Agripro Dicamba DMA) per acre per application.
- Do not apply a total of 2 applications of this product per acre per year.

**Tank-Mix Treatments**

This product may be tank-mixed with one or more of, but not limited to the following herbicides for control of grasses or additional broadleaf weeds.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

**Broadcast Rate of this Product per Treated Acre for Annual Weed Control**

Apply 0.25 to 1 pint of this product with:

Active Ingredient
Atrazine*
Chlorsulfuron + Metsulfuron-methyl**
2,4-D
Glyphosate
Glyphosate + 2,4-D
Glyphosate + Dicamba
Metribuzin*
Metsulfuron-methyl**
Paraquat
Pronamide*
Triasulfuron**
*Tank-mixes of this product with these products may be subject to special restrictions. See the product label of the tank-mix partner for intended use rates, restrictions, and other precautions.
**When tank-mixing with Sulfonylurea herbicides, refer to the product label for rates and restrictions. Use a surfactant of at least 80% active ingredient at the rate of 2 to 4 pts./100 gals. of spray or not more than 0.25 to 0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank-mix and/or when treating more mature weeds or dense vegetative growth. Sulfonylurea-resistant weeds may not be controlled by tank-mixes of this product and a Sulfonylurea. Refer to this product's tank-mix section for alternative tank-mixes.

### Broadcast Rate of this Product per Treated Acre for Biennial and Perennial Weed Control

Apply 1 to 2 pints of this product with:

Active Ingredient
Clopyralid + 2,4-D
2,4-D
Glyphosate
Glyphosate + 2,4-D
Picloram

### SPECIAL TANK-MIX TREATMENTS

For suppression of perennial weeds, apply this product with Glyphosate.

For Wild garlic control, apply 1 pint of this product with labeled rate of 2,4-D LV Ester (4 lbs./gal.) per treated acre. Apply when Wild garlic is 4 to 8 inches tall.

For Canada thistle control, use this product alone, this product plus Clopyralid + 2,4-D or this product plus Glyphosate tank-mix treatments.

Application may be made during fallow periods for control of Bulbous bluegrass, Common rye, Downy brome, Jointed goatgrass, Volunteer barley and Volunteer wheat when they are actively growing. Use 1 pint of this product with labeled rate of pronamide. Fall seeded Wheat may be planted 9 months or more after application. For best performance, make application between mid-October and mid-December prior to soil freeze up.

During fallow periods, apply this product plus Glyphosate + 2,4-D or Glyphosate + Dicamba to give improved control of Canada thistle, Field bindweed, Kochia, Prickly lettuce and Wild buckwheat. Use 0.125 to 0.25 pint of this product plus labeled rate of Glyphosate + 2,4-D or Glyphosate + Dicamba for annual weed control or for suppression of perennial weeds.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

### CROPPING RESTRICTIONS

Corn, Sorghum and Soybeans may be planted in the Spring following applications made during the previous year. If less than 1 inch of rainfall occurs between application and first killing frost, cultivate treated areas to allow herbicide to come in contact with moist soil. Cultivation may take place before or immediately after ground thaw.

Soybean injury may occur if the interval between application and planting is less than specified. In areas with greater than 30 inches of rainfall, delay planting for 30 days per pint of this product per treated acre. In areas with less than 30 inches of rainfall, delay planting for 45 days per pint of this product per treated acre. Exclude days when ground is frozen.

Wheat may be planted in the Fall or Spring following applications. Also, spot application may be made any time prior to crop emergence if crop injury can be tolerated in treated areas. Wheat injury may occur if the interval between application and planting is less than specified.

**East of the Mississippi River:** The interval is 20 days per pint of this product per treated acre, or 1.25 days per 1 fluid ounce. Moisture is essential for degradation of this product. Exclude days when ground is frozen.

**West of the Mississippi River:** The interval is 45 days per pint of this product per treated acre or 3 days per 1 fluid ounce. Moisture is essential for degradation of this product. Exclude days when ground is frozen.

Following a normal harvest of Barley, Oats or Wheat, any rotational crop may be planted. **Use Restriction:** If the interval before harvest is shortened, such as when cover crops will be plowed under, do not follow up with the planting of a sensitive crop.

#### **SUPPRESSION / CONTROL OF PERENNIAL BROADLEAF WEEDS IN CROPLAND (SPOT APPLICATION ONLY) (ID, MT, NV, OR, UT, WA Only)**

Observe all precautions and restrictions. Read and follow mixing, application and cleaning instructions on this label.

##### **Use Restrictions:**

- Do not treat sub-irrigated cropland or areas where the soil remains saturated with water throughout the year.
- Make only one application of this product per year.

##### **WEEDS CONTROLLED**

This product, when applied at specified rates, will control many broadleaf weeds including:

Bindweed, Field Dock, Broadleaf (Bitterdock) Dock, Curly	Knapweed, Black Knapweed, Russian Ragwort, Tansy	Spurge, Leafy Thistle, Canada
--	--	----------------------------------

##### **Rates and Timings**

This product may be applied at any time following a crop harvest to stubble fallow or other cropland. Apply when weeds are actively growing and prior to a killing frost.

Apply 2 pints (1 lb. a.e.) of this product per treated acre. Application may be made up to one month prior to the planting of Wheat.

Do not use unless injury to Wheat or rotated Barley will be acceptable.

Annual or perennial grass crops, Barley, Corn, Oats, Sorghum (Milo) may be planted into treated areas one year after application. Crops grown for seed (other than perennial grass seed) should not be planted into treated areas until three years after application.

**Use Restriction:** Do not plant broadleaf crops including Alfalfa, Beans, Peas, Potatoes or Sugar beets into treated areas until two years after application.

In most cases, treatments will not kill perennial seedlings which germinate from seed one or two years after treatment. Once the effect of the chemical has been lost, a follow-up program for seedling control or other cultural practices should be instituted.

##### **Wiper Application Uses**

Observe all precautions and restrictions. Read and follow mixing, application and cleaning instructions on this label.

This product may be applied through wiper application equipment to control or suppress actively growing broadleaf weeds, brush, and vines. Use a solution containing 1 part of this product to 1 part water. **Use Restriction:** Do not contact desirable vegetation with herbicide solution.

Make wiper application to crops (including pastures) and non-cropland areas described in this label with the exception of Grain sorghum (Milo).

#### **RIGHTS-OF-WAY, UTILITY AND INDUSTRIAL AREAS, AND FENCEROWS**

Observe all precautions and restrictions. Read and follow the Mixing and Application section.

This product may be used for use on non-crop land areas including rights-of-way (including roadways, rest areas, utility, railroad, highway, pipelines, and rights-of-way that run through pasture and rangeland); utility facilities (including substations, pipelines, tank farms, pumping stations, parking and storage areas, fencerows, and non-irrigated ditch banks); brush control for forest site preparation or maintenance.

##### **RIGHTS-OF-WAY**

This product can be used to control many broadleaf weeds on rights-of-way. This use includes applications to roadsides, roadways, and highways; areas along utilities including cable and powerlines; railroad track and embankment; highways, highway medians, bridge abutments, pipelines, and rights-of-way that run through pasture and rangeland. Use controlled application techniques that minimize the risk of off-target movement.

##### **UTILITY AND INDUSTRIAL AREAS**

This product can be used to control many broadleaf weeds and brush in non-crop areas on or surrounding substations, pipelines, tank farms, pump stations, production facilities, and bare ground situations. It may also be used on parking and storage areas (see **Best Stewardship Practices** to avoid direct runoff from impervious surfaces).

##### **FENCEROWS**

This product can be used to control many broadleaf weeds and brush in fencerows.

**Mixing and Application**

Refer to the **Mixing and Application** under **PASTURE, HAY, RANGELAND AND FARMSTEAD (NON-CROPLAND)** section for directions.

**Weeds and Brush Controlled**

This product applied at specified rates will give control of many annual, biennial, and perennial broadleaf weeds, and many woody brush and vine species commonly found in non-crop land areas (refer to **WEED LIST** section). Noted (\*) perennial weeds may be controlled with lower rates of either this product or this product in tank -mix combinations. See **Rates And Timings** below.

**Rates and Timings**

Application rates and timings of this product are given below. Use the higher level of listed rate ranges when treating dense or tall vegetative growth.

Weed Stage and Type	Amount of This Product	Gallons of Spray Mixture per Acre <sup>1</sup>	Spray Concentration for Low Volume Application <sup>2</sup> (% vol/vol)
Annuals Small, actively growing Established weed growth	0.5 to 1 pt. 1 to 1.5 pts.	25 to 50 50 to 75	3 3
Biennials <sup>3</sup> Rosette diameter Less than 3 inches 3 inches or more Bolting	0.5 to 1 pt. 1 to 2 pts. 2 pts.	25 to 50 50 to 100 100 to 150	3 to 4 3 to 4 3 to 4
Perennials Suppression or Top growth control Noted (*) perennials Other perennials	0.5 to 1 pt. 2 pts. 2 pts.	50 to 100 100 to 200 200	4 4 5
Woody Brush and Vines <sup>4</sup> Top growth suppression Top growth stems and Roots	0.5 to 2 pts. 2 pts.	50 to 200 200	5 5
<sup>1</sup> Assuming typical application rate of 2 pts. of this product per 100 gals. <sup>2</sup> Low volume rates must not exceed 4 pts. of this product per acre per year. (5% vol/vol = 10 gals. maximum solution per acre per year). <sup>3</sup> For best performance, apply when biennial weeds are in the rosette stage. <sup>4</sup> Tank-mixes may be required for optimal control. Refer to <b>WEED LIST</b> section.			

**Use Restriction:**

- Do not exceed a total of 4 pints (2 lbs. a.e.) of this product per treated acre per year. Do not apply more than 1 lb. ae (32 oz. Agripro Dicamba DMA) per acre per application.
- Do not apply more than twice per year.

**Tank-Mix Treatments**

This product may be tank-mixed with one or more of, but not limited to the following herbicides for additional weed control in Rights-of-way, Utility and Industrial Areas and Fencerows.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

Active Ingredient
Bromacil
Chlorsulfuron
Clpyralid
2,4-D
Diquat
Diuron
Fosamine ammonium
Glufosinate
Glyphosate
Hexazinone
Imazapyr
Imazemethapyr
Metsulfuron-methyl
Norflurazon
Pendimethalin
Prodiamine
Simazine
Sulfometuron-methyl
Tebuthiuron

## Triclopyr

Due to variations that may occur between formulated products and specific use ingredients (e.g., water supplies), if compatibility is not known, a compatibility test (see **Compatibility Test** section) should be carried out prior to actual tank-mixing.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container. Keep container tightly closed. Avoid cross-contamination with other pesticides.

**PESTICIDE DISPOSAL:** To avoid waste, use all materials in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often, such programs are run by State or local governments or by industry).

**CONTAINER HANDLING:**

**Nonrefillable Container (rigid material; less than 5 gallons):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Container (rigid material; 5 gallons up to < 250 gallons):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Refillable Container (≥ 250 gallons & Bulk):** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Agripro USA Limited or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Agripro USA Limited and Seller harmless for any claims relating to such factors.

Agripro USA Limited warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Agripro USA Limited and to the extent consistent with applicable law, Buyer and User assume the risk of any such use. To the extent consistent with applicable law, AGRIPRO USA LIMITED MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Agripro USA Limited nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF AGRIPRO USA LIMITED AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF AGRIPRO USA LIMITED OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Agripro USA Limited and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Agripro USA Limited.

[All trademarks are the property of their respective owners.]