



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505T)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

103591-13

Date of Issuance:

11/13/25

NOTICE OF PESTICIDE:

☒ Registration  
☐ Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Prism Diquat

Name and Address of Registrant (include ZIP Code):

Rainbow Agrosiences, LLC  
c/o Pyxis Regulatory Consulting, Inc.  
535 E Plainfield Road, Suite E  
Willowbrook, IL 60527

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).


Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

*Continues page 2*

Signature of Approving Official:

  
for

Kevin Ulrich (for)  
Nathan Mellor, Chief  
Fungicide Herbicide Branch, Registration Division (7505T)

Date:

11/13/25

2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 103591-13."
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 09/30/2024

If you have any questions, please contact Hector Escobar at 202-566-1371 or at [escobar.hector@epa.gov](mailto:escobar.hector@epa.gov).

Enclosure: Stamped Label

{Note to Reviewer: Text in brackets [ ] denotes optional text. In instances where a word or phrase has multiple optional text options, at least one will be used to ensure that the entire statement is clear and understandable. Text in braces { } denotes explanatory text that will not be included on the final printed label.}

**{BOOKLET FRONT PANEL LANGUAGE}**

DIQUAT DIBROMIDE	GROUP	22	HERBICIDE
------------------	-------	----	-----------

**TO PREVENT ACCIDENTAL POISONING, NEVER PUT INTO FOOD, DRINK OR  
OTHER CONTAINERS AND USE STRICTLY IN ACCORDANCE WITH ENTIRE LABEL**

## Prism Diquat

Contains diquat dibromide, the active ingredient used in [Reglone®].

<b>ACTIVE INGREDIENT:</b>	<b>(% by weight)</b>
Diquat dibromide [6,7-dihydrodipyrido (1,2-a:2',1'-c) pyrazinediium dibromide] .....	37.3%
<b>OTHER INGREDIENTS:</b> .....	62.7%
<b>TOTAL</b> .....	100.0%

**Prism Diquat** is formulated as a soluble liquid containing 2 lb diquat cation per gal as 3.73 lb salt per gal.

### KEEP OUT OF REACH OF CHILDREN CAUTION

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If  
you DO NOT understand the label, find someone to explain it to you in detail.)**


[See] [below] [inside] [label] [booklet] [for] [additional] [First Aid,] [and] [Precautionary Statements] [and]  
[Directions for Use] [including] [storage and disposal instructions] [.]

**[Prism Diquat]** is not manufactured, or distributed by Syngenta Crop Protection, LLC, seller of Reglone® [Dessicant].

**EPA Reg. No.: 103591-xx**

**EPA Est. No.:**

Manufactured for:  
RAINBOW AGROSCIENCES, LLC  
535 E Plainfield Road, Suite E  
Willowbrook, IL 60527



## {LANGUAGE INSIDE BOOKLET}

<b>FIRST AID</b>	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Immediately call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>NOTE TO PHYSICIAN</b>	
<p>To be effective, treatment for diquat poisoning must begin IMMEDIATELY. Treatment consists of binding diquat in the gut with suspensions of activated charcoal or bentonite clay, administration of cathartics to enhance elimination, and removal of diquat from the blood by charcoal hemoperfusion or continuous hemodialysis.</p> <p>If in eyes, treat symptomatically. Symptoms may develop gradually. Severe damage may be caused by apparently trivial contact and healing may be delayed. Medical supervision should be continued until healing is complete.</p>	
<b>HOT LINE NUMBER</b>	
<p>Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. You may also contact the Poison Control Center at 1-800-222-1222 for emergency medical treatment information. For non-emergency information concerning this product, call the National Pesticide Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8:00AM to 12:00 PM Pacific Time, or at <a href="https://npic.orst.edu">https://npic.orst.edu</a>.</p>	

**For Chemical Emergency:**  
**Spill, Leak, Fire, Exposure, or Accident,**  
**Call CHEMTREC Day or Night**  
**Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

### **PRECAUTIONARY STATEMENTS** **Hazards to Humans and Domestic Animals** **CAUTION**

Harmful if swallowed. Harmful if inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

## **Human Flagging is Prohibited**

### **Personal Protective Equipment (PPE)**

#### **Applicators, mixers, loaders, and other handlers must wear:**

- Coveralls worn over long-sleeved shirt and long pants
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading

### **Occupational Handler PPE**

In addition to the above PPE, the following users must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R, or P filter; OR a NIOSH-approved powered air-purifying respirator with a HE filter.

- Handlers supporting applications to aquatic areas for the control of submersed weeds via trailing or submersed hoses, during the time in which they are mixing and/or loading the diquat-containing material.
  - Alternatively, such mixer/loaders may use closed mixing/loading systems without a respirator. A closed system is one which removes the pesticide from its original container and transfers the pesticide product through connecting hoses, pipes and couplings that are sufficiently tight to prevent exposure of handlers to the pesticide product, except for the negligible escape associated with normal operation of the system (40 CFR § 170.607(d)).
- Mixers/loaders supporting applications to aquatic areas with mechanically-pressurized handguns AND mixers/loaders/applications using mechanically-pressurized handguns to treat aquatic areas, during mixing and loading operations ONLY.
  - Alternatively, these aquatic handlers may use closed mixing/loading systems without a respirator.
- Mixers/loaders/applicators using mechanically-pressurized handguns on industrial and commercial areas, alfalfa (seed crop only), clover (seed crop only), or non-bearing tree, vine, small fruit, vegetable crops.
- Mixers/loaders supporting groundboom applications to canola and potato.
- Applicators using groundboom equipment for all crop uses.

### **Respirator fit testing, medical qualification, and training**

Using a program that conforms to OSHA's requirements (see 29 CFR Part 1910.134), employers must verify that any handler who uses a respirator is:

- Fit-tested and fit-checked,
- Trained, and
- Examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn. A qualified medical practitioner is a physician or other licensed health care professional who will evaluate the ability of a worker to wear a respirator. The initial evaluation consists of a questionnaire that asks about medical conditions (such as a heart condition) that would be problematic for respirator use. If concerns are identified, then additional evaluations, such as a physical exam, might be necessary. The initial evaluation must be done before respirator use begins. Handlers must be reexamined by a qualified medical practitioner if their health status or respirator style or use-conditions change.

Upon request by local/state/federal/tribal enforcement personnel, employers must provide documentation demonstrating how they have complied with these requirements.

### **User Safety Requirements**

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **Engineering Controls**

Mixers and loaders supporting aerial applications are required to use closed systems. The closed system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4)). When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS. Mixers, loaders, and applicators using closed systems who meet these requirements may wear: long-sleeved shirt and long pants; protective eyewear; waterproof gloves; shoes plus socks; and a chemical-resistant apron when mixing, loading, or cleaning equipment. If handling tasks are performed from inside an enclosed cab or aircraft with enclosed cockpits that meet these requirements may wear: long-sleeved shirt, long pants, shoes, and socks for the labeling-specified PPE. All labeling-specified PPE must be immediately available for use in an emergency. All applicable requirements as specified in 40 CFR 170.240(d)(4-6) must be followed.

### **User Safety Recommendations**

#### **Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **Environmental Hazards (Terrestrial and Aquatic Uses)**

This pesticide is toxic to aquatic invertebrates. **For Terrestrial Uses, DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate. **For Aquatic Uses, DO NOT** apply directly to water except as specified on this label. Treatment of dense weed areas may result in oxygen loss from decomposition of dead weeds. This loss of oxygen may cause fish suffocation. Therefore, treat only 1/3 – 1/2 of the water body area at one time, especially if dense areas of weeds and/or algae exist and wait 14 days between treatments.

Necessary approval and/or Permits must be obtained prior to application if required. Consult the responsible State Agencies (i.e., Fish and Game agencies or Department of Natural Resources) before making applications to public waters.

### **NON-TARGET ORGANISM ADVISORY**

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

**Physical-Chemical Hazards**

**DO NOT** mix or allow coming in contact with oxidizing agents. Hazardous chemical reactions may occur.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**DO NOT** apply this product via backpack sprayer.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

**DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.**

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas or vicinity where there may be drift.

**DO NOT allow people or pets to touch treated plants until the sprays have dried.**

**For terrestrial uses, DO NOT** allow entry of maintenance workers into treated areas, or allow contact with treated vegetation wet with spray, dew, or rain, without appropriate protective clothing until spray has dried.

**For aquatic uses, DO NOT** enter treated areas while treatments are in progress.

### PRODUCT INFORMATION

**Prism Diquat** is a nonvolatile herbicide for use as a preharvest aid to desiccate certain crops in order to facilitate harvesting. **Prism Diquat** is also advised for use as a general herbicide to control weeds in noncrop areas, nonbearing crops and aquatic areas. **Prism Diquat** is a contact-type herbicide and requires actively growing green plant tissue to function. Thorough coverage of all green plant tissue is essential for effective control.

**Prism Diquat** is rapidly absorbed by green plant tissue and interacts with the photosynthetic process to produce compounds which destroy plant cells. Herbicidal activity is usually quite rapid, with effects visible in a few days.

### AGRICULTURAL USE DIRECTIONS

#### APPLICATION

Since **Prism Diquat** is a contact-type herbicide, it is essential to obtain complete coverage of the target weed or crop to achieve effective results. Improper application technique and/or application to large, stressed, or mowed weeds will generally result in unacceptable control.

Complete coverage is also essential for effective performance in harvest aid applications. See details below for additional information.

#### Nozzle Selection

The use of flat fan nozzles will result in the most effective application of **PRISM DIQUAT**. The use of nozzles other than flat fans may result in reduced performance due to inadequate coverage.

#### Spray Volume

Follow specified minimum spray volumes listed for each use of **PRISM DIQUAT**. These are **minimum** volumes only, and spray volumes must be increased as necessary to obtain complete coverage of the target weed or plant without runoff from the foliage. When spraying less than 20 gal of spray carrier per acre, target weeds must not exceed 6 inches in height.

### SPRAY ADJUVANTS

#### Always Add One of the Following:

##### Nonionic Surfactant (NIS)

Add a NIS containing 75% or greater surface active agent at 0.06–0.5% v/v (1/2–4 pt per 100 gal) of the finished spray volume.



## Other Adjuvants

Adjuvants other than NIS may be used providing the product meets the following criteria:

- Contains only EPA exempt ingredients.
- Is compatible in mixture. Compatibility may be established through a jar test.
- Is supported locally for use with **PRISM DIQUAT** through proven field trials and through university and extension directions.

When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification program is advised.

## RATES

Follow specified rates listed with each use of **Prism Diquat**. Use the higher label rates when weeds are large or dense. Also, use higher labeled rates for harvest aid when crop vegetation is dense.

## APPLICATION TIMING

**Prism Diquat must be applied to emerged weeds when they are small. Weeds 1 inch to 6 inches in height are the easiest to control.** When weeds have been grazed or mowed, thus removing much of the green foliage, allow the weeds to regrow to a height of 2–4 inches before spraying. For proper application timing of harvest aid applications, refer to each crop for specifications or directions.

Weeds emerging after application of **Prism Diquat** will not be controlled or suppressed.

## RAINFASTNESS

Because **Prism Diquat** is rapidly absorbed by green plant tissue, rain occurring 30 minutes after application will have no effect on the activity of **Prism Diquat**.

## ENVIRONMENTAL CONDITIONS

**Prism Diquat** is active over a wide range of environmental conditions. Cool weather (below 55°F) will slow the activity of **Prism Diquat**, as will cloudy, overcast weather, but will not affect performance.

In dry areas, dust stirred up by high winds or equipment tires can coat target surface and reduce **Prism Diquat** activity. Avoid applying **Prism Diquat** in extremely dusty conditions.

## **Mandatory Spray Drift Management**

### **Aerial Applications:**

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- If the windspeed is 10 miles per hour or less, applicators must use  $\frac{1}{2}$  swath displacement upwind at the downwind edge of the field. When the windspeed is between 11-15 miles per hour, applicators must use  $\frac{3}{4}$  swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions

### **Groundboom Applications:**

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions

### **Boomless Ground Applications:**

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions

## **SPRAY DRIFT MANAGEMENT**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.  
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

## **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### **Controlling Droplet Size – Groundboom**

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **Controlling Droplet Size – Aircraft**

- Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### **BOOM HEIGHT – Groundboom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **Boom-less Ground Applications:**

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

### **RELEASE HEIGHT – Aircraft**

Higher release heights increase the potential for spray drift.

### **Handheld Technology Applications:**

Take precautions to minimize spray drift

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

### **RESISTANCE MANAGEMENT RECOMMENDATIONS**

For resistance management, **Prism Diquat** is a Group 22 herbicide. Any weed population may contain or develop plants naturally resistant to **Prism Diquat** and other Group 22 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of **Prism Diquat** or other Group 22 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.

- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact a Rainbow Agrosiences, LLC representative. Report any incidence of non-performance of this product against a particular weed species to Rainbow Agrosiences, LLC, your local retailer or your local extension specialist. If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemical means to remove escapes, as practical, with the goal of preventing further seed production.

## SPECIFIC USE DIRECTIONS

The following table indicates use pattern, rates, minimum spray volumes, and preharvest interval for specific uses.

Crop	Use Pattern	Prism Diquat Rate per Acre	Minimum Total Spray Volume per Acre	Preharvest Interval (Days)	Directions
<b>Alfalfa</b> (seed crop only)	Preharvest desiccation broadcast	1 1/2–2 pt (0.38 – 0.5 lb ai) (see precautions section for additional rate information)	Ground: 15 gal Air: 5 gal	3	<b>Restrictions</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply through irrigation systems.</li> <li>• On thin stands of seed alfalfa use 1 pt (0.25 lb ai) per acre.</li> <li>• <b>DO NOT</b> graze or feed treated forage to livestock.</li> <li>• <b>DO NOT</b> use seed from treated plants for food, feed, or oil purposes.</li> <li>• <b>DO NOT</b> apply more than 2 pt (0.5 lb ai) per acre per application.</li> </ul>

Crop	Use Pattern	Prism Diquat Rate per Acre	Minimum Total Spray Volume per Acre	Preharvest Interval (Days)	Directions
					<ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 pt (2.0 lb ai) per acre per year.</li> <li>• <b>DO NOT</b> make more than 2 applications per crop cycle (2 crop cycles per year) and 4 applications per acre per year.</li> </ul> <b>Precautions</b> <ul style="list-style-type: none"> <li>• Desiccation is complete in 3–10 days.</li> </ul>
<b>Canola</b>	Preharvest desiccation broadcast	1 1/2–2 pt (0.38 – 0.5 lb ai)	Ground: 15 gal Air: 5 gal	7	<b>Restrictions</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply through irrigation systems.</li> <li>• <b>DO NOT</b> apply more than 2 pt (0.5 lb ai) per acre per application</li> <li>• <b>DO NOT</b> apply more than 2 pt (0.5 lb ai) per acre per year.</li> <li>• A maximum of one application per year is allowed.</li> <li>• Harvest no later than 10 days after application.</li> </ul>
<b>Clover</b> (seed crop only)	Preharvest desiccation broadcast	1 1/2–2 pt (0.38 – 0.5 lb ai)	Ground: 15 gal Air: 5 gal	3	<b>Restrictions</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply through irrigation systems.</li> <li>• <b>DO NOT</b> apply more than 2 pt (0.5 lb ai) per acre per application.</li> <li>• <b>DO NOT</b> graze or feed treated forage to livestock.</li> <li>• <b>DO NOT</b> use seed from treated plants for food, feed, or oil purposes.</li> <li>• <b>DO NOT</b> apply more than 8 pt (2.0 lb) per acre per year.</li> <li>• <b>DO NOT</b> make more than 2 applications per crop cycle (2 crop cycles per year) and 4 applications per acre per year.</li> </ul> <b>Precautions</b> <ul style="list-style-type: none"> <li>• Desiccation is complete in 3–10 days.</li> </ul>

Crop	Use Pattern	Prism Diquat Rate per Acre	Minimum Total Spray Volume per Acre	Preharvest Interval (Days)	Directions
Potato	Preharvest desiccation broadcast	1–2 pt (0.25 – 0.5 lb ai)	Ground: 20 gal Air: 5 gal	7	<b>Restrictions</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply through irrigation systems.</li> <li>• <b>DO NOT</b> apply more than 2 pt (0.5 lb ai) per acre per application.</li> <li>• <b>DO NOT</b> apply to drought-stressed potatoes.</li> <li>• <b>DO NOT</b> exceed a total of 4 pt (1 lb ai) per acre per year.</li> <li>• <b>DO NOT</b> make more than 2 applications per acre per year.</li> <li>• Retreatment Interval: 5 days.</li> </ul> <b>Precautions</b> <ul style="list-style-type: none"> <li>• Make a second application if necessary to obtain additional desiccation where vine growth is dense. For improved vine coverage, a 5-day interval is advised between applications.</li> </ul>
<b>Tree, Vine, Small Fruit, Vegetable Crops - Nonbearing</b> Acerola (West Indian Cherry) Almonds Apple Apricots Artichokes Asparagus Avocados Bananas Blackberry Blueberry Boysenberry Cherries Coffee Conifers Crabapple Cranberry Dates Dewberry Elderberry Figs Filberts Ginseng Gooseberry Grapes Grapefruit Guava Huckleberry	Directed spray	1 1/2–2 pt (0.38 – 0.5 lb ai)	Ground: 15 gal	<b>DO NOT</b> use for food or feed for one year after application	<b>Restrictions</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 2 pt (0.5 lb ai) per acre per application.</li> <li>• <b>DO NOT</b> apply more than 10 pt (2.5 lb ai) per acre per year.</li> <li>• <b>DO NOT</b> make more than 5 applications per acre per year.</li> <li>• <b>DO NOT</b> graze or treated areas.</li> </ul> <b>Precautions</b> <ul style="list-style-type: none"> <li>• PRISM DIQUAT can be used during site preparation prior to planting and up to 1 year of harvest.</li> <li>• Retreatment may be necessary for complete control of grasses and older established weeds.</li> <li>• <b>DO NOT</b> allow spray to contact green stems, foliage, or fruit as injury can occur.</li> </ul>

Crop	Use Pattern	Prism Diquat Rate per Acre	Minimum Total Spray Volume per Acre	Preharvest Interval (Days)	Directions
Jojoba Kiwi Lemons Limes Loganberry Macadamia Mango Nectarines Olives Oranges Papayas Passion Fruit Peaches Pears Pecans Persimmons Pistachios Plantains Plums Pomegranates Prunes Raspberry Tangelos Tangerines Walnuts					<ul style="list-style-type: none"> <li>Use a shield or wrap plant when spraying around young trees or vines.</li> </ul>

Other Uses	Use Pattern	Prism Diquat	Directions
Noncrop or Nonplanted Areas on Farms Fence Lines, Farmyards, Farm Buildings, Fuel Storage Areas, Barrier Strips, Equipment Areas, and Dry (non-flooded) Areas around ponds, lakes, and drainage ditches on farms	Broadcast	1–2 pt (0.25 – 0.5 lb ai) in a minimum of 15 gal water per acre	<b>Restrictions</b> <ul style="list-style-type: none"> <li><b>DO NOT</b> apply this product via backpack sprayer.</li> <li><b>DO NOT</b> apply more than 2 pt (0.5 lb ai) per acre per broadcast application.</li> <li><b>DO NOT</b> apply more than 20 pt (5 lb ai) per acre per year.</li> <li><b>DO NOT</b> make more than 10 applications per acre per year.</li> </ul> <b>Precautions</b> <ul style="list-style-type: none"> <li>Apply for full coverage and thorough weed contact.</li> <li>Retreatment may be necessary to control grasses and established weeds.</li> <li>Avoid spray contact with foliage of food crops or ornamental plants or other desirable vegetation.</li> <li>Add the labeled rate of 75% or greater nonionic surfactant to the finished spray volume.</li> </ul>
	Spot Treatment	1–2 quarts (0.5 – 1 lb ai) Plus the labeled rate of a 75% or greater nonionic surfactant per 100 gal water or 0.75 oz. (22 ml) Plus the labeled rate of a 75% or greater nonionic surfactant per 1 gal of water	

## AGRICULTURAL AQUATIC USE DIRECTIONS[\*]

[\*Not registered for use by California]

Necessary approval and/or permits must be obtained prior to application if required. Consult the responsible State Agencies (i.e., Fish and Game Agencies or Department of Natural Resources). Treatment of dense weed areas may result in oxygen loss from decomposition of dead weeds. This loss of oxygen may cause fish suffocation. Therefore, treat only  $\frac{1}{3}$ – $\frac{1}{2}$  of the water body area at one time and wait 14 days between treatments.

For application only to stillwater (e.g., farm ponds, farm lakes, and farm drainage ditches) where there is minimal or no outflow to public waters.

Treated water may be used according to the following table or until such time as an approved assay (example: PAM II Spectromatic Method) shows that the water does not contain more than the designated maximum contaminant level goal (MCLG) of 0.02 mg/l (ppm) of diquat dibromide (calculated as the cation).

Waters treated with this product may be hazardous to non-target aquatic organisms. Treatment of aquatic weeds and algae can result in oxygen loss from decomposition of dead biomass. This oxygen loss can cause fish and invertebrate suffocation. To minimize this hazard, do not use products containing diquat dibromide to treat more than ½ of the water body (excluding water infrastructure and constructed conveyances such as drainage canals, ditches and pipelines or intakes and aqueducts for drinking water or irrigation use) to avoid depletion of oxygen due to decaying vegetation. Wait at least 7 days between diquat dibromide applications to adjacent treatment areas within the same waterbody. Begin treatment along the shore and proceed outward in bands to allow fish to move into untreated areas. Consult with the state or local agency with primary responsibility for regulating pesticides before applying this product to public waters to determine if a permit is required.

#### Water Use Restrictions Following Application with Prism Diquat (Days)

Application Rate	Drinking	Fishing and Swimming	Livestock Consumption	Spray Tank Applications** and Irrigation to Turf and Ornamentals	Spray Tank Applications** and Irrigation to Food Crops
2 gal/surface acre	3 days	0	1 day	3 days	5 days
1 gal/surface acre	2 days	0	1 day	2 days	5 days
0.75 gal/surface acre	2 days	0	1 day	2 days	5 days
0.50 gal/surface acre	1 day	0	1 day	1 day	5 days
Spot Spray* (< 0.5 gal/surface acre)	1 day	0	1 day	1 day	5 days

\* Rates refer to total surface area.

\*\* For preparing agricultural sprays for food crops, turf or ornamentals (to prevent phytotoxicity), **DO NOT** use water treated with **Prism Diquat** before the specified time periods. When the contents of more than one spray tank is necessary to complete a single aquatic application, no water holding restrictions apply between the consecutive spray tanks.

#### Restrictions:

- **DO NOT** make applications in areas where commercial processing of fish, resulting in the production of fish protein concentrate or fish meal, is practiced. Before application, coordination and approval of local and/or State authorities must be obtained.
- **DO NOT** apply this product at a rate greater than 2.5 lb ai/surface-acre per application for the control of floating, emergent, or marginal vegetation.
- **DO NOT** apply this product at a rate greater than 1 lb ai/acre-foot per application for the control of submersed weeds
- **DO NOT** apply this product on cattails (*Typha* spp.) as a target weed species.
- The use of a vehicle-mounted boom sprayer to apply this product above the water surface for treatment of floating, emergent, or marginal vegetation, except for applications made via aircraft,



is prohibited. This prohibition does not apply to applications made below the water surface via submersed or trailing hoses

**Apply Prism Diquat in accordance with the following table:**

Weed Species	Subsurface or Bottom Placement (gal/surface acre)*	Surface Application (gal/surface acre)*
Bladderwort ( <i>Utricularia</i> spp.)	1–2	2
Coontail ( <i>Ceratophyllum demersum</i> )	2	2
Elodea ( <i>Elodea</i> spp.)	2	2
Naiad ( <i>Najas</i> spp.)	1–2	2
Pondweeds <sup>1</sup> ( <i>Potamogeton</i> spp.)	2	2
Watermilfoils ( <i>Myriophyllum</i> spp.)	1–2	2
Hydrilla ( <i>Hydrilla verticillata</i> )	2	2
Waterlettuce <sup>2</sup> ( <i>Pistia Stratiotes</i> )	NA	0.5–0.75
Waterhyacinth <sup>2</sup> ( <i>Eichhornia crassipes</i> )	NA	0.5–0.75
Pennywort <sup>3</sup> ( <i>Hydrocotyle</i> spp.)	NA	0.5–0.75
Frog's Bit <sup>6</sup> ( <i>Limnobium spongia</i> )	NA	0.5–0.75
Salvinia <sup>2</sup> ( <i>Salvinia</i> spp.)	NA	0.5–0.75
Duckweed <sup>4</sup> ( <i>Lemna</i> spp.)	NA	1
Algae <sup>5</sup> ( <i>Spirogyra</i> spp. & <i>Pithophora</i> spp.)	1–2	2

\* For water less than or equal to 2 ft in average depth of treatment area, use a maximum of 1 gal of **Prism Diquat** per surface acre. Lowest rates must be used in shallow areas where the water depth is considerably less than the coverage depth of the entire treatment area, for example, shallow shoreline area. At water temperatures below 50°–60°F efficacy and immediacy of results may be reduced.

<sup>1</sup> **Prism Diquat** controls *Potamogeton* species except Richardson's pondweed (*P. richardsonii*). For control of *P. robbinsii*, applications must be made when the plants are in the early stages of growth including in Spring and early Summer.

<sup>2</sup> For salvinia, waterlettuce and waterhyacinth, use the labeled rate of **PRISM DIQUAT** in 75–200 gal water plus the labeled rate of a 75% or greater nonionic surfactant per acre for surface sprays and for aerial application for waterlettuce and waterhyacinth control, apply the labeled rate of **PRISM DIQUAT** in 10–24 gal water plus the labeled rate of a 75% or greater nonionic surfactant per acre.

<sup>3</sup> For pennywort control, apply in 50–150 gal of water plus the labeled rate of a 75% or greater nonionic surfactant per acre for full coverage and thorough weed contact. Repeat treatments may be necessary to control regrowth.

<sup>4</sup> For duckweed control, apply as an overall spray in 50–150 gal. of water plus the labeled rate of a 75% or greater nonionic surfactant per acre. Retreatment may be necessary for plants missed in previous applications and regrowth.

<sup>5</sup> For suppression of certain filamentous algae species including *Spirogyra* and *Pithophora*, apply according to the submersed use directions.

<sup>6</sup> **Not registered for use by California.**

## Application

In mixed weed populations, use the high rate of application as indicated by weeds present.

## Subsurface Applications

Where the submersed weed growth, especially Hydrilla, has reached the water surface, apply either in a water carrier or an invert emulsion through boom trailing hoses carrying nozzle tips to apply the dilute spray below the water surface to insure adequate coverage.

### Bottom Placement

Where the submersed weeds, especially Hydrilla, Bladderwort, and Coontail growth have reached the water surface or where water is slowly moving through the submersed weed growth that has reached the water surface, especially Hydrilla, Bladderwort, and Coontail, control may be enhanced when applied in an invert emulsion carrier injecting diluted **Prism Diquat** near the bottom with weighted hoses. The addition of a copper-based algaecide will improve control. Where algae are present along with the submersed weeds, pretreatment with copper- based algaecide at specified rates is advised for best results.

### Surface Application

For submerged aquatic weeds, apply **Prism Diquat** either as concentrate slowly poured directly from the container in strips[\*] or as a spray in sufficient carrier. Applications must be made to ensure complete coverage of the weed areas. In mixed weed populations, use the high rate of application as indicated by weeds present.

[\*Not registered for use by California]

### Limitations and Precautions (Terrestrial and Aquatic Uses)

- Direct spray contact or drift of **Prism Diquat** will cause severe plant injury or death. Avoid contact of desirable vegetation.
- Weeds emerging after application of **Prism Diquat** will not be controlled or suppressed. Retreatment may be necessary to control large weeds or established weeds.
- Application to muddy water may result in reduced control. Minimize creating muddy water during application. **(Aquatic Uses)**
- Avoid applying under conditions of high wind, water flow, or wave action. **(Aquatic Uses)**
- **DO NOT** apply this product through any type of irrigation system. Rinse all spray equipment thoroughly with water after use.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage or disposal. TO PREVENT ACCIDENTAL POISONING, NEVER STORE THIS PRODUCT IN FOOD, DRINK, OR UNLABELED CONTAINERS.

**PESTICIDE STORAGE:** Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

**PESTICIDE DISPOSAL:** Pesticide spray mixture or rinsate that cannot be used must be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING:

**[For plastic containers ≤ 5 gallons: Nonrefillable Container: DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or

puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

**[For plastic containers > 5 gallons: Nonrefillable container. DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

**[NON-REFILLABLE PLASTIC CONTAINER [(E.G., INTERMEDIATE BULK CONTAINERS [IBC]) (SIZE OR SHAPE TOO LARGE TO BE TIPPED, ROLLED OR TURNED UPSIDE DOWN)]][:]**

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into formulation equipment and before final disposal using the following pressure rinsing procedure: Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure spray duration and/or spray volume. If the manufacturer's instructions are not available pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain pour or pump rinsate into formulation equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

**[REFILLABLE PLASTIC CONTAINER [(E.G., INTERMEDIATE BULK CONTAINERS [IBC]) (SIZE OR SHAPE TOO LARGE TO BE TIPPED, ROLLED OR TURNED UPSIDE DOWN)]][:]**

Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. If not returned to the point of purchase or to the designated location, triple rinse emptied container and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

When this container is empty, replace the cap and seal all openings that have been opened during use; and return the container to the point of purchase or to a designated location named at the time of purchase of this product in a bulk container. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

### **LIMITATION OF WARRANTY AND LIABILITY**

RAINBOW AGROSCIENCES, LLC, warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use. The Directions For Use are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of RAINBOW AGROSCIENCES, LLC, or the SELLER. To the extent consistent with applicable law RAINBOW AGROSCIENCES, LLC, shall not be liable for the consequential, special or indirect damages resulting from the handling or use of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Except as expressly provided herein, RAINBOW AGROSCIENCES, LLC, makes no warranties, guarantees, or representations of any kind, either express or implied, or by usage of trade, statutory or

otherwise, with regard to the product sold, including but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage

[[Reglone(R)] is a registered trademark of Syngenta Group Company.]

[EPA APPROVAL DATE]

(BASE LABEL)

DIQUAT DIBROMIDE	GROUP	22	HERBICIDE
------------------	-------	----	-----------

TO PREVENT ACCIDENTAL POISONING, NEVER PUT INTO FOOD, DRINK OR OTHER CONTAINERS AND USE STRICTLY IN ACCORDANCE WITH ENTIRE LABEL

## Prism Diquat

Contains diquat dibromide, the active ingredient used in [Reglone®]

ACTIVE INGREDIENT: (% by weight)

Diquat dibromide [6,7-dihydrodipyrido (1,2-a:2',1'-c) pyrazinedium dibromide] 37.3%

OTHER INGREDIENTS: 62.7%

TOTAL 100.0%

Prism Diquat is formulated as a soluble liquid containing 2 lb diquat cation per gal as 3.73 lb salt per gal.

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• DO NOT induce vomiting unless told to do so by the poison control center or doctor.</li><li>• DO NOT give anything by mouth to an unconscious person.</li></ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Immediately call a poison control center or doctor for further treatment advice.</li></ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>NOTE TO PHYSICIAN</b> <p>To be effective, treatment for diquat poisoning must begin IMMEDIATELY. Treatment consists of binding diquat in the gut with suspensions of activated charcoal or bentonite clay, administration of cathartics to enhance elimination, and removal of diquat from the blood by charcoal hemoperfusion of continuous hemodialysis.</p> <p>If in eyes, treat symptomatically. Symptoms may develop gradually. Severe damage may be caused by apparently trivial contact and healing may be delayed. Medical supervision should be continued until healing is complete.</p>	
<b>HOT LINE NUMBER</b> <p>Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. You may also contact the Poison Control Center at 1-800-222-1222 for emergency medical treatment information. For non-emergency information concerning this product, call the National Pesticide Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8:00AM to 12:00 PM Pacific Time, or at <a href="https://npic.orst.edu">https://npic.orst.edu</a>.</p>	

### For Chemical Emergency:

Spill, Leak, Fire, Exposure, or Accident,

Call CHEMTREC Day or Night

Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

### PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

### CAUTION

Harmful if swallowed. Harmful if inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. **Human Flagging is Prohibited**

### Environmental Hazards (Terrestrial and Aquatic Uses)

This pesticide is toxic to aquatic invertebrates. For Terrestrial Uses, DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment washwaters or rinsate. For Aquatic Uses, DO NOT apply directly to water except as specified on this label. Treatment of dense weed areas may result in oxygen loss from decomposition of dead weeds. This loss of oxygen may cause fish suffocation. Therefore, treat only 1/3 – 1/2 of the water body area at one time, especially if dense areas of weeds and/or algae exist and wait 14 days between treatments.

Necessary approval and/or Permits must be obtained prior to application if required. Consult the responsible State Agencies (i.e., Fish and Game agencies or Department of Natural Resources) before making applications to public waters.

### NON-TARGET ORGANISM ADVISORY

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

### Physical-Chemical Hazards

DO NOT mix or allow coming in contact with oxidizing agents. Hazardous chemical reactions may occur.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage or disposal. TO PREVENT ACCIDENTAL POISONING, NEVER STORE THIS PRODUCT IN FOOD, DRINK, OR UNLABELED CONTAINERS.

**PESTICIDE STORAGE:** Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

**PESTICIDE DISPOSAL:** Pesticide spray mixture or rinsate that cannot be used must be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING:

**[For plastic containers ≤ 5 gallons: Nonrefillable Container: DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

**[For plastic containers > 5 gallons: Nonrefillable container. DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

**[NON-REFILLABLE PLASTIC CONTAINER [(E.G., INTERMEDIATE BULK CONTAINERS [IBC]) (SIZE OR SHAPE TOO LARGE TO BE TIPPED, ROLLED OR TURNED UPSIDE DOWN)]][:]** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into formulation equipment and before final disposal using the following pressure rinsing procedure: Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure spray duration and/or spray volume. If the manufacturer's instructions are not available pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain pour or pump rinsate into formulation equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

**[REFILLABLE PLASTIC CONTAINER [(E.G., INTERMEDIATE BULK CONTAINERS [IBC]) (SIZE OR SHAPE TOO LARGE TO BE TIPPED, ROLLED OR TURNED UPSIDE DOWN)]][:]**

Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. If not returned to the point of purchase or to the designated location, triple rinse emptied container and offer for recycling. Disposal of this container must be in compliance with state and local regulations. When this container is empty, replace the cap and seal all openings that have been opened during use; and return the container to the point of purchase or to a designated location named at the time of purchase of this product in a bulk container. Prior to refilling, inspect

carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

[See] [inside] [label] [booklet] [side] [panel] [for] [First Aid][.] [additional]  
[Precautionary Statements][.] [and] [Directions for Use] [including] [Storage  
and Disposal] [instructions][.]

[Prism Diquat is not manufactured, or distributed by Syngenta Crop Protection,  
LLC seller of Reglone®] [Dessicant]

EPA Reg. No. 103591-xx

EPA Est. No.

Manufactured for:

NET CONTENTS

RAINBOW AGROSCIENCES, LLC

535 E Plainfield Road, Suite E

Willowbrook, IL 60527 [Lot][Batch][No.][:]\_\_\_\_\_