



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

October 30, 2024

John E. Inouye, Senior Environmental Scientist
California Department of Pesticide Regulation
1001 I Street
P.O. Box 4015
Sacramento, California 95812-4015

Subject: FIFRA 24(c) Registration - SLN Amendment Acknowledgment Letter
SLN Registration Number: CA130010
EPA Decision Number: 596235
SLN Expiration Date: May 31, 2029
Parent EPA Reg. No.: 59639-77
Product Name: Tame 2.4 EC Spray

Dear Mr. Inouye,

This letter acknowledges receipt of an amendment for the above-listed Special Local Need (SLN) registration pursuant to Section 24(c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. The SLN registration permits use of the product Tame 2.4 EC Spray Insecticide-Miticide (fenpropathrin) on listed bearing citrus, bearing fruiting vegetables, bearing grape, bearing tree nuts, and bearing tropical and subtropical fruit (inedible peel) grown in outdoor nurseries to control glassy-winged sharpshooter and Asian citrus psyllid. This amendment extends the expiration date to May 31, 2029.

The Agency has completed a review of the amendment for this use and acknowledges the State's Special Local Need registration, as amended. Please inform the registrant that we have placed a copy of the amended SLN label in our files. If you have any questions, please contact Stacey Groce via email at Groce.Stacey@epa.gov.

Sincerely,

Jennifer Gaines, Senior Regulatory Specialist
Emergency Response Team
Minor Use and Emergency Response Branch
Registration Division/Office of Pesticide Programs

cc: Fabiola Estrada, USEPA Region 9 Coordinator

RESTRICTED USE PESTICIDE

DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS.

For retail sale to and use by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

FIFRA 24(c) Special Local Need Label (SLN)

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

For use on Listed Bearing Citrus, Bearing Fruiting Vegetables, Bearing Grape, Bearing Tree Nuts and Bearing Tropical and Sub Tropical Fruit (inedible peel) Grown in Outdoor Nursery

Not for use in Commercial Production Orchards

Tame®

2.4 EC SPRAY

INSECTICIDE - MITICIDE

EPA Reg. No. 59639-77

EPA SLN No. CA-130010

	By Wt.
Active Ingredient	
Fenpropathrin*	30.9%
Other Ingredients	69.1%
Total	100.0%
*alpha-cyano-3-phenoxybenzyl 2,2,3,3-tetramethylcyclopropanecarboxylate	

This label expires and must not be distributed or used in accordance with this SLN registration after May 31, 2029.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the EPA registered label for Tame® 2.4 EC Spray (EPA Reg. No. 59639-77) and this label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Plants treated under this 24(c) Special Local Need registration cannot be shipped until the pre-harvest interval (PHI) for those plants has expired.

BEARING CITRUS		
PESTS	PRODUCT RATE	SPECIAL INSTRUCTIONS
Glassy-winged Sharpshooter Asian Citrus Psyllid	16 fl oz/A (0.3 lb ai/A) 16 fl oz/100 gallons	<p>Apply with ground equipment as a full coverage spray in sufficient water for thorough coverage. Begin applications when pest activity is first noted and repeat as needed to maintain control but not more often than every 10 days.</p> <p>Resistance Management: It is generally recommended that no more than one <i>Tame 2.4 EC</i> Spray application be made per season as part of a Resistance Management Program. Use non-pyrethroid products at other timings to control pests.</p>
<ul style="list-style-type: none"> • Best results are achieved if applications are made when temperatures are moderate (usually early morning or evening). • Avoid making applications if the temperature is 94°F or above. • Preharvest Interval (PHI) – 24 hours. • Do not ship product within 24 hours after application. • Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) total application of <i>Tame 2.4 EC</i> Spray per acre per season. • Do not feed or graze livestock on treated grove cover crops. • Additional pesticide sprays may be required to obtain season-long pest control. See Resistance Management Statement. 		

BEARING FRUITING VEGETABLES		
Bell Pepper, Chili Pepper, Cooking Pepper, Eggplant, Ground Cherry (<i>Physalis</i> spp.), Pepino, Pimento, Sweet Pepper, Tomatillo, Tomato		
PESTS	PRODUCT RATE	SPECIAL INSTRUCTIONS
Glassy-winged Sharpshooter	10-2/3 fl oz/A (0.2 lb ai/A) 10-2/3 fl oz/100 gallons	<p>Apply with ground equipment in adequate water for uniform coverage. Begin applications when first pest activity is noticed and repeat as needed to maintain control but not more often than every 7 days.</p> <p>DO NOT tank mix with any copper formulations.</p> <p><i>Tame 2.4 EC</i> Spray may be mixed with and/or alternated with commonly used insecticides and miticides to comply with local IPM programs.</p>
<ul style="list-style-type: none"> • Preharvest Interval (PHI) – 3 days. • Do not ship product within 3 days following application. • Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) total application of <i>Tame 2.4 EC</i> Spray per acre per season. 		

BEARING GRAPE		
PESTS	PRODUCT RATE	SPECIAL INSTRUCTIONS
Glassy-winged Sharpshooter	16 fl. oz/A (0.3 lb ai/A) 16 fl. oz/100 gallons	<p>Apply as a full coverage spray. Start sprays when pest activity begins and repeat as needed to maintain control but not more often than every 7 days. When pest pressure is moderate or severe use the higher recommended rate.</p> <p>Resistance Management: It is generally recommended that no more than 2 <i>Tame</i> 2.4 EC Spray applications be made per season as part of a Resistance Management Program. Use non-pyrethroid products at other timings to control pests.</p> <p><i>Tame</i> 2.4 EC Spray may be tank mixed and/or alternated with other commonly used insecticides and miticides to comply with local IPM programs.</p>
<ul style="list-style-type: none"> Preharvest Interval (PHI) – 21 days. Do not ship product for 21 days after application. Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) total application of <i>Tame</i> 2.4 EC Spray per acre per season. 		

BEARING TREE NUTS		
Almond, Beechnut, Brazil, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory, Macadamia, Pecan, Pistachio, Walnut		
PESTS	PRODUCT RATES	SPECIAL INSTRUCTIONS
Glassy-winged Sharpshooter	10-2/3 to 16 fl oz/A (0.2 to 0.3 lb ai/A) 10-2/3 to 16 fl oz/100 gallons	<p>Apply by as a full coverage spray and repeat as needed to maintain control but not more often than every 10 days. Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will assure better coverage.</p> <p>Begin applications as common to the production area and the target pest.</p> <p>Resistance Management: It is generally recommended that no more than two (2) <i>Tame</i> 2.4 EC Spray applications be made per season as part of a Resistance Management program. Use non-pyrethroid products at other timings to control pests.</p> <p>Make pyrethroid insecticide applications according to local Integrated Pest Management (IPM) programs and guidelines.</p>
<ul style="list-style-type: none"> Preharvest Interval (PHI) – 3 days. Do not ship product within 3 days of application. Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) total application of <i>Tame</i> 2.4 EC Spray per acre per season. Do not apply as an Ultra Low Volume (ULV) spray. Do not feed or allow livestock to graze on cover crops from treated orchards. 		

BEARING TROPICAL AND SUB TROPICAL FRUIT (Inedible Peel) Avocado, Canistel, Mango, Papaya, Sapodilla, Sapote, Black Mamey, Star Apple		
PESTS	PRODUCT RATES	SPECIAL INSTRUCTIONS
Glassy-winged Sharpshooter	16 to 21-1/3 fl oz/A (0.3 to 0.4 lb ai/A)	Apply as a ground application in a minimum of 100 gals of water per acre. Make applications beginning when first pest activity is noticed and repeat as needed but not more often than every 14 days. Resistance Management Avocado - Do not make more than one pyrethroid application per season as part of a Resistance Management Program. Other crops - Do not make more than two <i>Tame</i> 2.4 EC Spray applications per season as part of a Resistance Management Program. Use non-pyrethroid products at other timings to control pests.
<ul style="list-style-type: none"> Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) of <i>Tame</i> 2.4 EC Spray per acre per season. Preharvest Interval (PHI) – 24 hours. Do not ship product within 24 hours of application. 		

FIFRA §24(c) registrant: Valent U.S.A. LLC
 P.O. Box 5075
 San Ramon, CA 94583
 800-89-VALENT (898-2536)

Tame is a registered trademark of Valent U.S.A. LLC

