

RESTRICTED USE PESTICIDE

DUE TO HAZARDS TO NONTARGET SPECIES

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

FIFRA SECTION 24(c) SPECIAL LOCAL NEED LABEL

FOR DISTRIBUTION AND USE ONLY WITHIN IDAHO

This product may only be used to control voles along irrigation canal levees by ground application only.

This label is valid until December 31, 2029, or until otherwise amended, withdrawn, canceled, or suspended.

ZINC PHOSPHIDE ON WHEAT

Alternate Brand Name: **ZINC PHOSPHIDE ON WHEAT – ID**

EPA Reg. No. 56228-3

EPA SLN No. ID-130003

ACTIVE INGREDIENT

Zinc Phosphide..... 1.82%

OTHER INGREDIENTS..... 98.18%

TOTAL..... 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Misuse may result in civil or criminal enforcement action.

READ THIS LABEL: Read this entire label and the label for EPA Registration Number 56228-3. This product must be used strictly in accordance with this label's precautionary statements and use directions on both labels, as well as with all applicable State and Federal laws and regulations.

This state-specific Section 24(c) labeling and the Section 3 labeling for EPA Reg. No. 56228-3 must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this label and the EPA-registered label for EPA Reg. No. 56228-3.

Before using this product, contact the U.S. Fish and Wildlife Service and the applicable State wildlife agency and obtain all necessary kill or collecting permits. Use only for the sites, pests, and application methods described on this label.

Do not use this product within 0.5 miles of habitat of the Northern Idaho Ground Squirrel (*Urocitellus brunneus*, formerly *Spermophilus brunneus*) in Adams, Gem, Idaho, Valley, and Washington counties, Idaho.

APPLICATION DIRECTIONS

PREBAITING:

Prebaiting is recommended. Two or 3 days prior to scheduled use of toxic bait, prebait along irrigation canal levees with untreated wheat at a rate per acre similar to the intended application rate of the toxic bait. If prebaiting, apply toxic bait where prebait has been consumed using a rate similar to the prebait rate.

GROUND APPLICATIONS:

Apply toxic bait only in areas of current vole activity at a rate of 2.5 to 6 pounds per acre. Do not exceed 4 applications per year. Do not exceed 10 pounds (0.18 lb active ingredient) of toxic bait per treated acre per year.

Treatments may be applied by hand baiting, or drop-tubes fixed to ground-based (such as pickup truck, ATV, tractor) bait-dispensing equipment.

Apply bait only along irrigation canal levees. Do not treat the canal side of the levee.

FOLLOW-UP:

Wearing waterproof gloves, collect and dispose of dead, exposed animals. Dispose of leftover bait in accordance with **PESTICIDE DISPOSAL** instructions under **STORAGE AND DISPOSAL**. If reinfestation does occur, repeat treatment.

24(c) Registrant:

U.S. Department of Agriculture
Animal and Plant Health Inspection Service
4700 River Road, Unit 149
Riverdale, MD 20737

Local Contact:

USDA APHIS Wildlife Services
9134 West Blackeagle Drive
Boise, ID 83709

Net Contents: _____

Batch Code: _____