



## **RESTRICTED USE PESTICIDE**

May injure (phytotoxic) susceptible non-target plants.

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial and certified applicators must ensure that all persons involved in these activities are informed of the precautionary statements.

### ***FIFRA Section 24(c) Special Local Need Label***

#### **FOR DISTRIBUTION AND USE ONLY WITHIN KANSAS**

**For Fall or Spring Applications to Control Kochia, Puncturevine and Russian Thistle in Fallow or Eco-Fallow Fields**

## **Corvus<sup>®</sup> Herbicide**

EPA Reg. No. 264-1066

SLN. No. KS-140004

**THIS LABEL IS VALID UNTIL DECEMBER 31, 2029 UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.**

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

<b>Product</b>	<b>Rate</b>	<b>Crop</b>	<b>Weeds</b>
Corvus <sup>®</sup> Herbicide	3-4 fl. oz/acre	Fallow, Eco-Fallow	For enhanced control of kochia, Russian thistle, puncturevine

- Apply Corvus<sup>®</sup> Herbicide in the fall or spring to stubble of harvested crops or fallow/eco-fallow fields for control of weeds prior to emergence in no-till or conservation tillage systems.
- Do not apply to frozen soils.
- Apply as a broadcast spray with ground equipment using a minimum of 10 gallons of spray mixture per acre.
- Corvus Herbicide should be tank mixed with additional herbicides labeled for use in fallow or eco-fallow applications in order to achieve improved residual weed control. Recommended tank mix partners include atrazine (0.5-1.0 lb. ai/acre) or metribuzin (0.19-0.375 lb. ai/acre)
- For enhanced burndown of emerged broadleaf or grasses weeds, tank mix the Corvus Herbicide + atrazine or metribuzin mixture with a labelled burndown herbicide such as dicamba, 2,-4-D, Gramoxone<sup>®</sup> or glyphosate (Note: glyphosate will not add additional control of glyphosate-resistant weeds). Always include an appropriate adjuvant (COC, MSO, HSOC) and nitrogen source (UAN,AMS) for enhanced burndown activity.
- Follow all label directions, rotational crop guidelines and use restrictions/precautions for Corvus Herbicide and any tank-mix partners and use the most restrictive guidelines.

## CORVUS® HERBICIDE USE RESTRICTIONS AND PRECAUTIONS

If the water table (i.e., level of saturation) is less than 25 feet below the ground surface, **DO NOT** apply to the following vulnerable loamy sand or sand soils. If the water table is unknown, **DO NOT** apply CORVUS Herbicide to a restricted soil.

Aline	Els	Haxtun	Lincoln	Thurman
Bankard	Elsmere	Inavale	Pratt	Tivoli
Dillwyn	Goltry	Kanza	Sarpy	Valent
Dix	Goodnight	Las Animas	Schamber	Valentine
Dwyer	Gracemore	Likes	Simeon	

## ROTATIONAL CROP RESTRICTIONS

Rotational crops vary in their response to low concentrations of CORVUS Herbicide remaining in the soil. The amount of CORVUS Herbicide that may be present in the soil depends on soil moisture, soil temperature, application rate, elapsed time since application and other environmental factors. When CORVUS Herbicide is used in combination with other products, always follow the most restrictive rotational crop requirements. The following rotational crops may be planted after applying CORVUS Herbicide:

Crop	Rotational Interval <sup>2</sup>	Minimum Precipitation Requirement <sup>1</sup>
Field Corn	0 Months	None
Wheat, Triticale, Cereal and rye	4 Months	None
Barley, Soybean, Sweet corn <sup>3</sup> , Popcorn <sup>3</sup>	9 Months	15 inches of cumulative precipitation from application to planting of rotational crop
Rice <sup>3</sup> , Cotton <sup>3</sup>	10 Months	15 inches of cumulative precipitation from application to planting of rotational crop
Peanuts <sup>3</sup>	11 Months	15 inches of cumulative precipitation from application to planting of rotational crop
Alfalfa, Green and Dry Beans, Oats, Sorghum <sup>4</sup> , Sunflower, Canola, Potato, Sugar beet and all other crops <sup>5</sup>	17 Months	30 inches of cumulative precipitation from application to planting of rotational crop

<sup>1</sup> The amount of cumulative precipitation required before planting a rotational crop is in addition to the required rotational interval given in months. Furrow or flood irrigation must not be included in total. No more than 7 inches of overhead irrigation must be included in total.


<sup>2</sup> Crop varieties planted back at intervals of one year or less must not have known acute sensitivity to ALS-inhibiting and/or SU herbicides.

<sup>3</sup> When soil pH is 7.5 or above, crop plant back needs to be delayed to 17 months and to 24 months for crops listed in the 17 month interval above.

<sup>4</sup> For CORVUS Herbicide used at 2.25 - 3.3 fl oz. per acre or less and the total of Thiencarbazone-methyl from all sources is 0.014 pounds active ingredient per acre or less, sorghum can be planted at the 9 month or longer interval.

<sup>5</sup> All other crops may be seeded only after the completion of a successful bioassay after a CORVUS Herbicide application. Refer to the "Field/Small Scale Bioassay" section. In the event of crop failure: If the corn crop treated with CORVUS Herbicide is lost, only field corn and corn grown for silage may be replanted immediately. **DO NOT** make an additional application of CORVUS Herbicide.

**Location of soil types listed on Page 2 in the eleven counties overlying the Great Bend Prairie and the Equus Beds Aquifers**, are shown in the map below. Before using CORVUS Herbicide in the shaded areas or surrounding counties with restricted soil types, identify the soil series of the treated field. Identification may be done by consulting the NRCS website (<http://websoilsurvey.nrcs.usda.gov/app/WebSoilSurvey.aspx>), a copy of your county soil survey, or visit or contact your local NRCS office or local extension agent. Use is only restricted on the named soils if the water table is less than 25 feet below the ground surface.



For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

Bayer CropScience, LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167

Corvus is a registered trademark of Bayer.