



FIFRA 24(c) Special Local Need Label (SLN)
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MISSISSIPPI

ISOCYCLOSERAM	GROUP	30	INSECTICIDE
---------------	-------	----	-------------

Vertento®

In-Furrow, At-Plant Application for Management of Rootworm Species in Peanut

EPA Reg. No. 100-1711
EPA SLN No. MS-260001

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2031

KEEP OUT OF REACH OF CHILDREN
CAUTION

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and main EPA-registered label for Vertento, EPA Reg. No. 100-1711 must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the main EPA-registered label for Vertento, EPA Reg. No. 100-1711.

SOIL APPLICATION

Target Pest	Rate (fl oz/A)	Application Timing	Use Directions
Rootworm species (larvae) including southern corn rootworm and banded cucumber beetle	2.6	At planting	In-Furrow Applications: Apply into the seed furrow through spray nozzles or micro- tubes, behind the planter furrow openers and in front of the press wheel. Apply this product diluted in a minimum volume of 2 gal/A.
Resistance Management: Refer to Section 3.2 of the Vertento Section 3 label.			
USE RESTRICTIONS			
1) Refer to Section 6.1 of the Vertento Section 3 label for additional product use restrictions. 2) Maximum Single Application Rate: 2.6 fl oz/A/application a. DO NOT exceed 0.0676 lb ai/A of isocycloseram-containing products. 3) Minimum Application Interval: N/A			



Target Pest	Rate (fl oz/A)	Application Timing	Use Directions
4) Maximum Annual Rate: 2.6 fl oz/A/year a. DO NOT exceed 0.0676 lb ai/A/year of isocycloseram-containing in-furrow products. b. DO NOT exceed 0.172 lb ai/A/year of isocycloseram-containing products including all application types (seed treatment, soil, and foliar). 5) Make only one soil application of Vertento per year. However, in the case of a failed peanut crop, another peanut crop may be planted at any time. Vertento may be applied to the replanted crop if the total maximum annual rate does not exceed 2.6 fl oz/A (0.0676 lb ai/A of isocycloseram) for both the initial and replanted peanut crop. 6) DO NOT allow livestock to graze in treated areas or harvest treated peanut plants to be used as livestock feed. 7) Pre-Harvest Interval (PHI): normal commercial harvest			

©2026 Syngenta

Vertento® and the Syngenta logo are trademarks of a Syngenta Group Company.

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: MS1711581AA0226