

FIFRA SECTION 24(c) SPECIAL LOCAL NEED (SLN)

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

ASULAM	GROUP	18	HERBICIDE
--------	-------	----	-----------

ASULAM HERBICIDE

EPA Reg. No. 34704-904 EPA SLN No. OR-100008

FOR POSTEMERGENCE WEED CONTROL IN SPINACH GROWN FOR SEED

This label is valid until December 31, 2030, or until otherwise amended, withdrawn, cancelled, or suspended.

ACTIVE INGREDIENT:

Sodium salt of asulam (methyl sulfanilylcarbamate)* 36.2%

OTHER INGREDIENTS: 63.8%

TOTAL: 100.0%

*Equivalent to 33.1% asulam or not less than 3.34 lbs. per gallon

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This Special Local Need (SLN) label and the label affixed to the main container of Asulam Herbicide must be in the possession of the user at the time of pesticide application. Follow all applicable directions, Worker Protection Standard requirements, restrictions and precautions on this label and the main label affixed to the container of Asulam Herbicide.

WEEDS	RATE/A	INSTRUCTIONS
Dog fennel Lambsquarters (suppression) Pigweed Poppies (suppression) Prickly lettuce Shepherd's purse	1.5 to 3.0 pints	One application per season: Use the 1.5 pt/A on spinach with 2 to 4 leaves, or 3.0 pt/A on spinach with 4 to 6 true leaves. Two applications per season: use 3.0 pt/A on spinach with 4 to 8 true leaves and 3.0 pt/A 7 days later (typically 8 to 12 true leaves). Some stunting of the spinach may occur, especially with two applications at 3.0 pt/A. A non-ionic surfactant at 0.25% v/v may be added to the spray to increase efficacy. Crop oils and other adjuvants such as urea ammonium nitrate are not recommended. Apply by ground application equipment in 20 gallons of water per acre. Maximum 2 applications per year. Maximum annual application 6 pt/A.

RESTRICTIONS

Do not apply Asulam Herbicide through irrigation systems of any type or by aerial application.

SPECIAL CROP USE RESTRICTIONS

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a nonfeed/nonfood use by the Oregon Department of Agriculture and is regulated by Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use on Crops Grown for Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed crop may be used or distributed for food or feed for 1 year (365 days) after the last application of this product. This restriction pertains to, but is not limited to, green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that the screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

"This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies".