



FIFRA Section 24(c)
SPECIAL LOCAL NEED LABEL

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

GROUP M2 FUNGICIDE

MICROTHIOL[®] DISPERSS[®]
MICRONIZED WETTABLE SULFUR

EPA Registration No. 70506-187

EPA SLN No. OR-240006

This label is valid until December 31, 2030 or until otherwise amended, withdrawn, cancelled, or suspended.

For Control of Bud Mites in Hazelnuts/Filberts

ACTIVE INGREDIENT:

Sulfur.....80.0%
OTHER INGREDIENTS20.0%
TOTAL100.0%

MICROTHIOL DISPERSS contains 0.8 pound of sulfur per pound of product.

KEEP OUT OF REACH OF CHILDREN
CAUTION

ENVIRONMENTAL HAZARDS

Follow all applicable directions, restrictions, and precautions in the Environmental Hazards section of the label affixed to the main container of Microthiol Disperss (EPA Reg. No. 70506-187).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This SLN label and the label affixed to the main container of Microthiol Disperss must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the label affixed to the main container of Microthiol Disperss

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

CROP	PEST	RATE (LBS/A)	DIRECTIONS
Hazelnuts/Filberts	Bud Mite	10-20	Apply in 50-400 gallons of spray volume. Begin application at dormant bud stage and repeat as necessary through shoot elongation stage. Repeat applications every 7 to 10 days.



FIFRA Section 24(c)
SPECIAL LOCAL NEED LABEL

Spray Coverage:

Foliage must be completely covered on both the upper and lower leaf surfaces for effective control. Dense canopies can limit complete coverage. Adequate spray volume for thorough coverage is essential.

Precautions:

Some crops may be damaged by sulfur under certain climatic conditions, such as when temperature is high. UPL NA, Inc. does not recommend application if temperature will exceed 90° F within the three days following spraying, due to risk of crop injury.

Restrictions:

- Maximum use rate per acre per year: 60 lbs (48 lbs ai).
- Maximum number of applications per year: 6 at the lowest use rate.
- Under this SLN, do not apply through any type of irrigation system.

Rev. 3/16/2026