

FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience

9330 Zionsville Road,

Indianapolis IN 46268

CYMOXANIL

GROUP

27

FUNGICIDE

Curzate® 60DF

EPA Reg. No. 352-592

EPA Section 24(c) Special Local Need Registration SLN No. WA-260001

For Control of Downy Mildew in Cabbage, Swiss Chard and Table Beet Grown for Seed

This SLN label for Curzate 60DF expires and must not be distributed or used in accordance with this SLN registration after December 31, 2030

For Distribution and Use Only in the State of Washington

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the EPA registered label.

DIRECTIONS FOR USE

For the use of Curzate® 60DF fungicide on, cabbage, swiss chard and table beet grown for seed for the control of downy mildew. Consult the Curzate 60DF EPA-registered label for additional requirements, restrictions and precautions.

HOW TO USE

Use Rate, tank mixes and timing

- Apply Curzate 60DF fungicide at 3.2 ounces per acre (0.12 lb cymoxanil/A) per application in sufficient water to obtain thorough spray coverage of the crop.
- Curzate 60DF must be applied in a tank-mix with a rate of protectant fungicide (EBDC, chlorothalonil, copper, etc.) labeled for downy mildew control. Refer to Tank Mixtures section on Section 3 label. DO NOT use Curzate 60DF alone. Always tank-mix with a labeled rate of protectant fungicide. Follow the more restrictive labeling of any tank mix partners. Do not tank mix with products that contain a prohibition on tank mixing.
- Tank mixing with products that contain copper will likely increase the risk of adverse effects on pollinators and aquatic organisms.
- Applications can be made prior to or after transplanting.
- Initial applications should start when local conditions indicate that downy mildew infection is imminent.
- Make applications on a 5 to 7-day interval as long as disease conditions are favorable.
- DO NOT exceed 5 applications of Curzate 60DF per acre per year.
- DO NOT apply more than 16 oz of Curzate 60DF (equivalent to 0.6 lb cymoxanil) per acre per year.
- For conventional ground application equipment, including tractor-mounted sprayers, boom or backpack sprayers, use a minimum of 20 gals/acre.
- For hard-to-wet foliage, a spray adjuvant may be used to help increase spray coverage and

retention.

Restrictions/Precautions:

- Do not apply by air.
- For use under this SLN label do not apply this product through any type of irrigation system.
- Potatoes, tomatoes, cucurbit crops, lettuce (head and leafy), peppers, spinach, hops, bulb crops and other crops in leafy greens subgroup 4A may be re-planted any time after Curzate 60DF applications. All other crops cannot be planted until 30 days after Curzate 60DF application.

- Crops Grown for Seed
- Cabbage, swiss chard and table beet seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site, and shall include the lot number, amount of material disposed of, the grower(s), and date of disposal.
- No portion of the cabbage, swiss chard and table beet seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
- Cabbage, swiss chard and table beet seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
- Cabbage, swiss chard and table beet seed may not be distributed for human consumption or animal feed.

WSDA Aquatic Advisory

This product is toxic to fish and aquatic invertebrates. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

WSDA Container Disposal Guidance:

Completely empty the contents of the pesticide container into application equipment and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

Read the Limitation of Warranty and Liability on Section 3 of the Federal product label before buying or using this product. If terms are not acceptable, return the unopened package at once to seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

™®Trademarks of Corteva Agriscience and its affiliated companies

R612-004
Accepted: 1/14/02026
Replaces R612-003