



PROTHIOCONAZOLE	GROUP	3	FUNGICIDE
-----------------	-------	---	-----------

FIFRA Section 24(c) Special Local Need (SLN) Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

Proline® 480 SC Fungicide

EPA REG No. 264-825

EPA SLN No. WA-260008

For Suppression of Fusarium Wilt (*Fusarium* spp.) in Spinach Grown For Seed

THIS LABEL FOR PROLINE 480 SC FUNGICIDE EXPIRES AND MUST NOT BE DISTRIBUTED OR USED IN ACCORDANCE WITH THIS SLN REGISTRATION AFTER DECEMBER 31, 2030.

ENVIRONMENTAL HAZARDS

This product is toxic to estuarine/marine invertebrates, and freshwater/estuaries/marine aquatic plants. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of prothioconazole and degradates from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Degradates of prothioconazole are known to leach through soil into groundwater under certain conditions as a result of label use. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the EPA registered label.

Spinach Grown for Seed Production	
PEST SUPPRESSION	RATE PER APPLICATION fluid ounces/acre
Fusarium Wilt (<i>Fusarium</i> spp.)	5.7 (0.178 lbs ai/A)
Recommended Application Timings for Spinach Seed Crops for Management of Fusarium Wilt: Apply Proline 480 SC as a banded spray or drench over the row at (or immediately after) planting. Make two repeat applications at 3-week intervals as a banded application over the rows in a large enough volume of water to drench the plants and wash the product into the soil around the roots.	

RESTRICTIONS / PRECAUTIONS

- **DO NOT** apply more than 3 applications per year with a **minimum 3-week interval** between applications.
- **DO NOT** apply within 14 days of harvest.
- **DO NOT** windrow within 14 days of last application.
- Maximum Proline 480 SC Fungicide allowed per year: **17.1 fl oz (0.534 lbs prothioconazole) per acre**
- **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
- **DO NOT** apply prothioconazole with mechanically pressurized handgun equipment.
- For use in accordance with this SLN label, **DO NOT** apply this product through any type of irrigation system.

- **Crops Grown for Seed Production:**

1. All spinach seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.
2. No portion of the spinach seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
3. Spinach seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
4. Spinach seed may not be distributed for human consumption or animal feed.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA website under the Waste Pesticide Program. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

IMPORTANT—READ BEFORE USE

PROLINE 480 SC FUNGICIDE, WHEN USED ON SPINACH GROWN FOR SEED UNDER THIS SECTION 24(C) LABEL HAS NOT BEEN TESTED UNDER ALL CONDITIONS AND IN ALL TANKMIX SCENARIOS. BAYER RECOMMENDS THAT THE USER AND/OR GROWER TEST PROLINE 480 SC FUNGICIDE IN ORDER TO DETERMINE ITS COMPATIBILITY WITH THE INTENDED USE. **RISK OF CROP INJURY, CROP YIELD REDUCTION, AND/OR CROP LOSS ARE BEYOND THE CONTROL OF BAYER, AND BAYER INTENDS THAT ALL SUCH USE IS AT USER'S AND/OR GROWER'S RISK, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW.**

THIS SPECIAL CONDITIONS AND DISCLAIMER STATEMENT IS REQUIRED BY BAYER AND IS NOT REQUIRED OR ENFORCED BY THE USEPA OR THE WSDA.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

**Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)**