

RESTRICTED USE PESTICIDE

TOXIC TO FISH AND AQUATIC ORGANISMS

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification

FIFRA Section 24(c) Special Local Need (SLN) Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

Bifender FC

For Soil Application to Established Plantings of Blueberry and Raspberry to Control Garden Symphylans.

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

EPA Reg. No. 89118-2

EPA SLN No. WA-260009

This label for Bifender FC expires and must not be distributed or used in accordance with this SLN registration after **December 31, 2027**.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of application.
- Read the label affixed to the container for Bifender FC before applying.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the EPA registered label.

SPECIFIC DIRECTIONS FOR USE

ESTABLISHED BLUEBERRY

BLUEBERRY SOIL DRENCH / BANDED APPLICATIONS

PEST	USE RATES	
	fl. oz. product/acre (lb. a.i./acre)	fl. oz. product/1000 row ft. (lb. a.i./1000 row ft.)
Garden Symphylans	5.9 - 7.4 (0.08 - 0.1)	0.34 - 0.42 (0.0046 - 0.0057)

Apply Bifender FC when garden symphylans are present.

Soil Drench and Banded Instructions:

- Apply as a soil-directed band or drench application in established plantings in 35 gallons per acre of final spray volume.

Chemigation Instructions:

- This product may be applied through irrigation systems. Refer to the EPA registered label (Bifender FC, EPA Reg.

No. 89118-2) and the WSDA Chemigation Guidance section on this SLN label for additional chemigation directions.

- Apply via drip irrigation, applying it through the middle third of the irrigation run.

Specific Use Restrictions:

- **DO NOT** make more than 5 foliar applications per year. A soil application under this SLN registration counts as one foliar application.
- **DO NOT** make applications less than 7 days apart.
- **DO NOT** apply more than 0.5 pound bifenthrin per acre per year including at-plant, PPI & PRE, and foliar applications of Bifender FC and other bifenthrin containing products.
- **DO NOT** apply within 1 day of harvest.
- Buffer zone requirement: **DO NOT** apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).
- Vegetative filter strip requirement: Refer to the “Vegetative Filter Strips” section of the federal label.
- For use in accordance with this SLN label do not apply this product by air.

ESTABLISHED RASPBERRY

RASPBERRY SOIL DRENCH / BANDED APPLICATIONS

PEST	USE RATES	
	fl. oz. product/acre (lb. a.i./acre)	fl. oz. product/1000 row ft. (lb. a.i./1000 row ft.)
Garden Symphylans	5.9 - 7.4 (0.08 - 0.1)	0.34 - 0.42 (0.0046 – 0.0057)

Apply Bifender FC when garden symphylans are present.

Soil Drench, and Banded Instructions:

- Apply as a soil-directed band or drench application in established planting in 35 gallons per acre of final spray volume.

Chemigation Instructions:

- This product may be applied through irrigation systems. Refer to the EPA registered label (Bifender FC, EPA Reg. No. 89118-2) and the WSDA Chemigation Guidance section on this SLN label for additional chemigation directions.
- Apply via drip irrigation, applying it through the middle third of the irrigation run.

Specific Use Restrictions:

- **DO NOT** make more than 2 foliar applications per year. A soil application under this SLN registration counts as one foliar application.
- **DO NOT** apply more than 0.2 pound active ingredient per acre per year including at-plant, PPI & PRE, and foliar applications of Bifender FC and other bifenthrin containing products.
- **DO NOT** apply within 3 days of harvest.
- Buffer zone requirement: **DO NOT** apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).
- Vegetative filter strip requirement: Refer to the “Vegetative Filter Strips” section of the federal label.
- For use in accordance with this SLN label do not apply this product by air.

WSDA Chemigation Guidance

Application off-site is prohibited. The chemigation application must be continuously observed whenever sensitive areas as defined in WAC 16-202-110(47) (including but not limited to schools, parks, dwellings, occupied buildings or structures, public roadways, and waters of the state) are at risk of being exposed to drift, runoff, or overspray. In order to minimize the

potential for application off-site, WSDA recommends that the product only be applied through low-pressure irrigation systems (defined as 5 to 30 pounds per square inch measured at the nozzle) with a nozzle release height no higher than 3 feet above the target crop, and that end guns be disabled throughout the application.

An inspection port or a direct access point is required, and it must be positioned immediately upstream of the irrigation mainline check valve and be of sufficient size to allow visual and manual inspection of the check valve and low-pressure drain. The inspection port or access point must have a minimum diameter of 4 inches, unless an alternative access system is approved by WSDA (WAC 16-202-310(3)(b)).

The chemigation application tank cannot be placed within 20 feet of and must be down gradient from wellheads or other sensitive areas. Mixing or loading activities cannot occur within 20 feet of the wellhead or other sensitive areas (WAC 16-202-250(2)).

The operator of the chemigation system must read and understand the WA State Chemigation Rules (WAC 16-202-100 through WAC 16-202-410) prior to conducting a chemigation application and must adhere to the requirements contained therein. The rules can be accessed from the Washington State Legislature website (<https://app.leg.wa.gov/wac/default.aspx>).

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA website under the Waste Pesticide Program. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

24(c) Registrant:
Vive Crop Protection
500 Westover Dr., 10198
Sanford, NC 27330