

APPENDIX K: GIS Maps and Spatial Analyses

I. Labeled Uses for chlorpyrifos

The following use list is derived from label use information. It is used as a basis for the spatial analysis of chlorpyrifos.

Table 1. Use list from labels	
NLCD Category	Use
Cultivated crops	Alfalfa, sorghum (milo), soybeans, asparagus, beans (asparagus, road, dried, hyacinth, moth, mung, rice, lima, snap, tepary, urd), broccoli (raab, Chinese), brussels sprouts, cabbage (Mustard, Chinese), pea (Catjang/Jerusalem/Marble, cowpea, blackeye, dried, early dwarf, field, pigeon, southern, succulent), cauliflower, Christmas tree plantations, cole crops, collards, corn (field, sweet), cotton, cranberry, cucumber, garbanzos, kale, kohlrabi, legume vegetables, lentils, lupine (grain), mint, onion, peanut, pumpkin, radish, rutabaga, strawberry, sugar beet, sunflower, sweet potato, tobacco, turnip (greens, root), wheat, guar
Turf	Commercial/industrial lawns, industrial areas, golf course, ornamental grasses, ornamental herbaceous plants, ornamental non-flowering plants, ornamental sod farm
Rights-of-way	Non-agricultural rights-of-way, fencerows
Orchard/vineyard	Almond, apple, calamondin, cherry, citron, citrus (hybrids), deciduous fruit trees, filbert (hazelnut), fruits, grapefruit, grapes, kumquat, lemon, lime, nectarine, orange, orchards, peach, conifers (seed orchard), ornamental and/or shade trees, ornamental woody shrubs and vines, fig, seed orchard trees, pear, pecan, plum, prune, pummelo (shaddock), tangelo, walnut (English/black)

II. Initial Area of Concern

After determining which uses will be assessed from label information, an evaluation of the potential ‘footprint’ of use patterns is determined. The ‘footprint’ includes all areas within the state of California where the pesticide could be applied (**Figure 1**). The footprint of potential use represents the chemical’s initial area of concern, and is typically based on available land cover data. Uses that are not represented through available land cover data are not displayed spatially, as their extent cannot be defined using existing categories of land cover classes (*e.g.* dumpsters). The initial area of concern represents the starting point from which the action area is defined.

A. Land Cover

Base mapping land cover layers for the initial area of concern analysis were obtained from the National Land Cover Dataset (NLCD 2001) for the majority of land use types. The NLCD was

released as a nationally consistent, regionally indexed dataset in January 2007. California GAP data from the Biogeography Lab from UCLA-Santa Barbara (1998) were obtained for the orchard and vineyard uses. These raster files were converted to vectors using simplification and majority filter routines, and used in the analysis. The rights-of-way land cover layer was derived by combining road and rail information from TeleAtlas (2007) with U.S. Department of Transportation's National Pipeline Mapping System (1999). **Table 2** shows the land-cover sources used for the initial area of concern analysis.

Table 2. Land Cover Data Sources


Layer name	Base source	Description
Cultivated Crops	NLCD	Grid code 82: Areas used for the production of annual crops, such as corn, soybeans, vegetables, tobacco, and orchards/vineyards, and also perennial woody crops such as orchards and vineyards. Crop vegetation accounts for greater than 20 percent of total vegetation. This class also includes all land being actively tilled.
Developed, High Intensity	NLCD	Grid code 24: Includes highly developed areas where people reside or work in high numbers. Examples include apartment complexes, row houses and commercial/industrial. Impervious surfaces account for 80 to 100 percent of the total cover.
Developed, Low Intensity	NLCD	Grid code 22: Includes areas with a mixture of constructed materials and vegetation. Impervious surfaces account for 20-49 percent of total cover. These areas most commonly include single-family housing units.
Developed, Medium Intensity	NLCD	Grid code 23: Includes areas with a mixture of constructed materials and vegetation. Impervious surfaces account for 50-79 percent of the total cover. These areas most commonly include single-family housing units.
Developed, Open Space	NLCD	Grid code 21: Includes areas with a mixture of some constructed materials, but mostly vegetation in the form of lawn grasses. Impervious surfaces account for less than 20 percent of total cover. These areas most commonly include large-lot single-family housing units, parks, golf courses, and vegetation planted in developed settings for recreation, erosion control, or aesthetic purposes.
Forest	NLCD	Grid codes 41, 42, 43: Deciduous, evergreen and mixed. Areas dominated by trees generally greater than 5 meters tall, and greater than 20% of total vegetation cover.
Open Water	NLCD	Grid code 11: All areas of open water, generally with less than 25% cover of vegetation or soil.
Orchards and vineyards	CA GAP	Grid codes 11210, 11211 and 11212. This is the only CA GAP reference.
Pasture/Hay	NLCD	Grid code 81: Areas of grasses, legumes, or grass-legume mixtures planted for livestock grazing or the production of seed or hay crops, typically on a perennial cycle. Pasture/hay vegetation accounts for greater than 20 percent of total vegetation.
Wetlands	NLCD	Grid codes 90, 95: Woody wetlands and emergent herbaceous.
Turf	NLCD	A derived NLCD class based on developed classes and the impervious surface layer with corrections applied.
Rights-of-way	US DOT; TeleAtlas	A derived class using road, rail, and pipeline coverages.

B. Initial Stream Reaches

In addition to the land cover classes described above, the initial area of concern includes the stream segments found within those land cover areas. The stream segments are obtained from the NHDPlus dataset. For each stream reach in the hydrography network, the data provide a tally of the total area in each NLCD land cover class for the upstream cumulative area contributing to the given stream reach. Using the cumulative land cover data provided by the NHDPlus (<http://www.horizon-systems.com/nhdplus/>), an aggregated use class is created based on the classes listed in Table 1. A cumulative percent cropped area (PCA) is calculated for each stream reach based on the aggregate use class (divided by the total upstream contribution area). Pesticide exposures in the streams within the initial area of concern are conservatively assumed to be represented by the estimated environmental concentrations used in RQ calculation.

Figure 1.

Initial Area of Concern for Chlorpyrifos



III. Spatial Extent of the Effects Determination

Based on the results of the Agency's endangered species risk assessment for chlorpyrifos, a Likely to Adversely Affect (LAA) and modification to critical habitat determination was concluded for the California red-legged frog (CRLF). A Likely to Adversely Affect (LAA) and modification to critical habitat determination was also concluded for the Delta smelt (DS), California tiger salamander (CTS), Bay checkerspot butterfly (BCB), and the Valley elderberry longhorn beetle (VELB). A Likely to Adversely Affect (LAA) determination was concluded for the San Francisco garter snake (SFGS), California clapper rail, (CCR), Salt marsh harvest mouse (SMHM), San Joaquin kit fox (SJKF), and California freshwater shrimp (CFS). The spatial extent of the effects determination is based on the initial area of concern for application of chlorpyrifos on cultivated crops and developed land, and expanded to include the total area where there is potential for direct or indirect effects to occur via off-site transport mechanisms. The extent of potential off-site transport is determined by deriving the spray drift area and the run-off area using the downstream dilution model and AgDRIFT.

The identified direct and indirect effects are anticipated to occur only for those currently occupied core areas, CNDDDB occurrence sections, and areas of designated critical habitat for the CRLF, and occurrence sections and areas of designated critical habitat for the San Francisco Bay Species (DS, CTS, BCB, VELB, SFGS, CCR, SMHM, SJKF and CFS) that are located 1,000 feet from legal use sites where chlorpyrifos is applied to cultivated crops and developed land. AgDRIFT and/or AgDISP spray drift models are used to determine the distance from the initial area of concern where no direct or indirect effects are expected to occur due to exposure to spray drift. It is assumed that lentic (*i.e.*, non-flowing) waterbodies (or potential CRLF habitat) are included within the spray drift area. Spray drift analysis is discussed in **Section 5.4**. The downstream extent analysis indicates the furthest distance that could be added downstream is 285 kilometers. This distance is representative of the maximum continuous downstream dilution from the edge of the initial area of concern where direct/indirect effects and/or critical habitat modification may occur.

There are several types of habitat areas: critical habitat (CH), currently occupied core areas (for the CRLF only), the California Natural Diversity Database (CNDDDB) occurrence sections identified from *Center for Biological Diversity (CBD) vs. EPA et al.* (Case No. 07-2794-JCS), (shown in yellow, **Figures 2-10**), and recovery plan locations depicting either habitat point locations (**Figures 3,4,6,8, and 10**) or river segments (**Figure 7**).

The overlap of land cover corresponding to chlorpyrifos use patterns that result in an LAA determination and CRLF habitat for the entire state of California is shown in Figures 11-20. The overlap maps provides a depiction of land cover that corresponds to "LAA" chlorpyrifos use patterns, overlapped with habitat; however, the actual area of overlap is expected to be greater when off-site transport via spray drift is included for each land cover type. The buffers may be different for each land cover type due to varying application rates and/or methods for different use patterns. Further analysis of the extent of drift for each land cover type and the overlap with habitat can be included as part of the consultation process, if needed.

Figure 2.

Delta Smelt Habitat




Figure 3

California Clapper Rail Habitat




Figure 4

Salt Marsh Harvest Mouse Habitat




Figure 5

California Tiger Salamander Habitat




Figure 6

SF Garter Snake Habitat




Figure 7

California Freshwater Shrimp Habitat




Figure 8

San Joaquin Kit Fox Habitat




Figure 9

Valley Elderberry Longhorn Beetle Habitat




Figure 10

Bay Checkerspot Butterfly Habitat




Figure 11

Delta Smelt with Potential Chlorpyrifos Use




Figure 12

CA Clapper Rail with Potential Chlorpyrifos Use




Figure 13

Salt Marsh Harvest Mouse with Potential Chlorpyrifos Use




Figure 14

California Tiger Salamander with Potential Chlorpyrifos Use




Figure 15

San Francisco Garter Snake with Potential Chlorpyrifos Use




Figure 16

CA Freshwater Shrimp with Potential Chlorpyrifos Use




Figure 17

San Joaquin Kit Fox with Potential Chlorpyrifos Use




Figure 18

Valley Elderberry Longhorn Beetle with Potential Chlorpyrifos Use




Figure 19

Bay Checkerspot Butterfly with Potential Chlorpyrifos Use





Figure 20

CA Red-legged Frog and Potential Chlorpyrifos Use Overlap



IV. A Note on Limitations and Constraints of Tabular and Geospatial Sources

The geographic data sets used in this analysis are limited with respect to their accuracy and timeliness. The National Land Cover Data Set (NLCD 2001) represents the most current and comprehensive collection of national land use and land cover information for the United States and represents land cover data obtained between 1994-1998. Two additional data sets were included to account for uses not clearly defined by the NLCD. These supplemental data include orchard and vineyard land cover data from the California Gap Analysis Project data (CaGAP 1998), and rights-of-way data derived from the Teleatlas (2007).

Hydrographic data are from the NHDPlus data set (<http://www.horizon-systems.com/nhdplus/>). NHDPlus contains the most current and accurate nationwide representation of hydrologic data. At a spatial scale of 1:100,000, the NHDPlus might omit the smallest streams and waterbodies. In addition, in some isolated instances, there are errors in the data including missing or disconnected stream segments and incorrect assignment of flow direction.

The relatively coarse spatial scale and general classification categories of these data sets preclude use of the data for highly localized study. Additionally, some labeled uses are not possible to map precisely due to the lack of appropriate spatial data in NLCD on the location of these areas.

OPP will continue to endeavor to identify and incorporate (as appropriate) additional land cover data sets for other land classes not captured in this assessment. In addition, as new updates to existing data occur, these will also be evaluated and incorporated as appropriate.

V. References for GIS Maps

Crop Maps

GAP. Gap Analysis. National Biological Information Infrastructure. www.nbii.gov

MRLC, 2001. Multiresolution Land Characteristics (MRLC) www.mrlc.gov

EPA Environmental Fate and Effects Division NLCD-derived turf.

EPA Environmental Fate and Effects Division road-derived rights-of-way.

EPA Environmental Fate and Effects Division turf layer.

Physical Maps

ESRI, 2002. Detailed Counties, ESRI data and maps. (1:24,000) www.esri.com

NHD Plus Region 18 (v01), National Hydrography Dataset, nhd.usgs.gov

Habitat Maps

Center for Biological Diversity vs EPA Case No. 07-2794-JCS.

US FWS 2002 California red-legged frog General Recovery Zones

US FWS 2002 California red-legged frog Core Areas

US FWS 2005 Final Critical Habitat for California red-legged frog

California Red-legged Frog CNDDB Occurrence Sections – California Natural Diversity Database <http://www.dfg.ca.gov/bdb/html/cnddb.html>

Critical habitat maps (Delta Smelt, Calif. Tiger Salamander, Valley Elderberry Longhorn Beetle, Bay Checkerspot Butterfly, Alameda Whipsnake, Tidewater Goby) from crithab.fws.gov

Delta Smelt Recovery Plan, 1996. ecos.fws.gov

California Clapper Rail Recovery Plan, 1984. ecos.fws.gov

Salt Marsh Harvest Mouse Recovery Plan, 1984. ecos.fws.gov

San Francisco Garter Snake, 1985. ecos.fws.gov

California Freshwater Shrimp, 1998. ecos.fws.gov

San Joaquin Kit Fox, 1998. ecos.fws.gov

Bay Checkerspot Butterfly, 1998. ecos.fws.gov