

## **APPENDIX E: The Risk Quotient Method and Levels of Concern**

The Risk Quotient Method is the means used by EFED to integrate the results of exposure and ecotoxicity data. For this method, Risk Quotients (RQs) are calculated by dividing exposure estimates by the acute and chronic ecotoxicity values (i.e.,  $RQ = \text{EXPOSURE}/\text{TOXICITY}$ ). These RQs are then compared to OPP's levels of concern (LOCs). These LOCs are criteria used by OPP to indicate potential risk to non-target organisms and the need to consider regulatory action. EFED has defined LOCs for acute risk, potential restricted use classification, and for endangered species.

The criteria indicate that a pesticide used as directed has the potential to cause adverse effects on non-target organisms. LOCs currently address the following risk presumption categories:

- (1) acute - there is a potential for acute risk; regulatory action may be warranted in addition to restricted use classification;
- (2) acute restricted use - the potential for acute risk is high, but this may be mitigated through restricted use classification;
- (3) acute endangered species - the potential for acute risk to endangered species is high, regulatory action may be warranted; and
- (4) chronic risk - the potential for chronic risk is high, regulatory action may be warranted.

Currently, EFED does not perform assessments for chronic risk to plants, acute or chronic risks to non-target insects, or chronic risk from granular/bait formulations to mammalian or avian species.

The ecotoxicity test values (i.e., measurement endpoints) used in the acute and chronic RQs are derived from required studies. Examples of ecotoxicity values derived from short-term laboratory studies that assess acute effects are: (1)  $LC_{50}$  (fish and birds), (2)  $LD_{50}$  (birds, mammals and terrestrial invertebrates), (3)  $EC_{50}$  (aquatic plants and aquatic invertebrates), and (4)  $EC_{25}$  (terrestrial plants). Examples of toxicity test effect levels derived from the results of long-term laboratory studies that assess chronic effects are: (1) the Lowest Observed Adverse Effect Concentration (LOAEC) (birds, fish, and aquatic invertebrates), and (2) the No Observed Adverse Effect Concentration (NOAEC) (birds, fish and aquatic invertebrates). The NOAEC is generally used as the ecotoxicity test value in assessing chronic effects.

Table A Risk Presumptions and LOCs		
Risk Presumption	RQ	LOC
Birds <sup>1</sup>		
Acute Risk	EEC/LC <sub>50</sub> or LD <sub>50</sub> /sqft or LD <sub>50</sub> /day	0.5
Acute Restricted Use	EEC/LC <sub>50</sub> or LD <sub>50</sub> /sqft or LD <sub>50</sub> /day (or LD <sub>50</sub> < 50 mg/kg)	0.2
Acute Endangered Species	EEC/LC <sub>50</sub> or LD <sub>50</sub> /sqft or LD <sub>50</sub> /day	0.1
Chronic Risk	EEC/NOEC	1
Wild Mammals <sup>1</sup>		
Acute Risk	EEC/LC <sub>50</sub> or LD <sub>50</sub> /sqft or LD <sub>50</sub> /day	0.5
Acute Restricted Use	EEC/LC <sub>50</sub> or LD <sub>50</sub> /sqft or LD <sub>50</sub> /day (or LD <sub>50</sub> < 50 mg/kg)	0.2
Acute Endangered Species	EEC/LC <sub>50</sub> or LD <sub>50</sub> /sqft or LD <sub>50</sub> /day	0.1
Chronic Risk	EEC/NOEC	1
Terrestrial Invertebrates		
Acute Endangered Species Risk	EEC/LD <sub>50</sub>	0.05
Aquatic Animals <sup>2</sup>		
Acute Risk	EEC/LC <sub>50</sub> or EC <sub>50</sub>	0.5
Acute Restricted Use	EEC/LC <sub>50</sub> or EC <sub>50</sub>	0.1
Acute Endangered Species	EEC/LC <sub>50</sub> or EC <sub>50</sub>	0.05
Chronic Risk	EEC/NOEC	1
Terrestrial and Semi-Aquatic Plants		
Acute Risk	EEC/EC <sub>25</sub>	1
Acute Endangered Species	EEC/EC <sub>05</sub> or NOEC	1
Aquatic Plants <sup>2</sup>		
Acute Risk	EEC/EC <sub>50</sub>	1
Acute Endangered Species	EEC/EC <sub>05</sub> or NOEC	1

<sup>1</sup> LD<sub>50</sub>/sqft = (mg/sqft) / (LD<sub>50</sub> \* wt. of animal)

LD<sub>50</sub>/day = (mg of toxicant consumed/day) / (LD<sub>50</sub> \* wt. of animal)

<sup>2</sup> EEC = (ppm or ppb) in water