

# Effect of Connecticut Yankee Power Plant on Population Dynamics of Asiatic Clams and Their Interactions with Native Bivalves

DOUGLAS E. MORGAN\*, MILAN KESER, JOHN T. SWENARTON, AND JAMES F. FOERTCH

*Millstone Environmental Laboratory, Dominion Nuclear Connecticut, Inc.  
Waterford, Connecticut 06385, USA*

*Abstract.*—Asian clam *Corbicula fluminea*, first reported in the Connecticut River in 1990, was the dominant bivalve around Connecticut Yankee nuclear power station (CY) during the 10-year period (1991–2000) of this study. This population extended its range upriver at least 22 km from 1991 to 1992. *Corbicula* winter survival was positively correlated with winter water temperatures and negatively correlated with spring freshet river flows. For example, no *Corbicula* were found following the two coldest winters of the study (1993–1994 and 1995–1996). Regardless of the level observed for winter population survival, study sites were repopulated each year by August, attributed to passive downstream movement from populations in refugia from winter stresses (e.g., upstream thermal discharges at Middletown and South Meadow power stations, or groundwater seeps). Small clams (2-mm size-class) dominated the CY thermal discharge canal during power plant operation because, by June of each year, discharge temperatures (>35°C) exceeded the upper thermal tolerance for this clam. The CY discharge, therefore, was not an important winter refuge. In years following CY closure (1997–2000), a year-round *Corbicula* population was established in the discharge canal, but abundance at river sites around CY remained similar to levels observed during power plant operation. Our findings suggest that *Corbicula* has established a permanent population in the Connecticut River with little impact on native bivalves up to the last year of this study in 2000.

## Introduction

Asian clam *Corbicula fluminea* is a freshwater clam, native to Southeast Asia, now common in Europe, Africa, the Pacific Islands, and North and South America (McMahon 1982; Counts 1986). In its native range in Asia, *C. fluminea* (termed *Corbicula* throughout this report) is a commercial species collected and cultured for human consumption (Chen 1990; Phelps 1993). Once introduced to North America, this clam spread rapidly because of its high reproductive capacity, adult brooding of young to a free-living juvenile stage, fast growth, early maturity, effectiveness of all stages of development to readily disperse by water currents, and tolerance to a wide range of environmental stresses (Gardner et al. 1976; Eng 1979; Sickel 1979, 1986; McMahon 1983). Its

ability to clog industrial water systems makes *Corbicula* a very costly problem for water-dependent industries (Isom 1986; McMahon and Lutey 1988). By fouling water systems at electric generating stations, particularly nuclear power stations, *Corbicula* causes significant safety problems as well (Ingram 1959; Goss and Cain 1975; McMahon 1977; Goss et al. 1979; Mattice 1979; USNRC 1981; Johnson et al. 1986; Page et al. 1986).

*Corbicula* were initially identified at Connecticut Yankee nuclear power station (CY), which is located on the lower Connecticut River (Figure 1), during a cleaning of the service water system in April 1990 (Morgan et al. 1992, 2003). The discovery of *Corbicula* at CY necessitated the development of an effective control methodology. *Corbicula* is highly resistant to short-term application of conventional biocides because of its ability to tightly close its valves during treat-

\* Corresponding author: douglas\_e\_morgan@dom.com




Figure 1. *Corbicula fluminea* sampling sites used in surveys conducted in November 1991 and 1992 to determine the extent of up-river clam abundance.

ment (Mattice et al. 1982; Neitzel et al. 1984; Page et al. 1986). However, continuous chlorination is effective in controlling *Corbicula* (Isom 1971; Goss and Cain 1975; Cherry et al. 1986). The continuous low level (<0.1 ppm, Total Residual Chlorine) chlorination of the service water system at CY was approved by the Connecticut Department of Environmental Protection (CTDEP) and began in April 1991. As a condition of the permit, CTDEP required CY to initiate a monitoring study of the *Corbicula* population in the lower Connecticut River. This paper summarizes the results of studies from 1991 to 2000, including data for *Corbicula* distribution, growth, fecundity, and abundance. In addition, native clam and mussel abundance data are presented and examined for possible interactions with *Corbicula*.

## Methods

Surveys of the distribution of Connecticut River *Corbicula* population were conducted in November of 1991 and 1992. In 1991, sampling began at CY and continued upriver at intervals of 3–10 km (Middletown station, Wilcox Island Park, Gildersleeve Island, Glastonbury Ferry, Crow Point, and South Meadow station; Figure 1). In 1992, sampling was extended upriver to the Bulkeley Bridge and Hartford Landfill area. In these surveys, sediments were screened using a 3.4 mm, U.S.A. Standard Testing Sieve, which retained clams larger than approximately 6 mm in shell length. All sampling sites were located in the tidal freshwater areas of the Connecticut River.

*Corbicula* abundance, growth, reproduction and recruitment were studied at five sites in the vicinity of CY from August 1991 to November 2000. Four sampling sites were located in the mainstem river (Figure 2): Haddam Island (A); vicinity of CY-intakes (B); down-river of CY-intakes (C); and in the vicinity of the CY discharge canal outfall (D). A fifth site was located at the head of the CY discharge canal (E). Water depth at sampling sites ranged from 1 to 6 m at mean low water.

Sampling began in August 1991 and continued annually in May, August, and November through November 2000. Sampling at site D did not start until August 1993. Five 0.1-m<sup>2</sup> bottom samples were obtained at each site using a




Figure 2. Location of the sampling sites (A–E) in the vicinity of CY.

weighted Peterson Grab. Sample processing techniques were similar to those of Gardner et al. (1976). Grab samples were sieved in the field by passing the sample through a series of three screens (6.3-, 2.0-, and 1.0-mm mesh sizes). Clams and sediment retained on the 1-mm screen were subsampled in the field by placing a well-mixed 1-L sample in an elutriator (Magdych 1981) for 3 min at a water flow of 20–30 L/min. The overflow from the elutriator was collected on a 1.0-mm mesh sieve and sorted in the laboratory under a dissecting microscope (10 $\times$ ). Sediment and clams retained on the 6.3- and 2.0-mm screens were taken to the laboratory and washed through a series of five U.S.A. Standard Testing Sieves (19.0-, 12.5-, 6.3-, 3.4-, and 2.0-mm mesh sizes). Size-classes were based on the sieve size in which clams were retained. Clams retained on the 1.0-mm sieve averaged 2.0 mm in shell length; on the 2.0-mm sieve, 4.1 mm; on the 3.4-mm sieve, 6.7 mm; on the 6.3-mm sieve, 14.1 mm; on the 12.5-mm sieve, 19.3 mm; and on the 19-mm sieve, 31.1 mm. The calculated densities (clams/m<sup>2</sup>) are used in this study as a measure of abundance. Native bivalve species collected during the processing of these sediment samples were reported only at the total abundance level. Winter survival of clams was defined as the ratio of May clam abundance to November abundance from the previous year, expressed as a percentage.

Several hundred adult clams (>8.0 mm in shell length) were collected from river sites in June 1991 and in May 1992–1994 and held in two lantern-nets, one in the river and the other in the CY discharge canal. Clam growth was monitored monthly from 1991 to 1994 by measuring shell lengths of 12 randomly selected clams from these nets. Each year, monthly measurements were made until total winter mortality occurred in the river (December–February) or summer mortality occurred in the discharge canal (usually June).

Clam fecundity was monitored monthly using techniques of Aldridge and McMahon (1978). Fecundity was assessed on the 12 clams selected for growth estimates in each month that living clams could be found. Each clam was brought to the laboratory and held under static conditions at 20°C for 24 h in a 100-mL beaker filled with filtered Connecticut River water. The number of juveniles released during this period was recorded

and used as an index of spawning activity. Additional fecundity assessments were made by dissecting these clams and noting the presence of brood progeny. Maturity of gametes was assessed by removing egg and sperm cells from the gonadal tissues and examining the cells under a compound microscope (400 $\times$ ).

To estimate *Corbicula* recruitment to bottom sediments in the Connecticut River and CY discharge canal, plastic trays (34.5  $\times$  47.0  $\times$  12.5 cm, inside dimensions) were filled with defaunated marine beach sand (<1 mm in grain size). Two sets of four sediment trays were enclosed within wooden frames (84 cm  $\times$  112 cm, outside dimensions) partitioned into four compartments (37.0 cm wide  $\times$  48.0 cm long  $\times$  14.0 cm high) and anchored to the bottom near the mouth of the discharge canal on 30 May 1991 and in the Connecticut River just upriver of the CY intakes (site B) on 6 June 1991. At each site, four trays were placed in each frame; two trays were covered with 6.4  $\times$  6.4 mm plastic mesh screening, and two were not covered. The screen-covered trays were stocked with 50 adult clams (>8 mm), and the uncovered trays were left barren. One replicate tray for each treatment (screen-covered and uncovered) was collected and screened for *Corbicula* during river surveys in August and November 1991. Nearly all clams were dead in May 1992 and sampling was terminated.

Connecticut River flow rates were obtained from the USGS Thompsonville CT monitoring station (USGS 2002). Water temperature was recorded every 15 min at the CY intake and discharge using the Northeast Utilities Environmental Data Acquisition Network. Statistical analyses were performed using SAS version 8 software (SAS Inc., Cary, North Carolina). Abundance data in tables are presented using arithmetic means. Statistical testing (SAS General Linear models Procedure) of abundance data were performed after log transformation. *T*-tests were performed on log-transformed abundance data, and depending on the results for the equality of variances test, either a pooled or Satterthwaite method was used to determine degrees of freedom. The relationships between winter clam survival and temperature or flow indices were tested using the rank-order Spearman correlation statistic. Growth and reproduction data were not transformed prior to analysis and are presented as arithmetic means.

## Results

### River Water Flow and Temperature

Annual Connecticut River water flow averaged 509 m<sup>3</sup>/s over the 10-year period from 1991 to 2000. Mean monthly flow rates peaked during April in 9 of 10 years with the average April flow rate ranging from a low of 422 m<sup>3</sup>/s in 1995 to a high of 2,099 m<sup>3</sup>/s in 1993 (Figure 3). Similarly, maximal flow rates for daily averages usually occurred in April and often exceeded 1,700 m<sup>3</sup>/s. These values, which were collected at the Thompsonville Connecticut gaging station, represent flows beyond any tidal effects. Semidiurnal tides affect river flow rates and bring an average 425 m<sup>3</sup>/s of additional flow to the lower Connecticut River in the vicinity of CY (Merriman and Thorpe 2004 [1976], this volume), causing periodic fluctuations in river height of ~1 m (NSI 1995; Rozsa 2001). Tidal influences are large in relation to natural flow during periods of low river discharge and are absent or nearly absent during freshet conditions (Rozsa 2001; Boyd 2004

[1976], this volume). Daily average ambient water temperatures ranged between -1.7°C and 30.6°C during the 10-year study period (Figure 4). Discharge water temperatures at CY during plant operation were 8–12°C above ambient river temperatures. The maximum discharge flow of 25 m<sup>3</sup>/s was directed through a man-made canal 1 km long before mixing with the ambient Connecticut River waters.

### Distribution

In November 1991, *Corbicula* was observed as far north in the Connecticut River as Gildersleeve Island, where density averaged 168/m<sup>2</sup> (Figure 1; Table 1). Gildersleeve Island is approximately 16 km upstream of CY, where density averaged 531/m<sup>2</sup>. Between these sites, clam densities were 150/m<sup>2</sup> at Middletown station and 12/m<sup>2</sup> at Wilcox Island Park. In November 1992, clams were documented 22 km further north at South Meadow station (30/m<sup>2</sup>), which is just south of Hartford,




Figure 3. Monthly mean flow of the Connecticut River as measured at the Thompsonville station from January 1991 through December 2000. Black bars are April mean flows.




Figure 4. Intake (—) and discharge (---) water temperatures at CY from January 1, 1991 to January 1, 2001. Horizontal reference lines represent upper and lower lethal temperature limits for *Corbicula fluminea*. CY shut down in July 1996, as revealed by discharge temperatures tracking intake temperatures.

Connecticut. Clam densities were higher in 1992 than in 1991 at all sites, from 28/m<sup>2</sup> at Crow Point, approximately 3 km downstream of South Meadow station, to 9,169/m<sup>2</sup> at Middletown station.

Table 1. Mean abundance of *Corbicula fluminea* during the November 1991 and 1992 Connecticut River surveys.

| Location                | Abundance <sup>a</sup>       |                              |
|-------------------------|------------------------------|------------------------------|
|                         | 1991<br>clams/m <sup>2</sup> | 1992<br>clams/m <sup>2</sup> |
| Connecticut Yankee (CY) | 531                          | 2,182                        |
| Middletown station      | 150                          | 9,169                        |
| Wilcox Island Park      | 12                           | — <sup>b</sup>               |
| Gildersleeve Island     | 168                          | 3,300                        |
| Glastonbury Ferry       | 0                            | —                            |
| Crow Point              | 0                            | 28                           |
| South Meadow station    | 0                            | 30                           |
| Bulkeley Bridge         | —                            | 0                            |
| Hartford landfill       | —                            | 0                            |

<sup>a</sup> Abundance estimates for clams > 5.6 mm in shell length (3.4-mm screen).

<sup>b</sup> Site was not sampled.

## Abundance

Mean annual total abundance of *Corbicula* at CY sites from 1991 to 2000 was highly variable with significantly ( $P < 0.05$ ) higher abundance in 1992, 1995, and 1999 than any other year of the study, ranging from 1,158/m<sup>2</sup> in 1995–2,610/m<sup>2</sup> in 1992 (Table 2). Annual abundance in 1998 and 2000 was 326 and 286/m<sup>2</sup>, respectively. These abundances were significantly lower ( $P < 0.05$ ) than those in 1992, 1995 and 1999, but significantly higher ( $P < 0.05$ ) than those in 1993, 1994, 1996, and 1997, which ranged from 45/m<sup>2</sup> in 1996 to 138/m<sup>2</sup> in 1997.

Seasonal abundance (i.e., abundance in May, August, and November collections) was also highly variable (Table 2). May abundance was consistently the lowest value observed during the year; 10-year averages for May abundance ranged from 24/m<sup>2</sup> at site B to 86/m<sup>2</sup> at site A. Averages for all sites combined show November abundance was highest in 8 of 10 years. November abundance was lowest at site D (365/m<sup>2</sup>) and highest at site A (2,299/m<sup>2</sup>).

Table 2. Mean total abundance (clams/m<sup>2</sup> ± 95% CI) of *Corbicula fluminea* at each sampling site near the Connecticut Yankee power plant in the Connecticut River from August 1991 to November 2000.

| Site                         | Year        |                |           |           |               |           |           |             |               |             |             |  | All years |  |
|------------------------------|-------------|----------------|-----------|-----------|---------------|-----------|-----------|-------------|---------------|-------------|-------------|--|-----------|--|
|                              | 1991        | 1992           | 1993      | 1994      | 1995          | 1996      | 1997      | 1998        | 1999          | 2000        | 2000        |  |           |  |
| <b>A</b>                     |             |                |           |           |               |           |           |             |               |             |             |  |           |  |
| May                          | –           | 124 ± 115      | 12 ± 33   | 0         | 202 ± 161     | 0         | 10 ± 21   | 104 ± 147   | 146 ± 44      | 174 ± 284   | 86 ± 36     |  |           |  |
| Aug.                         | 640 ± 290   | 5,552 ± 3,980  | 8 ± 10    | 54 ± 59   | 3,052 ± 2,369 | 44 ± 52   | 118 ± 120 | 630 ± 440   | 4,076 ± 1,500 | 650 ± 477   | 1,482 ± 633 |  |           |  |
| Nov.                         | 1,642 ± 724 | 11,158 ± 4,315 | 332 ± 347 | 576 ± 341 | 2,452 ± 1,768 | 60 ± 50   | 648 ± 373 | 1,498 ± 168 | 3,552 ± 2,079 | 1,036 ± 238 | 2,299 ± 959 |  |           |  |
| Average <sup>a</sup>         | 1,141       | 5,611          | 117       | 210       | 1,902         | 35        | 271       | 744         | 2,591         | 620         | 1,331       |  |           |  |
| <b>B</b>                     |             |                |           |           |               |           |           |             |               |             |             |  |           |  |
| May                          | –           | 20 ± 26        | 0         | 0         | 30 ± 37       | 0         | 10 ± 28   | 14 ± 19     | 86 ± 60       | 54 ± 41     | 24 ± 11     |  |           |  |
| Aug.                         | 288 ± 99    | 982 ± 346      | 18 ± 31   | 40 ± 49   | 1,574 ± 759   | 32 ± 28   | 38 ± 27   | 196 ± 54    | 1,626 ± 346   | 196 ± 46    | 499 ± 186   |  |           |  |
| Nov.                         | 272 ± 215   | 2,688 ± 659    | 138 ± 102 | 110 ± 97  | 1,874 ± 1,829 | 80 ± 147  | 204 ± 73  | 326 ± 129   | 1,604 ± 595   | 278 ± 295   | 757 ± 289   |  |           |  |
| Average                      | 280         | 1,230          | 52        | 50        | 1,159         | 37        | 84        | 179         | 1,105         | 176         | 441         |  |           |  |
| <b>C</b>                     |             |                |           |           |               |           |           |             |               |             |             |  |           |  |
| May                          | –           | 20 ± 26        | 0         | 0         | 164 ± 200     | 0         | 8 ± 22    | 26 ± 46     | 136 ± 114     | 26 ± 45     | 42 ± 25     |  |           |  |
| Aug.                         | 270 ± 208   | 1,170 ± 645    | 40 ± 37   | 84 ± 52   | 1,934 ± 1,216 | 100 ± 64  | 42 ± 34   | 260 ± 199   | 1,590 ± 742   | 376 ± 381   | 587 ± 220   |  |           |  |
| Nov.                         | 508 ± 180   | 1,780 ± 1,078  | 140 ± 64  | 84 ± 71   | 1,184 ± 576   | 34 ± 27   | 220 ± 350 | 330 ± 121   | 956 ± 213     | 144 ± 152   | 538 ± 181   |  |           |  |
| Average                      | 389         | 990            | 60        | 56        | 1,094         | 45        | 90        | 205         | 894           | 182         | 401         |  |           |  |
| <b>D</b>                     |             |                |           |           |               |           |           |             |               |             |             |  |           |  |
| May                          | –           | –              | –         | 0         | 98 ± 73       | 0         | 4 ± 111   | 8 ± 22      | 8 ± 16        | 58 ± 71     | 25 ± 16     |  |           |  |
| Aug.                         | –           | –              | 74 ± 110  | 94 ± 61   | 754 ± 283     | 48 ± 55   | 30 ± 23   | 78 ± 54     | 1,300 ± 447   | 242 ± 161   | 328 ± 147   |  |           |  |
| Nov.                         | –           | –              | 214 ± 306 | 130 ± 143 | 580 ± 339     | 146 ± 105 | 294 ± 252 | 442 ± 56    | 922 ± 166     | 192 ± 116   | 365 ± 96    |  |           |  |
| Average                      | –           | –              | 144       | 75        | 477           | 65        | 109       | 176         | 743           | 164         | 249         |  |           |  |
| <b>All sites<sup>b</sup></b> |             |                |           |           |               |           |           |             |               |             |             |  |           |  |
| May                          | –           | 55 ± 40        | 4 ± 9     | 0         | 124 ± 56      | 0         | 8 ± 8     | 38 ± 33     | 94 ± 35       | 78 ± 58     | 45 ± 13     |  |           |  |
| Aug.                         | 399 ± 132   | 2,568 ± 1,548  | 35 ± 24   | 68 ± 22   | 1,828 ± 624   | 56 ± 22   | 57 ± 28   | 291 ± 130   | 2,148 ± 619   | 366 ± 139   | 745 ± 192   |  |           |  |
| Nov.                         | 807 ± 390   | 5,209 ± 2,647  | 206 ± 91  | 225 ± 119 | 1,522 ± 567   | 80 ± 38   | 350 ± 137 | 649 ± 273   | 1,758 ± 637   | 412 ± 189   | 1,023 ± 284 |  |           |  |
| Average                      | 603         | 2,610          | 89        | 98        | 1,158         | 45        | 138       | 326         | 1,334         | 286         | 625         |  |           |  |

<sup>a</sup> Annual averages (clams/m<sup>2</sup>) pooling the May, August, and November samples.

<sup>b</sup> Average abundance of clams, pooling the four sites.

Examination of *Corbicula* population structure from 1991 to 2000 shows juvenile clams (2- and 4-mm size-classes) significantly more abundant ( $P < 0.05$ ) than first-year adults (7- and 14-mm size-classes) or clams older than 1 year of age (19- and 31-mm size-classes, Tables 3, 4, and 5). The overall annual average for older size-classes was only 6/m<sup>2</sup> as compared to 282 and 336/m<sup>2</sup> for first-year adult and juvenile size-classes, respectively.

Abundance was also highly variable from site to site (Table 2). Mean abundance at site A (1,331/m<sup>2</sup>) was significantly higher than at other river sites. Average annual abundances (clams/m<sup>2</sup>) at sites B (441), C (401), and D (249) were not significantly ( $P < 0.05$ ) different from one another. Spatially, mean annual abundance decreased from site A (the northernmost site) to site D, which was the closest to the CY discharge canal. Mean annual abundance of *Corbicula* within the CY discharge canal (Site E) was substantially less than that at river sites and ranged from 25/m<sup>2</sup> in 1993–4,091/m<sup>2</sup> in 1992 (Table 6). Seasonally within the discharge canal, May abundance (0–92/m<sup>2</sup>) was significantly ( $P < 0.05$ ) lower than August (0–12,174/m<sup>2</sup>) and November (24–880/m<sup>2</sup>); November abundance was usually higher than in August.

Mean *Corbicula* abundance for all river sites combined was not significantly different ( $P > 0.05$ ) between operational (1991–1996) and postoperational (1997–2000) periods. Similarly, the difference between total *Corbicula* abundance in the discharge canal during operational and post-operational years was not significant ( $P > 0.05$ ). However, abundance of  $\geq 7$  mm size-class clams was significantly higher ( $P < 0.05$ ) in August and November samples during the post-operational period versus the operational period.

## Recruitment

Uncovered trays placed in the river without *Corbicula* in June 1991 were colonized by 72 clams (444/m<sup>2</sup>) in August and by 133 clams (820/m<sup>2</sup>) in November. Size of recruited clams in August ranged from 1.5 to 17.2 mm with a mean shell length of 6.8 mm, and 1.5–22.3 mm in November with a mean shell length of 9.4 mm. Screen-covered trays seeded with 50 adult-size clams/tray (14.1-mm average size) collected upstream of CY sites had 587 clams (3,620/m<sup>2</sup>) in August and 1,511 (9,318/m<sup>2</sup>) in November. In these trays,

clams in August ranged from 1.5 to 19.3 mm with a mean shell length of 7.8, and in November, the size range was 1.5–22.6 mm with a mean shell length of 11.4 mm. In May 1992, 2 live and 145 dead (shell only) clams were recovered from both uncovered trays. In the screened trays, 1,877 clams (all dead) were recovered.

## Survival

Average survival of *Corbicula* in the river from November to May for 1992–2000 was 9.9% at site A, 9.3% at site B, 10.4% at site C, and 12.7% at site D (Table 7). Overall winter survival of *Corbicula* (all river sites combined) was positively correlated with average winter water temperature (December–April;  $r_s = 0.87$ ,  $P = 0.004$ ) and negatively correlated with the April flow index ( $r_s = -0.91$ ,  $P = 0.002$ ; Table 8). In addition, winter survival was also negatively correlated to the number of days that water temperatures averaged less than or equal to 1°C ( $r_s = -0.73$ ,  $P = 0.04$ ). *Corbicula* winter survival was not significantly correlated with the number of days that water temperature was less than or equal to 2°C ( $r_s = -0.65$ ,  $P = 0.08$ ).

However, spatial differences among sites were evident between environmental stress indices and winter survival. At site A, the site furthest from the CY, all four indices were significantly correlated with *Corbicula* winter survival: average winter water temperatures ( $r_s = 0.92$ ,  $P = 0.001$ ), peak flows in April ( $r_s = -0.74$ ,  $P = 0.036$ ), the number of days that were less than or equal to 1°C ( $r_s = -0.87$ ,  $P = 0.004$ ) and less than or equal to 2°C ( $r_s = -0.82$ ,  $P = 0.013$ ). At the B site near the plant intakes, winter survival was significantly correlated with average winter water temperature ( $r_s = 0.73$ ,  $P = 0.04$ ) and peak flows in April ( $r_s = -0.77$ ,  $P = 0.02$ ). At C, only average winter water temperature ( $r_s = 0.85$ ,  $P = 0.02$ ) was significantly correlated with survival. At D, the closest site in the river to the CY discharge, winter survival was not significantly correlated with any of the four indices. *T*-tests comparing percentage winter survival during operational and post-operational years showed no differences at any of the four river sites.

## Reproduction

Clams within lantern-nets in the Connecticut River showed the presence of eggs and sperm in

Table 3. Mean total abundance (clams/m<sup>2</sup> ± 95% CI) of *Corbicula fluminea* in the 2- to 4-mm shell-length size-classes at each sampling site near the Connecticut Yankee power plant in the Connecticut River from August 1991 to November 2000.

| Site                   | Year      |               |           |           |               |          |           |           |               |           |             |     | All years |
|------------------------|-----------|---------------|-----------|-----------|---------------|----------|-----------|-----------|---------------|-----------|-------------|-----|-----------|
|                        | 1991      | 1992          | 1993      | 1994      | 1995          | 1996     | 1997      | 1998      | 1999          | 2000      | 2000        | All |           |
| A                      |           |               |           |           |               |          |           |           |               |           |             |     |           |
| May                    | -         | 10 ± 18       | 12 ± 33   | 0         | 176 ± 150     | 0        | 8 ± 22    | 104 ± 147 | 102 ± 51      | 6 ± 11    | 46 ± 24     |     |           |
| Aug.                   | 302 ± 236 | 4,246 ± 3,060 | 0 ± 0     | 18 ± 25   | 1,883 ± 1,937 | 44 ± 52  | 72 ± 88   | 488 ± 449 | 1,998 ± 547   | 352 ± 312 | 940 ± 447   |     |           |
| Nov.                   | 520 ± 313 | 5,924 ± 2,463 | 232 ± 265 | 456 ± 328 | 758 ± 1,040   | 38 ± 37  | 506 ± 305 | 710 ± 467 | 1,228 ± 1,241 | 292 ± 174 | 1,066 ± 513 |     |           |
| Average <sup>a</sup>   | 411       | 3,393         | 81        | 158       | 939           | 27       | 195       | 434       | 1,109         | 217       | 706         |     |           |
| B                      |           |               |           |           |               |          |           |           |               |           |             |     |           |
| May                    | -         | 4 ± 11        | 0         | 0         | 28 ± 33       | 0        | 10 ± 28   | 12 ± 20   | 80 ± 59       | 26 ± 35   | 18 ± 10     |     |           |
| Aug.                   | 170 ± 52  | 768 ± 302     | 0         | 16 ± 44   | 866 ± 733     | 14 ± 14  | 16 ± 19   | 156 ± 76  | 1,104 ± 322   | 24 ± 31   | 323 ± 127   |     |           |
| Nov.                   | 186 ± 123 | 2,162 ± 710   | 100 ± 90  | 82 ± 67   | 854 ± 1,149   | 64 ± 150 | 152 ± 82  | 222 ± 118 | 680 ± 419     | 74 ± 64   | 457 ± 202   |     |           |
| Average <sup>a</sup>   | 178       | 978           | 33        | 33        | 583           | 26       | 59        | 130       | 621           | 75        | 275         |     |           |
| C                      |           |               |           |           |               |          |           |           |               |           |             |     |           |
| May                    | -         | 4 ± 11        | 0         | 0         | 156 ± 197     | 0        | 6 ± 17    | 26 ± 46   | 114 ± 105     | 14 ± 32   | 36 ± 24     |     |           |
| Aug.                   | 116 ± 123 | 914 ± 599     | 18 ± 38   | 22 ± 20   | 996 ± 851     | 88 ± 59  | 12 ± 27   | 184 ± 196 | 814 ± 488     | 284 ± 306 | 345 ± 135   |     |           |
| Nov.                   | 122 ± 130 | 994 ± 1,052   | 74 ± 73   | 26 ± 24   | 224 ± 242     | 14 ± 19  | 192 ± 348 | 210 ± 149 | 222 ± 150     | 74 ± 88   | 215 ± 109   |     |           |
| Average <sup>a</sup>   | 119       | 637           | 31        | 16        | 459           | 34       | 70        | 140       | 383           | 124       | 204         |     |           |
| D                      |           |               |           |           |               |          |           |           |               |           |             |     |           |
| May                    | -         | -             | -         | 0         | 84 ± 67       | 0        | 0         | 8 ± 22    | 8 ± 16        | 38 ± 53   | 20 ± 13     |     |           |
| Aug.                   | -         | -             | 50 ± 108  | 40 ± 34   | 108 ± 73      | 36 ± 44  | 10 ± 22   | 30 ± 28   | 686 ± 314     | 190 ± 149 | 144 ± 76    |     |           |
| Nov.                   | -         | -             | 154 ± 290 | 92 ± 97   | 272 ± 87      | 88 ± 83  | 252 ± 239 | 292 ± 82  | 104 ± 73      | 88 ± 68   | 168 ± 44    |     |           |
| Average <sup>a</sup>   | -         | -             | 102       | 44        | 155           | 41       | 187       | 110       | 266           | 105       | 114         |     |           |
| All sites <sup>b</sup> |           |               |           |           |               |          |           |           |               |           |             |     |           |
| May                    | -         | 6 ± 6         | 4 ± 9     | 0 ± 0     | 111 ± 53      | 0 ± 0    | 6 ± 7     | 38 ± 33   | 76 ± 30       | 21 ± 14   | 30 ± 10     |     |           |
| Aug.                   | 196 ± 79  | 1,976 ± 1,185 | 17 ± 22   | 24 ± 12   | 963 ± 491     | 46 ± 20  | 28 ± 21   | 214 ± 118 | 1,150 ± 287   | 238 ± 90  | 454 ± 132   |     |           |
| Nov.                   | 276 ± 132 | 3,027 ± 1,375 | 140 ± 76  | 164 ± 102 | 527 ± 304     | 51 ± 33  | 276 ± 113 | 358 ± 132 | 558 ± 312     | 132 ± 58  | 493 ± 152   |     |           |
| Average <sup>a</sup>   | 236       | 1,670         | 58        | 63        | 534           | 32       | 103       | 204       | 595           | 130       | 336         |     |           |

<sup>a</sup> Annual averages (clams/m<sup>2</sup>) pooling the May, August, and November samples.

<sup>b</sup> Average abundance of clams, pooling the four sites.



Table 4. Mean abundance (clams/m<sup>2</sup> ± 95% CI) of *Corbicula fluminea* in the 7 to 14 mm shell length size-classes at each sampling site near the Connecticut Yankee power plant in the Connecticut River from August 1991 to November 2000.

| Site                   | Years                |               |               |          |           |             |         |           |               |               |           |             | All years |
|------------------------|----------------------|---------------|---------------|----------|-----------|-------------|---------|-----------|---------------|---------------|-----------|-------------|-----------|
|                        | 1991                 | 1992          | 1993          | 1994     | 1995      | 1996        | 1997    | 1998      | 1999          | 2000          | 2000      |             |           |
| A                      | May                  | -             | 112 ± 98      | 0        | 0         | 26 ± 30     | 0       | 0         | 44 ± 31       | 168 ± 286     | 39 ± 28   |             |           |
|                        | Aug.                 | 338 ± 151     | 1,284 ± 926   | 8 ± 10   | 36 ± 38   | 1,154 ± 708 | 0       | 46 ± 38   | 2,068 ± 1,185 | 290 ± 309     | 537 ± 224 |             |           |
|                        | Nov.                 | 1,092 ± 584   | 5,172 ± 2,417 | 100 ± 97 | 120 ± 65  | 1,552 ± 786 | 22 ± 34 | 176 ± 104 | 786 ± 458     | 2,290 ± 1,484 | 736 ± 219 | 1,204 ± 472 |           |
|                        | Average <sup>a</sup> | 715           | 2,189         | 36       | 52        | 911         | 7       | 74        | 309           | 1,467         | 398       | 612         |           |
| B                      | May                  | -             | 16 ± 29       | 0        | 0         | 2 ± 6       | 0       | 0         | 6 ± 11        | 28 ± 24       | 6 ± 4     |             |           |
|                        | Aug.                 | 118 ± 75      | 198 ± 96      | 18 ± 31  | 24 ± 7    | 702 ± 51    | 18 ± 16 | 22 ± 14   | 520 ± 156     | 72 ± 34       | 173 ± 67  |             |           |
|                        | Nov.                 | 72 ± 91       | 518 ± 68      | 36 ± 17  | 24 ± 41   | 926 ± 962   | 16 ± 17 | 52 ± 10   | 918 ± 419     | 204 ± 240     | 287 ± 122 |             |           |
|                        | Average              | 95            | 244           | 18       | 16        | 543         | 11      | 25        | 49            | 481           | 101       | 161         |           |
| C                      | May                  | -             | 16 ± 29       | 0        | 0         | 8 ± 6       | 0       | 2 ± 6     | 22 ± 14       | 12 ± 16       | 6 ± 4     |             |           |
|                        | Aug.                 | 154 ± 87      | 226 ± 60      | 22 ± 10  | 62 ± 39   | 928 ± 497   | 12 ± 14 | 30 ± 9    | 770 ± 315     | 92 ± 76       | 237 ± 98  |             |           |
|                        | Nov.                 | 368 ± 60      | 724 ± 145     | 66 ± 19  | 58 ± 58   | 926 ± 467   | 20 ± 15 | 28 ± 22   | 730 ± 334     | 70 ± 77       | 310 ± 104 |             |           |
|                        | Average              | 261           | 322           | 29       | 40        | 621         | 11      | 20        | 65            | 507           | 58        | 191         |           |
| D                      | May                  | -             | -             | -        | 0         | 14 ± 27     | 0       | 4 ± 11    | 0             | 20 ± 25       | 5 ± 5     |             |           |
|                        | Aug.                 | -             | -             | 24 ± 11  | 54 ± 35   | 646 ± 258   | 12 ± 16 | 18 ± 16   | 608 ± 159     | 52 ± 24       | 183 ± 87  |             |           |
|                        | Nov.                 | -             | -             | 48 ± 32  | 36 ± 47   | 292 ± 362   | 58 ± 32 | 42 ± 27   | 804 ± 155     | 96 ± 73       | 190 ± 87  |             |           |
|                        | Average              | -             | -             | 36       | 30        | 317         | 23      | 21        | 66            | 471           | 56        | 132         |           |
| All sites <sup>b</sup> |                      |               |               |          |           |             |         |           |               |               |           |             |           |
| May                    | -                    | 48 ± 36       | 0             | 0        | 12 ± 8    | 0           | 2 ± 2   | 1 ± 1     | 18 ± 10       | 57 ± 59       | 15 ± 8    |             |           |
| Aug.                   | 203 ± 71             | 569 ± 365     | 18 ± 7        | 44 ± 13  | 858 ± 184 | 10 ± 6      | 29 ± 9  | 76 ± 25   | 991 ± 371     | 126 ± 72      | 288 ± 70  |             |           |
| Nov.                   | 510 ± 284            | 2,138 ± 1,360 | 62 ± 22       | 60 ± 26  | 924 ± 320 | 29 ± 12     | 74 ± 34 | 290 ± 160 | 1,186 ± 412   | 276 ± 142     | 514 ± 141 |             |           |
| Average                | 357                  | 918           | 29            | 34       | 598       | 13          | 35      | 122       | 732           | 153           | 282       |             |           |

<sup>a</sup> Annual averages (clams/m<sup>2</sup>) pooling the May, August, and November samples.

<sup>b</sup> Abundance of clams, pooling the four sites.

Table 5. Mean abundance (clams/m<sup>2</sup> ± 95% CI) of *Corbicula fluminea* in the 19 to 31 mm shell length size-classes at each sampling site near the Connecticut Yankee power plant in the Connecticut River from August 1991 to November 2000.

| Site<br>Month                | Years   |           |         |       |           |      |         |       |         |        | All<br>years |           |
|------------------------------|---------|-----------|---------|-------|-----------|------|---------|-------|---------|--------|--------------|-----------|
|                              | 1991    | 1992      | 1993    | 1994  | 1995      | 1996 | 1997    | 1998  | 1999    | 2000   |              |           |
| <b>A</b>                     |         |           |         |       |           |      |         |       |         |        |              |           |
| May                          | -       | 2 ± 6     | 0       | 0     | 0         | 0    | 2 ± 6   | 0     | 0       | 0      | 0            | 0.4 ± 0.6 |
| Aug.                         | 0       | 22 ± 27   | 0       | 0     | 16 ± 26   | 0    | 0       | 0     | 10 ± 12 | 8 ± 10 | 0            | 6 ± 3     |
| Nov.                         | 30 ± 51 | 62 ± 58   | 0       | 0     | 142 ± 184 | 0    | 2 ± 6   | 2 ± 6 | 34 ± 68 | 8 ± 10 | 0            | 28 ± 18   |
| Average <sup>a</sup>         | 15      | 29        | 0       | 0     | 53        | 0    | 1       | 1     | 15      | 5      | 0            | 12        |
| <b>B</b>                     |         |           |         |       |           |      |         |       |         |        |              |           |
| May                          | -       | 0         | 0       | 0     | 0 ± 0     | 0    | 0       | 0     | 0       | 0      | 0            | 0         |
| Aug.                         | 0       | 16 ± 21   | 0       | 0     | 6 ± 8     | 0    | 0       | 0     | 2 ± 6   | 0      | 0            | 2 ± 2     |
| Nov.                         | 14 ± 27 | 8 ± 16    | 2 ± 6   | 4 ± 7 | 94 ± 22   | 0    | 0       | 0     | 6 ± 7   | 0      | 0            | 13 ± 8    |
| Average                      | 7       | 8         | 1       | 1     | 33        | 0    | 0       | 0     | 3       | 0      | 0            | 5         |
| <b>C</b>                     |         |           |         |       |           |      |         |       |         |        |              |           |
| May                          | -       | 0         | 0       | 0     | 0         | 0    | 0       | 0     | 0       | 0      | 0            | 0         |
| Aug.                         | 0       | 30 ± 23   | 0       | 0     | 10 ± 9    | 0    | 0       | 0     | 6 ± 11  | 0      | 0            | 5 ± 3     |
| Nov.                         | 18 ± 16 | 62 ± 16   | 0       | 0     | 34 ± 37   | 0    | 0       | 2 ± 6 | 4 ± 7   | 0      | 0            | 12 ± 6    |
| Average                      | 9       | 31        | 0       | 0     | 15        | 0    | 0       | 1     | 3       | 0      | 0            | 6         |
| <b>D</b>                     |         |           |         |       |           |      |         |       |         |        |              |           |
| May                          | -       | -         | -       | 0     | 0         | 0    | 0       | 0     | 0       | 0      | 0            | 0         |
| Aug.                         | -       | -         | 0       | 0     | 0         | 0    | 2 ± 6   | 0     | 6 ± 11  | 0      | 0            | 1 ± 1     |
| Nov.                         | -       | -         | 12 ± 10 | 2 ± 6 | 16 ± 19   | 0    | 0       | 0     | 14 ± 7  | 8 ± 6  | 0            | 7 ± 3     |
| Average                      | -       | -         | 6       | 1     | 5         | 0    | 1       | 0     | 7       | 3      | 0            | 3         |
| <b>All sites<sup>b</sup></b> |         |           |         |       |           |      |         |       |         |        |              |           |
| May                          | -       | 0.7 ± 1.4 | 0       | 0     | 0         | 0    | 0.5 ± 1 | 0     | 0       | 0      | 0            | 0.1 ± 0.2 |
| Aug.                         | 0       | 23 ± 10   | 0       | 0     | 8 ± 6     | 0    | 0.5 ± 1 | 0     | 6 ± 4   | 2 ± 2  | 0            | 4 ± 1     |
| Nov.                         | 21 ± 15 | 44 ± 21   | 4 ± 3   | 2 ± 2 | 72 ± 41   | 0    | 0.5 ± 1 | 1 ± 1 | 15 ± 13 | 4 ± 13 | 0            | 15 ± 6    |
| Average                      | 10      | 22        | 1       | 1     | 26        | 0    | 1       | <1    | 7       | 2      | 0            | 6         |

<sup>a</sup> Annual averages (clams/m<sup>2</sup>) pooling the May, August, and November samples.

<sup>b</sup> Abundance of clams, pooling the four sites.

Table 6. Mean abundance (clams/m<sup>2</sup> ± 95% CI) of *Corbicula fluminea* in the CY discharge canal (Site E) by size class from August 1991 to November 2000.

| Year | Month          | Abundance by size-class |         |          | Total abundance |
|------|----------------|-------------------------|---------|----------|-----------------|
|      |                | 2–4 mm                  | 7–14 mm | 19–31 mm |                 |
| 1991 | May            | –                       | –       | –        | –               |
|      | Aug.           | 34±63                   | 0       | 0        | 34±63           |
|      | Nov.           | 172±181                 | 6±17    | 0        | 178±195         |
|      | Year's average | 103                     | 3       | 0        | 106             |
| 1992 | May            | 2±6                     | 0       | 0        | 2±6             |
|      | Aug.           | 12,174±32,688           | 0       | 0        | 12,174±32,688   |
|      | Nov.           | 96±131                  | 0       | 0        | 96±131          |
|      | Year's average | 4,091                   | 0       | 0        | 4,091           |
| 1993 | May            | 8±10                    | 24±29   | 0        | 32±24           |
|      | Aug.           | 0                       | 0       | 0        | 0               |
|      | Nov.           | 32±33                   | 10±18   | 0        | 42±28           |
|      | Year's average | 13                      | 11      | 0        | 25              |
| 1994 | May            | 0                       | 0       | 0        | 0               |
|      | Aug.           | 36±36                   | 24±29   | 0        | 60±59           |
|      | Nov.           | 816±1,303               | 58±57   | 6±17     | 880±1,354       |
|      | Year's average | 284                     | 27      | 2        | 313             |
| 1995 | May            | 24±31                   | 0       | 0        | 24±31           |
|      | Aug.           | 38±54                   | 0       | 0        | 38±54           |
|      | Nov.           | 90±202                  | 0       | 0        | 90±202          |
|      | Year's average | 51                      | 0       | 0        | 51              |
| 1996 | May            | 0                       | 8±6     | 0        | 8±6             |
|      | Aug.           | 48±81                   | 0       | 0        | 48±81           |
|      | Nov.           | 10±12                   | 34±81   | 0        | 44±75           |
|      | Year's average | 19                      | 14      | 0        | 33              |
| 1997 | May            | 60±79                   | 32±31   | 0        | 92±66           |
|      | Aug.           | 0                       | 6±17    | 0        | 6±17            |
|      | Nov.           | 4±7                     | 18±10   | 2±6      | 24±7            |
|      | Year's average | 21                      | 19      | 1        | 41              |
| 1998 | May            | 18±20                   | 2±6     | 0        | 20±23           |
|      | Aug.           | 40±81                   | 16±32   | 6±11     | 62±89           |
|      | Nov.           | 106±213                 | 96±77   | 10±12    | 212±245         |
|      | Year's average | 55                      | 38      | 5        | 98              |
| 1999 | May            | 74±35                   | 2±6     | 0        | 76±40           |
|      | Aug.           | 4±7                     | 216±196 | 14±17    | 234±186         |
|      | Nov.           | 0                       | 124±148 | 86±92    | 210±79          |
|      | Year's average | 26                      | 114     | 33       | 173             |

Table 6. Continued

| Year            | Month          | Abundance by size-class |         |          | Total abundance |
|-----------------|----------------|-------------------------|---------|----------|-----------------|
|                 |                | 2–4 mm                  | 7–14 mm | 19–31 mm |                 |
| 2000            |                |                         |         |          |                 |
|                 | May            | 0                       | 4±11    | 0        | 4±11            |
|                 | Aug.           | 20±34                   | 8±16    | 2±6      | 30±38           |
|                 | Nov.           | 162±125                 | 96±100  | 10±22    | 268±195         |
|                 | Year's average | 61                      | 36      | 4        | 101             |
| 10-year average |                |                         |         |          |                 |
|                 | May            | 21±11                   | 8±5     | 0        | 29±12           |
|                 | Aug.           | 1,239±2,380             | 27±22   | 2±2      | 1,269±2,378     |
|                 | Nov.           | 149±111                 | 44±19   | 11±10    | 204±116         |
|                 | Annual average | 485                     | 27      | 5        | 517             |

Table 7. Average percentage winter survival of Asiatic clams (*Corbicula*), native clams (Sphaeridae), and native mussels (Unionidae) at four CY sampling sites.

| Site    | <i>Corbicula</i> | Sphaeridae | Unionidae |
|---------|------------------|------------|-----------|
| A       | 9.9%             | 47.0%      | 17.5%     |
| B       | 9.3%             | 65.6%      | 51.3%     |
| C       | 10.4%            | 28.5%      | 33.3%     |
| D       | 12.7%            | 28.4%      | –         |
| Average | 10.6%            | 42.4%      | 34.0%     |

gonadal tissues from May through January (Figure 5). The frequency of clams with eggs ranged from 75% to 100% and with sperm ranged from 52% to 100%. The presence of young in the demibranchs (brooding) and the active releasing

of juveniles occurred from June to October. The frequency of clams brooding during this period ranged from 4% in October to 62% in August (Figure 5). In the 4 years of the reproduction study, only one brooding clam was observed in October and three in January. The periods of brooding and release of juvenile clams (spawning) occurred at about the same time, from June to September. Spawning peaked in August with an average release of 2,861 juveniles/clam/d.

During power plant operation, the clams were able to survive in the CY discharge canal from November through May; eggs and sperm were continually present in gonadal tissues (Figure 6). Eggs were present in 100% of clams surveyed and sperm frequency ranged from 8% in Decem-

Table 8. Spearman correlation coefficients of percentage winter survival of *Corbicula fluminea* at CY sites versus four indices of winter severity for the Connecticut River from 1992 through 2000<sup>a</sup>.

| Winter severity index<br>(Avg. ± SE)                | Spearman correlation coefficients by site |         |                     |                     | All sites<br>combined |                     |
|---|---|---------|---------------------|---------------------|-----------------------|---------------------|
|   | A   | B       | C                   | D                   |                       |                     |
| Average<br>winter temp. <sup>b</sup>                | (2.93°C ±0.37)                            | +0.92** | +0.73*              | +0.84*              | +0.63 <sup>ns</sup>   | +0.87**             |
| No. days<br>≤1°C                                    | (54.9 d ±8.75)                            | –0.87** | –0.63 <sup>ns</sup> | –0.74 <sup>ns</sup> | –0.56 <sup>ns</sup>   | –0.73*              |
| No. days<br>≤2°C                                    | (70.6 d ±8.12)                            | –0.82*  | –0.52 <sup>ns</sup> | –0.62 <sup>ns</sup> | –0.67 <sup>ns</sup>   | –0.65 <sup>ns</sup> |
| April flow<br>≥1,700 m <sup>3</sup> /s <sup>c</sup> | (6.4 d ±1.54)                             | –0.74*  | –0.78*              | –0.73 <sup>ns</sup> | –0.69 <sup>ns</sup>   | –0.91**             |

<sup>a</sup> 1993 data were omitted because of the mortality caused by the March storm Joshua (see text).

<sup>b</sup> Average winter temperature = the annual Dec.–Apr. mean daily Connecticut River temperature at CY.

<sup>c</sup> Number of days in April when the Connecticut River flow equaled or exceeded a daily average of 1,700 m<sup>3</sup>/s.

\*Significance levels: \*\* =  $P < 0.01$ ; \* =  $P < 0.05$ ; ns =  $P \geq 0.05$ .




Figure 5. Average seasonal reproductive status of *Corbicula fluminea* (1991-1994) at ambient river temperatures. The percentage of adult clams (e"8 mm) with mature gametes or brooding young is plotted using the left axis. Active spawning is plotted using the right axis and is defined as the average number juveniles released over a 24-hour period by 12 randomly selected adult clams.




Figure 6. Reproductive status of *Corbicula fluminea* held in the CY discharge canal from November 1992 to May 1993, when water temperatures were 8°C to 12°C above ambient river temperatures. The percentage of adult clams (≥8 mm) with mature gametes or brooding young is plotted using the left axis. Active spawning is plotted using the right axis and is defined as the number juveniles released over a 24-h period by 12 randomly selected adult clams.

ber to 100% in May. Continuous brooding and spawning occurred from March through May with brooding adults ranging from 8% to 100%, when water temperature averaged  $\geq 17^{\circ}\text{C}$ . Brooding and spawning were also observed in November (92% brooding), when temperatures averaged  $18.3^{\circ}\text{C}$ . The sharp decrease in spawning activity in May was the result of power plant shut down beginning on 15 May 1993, which dropped cooling water temperatures from  $30\text{--}18^{\circ}\text{C}$  in a single day. Brooding and spawning peaked in June and July as the power outage continued and reproduction of discharge clams followed the pattern observed at ambient river temperatures ( $17\text{--}21^{\circ}\text{C}$ ). On 21 July 1993 the power plant restarted and temperatures increased to  $>35^{\circ}\text{C}$  in 4 d. All clams held in the CY discharge canal were dead by 18 August 1993.

## Growth

Quadratic regression models fit to captive clam growth data had slope parameters ( $b_1$  and  $b_2$ ) that were significantly different from zero ( $P < 0.05$ ; Figure 7). The 1991 growth data showed the best fit to the model ( $R^2 = 0.96$ ) and had the highest peak growth rate of 1.08 mm/week. Peak growth rates in 1992 and 1993 were 0.68 mm/week and 0.44 mm/week, respectively. *Corbicula* growth under ambient conditions during 1991–1993 exhibited seasonal cycles. For example, growth in 1991 between June and July was 1.08 mm/week, between July and August was 0.94 mm/week, and between August and September was 0.60 mm/week. Growth was minimal from September to October (0.10 mm/week) and October to November (0.06 mm/week).

Clam growth in the CY discharge canal was




Figure 7. Growth of *Corbicula fluminea* at ambient river conditions in 1991, 1992 and 1993 (Panels A–C), as compared to growth at CY discharge (Panel D) conditions from November 1992 to August 1993. The quadratic regression models (1991, 1992, and 1993) are of the form,  $\text{Length} = a + b_1(\text{Day}) + b_2(\text{Day})^2$ , where “a” is the intercept, “ $b_1$ ” and “ $b_2$ ” are the slopes, and “Day” references the Julian day of the year for 1991, 1992, and 1993. The linear regression model (1992–1993) is of the form,  $\text{Length} = a + b(\text{Day})$ , where “Day” references a cumulative Julian day over the 2-year period 1992–1993. Regression lines are bracketed by 95% confidence limits, and the vertical bars indicate the minimum, maximum, and average shell length by horizontal bars ( $n = 12$ ).

0.05 mm/week from November 1992 to February 1993, when water temperatures ranged from 12–21°C (10–12°C above ambient river temperatures; Figure 7). Increased growth rates were observed from February to May (0.14 mm/week), when water temperatures ranged from 12–30°C. Maximum growth rates at this site occurred from May to July (0.36 mm/week), when canal temperatures were similar to those at ambient river sites because of a power plant shut down. All clams died after the power plant restarted in July 1993 and discharge water temperatures exceeded 37°C.

## Native Bivalve Abundance

Clams of the family Sphaeriidae comprised more than 90% of the native bivalves collected and were not identified below this taxonomic level, while adult mussels (Unionidae) were identified to species. Four mussel species were documented

at the Connecticut River sites: eastern elliptio *Elliptio complanata*, alewife floater *Anodonta imbecilis*, tidewater mucket *Ligumia ochracea*, and eastern pondmussel *L. nasuta*. Of these, *E. complanata* comprised more than 90% of all mussels collected. *Ligumia ochracea* and *L. nasuta*, the rarest mussels, were not collected until 1997 and 1999, respectively.

Mean annual abundances (individuals/m<sup>2</sup>) of native clams and mussels were highly variable with clams ranging from 16 to 242 and mussels ranging from 1 to 5 (Figure 8). Abundances of both mussels and clams were highest in 1991. The lowest abundances for clams occurred in 1993, 1996, and 2000, while those for mussels occurred in 1995 and 2000. Mean annual abundances of native clams (Sphaeriidae) averaged six times lower than those for *Corbicula*.

Winter survival was considerably greater for native clams and mussels than for *Corbicula* (Table 7). Winter survival of Sphaeriidae ranged




Figure 8. Annual abundance ((#/m<sup>2</sup>)+1) of native clams (Sphaeriidae) and mussels (Unionidae) from 1991 through 2000 at Connecticut River sites in the vicinity of CY. Annual abundance estimates for *Corbicula* are given in numeric values (#/m<sup>2</sup>) above each year.

from 28.4% at D to 65.6% at B, averaging 42% for all sites. Unionidae survival ranged from 17.5% at A to 51.3% at B, averaging 34% for sites A–C. Mussels were not collected at D. In comparison, *Corbicula* survival ranged from 9.3% at B to 12.7% at D, averaging 11% overall.

Statistical trend analyses on the log transformed abundance data using a Mann-Kendall statistic showed no significant ( $P < 0.05$ ) trends in abundance for either native clam or mussels from 1991 to 2000. *T*-tests of native clam and mussel seasonal abundance during the CY operational period (1991–1996) versus the post-operational period (1997–2000) showed no significant differences, except for mussels in November (Table 9).

## Discussion

*Corbicula* has been a major component of macrobenthic communities in the lower Connecticut River following its initial discovery in 1990. *Corbicula* distribution in 1991 extended northward from CY to Gildersleeve Island, a 16 km section of the river, and it spread 22 km further upriver in 1992. Savoy and Benway (2004, this volume) collected sediment grab samples during 2000–2002 approximately 5–10 km upriver from the northern extent reported for *Corbicula* in 1992 (South Meadow station) and did not report the presence *Corbicula*. Upstream movements of *Corbicula* are not unusual and can happen without anthropogenic intervention (Voelz et al. 1998). Rodgers et al. (1977) reported movement of 222

km in 15 years. Once a *Corbicula* population is established upriver, downriver dispersal is more rapid; the clams are able to free themselves from the bottom and are aided by water currents in their movements (Sinclair and Isom 1963; Britton and Morton 1979; Eng 1979; Prezant and Chalermwat 1984; French and Schloesser 1996).

*Corbicula* winter survival in the Connecticut River was correlated positively with water temperature from December through April, and negatively with high-water flows in spring. No clams survived at any of the sites during the coldest winters of 1993–1994 and 1995–1996, when temperatures were below 2°C for 12–15 weeks. Similarly low survival due to coldwater temperatures was reported by a number of other researchers (Horning and Keup 1964; Bickel 1966; Dreier 1977; Rodgers et al. 1979; Cherry et al. 1980; Graney et al. 1980; Cairns and Cherry 1983). *Corbicula* survival at individual stations was also influenced by distance from the thermal plume discharge. Winter survival at site D, the river site nearest to where the CY discharge mixes with the river, was not significantly correlated with winter water temperature indices. The sites between the CY intakes and discharge canal (sites B and C) were significantly correlated with average winter water temperatures, but not correlated with the number of days water temperatures below 1°C or 2°C. The reason for the poor correlation of these temperature indices ( $\leq 2^\circ\text{C}$  and  $\leq 1^\circ\text{C}$ ) at sites B, C, and D might be related to the thermal plume mixing with ambient temperature river water during tidal incursions into our sampling sites or the

Table 9. Total abundance of *Corbicula fluminea*, native clams (Sphaeriidae), and native mussels (Unionidae) in relation to the CY power plant's operational status.

| Month    | Time                          | <i>Corbicula fluminea</i> |        | Sphaeriidae |       | Unionidae |      |
|----------|-------------------------------|---------------------------|--------|-------------|-------|-----------|------|
|          |                               | Mean <sup>a</sup>         | S.E.   | Mean        | S.E.  | Mean      | S.E. |
| May      | Operational <sup>b</sup>      | 37.22                     | 8.48   | 39.11       | 13.00 | 1.22      | 0.47 |
|          | Post-operational <sup>c</sup> | 54.50                     | 9.70   | 45.75       | 11.49 | 2.12      | 0.58 |
| August   | Operational                   | 733.33                    | 142.29 | 147.54      | 21.90 | 3.18      | 0.83 |
|          | Post-operational              | 715.50                    | 120.76 | 119.00      | 23.50 | 1.75      | 0.50 |
| November | Operational                   | 1190.09                   | 235.71 | 110.46      | 25.77 | **0.91    | 0.42 |
|          | Post-operational              | 792.62                    | 106.96 | 94.25       | 16.56 | **4.00    | 1.15 |

<sup>a</sup>Mean = #/m<sup>2</sup> is calculated by averaging the total number of individuals per grab at the A, B, C, and D for each sample month from 1991 to 2000 and converting that value to #/m<sup>2</sup> ((#/grab) × 10).

<sup>b</sup>CY operational years (1991–1996).

<sup>c</sup>Post-operational years (1997–2000).

\*\* Difference in abundance between time periods is highly significant; Satterthwaite *t*-test,  $P < 0.01$ .



recruitment of some CY discharge canal clams to these sites closest to the discharge outfall. However, *t*-tests of winter survival data at each site between operation and postoperational years indicated no differences. Although twice each day during low flow conditions the river reverses flow and thermal plume water moves upriver where our sites are located (Rozsa 2001; Boyd 2004 [1976]), differences in site correlations to winter severity indices could be caused by site locations in relation to river conditions other than CY discharge effects. The significant correlations observed at site A between river temperature indices and winter survival of clams indicates no thermal plume effects at this station. During periods of high river flow, usually in the spring and other times of flooding, the flow at CY is unidirectional and not influenced by tidal flows. The *Corbicula* population at CY sites was able to recover after high winter mortalities by August and usually peaked in November. This recovery was driven by passive downriver recolonization from populations that were able to survive the cold winter conditions in refugia (e.g., those known to exist near thermal plumes at Middletown and possibly South Meadow power stations or natural refugia, such as groundwater seeps)(Morgan et al. 2003).

The importance of CY thermal discharge as a refuge for *Corbicula* survival in the Connecticut River during cold winters appears minimal. The clams were not able to maintain year-round populations in the discharge canal because of total mortality due to high discharge temperatures from June through October. Most *Corbicula* that could colonize the discharge canal in winter and spring, therefore, were small and not reproductively mature. If movement of clams to the river sites occurred during March through May, it would occur when conditions were poor for their survival (e.g., low spring water temperatures [ $<12^{\circ}\text{C}$ ] and high, unidirectional spring flows).

Additional evidence that the CY discharge was not necessary for survival of *Corbicula* populations in the Connecticut River is apparent when *Corbicula* abundance during CY operation (1991–1996) was compared to abundance following the plant closure (1997–2000). Following closure of the CY power plant in 1996, the abundance of *Corbicula* at all sites was not significantly different than during the operational period. In addition, following CY closure, clams were able to survive in the discharge canal throughout the year

and significant increases in the abundance of larger size clams were observed.

The growth patterns of *Corbicula* in the Connecticut River are similar to those reported by other researchers. The highest growth rate occurred in the summer at temperatures approximating  $25^{\circ}\text{C}$ , and minimal growth occurred at temperatures below  $15^{\circ}\text{C}$ . Others have reported  $10$ – $15^{\circ}\text{C}$  to be the lower threshold for *Corbicula* growth (Joy 1985; Mattice and Wright 1986; McMahon and Williams 1986a; Stites et al. 1995).

The high *Corbicula* growth rate in 1991 was due to small-sized individuals used in the study. Other researchers have also reported that smaller clams have higher growth rates (O’Kane 1976; Britton et al. 1979; Mattice 1979; Dreier and Tranquilli 1981; Dauble et al. 1985; Joy 1985; Mattice and Wright 1986; McMahon and Williams 1986a, 1986b; Foe and Knight 1986; Morgan et al. 2003). The total growth of *Corbicula* in the Connecticut River was in the range of growth reported by others (Britton et al. 1979; McMahon 1983; Welch and Joy 1984; Mattice and Wright 1986; McMahon and Williams 1986a; French and Schloesser 1991). The growth of *Corbicula* at the CY discharge started in the fall and continued modestly until discharge water temperatures exceeded  $37^{\circ}\text{C}$  in June or July, when all clams died. Others reported that the upper temperature tolerance threshold for this clam between  $34$ – $37^{\circ}\text{C}$  (Mattice and Dye 1976; Dreier 1977; Mattice 1979; Cairns and Cherry 1983; McMahon 1983; McMahon and Williams 1986b).

Brooding and spawning of *Corbicula* in the Connecticut River and CY discharge canal occurred at water temperatures between  $15^{\circ}\text{C}$  and  $31^{\circ}\text{C}$ . Ambient river temperatures remained within this range from June to September, and CY discharge conditions remained within this range in November and April to mid-May. Peak reproduction, as described by brooding and spawning, occurred in August at river temperatures ( $22$ – $28^{\circ}\text{C}$ ) and in April at CY discharge temperatures ( $13$ – $24^{\circ}\text{C}$ ). The survival of juveniles released in the discharge canal is questionable, particularly during times of high water flow. Potential recruitment to the river from the discharge canal comes when temperature in the river is low for successful growth and river flow rates are at yearly optimal levels for downriver transport by the swift, unidirectional currents of spring freshets. As a result, they would be exposed to

estuarine waters of Long Island Sound and perish due to high salinity.

While *Corbicula* quickly established itself as the dominant bivalve in the Connecticut River, there was little change in native bivalve abundance found in the same sediments. We believe this is due to relatively high winter survival of native clams and mussels when compared to that for *Corbicula*. Similarly, Fuller and Imlay (1976) questioned whether there was any displacement of native Unionidae species following *Corbicula* colonization. Diaz (1994) concluded that *Corbicula* invasions of Northern American tidal freshwater systems in the late 1960s did not displace tidal freshwater species, but rather, these nonnative clams took advantage of underutilized benthic resources. Others suggest filtration activities by high densities of *Corbicula* can lead to alteration of nutrient dynamics in aquatic systems, resulting in negative impacts on native bivalves (Kraemer 1979; Lauritsen 1986; Leff et al. 1990; Phelps 1994). Karatayev et al. (2003) found no correlations between the densities of *Corbicula* and other benthic invertebrates in an East Texas lake.

Low densities of native clams and mussels in the Connecticut River may suggest availability of an unoccupied niche for *Corbicula* to colonize. In addition, high water flow of the Connecticut River may provide a steady supply of nutrients and food for filter feeders, further limiting the effects of *Corbicula* on native species.

## Conclusion

Within 2 years of their discovery in the Connecticut River, *Corbicula* colonized 38 km of river upstream of the CY plant. The *Corbicula* population in the vicinity of the CY plant was highly variable within and between years. However, annual densities during plant operation in 1991–1996 were not significantly different from those following the plant closure in 1997–2000. This suggests that the CY thermal discharge did not serve as an important refuge area for *Corbicula* overwintering in the vicinity of the plant. Continuous chlorination (at <0.10 ppm) was effective in preventing *Corbicula* from fouling service water systems at CY. The lack of correlation between presence of *Corbicula* and abundance of native clams and mussels suggest no detrimental effect of *Corbicula* on native species in the Con-

necticut River during these studies. Finally, the success of *Corbicula* in more northerly ranges of New England may not always be associated with the thermal plumes from industrial facilities, but could also result from other thermal refugia such as groundwater seeps that are above 2°C.

## Acknowledgments

This research was conducted by Northeast Utilities Service Company (NU) as a Connecticut State Department of Environmental Protection permit requirement for the operation of the Connecticut Yankee nuclear power plant. The authors' gratitude is extended to John Tietjen, Robert Wilce, Guy Hoffman, Ernesto Lorda, Douglas G. Smith, and Edward Roseman for their critical reviews of this paper. Special appreciation is given to Bob Parker, Bill McDermott, and all the NU Production Operation Services Laboratory personnel for their many hours of boat work and sampling. We also extend this appreciation to our colleagues at the Millstone Environmental Laboratory for their support during sample collection and processing.

## References

- Aldridge, D. W., and R. F. McMahon. 1978. Growth, fecundity, and bioenergetics in a natural population of freshwater clam, *Corbicula manilensis* Philippi, from north central Texas. *Journal of Molluscan Studies* 44:49–70.
- Bickel, D. 1966. Ecology of *Corbicula manilensis* in the Ohio River at Louisville, Kentucky. *Sterkiana* 23:19–24.
- Boyd, W. A. 2004. Hydrology. Pages xxx–xxx in P. M. Jacobson, D. A. Dixon, W. C. Leggett, B. C. Marcy, Jr., and R. R. Massengill, editors. *The Connecticut River Ecological Study (1965–1973) revisited: ecology of the lower Connecticut River 1973–2003*. American Fisheries Society, Monograph 9, Bethesda, Maryland. (Originally published in 1976)
- Britton, J. C., D. R. Coldiron, L. P. Evans, Jr., C. Golightly, K. D. O'Kane, and J. R. TenEyck. 1979. Reevaluation of the growth pattern in *Corbicula fluminea* (Müller). Pages 177–192 in J. C. Britton, editor. *Proceedings, First International Corbicula Symposium*. October 13–15, 1977. Texas Christian University, Fort Worth.
- Britton, J. C., and B. Morton. 1979. *Corbicula* in North America: the evidence reviewed and evaluated. Pages 249–287 in J. C. Britton, editor. *Proceedings, First International Corbicula Symposium*.

- October 13–15, 1977. Texas Christian University, Fort Worth.
- Cairns, J., Jr., and D. S. Cherry. 1983. A site-specific field and laboratory evaluation of fish and Asiatic clam population responses to coal-fired power plant discharges. *Water Science and Technology* 15:10–37.
- Chen, L. 1990. *Aquaculture in Taiwan*. Fishing News Books, Cambridge, Massachusetts.
- Cherry, D. S., J. H. Rodgers, Jr., R. L. Graney, and J. Cairns, Jr. 1980. Dynamics and control of the Asiatic clam in the New River. Virginia. *Bulletin of Virginia Water Resources Research Center* 123:1–72.
- Cherry, D. S., R. L. Roy, R. A. Lechleitner, P. A. Dunhardt, G. T. Peters, and J. Cairns, Jr. 1986. *Corbicula* fouling and control measures at the CELCO plant, Virginia. Pages 69–81 in J. C. Britton and R. S. Prezant, editors. Second International *Corbicula* Symposium. June 21–24, 1983, Hattiesburg, Mississippi. *American Malacological Bulletin Special Edition* 2.
- Counts, C. L., III. 1986. The zoogeography and history of the invasion of the United States by *Corbicula fluminea*. Pages 7–39 in J. C. Britton and R. S. Prezant, editors. Second International *Corbicula* Symposium. June 21–24, 1983, Hattiesburg, Mississippi. *American Malacological Bulletin Special Edition* 2.
- Dauble D. D., D. S. Daly, and C. S. Abernethy. 1985. Factors affecting growth and survival of the Asiatic clam, *Corbicula* sp., under controlled laboratory conditions. Pages 134–144 in R. D. Cardwell, R. Purdy, and R. C. Bahner, editors. *Aquatic Toxicology and Hazard Assessment: Seventh Symposium*, ASTM, STP854. American Society for Testing and Materials, Philadelphia.
- Dreier, H. 1977. Study of *Corbicula* in Lake Sangchris. Pages 1–52 in *The Annual Report for Fiscal Year 1976, Lake Sangchris Project. Section 7. Illinois Natural History Survey, Urbana*.
- Diaz, R. J. 1994. Response of tidal freshwater macrobenthos to sediment disturbance. *Hydrobiologia* 278:201–212.
- Dreier, H., and J. A. Tranquilli. 1981. Reproduction, growth, distribution and abundance of *Corbicula* in an Illinois cooling lake. Pages 378–393 in R. W. Lawrimore and J. A. Tranquilli, editors. *The Lake Sangchris study: case history of an Illinois cooling lake. Bulletin Illinois Natural History Survey* 32.
- Eng, L. L. 1979. Population dynamics of the Asiatic clam, *Corbicula fluminea* (Müller), in the concrete-lined Delta-Mendota Canal of Central California. Pages 39–68 in J. C. Britton, editor. *Proceedings, First International Corbicula Symposium*. October 13–15, 1977. Texas Christian University, Fort Worth.
- Foe, C., and A. Knight. 1986. A thermal energy budget for juvenile *Corbicula fluminea*. *American Malacological Bulletin* 2:143–150.
- French, J. R. P., III, and D. W. Schloesser. 1991. Growth and over-winter survival of the Asiatic clam, *Corbicula fluminea*, in the St Clair River, Michigan. *Hydrobiologia* 219:165–170.
- French, J. R. P., III, and D. W. Schloesser. 1996. Distribution and winter survival health of Asian clams, *Corbicula fluminea*, in the St Clair River, Michigan. *Journal of Freshwater Ecology* 11:183–192.
- Fuller, S. L. H., and M. J. Imlay. 1976. Spatial competition between *Corbicula manilensis* (Philippi), the Chinese clam (Corbiculidae), and fresh-water mussels (Unionidae) in the Waccamaw River Basin of the Carolinas (Mollusca: Bivalvia). *Association of Southeastern Biologists Bulletin* 23:60.
- Gardner, J. A., Jr., W. R. Woodall, Jr., A. A. Staats, Jr., and J. F. Napoli. 1976. The invasion of the Asiatic clam (*Corbicula manilensis* Philippi) in the Altamaha River, Georgia. *Nautilus* 90:117–125.
- Goss, L. B., and C. Cain, Jr. 1975. Power plant condenser and service water system fouling by *Corbicula*, the Asiatic clam. Presented at the Biofouling Workshop held at Johns Hopkins University, Baltimore, Maryland, June 16–19, 1975.
- Goss, L. B., J. M. Jackson, H. B. Flora, B. G. Isom, C. Gooch, S. A. Murray, C. G. Burton, and W. S. Bain. 1979. Control studies on *Corbicula* for steam-electric generating plants. Pages 139–151 in J. C. Britton, editor. *Proceedings, First International Corbicula Symposium*. October 13–15, 1977. Texas Christian University, Fort Worth.
- Graney, R. L., D. S. Cherry, J. H. Rodgers, Jr., and J. Cairns, Jr. 1980. The influence of thermal discharges and substrate composition on the population structure and distribution of the Asiatic clam, *Corbicula fluminea*, in the New River, Virginia. *Nautilus* 94:130–135.
- Horning, W. B., and L. Keup. 1964. Decline of the Asiatic clam in the Ohio River. *Nautilus* 78:29–30.
- Ingram, W. M. 1959. Asiatic clams as potential pests in California water supplies. *Journal of the American Water Works Association* 51:363–370.
- Isom, B. G. 1971. Evaluation and control of macroinvertebrate nuisance organisms in freshwater industrial supply system. Presented at the 24th Annual Meeting of the Midwest (N. A.) Benthological Society.
- Isom, B. G. 1986. Historical review of the Asiatic clam (*Corbicula*) invasion and biofouling of waters and industries in the Americas. Pages 1–5 in J. C. Britton

- and R. S. Prezant, editors. Second International *Corbicula* Symposium. June 21–24, 1983, Hattiesburg, Mississippi. American Malacological Bulletin Special Edition No. 2.
- Johnson, K. I., C. H. Henager, T. L. Page, and P. F. Hayes. 1986. Engineering factors influencing *Corbicula* fouling in nuclear service water systems. Pages 47–52 in J. C. Britton and R. S. Prezant, editors. Second International *Corbicula* Symposium. June 21–24, 1983, Hattiesburg, Mississippi. American Malacological Bulletin Special Edition 2.
- Joy, J. E. 1985. A 40-week study on growth of the Asian clam, *Corbicula fluminea* (Müller), in the Kanawha River, West Virginia. *Nautilus* 99:110–116.
- Karatayev, A. Y., L. E. Burlakova, T. Kesterson, and D. K. Padilla. 2003. Dominance of the Asiatic clam, *Corbicula fluminea* (Müller), in the benthic community of a reservoir. *Journal of Shellfish Research* 22:487–493.
- Kraemer, L. S. 1979. Juvenile *Corbicula*: their distribution in the Arkansas River benthos. Pages 89–97 in J. C. Britton, editor. Proceedings, First International *Corbicula* Symposium. October 13–15, 1977. Texas Christian University, Fort Worth.
- Lauritsen, D. D. 1986. Assimilation of radiolabeled algae by *Corbicula*. Pages 219–222 in J. C. Britton and R. S. Prezant, editors. Second International *Corbicula* Symposium. June 21–24, 1983, Hattiesburg, Mississippi. American Malacological Bulletin Special Edition No. 2.
- Leff, L. G., J. L. Burch, and J. V. McArthur. 1990. Spatial distribution, seston removal and potential competitive interactions of the bivalves *Corbicula fluminea* and *Elliptio complanata*, in a coastal plain stream. *Freshwater Biology* 24:409–416.
- Magdych, W. P. 1981. An efficient, inexpensive elutriator design for separating benthos from sediment samples. *Hydrobiologia* 85:157–159.
- Mattice, J. S. 1979. Interactions of *Corbicula* sp. with power plants. Pages 119–138 in J. C. Britton, editor. Proceedings, First International *Corbicula* Symposium. October 13–15, 1977. Texas Christian University, Fort Worth.
- Mattice, J. S., and L. L. Dye. 1976. Thermal tolerance of the adult Asiatic clam. Pages 130–135 in G. W. Esch and R. W. McFarlane, editors. Thermal ecology II. Conference 750425. National Technical Information Services, Springfield, Virginia.
- Mattice J. S., R. B. McLean, and M. B. Burch. 1982. Evaluation of short-term exposure to heated water and chlorine for control of the Asiatic clam (*Corbicula fluminea*). Oak Ridge National Laboratory, Environmental Services Division Publication No. 1748, Oak Ridge, Tennessee, operated by Union Carbide Corporation for the Department of Energy.
- Mattice, J. S., and L. L. Wright. 1986. Aspects of growth of *Corbicula fluminea*. Pages 167–178 in J. C. Britton and R. S. Prezant, editors. Second International *Corbicula* Symposium. June 21–24, 1983, Hattiesburg, Mississippi. American Malacological Bulletin Special Edition 2.
- McMahon, R. F. 1977. Shell size-frequency distributions of *Corbicula manilensis* from a clam-fouled steam condenser. *Nautilus* 91:54–59.
- McMahon, R. F. 1982. The occurrence and spread of the introduced Asiatic freshwater clam, *Corbicula fluminea*, in North America: 1924–1982. *Nautilus* 96:134–141.
- McMahon, R. F. 1983. Ecology of an invasive pest bivalve, *Corbicula*. Pages 505–561 in W. D. Russell-Hunter, editor. The mollusca. Volume 6, Ecology. Academic Press, Orlando, Florida.
- McMahon, R. F., and R. W. Lutey. 1988. Field and laboratory studies of the efficacy of Poly[oxyethylene(dimethyliminio)ethylene(dimethyliminio)ethylene dichloride] as a biocide against the Asian clam, *Corbicula fluminea*. Presented at Service Water System Reliability Improvement Seminar on October 17–19, 1988, Charlotte, North Carolina. Sponsored by Electric Power Research Institute, Palo Alto, California.
- McMahon, R. F., and C. J. Williams. 1986a. A reassessment of growth rate, life span, life cycles and population dynamics in a natural population and field caged individuals of *Corbicula fluminea* (Müller) (Bivalvia: Corbiculacea). Pages 151–166 in J. C. Britton and R. S. Prezant, editors. Second International *Corbicula* Symposium. June 21–24, 1983, Hattiesburg, Mississippi. American Malacological Bulletin Special Edition 2.
- McMahon, R. F., and C. J. Williams. 1986b. Growth, life cycle, upper thermal limit and downstream colonization rates in a natural population of the freshwater bivalve mollusc, *Corbicula fluminea* (Müller) receiving thermal effluent. Pages 231–239 in J. C. Britton and R. S. Prezant, editors. Second International *Corbicula* Symposium. June 21–24, 1983, Hattiesburg, Mississippi. American Malacological Bulletin Special Edition 2.
- Merriman, D., and L. M. Thorpe. 2004. Introduction. Pages xxx–xxx in P. M. Jacobson, D. A. Dixon, W. C. Leggett, B. C. Marcy, Jr., and R. R. Massengill, editors. The Connecticut River Ecological Study (1965–1973) revisited: ecology of the lower Connecticut River 1973–2003. American Fisheries Society, Monograph 9, Bethesda, Maryland. (Originally published in 1976)

- Morgan, D. E., M. Keser, J. F. Foertch, and J. M. Vozarik. 1992. The Asiatic clam, *Corbicula fluminea*, extends its North American distribution to the Connecticut River (abstract only). Page 523 in W. J. Blogoslawski, editor. Aquaculture and environmental stewardship: Milford Shellfish Biology Seminar - 1991. Environmental Management 16:521–529.
- Morgan, D. E., M. Keser, J. T. Swenarton, and J. F. Foertch. 2003. Population dynamics of the Asiatic clam, *Corbicula fluminea* (Müller) in the lower Connecticut River: establishing a foothold in New England. Journal of Shellfish Research 22:193–203.
- Neitzel, D. A., K. I. Johnson, T. L. Page, J. S. Young, and P. M. Daling. 1984. Bivalve fouling of nuclear power plant service water systems In Volume 1, Correlation of bivalve biological characteristics and raw water system design. U.S. Nuclear Regulatory Commission, NUREG/CR-4070, PNL-5300, Washington, D. C.
- NSI (Nautical Software, Inc.). 1995. Tides and currents for Windows, version 2.0. Beaverton, Oregon. Available: nauticalsw@aol.com.
- O’Kane, K. D. 1976. A population study of the exotic bivalve *Corbicula manilensis* (Philippi, 1841) in selected Texas reservoirs. Master’s thesis. Texas Christian University, Fort Worth.
- Page, T. L., D. A. Neitzel, M. A. Simmons, and P. F. Hayes. 1986. Biofouling of power plant service systems by *Corbicula*. Pages 41–45 in J. C. Britton and R. S. Prezant, editors. Second International *Corbicula* Symposium. June 21–24, 1983, Hattiesburg, Mississippi. American Malacological Bulletin Special Edition 2.
- Phelps, H. L. 1993. *Corbicula* production and marketing techniques in Taiwan. Report to the University of the District of Columbia Research Council, Washington, D.C.
- Phelps, H. L. 1994. The Asiatic clam (*Corbicula fluminea*) invasion and system-level ecological change in the Potomac River Estuary near Washington, D.C. Estuaries 17:614–621.
- Prezant, R. S., and K. Chalermwat. 1984. Flotation of the bivalve *Corbicula fluminea* as a means of dispersal. Science 225:1491–1493.
- Rodgers, J. H., Jr., D. S. Cherry, K. L. Dickson, and J. Cairns, Jr. 1977. Invasion of Asiatic clam, *Corbicula manilensis* in the New River, Virginia. Nautilus 9:43–46.
- Rodgers, J. H., Jr., D. S. Cherry, K. L. Dickson, and J. Cairns, Jr. 1979. Invasion population dynamics and elemental accumulation of *Corbicula fluminea* in the New River at Glen Lyn, Virginia. Pages 99–110 in J. C. Britton, editor. Proceedings, First International *Corbicula* Symposium. October 13–15, 1977. Texas Christian University, Fort Worth.
- Rozsa, R. 2001. Hydrology of the lower Connecticut River. Pages 18–22 in G. D. Dreyer and M. Caplis, editors. Living resources and habitats of the lower Connecticut River. December 2001. The Connecticut College Arboretum Bulletin 37, New London, Connecticut.
- Savoy, T. F., and J. Benway. 2004. Food habits of shortnose sturgeon collected in the lower Connecticut River from 2000 through 2002. Pages xxx–xxx in P. M. Jacobson, D. A. Dixon, W. C. Leggett, B. C. Marcy, Jr., and R. R. Massengill, editors. The Connecticut River Ecological Study (1965–1973) revisited: ecology of the lower Connecticut River 1973–2003. American Fisheries Society, Monograph 9, Bethesda, Maryland.
- Sickel, J. B. 1979. Population dynamics of *Corbicula* in the Altamaha River, Georgia. Pages 69–80 in J. C. Britton, editor. Proceedings, First International *Corbicula* Symposium. October 13–15, 1977. Texas Christian University, Fort Worth.
- Sickel, J. B. 1986. *Corbicula* population mortalities: factors influencing population control. Pages 89–94 in J. C. Britton and R. S. Prezant, editors. Second International *Corbicula* Symposium. June 21–24, 1983, Hattiesburg, Mississippi. American Malacological Bulletin Special Edition 2.
- Sinclair, R. M., and B. G. Isom. 1963. Further studies on the introduced Asiatic clam *Corbicula* in Tennessee. Tennessee Stream Pollution Control Board.
- Sites, D. L., A. C. Benke, and D. M. Gillespie. 1995. Population dynamics, growth, and production of the Asiatic clam, *Corbicula fluminea*, in a blackwater river Canadian Journal of Fisheries and Aquatic Sciences 52:425–437.
- USGS (United States Geological Survey). 2002. Water resources of the United States, surface-water data for Connecticut, August 15, 2002. Available: <http://waterdata.usgs.gov/ct/nwis/sw>.
- USNRC (United States Nuclear Regulatory Commission). 1981. Flow blockage of cooling water to safety system components by *Corbicula* sp (Asiatic clam) and *Mytilus* sp. (mussel). IE Bulletin no. 81–03, Office of Inspection and Enforcement, United States Nuclear Regulatory Commission, Washington, D.C.
- Voelz, N. J., J. V. McArthur, and R. B. Rader. 1998. Upstream mobility of the Asiatic clam *Corbicula fluminea*: identifying potential dispersal agents. Journal of Freshwater Ecology 13:39–45.
- Welch, K. J., and J. E. Joy. 1984. Growth rates of the Asiatic Clam, *Corbicula fluminea* (Müller), in the Kanawha River, West Virginia. Freshwater Invertebrate Biology 3:139–142.

