

17.7.46

Superfund Records Center
SITE: New Bedford
BREAK: 177
OTHER: 268621

Mutation Research, 220 (1989) 31-47
Elsevier



MTR 07264

SDMS DocID 268621

Polychlorinated biphenyls (PCBs): mutagenicity and carcinogenicity

Stephen Safe

*Department of Physiology and Pharmacology, College of Veterinary Medicine, Texas A & M University,
College Station, TX 77840 (U.S.A.)*

(Received 29 August 1988)

(Accepted 15 September 1988)

Keywords: Polychlorinated biphenyls; Mutagenicity; Carcinogenicity

Contents

1. Introduction	31
2. Mutagenic activity of PCBs	32
i. Metabolism and macromolecular binding	32
ii. Mutagenicity	34
3. Carcinogenicity of PCBs — laboratory animal studies	34
i. Cancer as a multi-stage process	34
ii. Chronic feeding studies	35
iii. PCBs as modulators of carcinogenesis	37
4. Carcinogenicity of PCBs — human studies	41
5. Summary and conclusions	42
Acknowledgements	42
References	43

(1) Introduction

Commercial polychlorinated biphenyls (PCBs) (Fig. 1) were widely used as industrial compounds with such diverse applications as plasticizers, heat transfer fluids, organic diluents, cutting oils, wax extenders, adhesives, flame retardants and dielectric fluids for capacitors and transformers (Hutzinger et al., 1974; Pomerantz et al., 1978). It has been estimated that up to 1.4×10^9 pounds of PCBs were produced in the U.S.A. PCBs were detected in the environment in the late 1960s (Risebrough et al., 1968) and within a short time

these compounds were reported as contaminants in almost every component of the global ecosystem including air, water, fish, wildlife and human blood, adipose tissue and milk (Bush et al., 1985, 1986; Buckley, 1982; Ballschmiter et al., 1981; Wasserman et al., 1979; Safe, 1982; Holdrinet et al., 1977; Kannan et al., 1988; Tanabe, 1988). In the 1970s, all 'open' uses for PCBs were discontinued, however, the 'closed' applications of PCBs as dielectric fluids in capacitors and transformers were continued until 1978. New transformers and capacitors now contain alternative dielectric fluids and as older PCB-containing electrical equipment is withdrawn from service or repaired, the PCB fluids are replaced.

Commercial PCBs are prepared by the chlorination of biphenyl and the products are graded and sold according to their chlorine content. The

Correspondence: Dr. Stephen Safe, Department of Physiology and Pharmacology, College of Veterinary Medicine, Texas A & M University, College Station, TX 77840 (U.S.A.).




Fig. 1. The structure of polychlorinated biphenyls and the numbering system.

lower chlorinated PCBs typified by Aroclor 1221 (21% by weight of Cl; 1.15 Cl/biphenyl) and the higher chlorinated products typified by Aroclor 1260 (60% by weight of Cl; 6.30 Cl/biphenyl) exhibit markedly different physical chemical properties and use patterns (Hutzinger et al., 1974). Moreover, it is apparent from gas chromatographic analysis of the commercial products that PCB mixtures also differ with respect to the individual congeners present in these mixtures and their relative concentrations (Mullin et al., 1984; Safe et al., 1985a; Albro et al., 1981; Alford-Stevens et al., 1986; Jensen and Sundstrom, 1974; Ballschmiter and Zell, 1980). Obviously there is considerable overlap between the individual PCB congeners present in mixtures which contain similar % chlorine compositions and the extent of PCB congener overlap decreases with increasing differences in the degree of chlorination of the PCB mixtures.

The introduction of commercial PCBs into the ecosystem results in their subsequent translocation and uptake into different environmental components (Safe et al., 1985a). Since individual PCBs exhibit different physicochemical properties and susceptibilities to environmental breakdown, it is not surprising that the composition of most PCB extracts from environmental samples do not resemble the commercial PCB mixtures (Safe et al., 1985a; Wolff et al., 1982, 1986; Hansen et al., 1983; Brown et al., 1985, 1987). Conventional

analysis of these environmental mixtures usually estimates 'total PCBs' using an appropriate commercial PCB cocktail as a surrogate quantitative standard (Webb and McCall, 1973). However, it is apparent from high-resolution gas chromatographic analyses that the congener composition and relative concentrations of the individual components in many PCB extracts from environmental samples differ markedly from the commercial PCBs (Safe et al., 1985a; Wolff et al., 1985, 1986; Jensen and Sundstrom, 1974). The differences in the composition of PCB mixtures are reflected in toxicity differences since the higher chlorinated products such as Aroclor 1254 and Aroclor 1260 are more toxic than the lower chlorinated PCBs (Safe, 1984; Fishbein, 1974; Nelson et al., 1972). This is not surprising since extensive structure-activity (including toxicity) studies with individual PCB congeners have demonstrated that the most toxic PCB congeners contain 5 or 6 Cl groups/biphenyl (Safe, 1984, 1985b; Parkinson et al., 1983; Leece et al., 1985). Since congener-specific analyses of few environmental mixtures have been determined, their composition and potential toxicities are therefore largely unknown.

(2) Mutagenic activity of PCBs

(i) Metabolism and macromolecular binding

A large number of PCB metabolites have been identified in both in vivo and in vitro studies in several animal species, microorganisms and also with microsome preparations (reviewed in Matthews and Dedrick, 1984; Safe, 1980; Schnellmann et al., 1985; Sundstrom et al., 1976). The major metabolic products are invariably phenolic compounds however, dihydroxy, dihydrodiol, phenol conjugate and thioether metabolites have also been identified. In addition, several studies have also identified hydroxylated/dechlorinated products as PCB metabolites (Hass et al., 1977; Safe et al., 1975a,b, 1976; Sundstrom et al., 1976). The metabolic pathways associated with PCB metabolism have been investigated using specifically labeled substrates (Schnellmann et al., 1984; Safe et al., 1975a) and induced microsomal enzymes which exhibit different monooxygenase enzyme activities (Kennedy et al., 1980, 1981; Kaminsky et al., 1981). Fig. 2 summarizes the pathways




Fig. 2

assoc:
2,2',5,
vole
of an
rearr.
addit
phase
al., 19
Melan
labeled
some
tion o
Allen,
synth
(Reich
solutic
Allen,
non-ar




Fig. 2. Metabolism of 2,2',5,5'-tetrachlorobiphenyl.

associated with the oxidative metabolism of 2,2',5,5'-tetrachlorobiphenyl. The initial step involves the cytochrome P-450-dependent formation of an unstable arene oxide intermediate which can rearrange to form phenolic metabolites or undergo additional metabolic processing by phase I and phase II drug-metabolizing enzymes (Norback et al., 1976; Hsu et al., 1975; Gardner et al., 1973; Melancon and Lech, 1976). Incubation of radio-labeled 2,2',5,5'-tetrachlorobiphenyl with microsome preparations has resulted in the identification of the arene oxide intermediate (Forgue and Allen, 1982) and it has also been reported that synthetic 3,4-oxo-2,2',5,5'-tetrachlorobiphenyl (Riech et al., 1978) spontaneously rearranges in solution to give the phenolic products (Forgue and Allen, 1982) (Fig. 2). It has also been reported that non-arene oxide pathways are prevalent in the

metabolism of 2,2',5,5'-tetrachlorobiphenyl using phenobarbital-induced microsomes (Preston et al., 1984). However, data from most studies suggest that oxidative metabolism of PCBs results in the initial formation of arene oxide intermediates which are known to be strong electrophiles and alkylating agents.

In vivo and in vitro studies with individual PCBs and PCB mixtures have demonstrated that these hydrocarbons can undergo metabolic activation and alkylate cellular macromolecules (Wong et al., 1979; Wyndham and Safe, 1978; Morales et al., 1979; Hesse and Wolff, 1977; Hesse et al., 1978; Shimada and Sato, 1978; Hargraves and Allen, 1979; Wyndham et al., 1976). For example, incubation of [^3H]4-chlorobiphenyl with Aroclor 1248-induced rat liver microsomes resulted in the formation of protein, RNA and DNA adducts

usually
e com-
titative
er. it is
omato-
osition
al com-
onmen-
mercial
. 1986;
nces in
cted in
rinated
or 1260
l PCBs
(1972).
re-ac-
ividual
e most
roups /
.. 1983;
nalyzes
en de-
toxici-

e been
dies in
id also
1 Mat-
ichnell-
5). The
henolic
). phe-
ve also
s have
1 prod-
Safe et
). The
etabo-
ifically
4; Safe
nzymes
enzyme
minsky
thways

with 4-chlorobiphenyl and incubation with homopolynucleotides indicated that poly G was more readily adducted than poly A, poly C or poly U (Wyndham and Safe, 1978). Comparable in vitro adduction of cellular macromolecules by other individual PCBs and PCB mixtures has also been reported. Morales and Matthews (1979) administered [^{14}C]2,2',4,4',5,5'- and [^{14}C]2,2',3,3',6,6'-hexachlorobiphenyl to mice and reported the in vitro incorporation of radioactivity into cellular protein and RNA for both compounds and into DNA only for the 2,2',3,3',6,6'-hexachlorobiphenyl isomer. The higher macromolecular binding (i.e., protein, RNA and DNA) by the latter isomer was consistent with its more rapid rate of metabolism (i.e., compared to the poorly metabolized 2,2',4,4',5,5'-hexachlorobiphenyl).

(iii) Mutagenicity

It was initially reported by Wyndham and coworkers (1976) that the 4-chlorobiphenyl and the lower chlorinated PCB mixture, Aroclor 1221, were mutagenic to *Salmonella typhimurium* strain TA1538 in the presence of an exogenous source of metabolic activation (S9). More highly chlorinated biphenyls (Aroclor 1254 and 2,2',5,5'-tetrachlorobiphenyl) were not mutagenic; these results were consistent with the facile metabolic activation of lower chlorinated biphenyls and the activity of 4-chlorobiphenyl to cause unscheduled DNA synthesis in Chinese hamster ovary cells in culture (Wong et al., 1979). However, subsequent studies in the authors' laboratory (S. Safe, unpublished results) failed to reproduce the original observations. Moreover, Schoeny and coworkers (1979; Schoeny, 1982) have reported the lack of bacterial mutagenicity of Aroclor 1254, 4-chlorobiphenyl, 3,3',4,4'- and 2,2',4,4'-tetrachlorobiphenyl and 2,2',4,4',6,6'-hexachlorobiphenyl. Hsia et al. (1978) have also shown that 2,2',5,5'-tetrachlorobiphenyl, 4-hydroxy-2,2',5,5'-tetrachlorobiphenyl (a metabolite, see Fig. 2) and 3,4-oxo-2,2',5,5'-tetrachlorobiphenyl (arene oxide intermediate, see Fig. 2) were inactive as bacterial mutagens in *Salmonella typhimurium* strain TA1537.

Heddle and Bruce (1977) reported that Aroclor 1254 did not cause cytogenic effects in mice (i.e., micronuclei and abnormal sperm production) and Aroclors 1242 and 1254 did not cause cytogenetic

damage in the bone marrow of rats. It was also concluded that Clophens A-30 and A-60 did not exhibit clastogenic effects in *Drosophila melanogaster* (Nilsson and Ramel, 1974), however, Peakall and coworkers (1972) observed some chromosomal aberrations in ring dove embryos treated with Aroclor 1254. These results suggested possible clastogenic effects of PCBs in ring dove embryos.

(3) Carcinogenicity of PCBs — laboratory animal studies

(i) Cancer as a multi-stage process

The development of cellular neoplasia is a highly complex process which has been investigated using several different model systems (Farber, 1984a,b; Schulte-Hermann, 1985; Farber and Sarma, 1987; Pitot et al., 1987; Slaga, 1984). In the liver, a primary target organ for PCBs, several stages in the sequential development of hepatocellular cancer have been identified or proposed (Fig. 3). The initiation step in cancer development is in itself highly complex in which only one of the steps is presumed to involve irreversible cellular (DNA) damage. For chemical carcinogens, this step in the initiation process is believed to be associated with covalent adduction of DNA. Cellular damage such as the formation of carcinogen-DNA adducts is readily repaired in most tissues and, in the liver, the development of initiated hepatocytes requires a round of cell proliferation following the initial damage. The conversion of the initiated cell populations into focal proliferations such as nodules (liver), papillomas (skin) and polyps (colon) is a complex process known as promotion. Several classes of organ-specific promoters have been identified and have been utilized to study the biochemical and cellular process associated with the development of this altered cell population. The subsequent conversion of initiated cells which have been promoted into cancer cells is known as progression. The specific steps associated with this multi-stage process are not well defined, however Fig. 3 (adapted from Farber, 1986) summarized a proposed model which describes the sequential development of liver cancer in which several of the steps have either been identified or inferred from experimental

CARCINO
+
LIVER C

INITIA
CELLS

HEPAT
NODUL

Fig. 3

studies
the de
act
ever, it
may
comple
cells
review
1987

(iii) Ch
Sev
effect
tration

TABLE
THE IN
SHERM
(1986)
(Kimm

Type of

Hepatic
Neoplas
Focal
cystic




Fig. 3. Some of the proposed steps in the development of hepatic cancer (from Farber, 1986).

studies. Most studies on the role of chemicals in the development of cancer have focused on their activity as initiators or as tumor promoters. However, it is conceivable that chemicals such as PCBs may act at more than one step in the highly complex conversion of normal cells into neoplastic cells and this possibility will be noted in this review and has been discussed in detail by Hayes (1987).

(ii) Chronic feeding studies

Several studies have reported the carcinogenic effects of commercial PCBs after chronic administration. In male and female Donryu rats, Kimura

TABLE 1

THE INCIDENCE AND TYPE OF LIVER LESIONS IN SHERMAN STRAIN FEMALE RATS FED AROCLOR 1260 (1000 ppm) FOR APPROXIMATELY 21 MONTHS (Kimbrough et al., 1975)

Type of lesion	Incidence	
	Controls	PCB-treated
Hepatocellular carcinomas	1/173	26/184
Neoplastic nodules	0/173	144/184
Foci or areas of cytoplasmic alteration	28/173	182/184

and Baba (1973) reported adenomatous nodule formation in the liver of female (6/10) but not male rats; however, it was apparent that the dose levels used in this study were toxic. Ito and co-workers (1974) investigated the effects of chronic feeding of several Kanechlor preparations to male Wistar rats and observed nodular hyperplasia in some of the treatment groups. Tables 1 and 2 summarize results of 2 studies in which Aroclor 1260 was utilized. Exposure of female Sherman rats to Aroclor 1260 (100 ppm) in the diet for approximately 21 months results in 14% (26/184) hepatocellular carcinoma and 78% neoplastic nodules (144/184) in the liver whereas only 0.58% (1/173) and 0% (0/173) were observed in the control animals (Kimbrough et al., 1975). In a second long-term study (Norback and Weltman, 1985) using Aroclor 1260 (Table 2), it was demonstrated that this PCB preparation caused hepatocellular trabecular carcinoma, adenocarcinoma and neoplastic nodules in Sprague-Dawley rats. Moreover, it was apparent that female rats were more highly susceptible than males to the hepatocarcinogenic effects of Aroclor 1260.

The National Cancer Institute (1978) investigated the effects of dietary administration of Aroclor 1254 on male and female F344 rats at dose levels of 0, 25, 50 and 100 ppm in the diet for 105

TABLE 2
EFFECTS OF AROCLOR 1260 ON HEPATOCELLULAR NEOPLASM FORMATION IN MALE AND FEMALE SPRAGUE-DAWLEY RATS (Norback and Weltman, 1984) ^a

Response	Treatment	% Incidence		
		Total (n)	Female (n)	Male (n)
Trabecular carcinoma ^a	Aroclor 1260	23 (21)	40 (19)	4 (2)
	Corn oil	0	0	0
Adenocarcinoma ^{b,c}	Aroclor 1260	26 (24)	51 (24)	0
	Corn oil	0	0	0
Neoplastic nodule only	Aroclor 1260	8 (7)	4 (2)	11 (5)
	Corn oil	1 (1)	2 (1)	0
Negative	Aroclor 1260	44 (41)	4 (2)	85 (39)
	Corn oil	99 (80)	98 (48)	100
Total number of animals	Aroclor 126	93	47	46
	Control	81	49	32

^a Included in the table are data from 8 male and 7 female rats in the group administered Aroclor 1260 and 8 male and 10 female rats in the control (corn oil) group that had received partial hepatectomy during the first 18 months of the study.

^b Animals containing neoplastic nodules plus carcinoma were only included in the carcinoma category.

^c Animals with trabecular carcinoma and adenocarcinoma were only placed in the adenocarcinoma category.

weeks. The results summarized in Table 3 illustrate a dose-dependent increase of hepatocellular carcinomas and hyperplastic nodules in male and female rats and a significant increase in intestinal metaplasia (male + female rats data combined) (Ward, 1985) at the 100 ppm dose level. In contrast to the results obtained using Aroclor 1260 (Norback and Weltman, 1985), Aroclor 1254 was equally carcinogenic in both male and female F344 rats. Moreover, the incidence of hepatocellular carcinomas was considerably lower

in the Aroclor 1254-treated animals than in the rats treated with Aroclor 1260. These results suggest a possible difference in the carcinogenic potencies of the two PCB mixtures and this must be related to differences in their composition.

The hepatocarcinogenicities of Clophens A60 and A30 were determined in male Wistar rats maintained on diets containing 100 ppm of the commercial PCBs for up to 832 days (Schaeffer et al., 1984). Table 4 summarizes the frequencies of the major lesions observed in the rats at the end of

TABLE 3
EFFECTS OF LONG-TERM FEEDING OF AROCLOR 1254 TO F344 RATS (NCI, 1978; Ward, 1985)

Lesion	Sex	Number of rats with lesions/number treated			
		Dose (ppm)			
		0	25	50	100
Hepatocellular carcinomas	M	0	0	1/24	3/24
	F	0	0	1/24	2/24
Hyperplastic nodules	M	0	5/34	8/24	12/24
	F	0	6/24	9/22	12/24
Intestinal metaplasia	M + F	3/47	4/48	5/48	15/48
Adenocarcinoma gastric	M + F	0/47	1/48	3/48	2/48

TABLE
COMPA
(Schaeff

Lesion

Hepatocellular
Neoplasia
Bile duct
Adenofibrosis
Thymoma
Other neoplasms
Wistar rats

* Signifi

the study
caused
level
hepatocellular
demonstrated
carcinomas
Clophens A60
the results
of the study
relatively
divided
these means
were

The
chronic
Administered
males
250 and
level
carcinomas
Ito et al.
and 10
hepatocellular
BALB/c
1254 (8)
was not
based on
reported
Clophens
the low
cantly
chlorin.

TABLE 4

COMPARATIVE EFFECTS OF CHRONIC FEEDING OF CLOPHEN A60 AND CLOPHEN A30 ON MALE WISTER RATS (Schaeffer et al., 1984)

Lesion	Number of rats with lesions/number treated (%)		
	Control	Clophen A30	Clophen A60
Hepatocellular carcinoma	1/53 (2)	3/87 (3)	52/85 (61) *
Neoplastic nodules	2/53 (4)	35/87 (40)	34/85 (40) *
Bile duct hyperplasia	7/53 (13)	28/87 (32)	62/85 (73)
Adenofibrosis	12/53 (23)	6/87 (7)	2/85 (2) *
Thymoma	9/53 (17)	12/87 (14)	0/85 (0) *
Other neoplasms	19/53 (36)	33/87 (38)	13/85 (15)
Wistar nephritis	21/53 (40)	7/87 (8)	0/85 (0) *

* Significantly different ($P < 0.05$) from control animals.

the study. Both Clophen A60 and Clophen A30 caused the formation of neoplastic nodules in the liver but only the former compound caused hepatocellular carcinomas. These results clearly demonstrated the markedly increased hepatocarcinogenic potency of the higher chlorinated Clophen A60 (60% by weight of Cl) compared to the lower chlorinated Clophen A30 (30% by weight of Cl). These differences must be related to the relative distribution and concentrations of the individual PCB isomers and congeners present in these mixtures (note: polychlorinated dibenzofurans were not detected in the Clophens used in this study).

The carcinogenicity of commercial PCBs after chronic feeding to mice has also been reported. Administration of Kanechlors 500, 400 and 300 to male dd mice for 32 weeks at dietary levels of 500, 250 and 100 ppm indicated that at the 500 ppm level, Kanechlor 500 caused hepatocellular carcinomas (5/12, 417%) (Nagasaki et al., 1972; Ito et al., 1973). In a second study (Kimbrough and Linder, 1974) there was an increase in hepatomas (9/22, 40%) observed in male BALB/CJ mice maintained on a diet of Aroclor 1254 (300 ppm) for 11 months whereas this lesion was not observed in the control mice. Moreover, based on the data noted above and the results reported by Schaeffer and coworkers (1984) for Clophen A30 and Clophen A60, it is apparent that the lower chlorinated PCB mixtures were significantly less potent as carcinogens than the higher chlorinated commercial products.

(iii) PCBs as modulators of carcinogenesis

PCBs are known to interact with structurally diverse carcinogens (initiators) and modulate their mutagenic and carcinogenic potencies. For example, PCB mixtures induce several phase I and phase II drug-metabolizing enzymes including glutathione *S*-transferases, epoxide hydrolase, glucuronyl transferases and several cytochrome P-450-dependent monooxygenases (reviewed in Safe, 1984). Commercial PCBs such as Aroclor 1254 and individual isomers and congeners induce a spectrum of cytochrome P-450 isozymes (cytochromes P-450a, P-450b, P-450c, P-450d, P-450e and P-450p) (Parkinson et al., 1983; Ryan et al., 1979a,b, 1981, 1982; Scheutz et al., 1986; Thomas et al., 1981, 1983) which in turn catalyze the oxidation of diverse compounds, including carcinogens such as benzo[*a*]pyrene and related polynuclear aromatic hydrocarbons, aflatoxin B₁ and acetylaminofluorene (Astrom and DePierre, 1985; Yang, 1988; Goldstein et al., 1984; McManus et al., 1984; Yoshizawa et al., 1982). Not surprisingly, PCBs can act as cocarcinogens or anticarcinogens which enhance or inhibit the tumorigenic activity of other carcinogenic compounds. For example Kanechlor 500 decreased the incidence of hepatic tumors in rats treated with the hepatocarcinogens 3'-methyl-4-dimethylaminoazobenzene, 2-acetylaminofluorene and diethylnitrosamine in the diet (Makiura et al., 1974). In a similar experiment, Kanechlor 400 also decreased the incidence of 3'-methyl-4-dimethylaminoazobenzene-induced hepatocarcinomas in

FEMALE

male rats

in the
its sug-
nogenic
is must
on.

ns A60
ar rats
of the
ffer et
cies of
end of

rats (Kimura et al., 1976).

The effects of PCBs on aflatoxin B₁-mediated hepatocarcinoma formation in rainbow trout have been extensively investigated. When 6 ppb of aflatoxin B₁ and Aroclor 1254 are simultaneously fed to rainbow trout, the PCBs inhibit the incidence of liver tumors (Hendricks et al., 1977). Similarly, PCB administration prior to aflatoxin B₁ treatment also decreased the liver tumor incidence (Shelton et al., 1983) whereas if Aroclor 1254 is fed after exposure of trout embryo to aflatoxin B₁ there is no effect on the formation of liver tumors. Table 5 summarizes the incidence of hepatic tumors in rainbow trout fed different dietary levels of aflatoxin B₁ (1, 4 and 8 ppb), Aroclor 1254 (50 ppm) and aflatoxin B₁ plus Aroclor 1254 (Shelton et al., 1984). The inhibition of aflatoxin B₁-mediated carcinogenicity was also correlated with the decreased bacterial mutagenicity of this compound in the presence of an Aroclor 1254-induced drug-metabolizing enzyme fraction from fish. The potential mechanisms of this interaction were further investigated (Shelton et al., 1986) by studying the effects of Aroclor 1254 on aflatoxin B₁ distribution, metabolism and DNA adduct formation. The results from the in

vivo studies showed that PCB treatment resulted in a marked increase in the metabolism of aflatoxin B₁ to aflatoxin M₁ and aflatoxicol M₁ and their glucuronide conjugates. Moreover, after 21 days, the DNA adduct levels in the PCB-treated fish were 48–69% lower than in the controls. The results strongly suggest that the anticarcinogenic activity of PCBs in the fish model using aflatoxin B₁ as the carcinogen is associated with the activity of Aroclor 1254 as an inducer of cytochrome P-450-dependent monooxygenases (e.g., such as aflatoxin-4-hydroxylase) (Shelton et al., 1986; Halvorson et al., 1985).

The anticarcinogenic activities of Aroclor 1254 and some PCB congeners have also been reported in the mouse skin model using polynuclear aromatic hydrocarbons as initiators (Berry et al., 1978; DiGiovanni et al., 1977, 1979, 1980). The results summarized in Table 6 show that Aroclor 1254 (100 µg/mouse) administered 18 h prior to the initiator, 7,12-dimethylbenz[*a*]anthracene (DMBA), significantly decreased the incidence of papilloma formation in female Charles River CD-1 mice. 2,2',4,4',5,5'-Hexachlorobiphenyl (625 µg/mouse), a PCB congener which resembles phenobarbital in its mode of incubation of drug-

TABLE 5
HEPATOCARCINOMA INCIDENCE FOLLOWING VARIOUS DIETARY CONCENTRATIONS OF AFB₁ AND AROCLOR 1254 (from Shelton et al., 1984)

Treatment diet	9-month sample		12-months sample	
	Hepatocarcinoma incidence *	%	Hepatocarcinoma incidence	%
1 ppb AFB ₁ ^b	2/52	3.8	27/121	22.3
4 ppb AFB ₁	5/51	9.8	68/126	54.0
8 ppb AFB ₁	32/62	51.6	98/118	83.1
1 ppb AFB ₁ + 50 ppm PCB ^{c,d}	-	-	14/120	11.7
4 ppb AFB ₁ + 50 ppm PCB	3/59	5.1	38/122	31.2
8 ppb AFB ₁ + 50 ppm PCB	12/58	20.7	88/118	74.6
4 ppb AFB ₁ + 5 ppm PCB	10/67	14.9	56/119	47.1
50 ppm PCB	0/74	0	0/120	0
Control	0/70	1.1	0/120	0

* The 9- and 12-month samples show pooled data from duplicate tanks. Less than 80 fish at 9 months and 120 fish at 12 months reflect mortalities during the experiment.

^b The 12-month AFB₁ dose-response curve was linear ($y = 8.60x + 15.84$, $r = 0.944$) with an ED₅₀ of 4.0 ppb.

^c The 12-month AFB₁ + 50 ppm PCB dose-response curve was linear ($y = 9.09x - 0.25$, $r = 0.0991$) with an ED₅₀ of 5.5 ppb.

^d Regression line for AFB₁ + PCB not significantly different from parallel ($0.75 > P > 0.5$) when compared with regression line for AFB₁ alone.

TABLE
EFFECT
INITIA

Pretreat
µg/m

Acetone
Aroclor

Aroclor

Aroclor

Aroclor

Acetone

3,4,3',4'

2,4,5,2',5'

Acetone

TCDD

* Pretre

metab
carcin
was m
Both
trachl
in the
(Parki
hydro
of DN
the PC
of pu
that th
skin t
tered
carcin
oxygen
Sex
PCBs
differ
PCBs
which
as the
N-hy
and 2-
et al.
1982;
1983.
treatr

TABLE 6

EFFECTS OF 2,3,7,8-TCDD, AROCLOR 1254 AND POLYCHLORINATED BIPHENYL CONGENERS ON SKIN TUMOR INITIATION BY DMBA IN CD-1 MICE (from DiGiovanni et al., 1979) ^a

Pretreatment ($\mu\text{g}/\text{mouse}$)	Weeks of promotion	Initiator	Pretreatment time	Papillomas per mouse	Percent of control
Acetone	20	DMBA	5 min	3.80	100
Aroclor 1254 (100)	20	DMBA	5 min	3.60	85
Aroclor 1254 (100)	20	DMBA	18 h	2.1	55
Aroclor 1254 (100)	20	DMBA	3 days	3.3	88
Aroclor 1254 (625)	20	DMBA	18 h	2.70	71
Acetone	24	DMBA	3 days	3.83	100
3,4,3',4'-TCB (625)	24	DMBA	3 days	0.40	10
2,4,5,2',4',5'-HCB (625)	24	DMBA	3 days	3.36	88
Acetone	20	DMBA	5 min	3.80	100
TCDD (1)	20	DMBA	3 days	0.34	9

^a Pretreated mice (30 per group) were initiated with DMBA as described (DiGiovanni et al., 1979).

metabolizing enzymes, did not act as an anticarcinogen, whereas 3,3',4,4'-tetrachlorobiphenyl was more active than Aroclor 1254 as an inhibitor. Both 3,3',4,4'-tetrachlorobiphenyl and 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) resemble MC in their mode of cytochrome P-450 induction (Parkinson et al., 1983) and both halogenated aryl hydrocarbons significantly decreased the number of DMBA-initiated papillomas/mouse. Although the PCB treatment did not modulate the incidence of papillomas caused by B[a]P, it was suggested that the anticarcinogenic effects of PCBs on mouse skin tumors initiated by DMBA were due to altered metabolism and DNA binding of the carcinogen by the PCB-induced skin monooxygenases (DiGiovanni et al., 1979).

Several studies have investigated the activity of PCBs as cancer promoters using a number of different bioassays and target organs. Commercial PCBs promote hepatocarcinogenesis in rodents which have been initiated with carcinogens such as the isomeric hexachlorocyclohexanes, *N*-ethyl-*N*-hydroxyethylnitrosamine, diethylnitrosamine and 2-acetylaminofluorene (Ito et al., 1973; Hirose et al., 1981; Kimura et al., 1976; Periera et al., 1982; Tatematsu et al., 1979; Anderson et al., 1983, 1986). Table 7 summarizes the effects of treatment of male Sprague-Dawley rats with Ar-

roclor 1254 for 18 weeks (100 ppm in the diet) following initiation with diethylnitrosamine (administered for 5 weeks in the drinking water). The results demonstrated that Aroclor 1254 treatment promoted diethylnitrosamine-initiated hepatocellular carcinomas in male rats. Moreover, the promoter activity was observed using either commer-

TABLE 7

INCIDENCE AND TYPE OF LIVER LESIONS IN MALE SPRAGUE-DAWLEY RATS TREATED WITH DIETHYLNITROSAMINE AND AROCLOR 1254 (Preston et al., 1981)

Treatment	Number of rats	Neoplastic nodules	Hepatocellular carcinoma (%)
Control	72	0	0 (9)
Diethylnitrosamine	32	6	5 (16)
Aroclor 1254 ^a	34	0	0 (9)
Diethylnitrosamine plus Aroclor 1254 ^a	33	10	21 (64)
Aroclor 1254-U ^b	34	0	0 (9)
Diethylnitrosamine plus Aroclor 1254-U ^b	33	5	27 (84)

^a Aroclor 1254 cleaned up to remove polychlorinated dibenzofurans.^b Aroclor 1254-U not cleaned up as noted above.

resulted of aflato-
M₁ and
after 21
B-treated
controls. The
cinogenic
aflatoxin
e activity
tochrome
such as
l., 1986;

lor 1254
reported
lynuclear
ry et al.,
(80). The

Aroclor
prior to
thracene
dence of
ver CD-1
625 $\mu\text{g}/$
s pheno-
of drug-

ROCLOR

2 months

ppb.
n line for

cial Aroclor 1254 or a sample that had undergone cleanup to remove polychlorinated dibenzofuran impurities. Anderson and coworkers (1986) showed that a single dose of Aroclor 1254 enhanced *N*-nitrosodimethylamine-initiated lung tumors in male Swiss mice, however, the effects on liver tumor formation were variable. It was also reported that 2 higher chlorinated biphenyls 2,2',4,4',5,5'-hexachlorobiphenyl and 2,2',3,4',5,5'-hexachlorobiphenyl were the dominant congeners persisting in the tissues of the PCB-treated mice. It should be noted that only the more highly chlorinated commercial PCBs (> 50% by weight) were reported as promoters of hepatocarcinogenesis in rodents and the promoting effects of the lower chlorinated PCBs and individual PCB congeners have not been determined.

The activity of PCBs as promoters has also been investigated utilizing short-term in vivo assays in which the end points are preneoplastic lesions which are presumed to progress into cancer cells. Most of these studies demonstrate that the higher chlorinated commercial PCB mixtures promote the formation of putative preneoplastic lesions (Anderson et al., 1983, 1986; Ito et al., 1973; Nishizumi, 1976, 1979; Preston et al., 1981; Deml and Oesterle, 1982; Oesterle and Deml, 1983, 1984; Deml et al., 1983; Pereira et al., 1982; Buchmann et al., 1986). These studies were carried out using different initiators, commercial PCBs (usually Aroclor 1254 or Clophen A50) or PCB congeners and different treatment protocols; however, the results clearly demonstrate the activity of PCBs as promoters of preneoplastic lesions in rodent liver. Buchmann and coworkers (1976) reported the effects of 3,3',4,4'-tetrachlorobiphenyl and 2,2',4,4',5,5'-hexachlorobiphenyl as promoters; although there were some problems with these studies (i.e., limited numbers of animals used and large standard deviations in some of the data) the results suggested that both congeners promoted diethylnitrosamine-initiated hepatocellular adenosine triphosphate-deficient focal lesions.

The effects of PCB mixtures and selected congeners have also been investigated (Hayes et al., 1985, 1986) using the resistant hepatocyte model developed by Farber and coworkers (Solt and Farber, 1976; Tsuda et al., 1980; Farber, 1984a,b, 1986). The PCBs used in these studies included

Aroclor 1254, a reconstituted mixture of PCBs with a defined congener composition, 2,2',5,5'-tetrachlorobiphenyl, 2,2',4,4'-tetrachlorobiphenyl and 2,2',4,4',5,5'-hexachlorobiphenyl. None of these compounds exhibited activity as initiators in this model (Hayes et al., 1985). Subsequent studies (Hayes et al., 1986) using the aforementioned compounds and 3,3',4,4'-tetrachlorobiphenyl (a prototypical MC-type inducer of cytochrome P-450-dependent monooxygenases) showed that these PCBs (50 μ moles/kg) given 10 days after a dose of the initiator, diethylnitrosamine, and 7 days before 2-acetylaminofluorene, all reduced the size of the 2-acetylaminofluorene-selected γ -glutamyltranspeptidase-positive nodules. These results show that in contrast to the previous studies, noted above, PCBs also exhibit 'anti-promoting' activities in this model which utilizes 2-acetylaminofluorene as a mitoinhibitory toxin.

Poland and coworkers (1982) investigated the activities of 2,3,7,8-TCDD, Aroclor 1254 and other halogenated aryl hydrocarbons as promoters of DMBA and *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine-initiated papilloma formation in HRS/J hairless mice. The structure-dependent effects of these compounds as promoters correlated with their affinities for the aryl hydrocarbon (Ah) receptor and thus supported the role of the Ah receptor in mediating this process. With the exception of this study, the structure-dependent activities of PCBs as promoters of rodent hepatocarcinogenesis have not been investigated. However, like Aroclor 1254 and Aroclor 1260, the polybrominated biphenyl (PBB) mixture, Fire-Master BP-6, exhibits comparable biologic and toxic effects (Safe, 1984; Parkinson et al., 1983) and is also a rodent hepatocarcinogen (Kimbrough et al., 1981; Gupta et al., 1983). Moreover, the structure-function relationships for PBB and PCB isomers and congeners are similar (Safe, 1984). Structure-activity studies utilizing 2,2',4,4',5,5'- and 3,3',4,4',5,5'-hexabromobiphenyl (HBB) as promoters of *N*-nitrosodiethylamine-initiated enhancement of γ -glutamyltranspeptidase-positive altered hepatic foci showed that 2,2',4,4',5,5'-HBB but not 3,3',4,4',5,5'-HBB enhanced the development of altered hepatic foci and hepatic nodules (Jensen and Sleight, 1986). A combination of the two congeners resulted in a synergistic tumor pro-

motor of the
gation
toxic
affinit
chloro
to the
al., 19
1,2,3,4
and N
neopla
parent
the dit
be furt
Ker
the ar
inocul.
Depen
PCBs
study,
crease
increas
regress
establi
Arocl
been r

(4) Ca

An
indiv
in the
ppm)
enviro
genic
ground
addit
exper
PCBs,
contan
incide
and oc
suffic
expos
this da
human
studies
involve
relative

motion response. The significance and mechanism of these interactive effects require further investigation. The negative results obtained with the toxic halogenated biphenyls which exhibit high affinity for the Ah receptor (i.e., 3,3',4,4'-tetrachlorobiphenyl and 3,3',4,4',5,5'-HBB) contrasted to the reported activity of 2,3,7,8-TCDD (Pitot et al., 1980), 2,3,4,7,8-pentachlorodibenzofuran and 1,2,3,4,7,8-hexachlorodibenzofuran (Nishizumi and Masuda, 1986) as promoters of hepatic preneoplastic lesions and tumors in rodents. The apparent dichotomy in these results may be due to the different model systems used and this should be further investigated.

Kerkvliet and Kimeldorf (1977) have reported the antitumor activity of Aroclor 1254 in rats inoculated with Walker 256 carcinosarcoma cells. Depending on the timing of the treatment with PCBs and the number of tumor cells used in the study, Aroclor 1254 inhibited tumor growth, increased the latency period for tumor development, increased the host survival time and caused tumor regression (if administered after the tumors were established). Additional studies on the effects of Aroclor 1254 on transplantable tumors have not been reported.

(4) Carcinogenicity of PCBs — human studies

Analytical studies have confirmed that most individuals carry significant body burdens of PCBs in their adipose tissues (in the range of 0.1–1.0 ppm) and these residues are derived from multiple environmental pathways. The potential carcinogenic (or anticarcinogenic) effect of these background PCB levels has not been determined. In addition, there are 2 major groups which have experienced exposure to relatively high levels of PCBs, namely, individuals who consumed PCB-contaminated rice oil in the Yusho or Yu Cheng incidents (Kuratsune, 1980; Masuda et al., 1982) and occupationally exposed workers. There is insufficient data to assess the carcinogenic effects of exposure to the contaminated rice oil, moreover, this data may not be appropriate for assessing the human carcinogenic potential of PCBs. Several studies have reported that the PCB industrial fluid involved in the rice oil contamination contained relatively high levels of polychlorinated dibenzo-

furans (Chen et al., 1985; Buser et al., 1978) and these compounds are likely the major etiologic agents of Yusho/Yu Cheng poisoning (Bandiera et al., 1984; Kunita et al., 1985).

Zack and coworkers (1979) examined a small cohort of workers (89) with occupational exposure to PCBs. No liver cancer were reported among the 30 deaths which occurred in this group. Gustavsson and coworkers (1986) did not observe any excesses in either overall mortality or cancer in a group of 142 male Swedish workers exposed to PCBs during the manufacture of capacitors. Both of these studies are limited because of the small size of the sample population.

Bertazzi et al. (1981, 1987) examined a group of 1310 workers employed (≥ 6 months) in a capacitor-manufacturing plant from 1946 to 1970. The updated study of these PCB-exposed individuals (Bertazzi et al., 1987) added 790 workers (total = 2100) and their data showed a statistically significant excess of cancer in males (14 observed vs. 7.6 expected) and females (12 observed vs. 5.3 expected). Excess (significant) lymphatic hematopoietic cancer (4 observed vs. 1.1 expected) was noted for females whereas significantly increased digestive cancers (6 observed vs. 2.2 expected) were seen in males. However, it was reported that there were no apparent correlations between cancer mortalities (individual or total cancers) and the duration of exposure and latency periods from the time of first exposure. The small number of deaths observed in this study do not yet permit any firm conclusions regarding the carcinogenicity of PCBs and additional follow-up studies on this group will be required.

The most comprehensive mortality and cancer studies of PCB-exposed workers were reported by Brown and Jones (1981) and Brown (1987). For the 2588 workers employed in 2 capacitor-manufacturing plants (Brown, 1987), the overall mortality of this group (295 deaths) was lower than expected (318) and the mortality for cancer deaths (62 observed) was also lower than expected (80). Inspection of the cancer mortality data among workers exposed to PCBs showed that the only category where there was a significant increase ($P < 0.05$) in deaths was associated with liver, gall bladder and biliary tract cancers (combined). A detailed analysis of the data with respect to latency

periods, length of employment and sex of the workers was carried out. There was a higher female susceptibility to these cancers in the PCB-exposed workers. Moreover, for the individuals employed in plant 2, the length of employment was also important and this correlated with potential exposure to the more highly chlorinated PCBs which were initially used in this plant. This observation also correlated with the animal data. It was interesting to note that based on pathology reports, it is likely that only 1 of the 5 individuals with liver/gall bladder/biliary tract cancer had primary liver cancer. Most of these neoplasms were originally bile duct or gall bladder cancers and PCB-mediated cancer at these sites has not been extensively investigated or reported in rodent cancer studies. Subsequent updates of the mortality data in this cohort and others will be useful in determining the potential role of PCBs as human carcinogens.

(5) Summary and conclusions

The potential mutagenicity and carcinogenicity of commercial PCBs has been investigated in both in vivo and in vitro systems and several conclusions can be drawn from these studies.

(1) PCBs can covalently adduct DNA both in vivo and in vitro (using a source of metabolic activation); the more highly chlorinated biphenyls are poorly metabolized and these compounds tend to exhibit very low binding to DNA. Based on the structure-activity relationships for PCBs (Safe, 1984) it is unlikely that the more toxic compounds such as 3,3',4,4',5-penta- and 3,3',4,4',5,5'-hexachlorobiphenyl, would form covalent adducts with DNA.

(2) PCB mixtures and individual compounds exhibit minimal mutagenic activity in most assay systems.

(3) The more highly chlorinated PCB mixtures (i.e. > 50% Cl by weight) are hepatocarcinogens in rodents whereas data from a limited number of studies suggest that the lower chlorinated mixtures are not carcinogenic.

(4) In some model systems, the higher chlorinated PCB mixtures act as promoters of preneoplastic lesions and hepatocellular

carcinomas in rodents treated with a variety of initiators.

(5) Aroclor 1254 acts as a promoter of skin papilloma formation in HRS/J hairless mice and structure-activity and genetic studies suggest that the Ah receptor is necessary but not sufficient for the activity of halogenated aryl hydrocarbons as promoters in hairless mice.

(6) Individual PCB congeners and higher chlorinated commercial mixtures also exhibit anticarcinogenic activity in the CD-1 mouse skin cancer model.

(7) Results from occupational studies suggest that individuals exposed to PCBs may have an excess of cancer at some sites, however, the most comprehensive study (Brown, 1987) suggests that there are no significant increases in the overall cancer rate in workers exposed to PCBs. Follow-up and continuing epidemiological studies on the PCB-exposed workers are required to further clarify the potential carcinogenic effects of PCBs on humans.

In several strains of rats and mice, there is a high incidence of hepatic preneoplastic lesions and carcinomas and these lesions can be induced by diverse promoting agents (Schulte-Hermann et al., 1983; Weinstein, 1984). Since PCBs are not mutagenic and do not readily form covalent adducts with cellular DNA, it is likely that the higher chlorinated biphenyls are not genotoxic and act as promoters of carcinogenesis in rodents. A comparable mechanism has been suggested for 2,3,7,8-TCDD (Shu et al., 1987; Weinstein, 1984). For PCBs, the role of the Ah receptor in mediating their activity as promoters has not been delineated. It is conceivable that PCB congeners (e.g., 2,2',4,4',5,5'-hexachlorobiphenyl) which possess many properties similar to the promoter, phenobarbital, may also contribute to the development of cancer in rodents. Future studies should focus on the structure-dependent effects of PCBs and more accurately define the class of higher chlorinated biphenyl congeners which are carcinogenic.

Acknowledgements

The financial assistance of the Texas Agricultural Experiment Station is gratefully acknowl-

edged.
apprec

Refere

Albro. F
tive
(Aro-
using
Alford-S
Bude
auto
rinat
2014
Andersc
Effe
tume
mune
Andersc
Rigg
biph
trate.
Astrom.
2-ace
toch
113
Ballschn
rnat
raph
clopt
20-3
Ballschn
stud:
323-
Bandier.
Banr
poly
mat
507
Berry. I
Glea
abili
mou
Path
Bertazzi.
Pesat
fema
sente
cupa
Bertazzi.
Zoch
turn
Brown. I
rinat
333-
Brown.
hygie
phen

variety of
ter of skin
s mice and
uggest that
fficient for
arbons as

nd higher
hibit anti-
nose skin

ies suggest
y have an
r. the most
ggests that
the overall
Follow-up
es on the
to further
ts of PCBs

there is a
esions and
duced by
ann et al.,
not muta-
it adducts
the higher
and act as
s. A com-
ested for
ein, 1984).
in mediat-
een delin-
eners (e.g.,
h possess
er, pheno-
velopment
ould focus
PCBs and
gher chlo-
: carcino-

Agricul-
acknowl-

edged. The assistance of Mrs. Laura Young is also appreciated.

References

- Albro, P.W., J.T. Corbett and J.L. Schroeder (1981) Quantitative characterization of polychlorinated biphenyl mixtures (Aroclors 1242, 1254 and 1260) by gas chromatography using capillary columns. *J. Chromatogr.*, 205, 103-111.
- Alford-Stevens, A.L., T.A. Bellar, J.W. Eichelberger and W.L. Budde (1986) Characterization of commercial Aroclors by automated mass spectrometric determination of polychlorinated biphenyls by level of chlorination. *Anal. Chem.*, 58, 2014-2022.
- Anderson, L.M., K. Van Havere and J.M. Budinger (1983) Effects of polychlorinated biphenyls on lung and liver tumors initiated in suckling mice by *N*-nitrosodimethylamine. *J. Natl. Cancer Inst.*, 71, 157-163.
- Anderson, L.M., J.M. Ward, S.D. Fox, H.J. Isaaq and C.W. Riggs (1986) Effects of a single dose of polychlorinated biphenyl to infant mice on *N*-nitrosodimethylamine-initiated lung and liver tumors. *Int. J. Cancer*, 38, 109-116.
- Astrom, A., and J.W. DePierre (1985) Metabolism of 2-acetylaminofluorene by eight different forms of cytochrome P-450 isolated from rat liver. *Carcinogenesis*, 6, 113-120.
- Ballschmiter, K., and M. Zell (1980) Analysis of polychlorinated biphenyls (PCB) by glass capillary gas chromatography. Composition of technical Aroclor- and clophen-PCB mixtures. *Fresenius Z. Anal. Chem.*, 302, 20-31.
- Ballschmiter, K., H. Buckert and S. Bihler (1981) Baseline studies of the global pollution. *Z. Anal. Chem.*, 306, 323-339.
- Bandiera, S., K. Farrell, G. Mason, M. Kelley, M. Romkes, R. Bannister and S. Safe (1984) Comparative toxicities of the polychlorinated dibenzofuran (PCB) and biphenyl (PCB) mixtures which persist in Yusho victims. *Chemosphere*, 13, 507-512.
- Berry, D.L., DiGiovanni, M.R. Juchau, W.M. Bracken, G.L. Gleason and T.J. Slaga (1978) Lack of tumor promoting ability of certain environmental chemicals in a two-stage mouse skin tumorigenesis assay. *Res. Commun. Chem. Pathol. Pharmacol.*, 20, 101-108.
- Bertazzi, P.A., C. Zochetti, S. Guercilena, M.D. Foglia, A. Pesatori and L. Riboldi (1981) Mortality study of male and female workers exposed to PCBs. Helsinki, Finland, presented at International Symposium on Prevention of Occupational Cancer.
- Bertazzi, P.A., L. Riboldi, A. Pesatori, L. Radice and C. Zochetti (1987) Cancer mortality of capacitor manufacturing workers. *Am. J. Ind. Med.*, 11, 165-176.
- Brown, D.P. (1987) Mortality of workers exposed to polychlorinated biphenyls — an update. *Arch. Environ. Health*, 42, 333-339.
- Brown, D.P., and M. Jones (1981) Mortality and industrial hygiene study of workers exposed to polychlorinated biphenyls. *Arch. Environ. Health*, 36, 120-129.
- Brown, Jr., J.F., D.L. Bedard, M.J. Brennan, J.C. Carnahan, H. Feng and E. Wagner (1987) Polychlorinated biphenyl dechlorination in aquatic sediments. *Science*, 236, 709-712.
- Brown, M.P., M.B. Werner, R.J. Sloan and K.W. Simpson (1985) Polychlorinated biphenyls in the Hudson River. *Environ. Sci. Technol.*, 19, 656-661.
- Buckley, E.H. (1982) Accumulation of airborne polychlorinated biphenyls in foliage. *Science*, 216, 520-522.
- Buckmann, A., W. Kunz, C.R. Wolf, F. Oesch and L.W. Robertson (1986) Polychlorinated biphenyls, classified as either phenobarbital or 3-methylcholanthrene-type inducers of cytochrome P-450 are both hepatic tumor promoters in diethylnitrosamine-initiated rats. *Cancer Lett.*, 32, 243-253.
- Buser, H.R., C. Rappe and A. Gara (1978) Polychlorinated dibenzofurans (PCDFs) found in Yusho oil and in used Japanese PCB. *Chemosphere*, 7, 439-449.
- Bush, B., J. Snow, S. Connor and R. Koblintz (1985) Polychlorinated biphenyl congeners (PCBs), p,p'-DDE and hexachlorobenzene in human milk in three areas of upstate New York. *Arch. Environ. Contam. Toxicol.*, 14, 443-450.
- Bush, B., A.H. Bennett and J.T. Snow (1986) Polychlorinated biphenyl congeners p,p'-DDE and sperm function in humans. *Arch. Environ. Contam. Toxicol.*, 15, 333-341.
- Chen, P.H., C.K. Wong, C. Rappe and M. Nygren (1985) Polychlorinated biphenyls, dibenzofurans and quaterphenyls in toxic rice-bran oil and in the blood and tissues of patients with PCB poisoning (Yu-Cheng) in Taiwan. *Environ. Health Perspect.*, 59, 59-65.
- Deml, E., and D. Oesterle (1982) Sex-dependent promoting effect of polychlorinated biphenyls on enzyme-altered islands induced by diethylnitrosamine in rat liver. *Carcinogenesis*, 3, 1449-1453.
- Deml, E., D. Oesterle and F.J. Wiebel (1983) Benzo[a]pyrene initiates enzyme-altered islands in the liver of adult rats following single pretreatment and promotion with polychlorinated biphenyls. *Cancer Lett.*, 19, 301-304.
- DiGiovanni, J., A. Viaje, D.L. Berry, T.J. Slaga and M.R. Juchau (1977) Tumor-initiating ability of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) and Aroclor 1254 in the two-stage system of mouse skin carcinogenesis. *Bull. Environ. Contam. Toxicol.*, 18, 552-557.
- DiGiovanni, J., D.L. Berry, T.J. Slaga and M.R. Juchau (1979) Studies on the relationship between induction of biotransformation and tumor-initiating activity of 7,12-dimethylbenz[a]anthracene in mouse skin, in: P.W. Jones and P. Leber (Eds.), *Polynuclear Aromatic Hydrocarbons*. Ann Arbor Science Publishers, Ann Arbor, MI, pp. 553-568.
- DiGiovanni, J., T.J. Slaga, D.L. Berry and M.R. Juchau (1980) Inhibitory effects of environmental chemicals on polycyclic hydrocarbon carcinogenesis. in: T.J. Slaga (Ed.), *Carcinogenesis*, Vol. 5, *Modifiers of Chemical Carcinogenesis*. Raven, New York, pp. 145-168.
- Farber, E. (1984a) Perspectives in cancer research. The multi-step nature of cancer development. *Cancer Res.*, 44, 4217-4223.
- Faber, E. (1984b) Cellular biochemistry of the step-wise development of cancer with chemicals. G.H. Clowes Memorial Lecture, *Cancer Res.*, 44, 5463-5474.

- Farber, E. (1986) Some emerging general principles in the pathogenesis of hepatocellular carcinoma, *Cancer Surv.* 5, 695-718.
- Farber, E., and D.S.R. Sarma (1987) Biology in disease: hepatocarcinogenesis, a dynamic cellular perspective. *Lab. Invest.*, 56, 4-222.
- Forgue, S.T., and J.R. Allen (1982) Identification of an arene oxide metabolite of 2,2',5,5'-tetrachlorobiphenyl by gas chromatography-mass spectroscopy. *Chem.-Biol. Interact.*, 40, 233-245.
- Gardner, A.M., J.T. Chen, J.A.G. Roach and E.P. Ragelis (1973) Polychlorinated biphenyls: hydroxylated urinary metabolites of 2,2',5,5'-tetrachlorobiphenyl identified in rabbits, *Biochem. Biophys. Res. Commun.*, 55, 1377-1384.
- Goldstein, J.A., R. Weaver and D.W. Sundheimer (1984) Metabolism of 2-acetylaminofluorene by two methylcholanthrene-inducible forms of rat liver cytochrome P-450. *Cancer Res.*, 44, 3768-3771.
- Gupta, B.N., E.E. McConnell, J.A. Moore and J.K. Haseman (1983) Effects of a polybrominated biphenyl mixture in the rat and mouse, *Toxicol. Appl. Pharmacol.*, 68, 19-35.
- Gustavsson, P., C. Hogstedt and C. Rappe (1986) Short-term mortality and cancer incidence in capacitor manufacturing workers exposed to polychlorinated biphenyls (PCBs). *Am. J. Ind. Med.*, 10, 341-344.
- Halvorson, M.R., T.D. Phillips, S.H. Safe and L.W. Robertson (1985) Metabolism of aflatoxin B₁ by rat hepatic microsomes induced by polyhalogenated biphenyl congeners. *Appl. Environ. Microbiol.*, 49, 882-886.
- Hansen, L.G., L.G.M.Th. Tuinstra, C.A. Kan, J.T.W.A. Strick and J.H. Koeman (1983) Accumulation of chlorobiphenyls in chicken fat and liver after feeding Aroclor 1254 directly or fat from swine fed Aroclor 1254, *J. Agric. Food Chem.*, 31, 254-260.
- Hargraves, W.A., and J.R. Allen (1979) The in vitro binding of 2,2',5,5'-tetrachlorobiphenyl metabolites to rat liver microsomal proteins. *Res. Commun. Chem. Pathol. Pharmacol.*, 25, 33-52.
- Hass, J.R., L.T. Jao, N.K. Wilson and H.B. Matthews (1977) Metabolism of 4-chlorobiphenyl and 4,4'-dichlorobiphenyl in the rat: qualitative and quantitative aspects, *J. Agric. Food Chem.*, 25, 1330-1333.
- Hayes, M.A. (1987) *Carcinogenic and Mutagenic Effects of PCBs in Polychlorinated Biphenyls (PCBs), mammalian and Environmental Toxicology*, Vol. 1, Springer, Heidelberg, pp. 77-95.
- Hayes, M.A., S.H. Safe, D. Armstrong and R.G. Cameron (1985) Influence of cell proliferation on initiating activity of pure polychlorinated biphenyls and complex mixtures in resistant hepatocyte in vivo assays for carcinogenicity, *J. Natl. Cancer Inst.*, 74, 1037-1041.
- Hayes, M.A., E. Roberts, S.H. Safe, E. Farber and R.G. Cameron (1986) Influences of different polychlorinated biphenyls on cytotoxic, mitoinhibitory and nodule-selecting activities of *N*-2-fluorenylacetylacetamide in rat liver, *J. Natl. Cancer Inst.*, 76, 683-691.
- Hedde, J.A., and W.R. Bruce (1977) Comparison of tests for mutagenicity or carcinogenicity using assays for sperm abnormalities, formation of micronuclei and mutations in Salmonella. in: H.H. Hiatt et al. (Eds.), *Origins of Human Cancer*, Cold Spring Harbor Conf. Cell Prolif., Vol. 4, Cold Spring Harbor Lab., Cold Spring Harbor, NY, pp. 1549-1557.
- Hendricks, D., T.P. Putnam, D.D. Bills and R.O. Sinnhuber (1977) Inhibitory effect of a polychlorinated biphenyl (Aroclor 1254) on aflatoxin B₁ carcinogenesis in rainbow trout (*Salmo gairdneri*). *J. Natl. Cancer Inst.*, 59, 1545-1551.
- Hesse, S., and T. Wolff (1977) In vitro interactions of di-, tetra- and hexachlorobiphenyl with rabbit liver monooxygenase. *Biochem. Pharmacol.*, 26, 2043-2047.
- Hesse, S., M. Metzger and T. Wolff (1978) Activation of [¹⁴C]-chlorobiphenyls to protein-binding metabolites by rat liver microsomes. *Chem.-Biol. Interact.*, 20, 355-365.
- Hirose, M., T. Shirai, H. Tsuca, S. Fukushima, T. Ogiso and N. Ito (1981) Effect of phenobarbital, polychlorinated biphenyl and sodium saccharin on hepatic and renal carcinogenesis in unilaterally nephrectomized rats given *N*-ethyl-*N*-hydroxyethylnitrosamine orally, *Carcinogenesis*, 2, 1299-1302.
- Holdrinet, M.V., H.E. Braun, R. Frank, G.J. Stopps, M.S. Smout and J.W. McWade (1977) Organochlorine residues in human adipose tissue and milk from Ontario residents, 1969-1974. *Can. J. Publ. Health*, 68, 74-80.
- Hsia, M.T.S., F.S.D. Lin and J.R. Allen (1978) Comparative mutagenicity and toxic effects of 2,5,2',5'-tetrachlorobiphenyl and its metabolites in bacterial and mammalian test systems. *Res. Commun. Chem. Pathol. Pharmacol.*, 21, 485-496.
- Hsu, I.C., J.P. Van Miller and J.R. Allen (1975) metabolic fate of [³H]-2,2,2',5'-tetrachlorobiphenyl in infant non-human primates. *Bull. Environ. Contam. Toxicol.*, 14, 233-240.
- Hutzinger, O., S. Safe and V. Zitko (1974) *The Chemistry of PCBs*. CRC Press, Cleveland, OH.
- Ito, N., H. Nagasaki, M. Arai, S. Makiura, S. Sugihara and K. Hirao (1973) Histopathologic studies on liver tumorigenesis induced in mice by technical polychlorinated biphenyls and its promoting effect on liver tumors induced by benzene hexachloride. *J. Natl. Cancer Inst.*, 51, 1637-1646.
- Ito, N., H. Nagasaki, S. Makiura and M. Arai (1974) Histopathological studies on liver tumorigenesis in rats treated with polychlorinated biphenyls. *Gann*, 65, 545-549.
- Jensen, R.K., and S. Sleight (1986) Sequential study on the synergistic effects of 2,2',4,4',5,5'-hexabromobiphenyl and 3,3',4,4',5,5'-hexabromobiphenyl on hepatic tumor promotion. *Carcinogenesis*, 7, 1771-1774.
- Jensen, S., and G. Sundstrom (1974) Structures and levels of most chlorobiphenyls in the technical PCB products and in human adipose tissue. *Ambio*, 3, 70-76.
- Kaminsky, L.S., M.W. Kennedy, S.M. Adams and F.P. Guengerich (1981) Metabolism of dichlorobiphenyls by highly purified isozymes of rat liver cytochrome P-450. *Biochemistry*, 20, 7379-7384.
- Kannan, N., S. Tanabe and R. Tastsukawa (1988) Potentially hazardous residues of non-ortho chlorine substituted coplanar PCBs in human adipose tissue. *Arch. Environ. Health*, 43, 11-14.
- Kenne
and
ph
Pha
Kenne
Kur
hep
col
Kerke
of
mo
Car
Kimbr
ade
by
Ins
Kimbr
R.J
tur
bip
145
Kimbr
Cl
ra
535
Kimur
rat
105
Kimur
nit
car
Kunita
PC
an
Kurats
H
ber
p.2
Leece
(19
vis
ship
Makiur
(19
lve
ylat
nit
Masud
Kur
age
osp
Matthe
PCI
McMa
E F
pro
cye
171
Melan

- Kennedy, M.W., N.K. Carpentier, P.O. Dymerski, S.M. Adams and L.S. Kaminsky (1980) Metabolism of monochlorobiphenyls by hepatic microsomal cytochrome P-450. *Biochem. Pharmacol.*, 29, 727-736.
- Kennedy, M.W., N.K. Carpentier, P.O. Dymerski and L.S. Kaminsky (1981) Metabolism of dichlorobiphenyls by hepatic microsomal cytochrome P-450. *Biochem. Pharmacol.*, 30, 577-588.
- Kerkvliet, N.I., and D.J. Kimeldorf (1977) Antitumor activity of a polychlorinated biphenyl mixture, Aroclor 1254 in rats inoculated with Walker 256 carcinosarcoma cells. *J. Natl. Cancer Inst.*, 59, 951-955.
- Kimbrough, R.D., and R.E. Linder (1974) Induction of adenofibrosis and hepatomas in the liver of Balb/c mice by polychlorinated biphenyls (Aroclor 1254). *J. Natl. Cancer Inst.*, 53, 547-552.
- Kimbrough, R.D., R.A. Squire, R.E. Linder, J.D. Strandberg, R.J. Montali and V.W. Burse (1975) Induction of liver tumors in Sherman strain female rats by polychlorinated biphenyls (Aroclor 1260). *J. Natl. Cancer Inst.*, 55, 1453-1459.
- Kimbrough, R.D., D.F. Groce, M.P. Korver and V.W. Burse (1981) Induction of liver tumors in female Sherman strain rats by polybrominated biphenyls. *J. Natl. Cancer Inst.*, 66, 535-542.
- Kimura, N.T., and T. Baba (1973) Neoplastic changes in the rat liver induced by polychlorinated biphenyls. *Gann*, 64, 105-108.
- Kimura, N.T., T. Kanematsu and T. Baba (1976) Polychlorinated biphenyl(s) as a promoter in experimental hepatocarcinogenesis in rats. *Z. Krebsforsch.*, 87, 257-266.
- Kunita, N., S. Hori, H. Obana et al. (1985) Biological effects of PCBs, PCQs and PCDFs present in the oil causing Yusho and Yu-Cheng. *Environ. Health Perspect.*, 59, 79.
- Kuratsune, M. (1980) Yusho. In: R.D. Kimbrough (Ed.), *Halogenated Biphenyls, Terphenyls, Naphthalenes, Dibenzodioxins and Related Products*. Elsevier, Amsterdam, p. 287.
- Leece, B., M.A. Denomme, R. Towner, S.M.A. Li and S. Safe (1985) Polychlorinated biphenyls: Correlation between in vivo and in vitro quantitative structure-activity relationships (QSARs). *J. Toxicol. Environ. Health*, 16, 379-388.
- Makiura, S., H. Aoe, S. Sugihara, K. Hiara, M. Arai and N. Ito (1974) Inhibitory effect of polychlorinated biphenyls on liver tumorigenesis in rats treated with 3'-methyl-4-dimethylaminobenzene, N-2-fluorenylacetamide and diethylnitrosamine. *J. Natl. Cancer Inst.*, 53, 1253-1257.
- Masuda, Y., H. Kuroki, T. Yamaryo, K. Haraguchi, M. Kuratsune and S.T. Hsu (1982) Comparison of causal agents in Taiwan and Fukuoka PCB poisonings. *Chemosphere*, 11, 199-206.
- Matthews, H.B., and R.L. Dedrick (1984) Pharmacokinetics of PCBs. *Annu. Rev. Pharmacol. Toxicol.*, 24, 85-103.
- McManus, M.E., R.F. Minchin, N. Sanderson, D. Schwartz, E.F. Johnson and S.S. Thorgeirsson (1984) Metabolic processing of 2-acetylaminofluorene and six highly purified cytochrome P-450 forms of rabbit liver. *Carcinogenesis*, 5, 1717-1723.
- Melancon, M.J., and J.J. Lech (1976) Isolation and identification of a polar metabolite of tetrachlorobiphenyl from bile of rainbow trout exposed to ^{14}C -tetrachlorobiphenyl. *Bull. Environ. Contam. Toxicol.*, 15, 181-188.
- Morales, N.M., and H.B. Matthews (1979) In vivo binding of 2,3,6,2',3',6'-hexachlorobiphenyl and 2,4,5,2',4',5'-hexachlorobiphenyl to mouse liver macromolecules. *Chem.-Biol. Interact.*, 27, 94-110.
- Mullin, M.D., C.M. Pochini, S. McCrindle, M. Romkes, S.H. Safe and L.M. Safe (1984) High-resolution PCB analysis. The synthesis and chromatographic properties of all 209 PCB congeners. *Environ. Sci. Technol.*, 18, 468-276.
- Nagasaki, H., S. Tomii, T. Mega, M. Marugami and N. Ito (1972) Hepatocarcinogenicity of polychlorinated biphenyls in mice. *Gann*, 63, 805.
- NCI (National Cancer Institute) (1978) Bioassay of Aroclor (trademark) 1254 for possible carcinogenicity. CAS No. 27343-18-8, NCI Carcinogenesis Tech. Rep. Ser. No. 38.
- Nelson, N.N., P.B. Hammon, I.C.T. Nisbet, A.F. Sarofim and W.H. Drury (1972) Polychlorinated biphenyls: environmental impact. *Environ. Res.*, 5, 249-362.
- Nilsson, B., and C. Ramel (1974) Genetic tests on *Drosophila melanogaster* with polychlorinated biphenyls (PCB). *Hereditas*, 77, 319-322.
- Nishizumi, M. (1976) Enhancement of diethylnitrosamine hepatocarcinogenesis in rats by exposure to polychlorinated biphenyls or phenobarbital. *Cancer Lett.*, 2, 11-16.
- Nishizumi, M. (1979) Effects of phenobarbital, dichlorodiphenyltrichlorethane, and polychlorinated biphenyls on diethylnitrosamine-induced hepatocarcinogenesis. *Gann*, 70, 835-837.
- Nishizumi, M., and Y. Masuda (1986) Enhancing effect of 2,3,4,7,8-pentachlorodibenzofuran and 1,2,3,4,7,8-hexachlorodibenzofuran on diethylnitrosamine hepatocarcinogenesis in the rat. *Cancer Lett.*, 33, 333-339.
- Norback, D.J., and R.H. Weltman (1985) Polychlorinated biphenyl induction of hepatocellular carcinoma in the Sprague-Dawley rat. *Environ. Health Perspect.*, 60, 97-105.
- Norback, D.H., J.L. Seymour, K.M. Knieriem, R.E. Peterson and J.R. Allen (1976) Biliary metabolites of 2,5,2',5'-tetrachlorobiphenyl in the rat. *Res. Commun. Chem. Pathol. Pharmacol.*, 14, 527-533.
- Oesterle, D., and E. Deml (1983) Promoting effects of polychlorinated biphenyls on development of enzyme-altered islands in livers of weaning and adult rats. *J. Cancer Res. Clin. Oncol.*, 105, 141-147.
- Oesterle, D., and E. Deml (1984) Dose-dependent promoting effects of polychlorinated biphenyls on enzyme-altered islands in livers of adult and weaning rats. *Carcinogenesis* 5, 351-355.
- Parkinson, A., P.E. Thomas, D.E. Ryan, L.M. Reik, S.H. Safe, L.W. Robertson and W. Levin (1983) Immunochemical quantitation of cytochrome P-450 isozymes and epoxide hydrolase in liver microsomes from polychlorinated and polybrominated biphenyls: A study of structure activity relationships. *J. Biol. Chem.*, 258, 5967-5976.
- Peakall, D.B., J.L. Lincer and S.E. Bloom (1972) Embryonic mortality and chromosomal alterations caused by Aroclor 1254 in ring doves. *Environ. Health Perspect.*, 1, 103-104.
- Pereira, M.A., S.L. Herren, A.L. Britt and M.M. Khoury

- (1982) Promotion by polychlorinated biphenyls of enzyme-altered foci in rat liver. *Cancer Lett.*, 15, 185-190.
- Pitot, H.C., T. Goldsworthy and H. Poland (1980) Promotion by 2,3,7,8-tetrachlorodibenzo-*p*-dioxin of hepatocarcinogenesis from diethylnitrosamine. *Cancer Res.*, 40, 3616-3620.
- Pitot, H.C., D.G. Beer and S. Hendrich (1987) Multi-stage carcinogenesis of the rat hepatocyte. in: *Banbury Report 25: Non Genotoxic Mechanism in Carcinogenesis*. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, pp. 41-53.
- Poland, A., D. Palen and E. Glover (1982) Tumor promotion by TCDD in skin of HRS/1 hairless mice. *Nature (London)*, 300, 271-273.
- Pomerantz, I., J. Burke, D. Firestone, J. McKinney, J. Roach and W. Trotter (1978) Chemistry of PCBs and PBBs. *Environ. Health Perspect.*, 24, 133-146.
- Preston, B.D., J.P. Van Miller, R.W. Moore and J.R. Allen (1981) Promoting effects of polychlorinated biphenyls (Aroclor 1254) and polychlorinated dibenzofuran-free Aroclor 1254 on diethylnitrosamine-induced tumorigenesis in the rat. *J. Natl. Cancer Inst.*, 66, 509-515.
- Preston, B.D., J.A. Miller and E.C. Miller (1984) Reactions of 2,2',5,5'-tetrachlorobiphenyl 3,4-oxide with methionine, cysteine and glutathione in relation to the formation of methylthio-metabolites of 2,2',5,5'-tetrachlorobiphenyl in the rat and mouse. *Chem.-Biol. Interact.*, 50, 289-312.
- Reich, H.J., I.L. Reich and S. Wollowitz (1978) Conversion of allyl alcohols to dienes by sulfoxide and selenoxide syn elimination: synthesis of PCB arene oxides. *J. Am. Chem. Soc.*, 100, 5981-5982.
- Risebrough, R.W., P. Rieche, S.G. Herman, D.B. Herman, D.C. Peakall and M.N. Kirven (1968) Polychlorinated biphenyls in the global ecosystem. *Nature (London)*, 220, 1098-1102.
- Ryan, D.E., P.E. Thomas, D. Korzeniowski and W. Levin (1979a) Separation and characterization of highly purified forms of liver microsomal cytochrome P-450 from rats treated with polychlorinated biphenyls, phenobarbital and 3-methylcholanthrene. *J. Biol. Chem.*, 254, 1365-1374.
- Ryan, D.E., P.E. Thomas, L.M. Reik and W. Levin (1979b) Purification, characterization and regulation of five rat hepatic cytochrome P-450 isozymes. *Xenobiotica*, 12, 727-744.
- Safe, S. (1980) Metabolism, uptake, storage and bioaccumulation of halogenated aromatic pollutants. in: R.D. Kimbrough (Ed.), *Halogenated Biphenyls, Terphenyls, Naphthalenes, Dibenzodioxins and Related Products*. Elsevier, Amsterdam, pp. 81-107.
- Safe, S. (1982) Halogenated hydrocarbons and aryl hydrocarbons identified in human tissues. *Toxicol. Environ. Chem.*, 5, 153-165 (1982).
- Safe, S. (1984) Polychlorinated biphenyls (PCBs) and polybrominated biphenyls (PBBs): biochemistry, toxicology and mechanism of action. *CRC Crit. Rev. Toxicol.*, 13, 319-395.
- Safe, S., O. Hutzinger and D. Jones (1975a) The mechanism of chlorobiphenyl metabolism. *J. Agric. Food Chem.*, 23, 851-853.
- Safe, S., N. Platonow, O. Hutzinger and W.D. Jamieson (1975b) Analysis of organochlorine metabolites in crude extracts by high resolution photoplate mass spectrometry. *Biomed. Mass Spectrom.*, 2, 201-203.
- Safe, S., D. Jones and O. Hutzinger (1976) Metabolism of 4,4'-dihalobiphenyls. *J. Chem. Soc. Perkin Trans.*, 1, 357-533.
- Safe, S., L. Safe and M. Mullin (1985a) Polychlorinated biphenyls (PBCs) — Congener-specific analysis of a commercial mixture and a human milk extract. *J. Agric. Food Chem.*, 33, 24-29.
- Safe, S., S. Bandiera, T. Sawyer, L. Robertson, A. Parkinson, P.E. Thomas, L.M. Reik, W. Levin, M.A. Denomme and T. Fujita (1985b) PCBs: Structure-function relationships and mechanism of action. *Environ. Health Perspect.*, 60, 47-56.
- Schaffer, E., H. Greim and W. Goessner (1984) Pathology of chronic polychlorinated biphenyl (PCB) feeding in rats. *Toxicol. Appl. Pharmacol.*, 75, 278-288.
- Scheutz, E.G., S.A. Wrighton, S.H. Safe and P.S. Guzelian (1986) Regulation of cytochrome P-450p by phenobarbital and phenobarbital-like inducers in adult rat hepatocytes in primary monolayer and in vivo. *Biochemistry*, 25, 1124-1133.
- Schnellmann, R.G., R.F. Volp, C.W. Putnam and I.G. Sipes (1984) The hydroxylation, dechlorination and glucuronidation of 4,4'-dichlorobiphenyl by human hepatic microsomes. *Biochem. Pharmacol.*, 33, 3503-3509.
- Schnellmann, R.G., E.M. Vickers and I.G. Sipes (1985) Metabolism and disposition of polychlorinated biphenyls. in: E. Hodgson, J.R. Bend and R.M. Philpot (Eds.), *Reviews in Biochemical Toxicology*, Vol. 7, Elsevier, Amsterdam, pp. 274-282.
- Schoeny, R. (1982) Mutagenicity testing of chlorinated biphenyl and chlorinated dibenzofurans. *Mutation Res.*, 101, 45-56.
- Schoeny, R.S., C.C. Smith and J.C. Loper (1979) Non-mutagenicity for Salmonella of the chlorinated hydrocarbons Aroclor 1254, 1,2,4-trichlorobenzene, nutrex and kepone. *Mutation Res.*, 68, 125-132.
- Schulte-Hermann, R. (1985) Tumor promotion in the liver. *Arch. Toxicol.*, 57, 147-158.
- Schulte-Hermann, R., I. Timmermann-Trosiener and J. Schuppler (1983) Promotion of spontaneous preneoplastic cells in rat liver as a possible explanation of tumor production of nonmutagenic compounds. *Cancer Res.*, 43, 839-844.
- Shelton, D.W., R.A. Coulombe, C.B. Pereira, J.L. Casteel and J.D. Hendricks (1983) Inhibitory effect of Aroclor 1254 on aflatoxin-initiated carcinogenesis in rainbow trout and mutagenesis using a Salmonella-trout hepatic activation system. *Aquat. Toxicol.*, 3, 229-238.
- Shelton, D.W., J.D. Hendricks, R.A. Coulombe and G.S. Bailey (1984) Effects of dose on the inhibition of carcinogenesis/mutagenesis by Aroclor 1254 in rainbow trout fed aflatoxin B₁. *J. Toxicol. Environ. Health*, 13, 659-667.
- Shelton, D.W., D.E. Goeger, J.D. Hendricks and G.S. Bailey (1986) Mechanisms of anticarcinogenesis: the distribution and metabolism of aflatoxin B₁ in rainbow trout fed Aroclor 1254. *Carcinogenesis*, 7, 1065-1071.

Shimada,
poly
Biochem.
Shu, H.I.
evalu
tum
2,3,7
Phar
Slaga, J
gene
Tum
gene
Solt, D
of ch
Sundstr
lism
267-
Tanabe,
haz
exp
Hea
Tatema
Hir
no
Inst
Thoma
Reg
ind
Thoma
Ind
chr
stru
454
Tsuda
no
V
Ward,
and
E

on (1975b) extracts by Biomed.

abolism of Trans., 1.

nated bi-a commer-gnc. Food

Parkinson, ime and T. nshups and 60, 47-56. thology of ng in rats.

Guzelian nobarbitol atocytes in istry., 25.

I.G. Sipes ucuronida-tic micro-

es (1985) biphenyls. Eds.), Re-r. Amster-

nated bi-Res., 101.

non-muta-trocabons d kepone.

the liver.

I.J. Schup-tic cells in duction of -844. asteel and r 1254 on trout and activation

G.S. Bai-rcinogene-trout fed -667.

i.S. Bailey istribution t fed Aro-

Shimada, T., and R. Sato (1978) Covalent binding in vitro of polychlorinated biphenyls to microsomal macromolecules. *Biochem. Pharmacol.*, 27, 585-593.

Shu, H.P., D.J. Paustenbach and F.J. Murray (1987) A critical evaluation of the use of mutagenesis, carcinogenesis and tumor promotion data in a cancer risk assessment of 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Regulat. Toxicol. Pharmacol.* 7, 57-88.

Slaga, T.J. (1984) Mechanisms involved in two-stage carcinogenesis in mouse skin; in: T.J. Slaga (Ed.), *Mechanisms of Tumor Promotion: Tumor Promotion and Skin Carcinogenesis*. Vol. 1, CRC Press, Boca Raton, FL, p. 1

Solt, D.B., and E. Farber (1976) New principle for the analysis of chemical carcinogenesis. *Nature (London)*, 263, 702-703.

Sundstrom, G., O. Hutzinger and S. Safe (1976) The metabolism of chlorobiphenyls — A review. *Chemosphere*, 5, 267-298.

Tanabe, N., S. Tanabe and R. Tastsukawa (1988) Potentially hazardous residues of non-ortho chlorine substituted coplanar PCBs in human adipose tissue. *Arch. Environ. Health*, 43, 11-14

Tatematsu, K., K. Nakanishi, G. Murasaki, Y. Miyata, M. Hirose and N. Ito (1979) Enhancing effect of inducers of liver microsomal enzymes on induction of hyperplastic liver nodules by N-2-fluorenylacetylamide in rats. *J. Natl. Cancer Inst.*, 63, 1411-1416

Thomas, P.E., L.M. Reik, D.E. Ryan and W. Levin (1981) Regulation of three forms of cytochrome P-450 and epoxide hydrolase in rat liver microsomes: effects of age, sex and induction. *J. Biol. Chem.*, 256, 1044-1052.

Thomas, P.E., L.M. Reik, D.E. Ryan and W. Levin (1983) Induction of two immunohistochemically related rat liver cytochrome P-450 isozymes cytochrome P-450c and P-450d, by structurally diverse xenobiotics. *J. Biol. Chem.*, 258, 4590-4598

Tsuda, H., G. Lee and E. Farber (1980) Induction of resistant hepatocytes as a new principle for a possible short-term in vivo test for carcinogens. *Cancer Res.*, 40, 1157-1164.

Ward, J.M. (1985) Proliferative lesions of the glandular stomach and liver in F344 rats fed diets containing Aroclor 1254. *Environ. Health Perspect.*, 60, 89-95.

Wasserman, M., D. Wasserman, S. Cucos and H.J. Miller (1979) World PCBs map: storage and effects in man and his biologic environment in the 1970s. *Ann. N.Y. Acad. Sci.*, 320, 69-124.

Webb, R.G., and A.C. McCall (1973) Quantitative PCB standards for electron capture gas chromatography. *J. Chromatogr. Sci.*, 11, 366-373.

Weinstein, I.B. (1984) Dioxins as carcinogenic promoters. in: W.W. Lowrance (Ed.), *Public Health Risks of the Dioxins*. Kaufmann, Los Altos, CA, pp. 155-159.

Wolff, M.S., J. Thornton, A. Fischbein, R. Lillis and I.J. Selikoff (1982) Disposition of polychlorinated biphenyl congeners in occupationally exposed persons. *Toxicol. Appl. Pharmacol.*, 62, 294-308

Wolff, M.S., A. Fischbein, K.D. Rosenman and S.M. Levin (1986) Pattern designation of PCBs in human samples. *Chemosphere*, 15, 301-307.

Wong, A., P.K. Basur and S. Safe (1979) The metabolically mediated DNA damage and subsequent repair by 4-chlorobiphenyl in Chinese hamster ovary cells. *Res. Commun. Chem. Pathol. Pharmacol.*, 24, 543-570

Wyndham, C., and S. Safe (1978) In vitro metabolism of 4-chlorobiphenyl by control and induced rat liver microsomes. *Biochemistry*, 17, 208-215.

Wyndham, C., J. Devenish and S. Safe (1976) The in vitro metabolism, macromolecular binding and bacterial mutagenicity of 4-chlorobiphenyl, a model PCB substrate. *Res. Commun. Chem. Pathol. Pharmacol.*, 12, 563-570.

Yang, S.K. (1988) Stereoselectivity of cytochrome P-450 isozymes and epoxide hydrolase in the metabolism of polycyclic aromatic hydrocarbons. *Biochem. Pharmacol.*, 37, 61-70.

Yoshizawa, H., R. Uchimaru, T. Kamataki, R. Kato and Y. Ueno (1982) Metabolism and activation of aflatoxin B₁ by reconstituted P-450 system at rat liver. *Cancer Res.* 42, 1120-1124.

Zack, T.A., and D.C. Musch (1979) Mortality of PCB workers at the Monsanto Plant in Sanget, Illinois. St. Louis, MO Monsanto International Report