Lower Duwamish Waterway Group

Port of Seattle / City of Seattle / King County / The Boeing Company

Lower Duwamish Waterway Remedial Investigation

PHASE 2 REMEDIAL INVESTIGATION REPORT

APPENDIX A:

BASELINE ECOLOGICAL RISK ASSESSMENT FINAL

For submittal to:

The US Environmental Protection Agency
Region 10
Seattle, WA

The Washington State Department of Ecology
Northwest Regional Office
Bellevue, WA

July 31, 2007

Prepared by: Windward environmental LLC

200 West Mercer Street, Suite 401 ♦ Seattle, Washington ♦ 98119

Table of Contents

List of Tables iv

List of Figures x

List of Maps x

Acronyms xi

Executive Summary ES-1
- ES.1 PROBLEM FORMULATION ES-1
- ES.2 EXPOSURE ASSESSMENT ES-3
- ES.3 EFFECTS ASSESSMENT ES-3
- ES.4 RISK CHARACTERIZATION AND UNCERTAINTY ANALYSIS ES-4

A.1.0 Introduction 1

A.2.0 Problem Formulation 3
- A.2.1 ENVIRONMENTAL SETTING 3
  - A.2.1.1 Site description 3
  - A.2.1.2 Habitat features 4
- A.2.2 RESOURCES POTENTIALLY AT RISK 7
  - A.2.2.1 State and federal threatened, endangered, and sensitive species in the LDW 7
  - A.2.2.2 Benthic invertebrate community 9
  - A.2.2.3 Fish 18
  - A.2.2.4 Wildlife 27
- A.2.3 RECEPTOR OF CONCERN SELECTION 35
  - A.2.3.1 Benthic invertebrate community 36
  - A.2.3.2 Fish 37
  - A.2.3.3 Wildlife 41
  - A.2.3.4 Summary of ROC selection 44
- A.2.4 DATA SELECTION, REDUCTION, AND SUITABILITY 47
  - A.2.4.1 Data selection and reduction 47
  - A.2.4.2 Suitability of data for risk assessment 54
- A.2.5 SELECTION OF CHEMICALS OF POTENTIAL CONCERN 58
  - A.2.5.1 Benthic invertebrates 59
  - A.2.5.2 Fish 75
  - A.2.5.3 Wildlife 89
- A.2.6 CONCEPTUAL SITE MODEL 104
  - A.2.6.1 Exposure pathways 105
  - A.2.6.2 Food web diagram 107
  - A.2.6.3 Assessment endpoints and measures of effect and exposure 109

A.3.0 Exposure and Effects Assessment: Benthic Invertebrates 114
A.3.1 BENTHIC INVERTEBRATE COMMUNITY EXPOSURE ASSESSMENT 114
A.3.1.1 Sediment exposure assessment 114
A.3.1.2 Porewater exposure assessment 117
A.3.1.3 TBT exposure assessment 118
A.3.2 BENTHIC INVERTEBRATE COMMUNITY EFFECTS ASSESSMENT 121
A.3.2.1 Sediment effects assessment 121
A.3.2.2 Site-specific toxicity tests assessment 125
A.3.2.3 Porewater effects assessment 133
A.3.2.4 TBT effects assessment 133
A.3.3 CRAB EXPOSURE ASSESSMENT 137
A.3.4 CRAB EFFECTS ASSESSMENT 139
A.3.5 SUMMARY OF BENTHIC INVERTEBRATE COMMUNITY ASSESSMENT 141
A.3.5.1 Exposure assessment 141
A.3.5.2 Effects assessment 141
A.3.6 SUMMARY OF CRAB ASSESSMENT 142
A.3.6.1 Exposure assessment 142
A.3.6.2 Effects assessment 142

A.4.0 Exposure and Effects Assessment: Fish 143
A.4.1 EXPOSURE ASSESSMENT 144
A.4.1.1 Critical tissue-residue exposure assessment 144
A.4.1.2 Dietary exposure 145
A.4.2 EFFECTS ASSESSMENT 152
A.4.2.1 COPCs evaluated using the critical tissue-residue approach 152
A.4.2.2 COPCs evaluated using a dietary approach 160
A.4.3 SUMMARY OF FISH ASSESSMENT 167
A.4.3.1 Exposure assessment 167
A.4.3.2 Effects assessment 167

A.5.0 Exposure and Effects Assessment: Wildlife 169
A.5.1 EXPOSURE ASSESSMENT 170
A.5.1.1 Approach 170
A.5.1.2 Exposure assumptions 172
A.5.1.3 Prey tissue, sediment, and water data 184
A.5.1.4 Estimated dietary doses 192
A.5.2 EFFECTS ASSESSMENT 194
A.5.2.1 TRVs for birds 195
A.5.2.2 TRVs for mammals 217
A.5.3 SUMMARY OF WILDLIFE ASSESSMENT 234
A.5.3.1 Exposure assessment 234
A.5.3.2 Effects assessment 234

A.6.0 Risk Characterization and Uncertainty Analysis 235

A.6.1 BENTHIC INVERTEBRATES 235
A.6.1.1 Benthic invertebrate community 236
A.6.1.2 Crabs 264
A.6.1.3 Summary of risk conclusions for benthic invertebrates 271
A.6.2 RISK CHARACTERIZATION FOR FISH 272
A.6.2.1 Juvenile chinook salmon 272
A.6.2.2 English sole 293
A.6.2.3 Pacific staghorn sculpin 311
A.6.2.4 Summary of risk conclusions for fish 324
A.6.3 RISK CHARACTERIZATION FOR WILDLIFE 325
A.6.3.1 Spotted sandpiper 325
A.6.3.2 Great blue heron 346
A.6.3.3 Osprey 350
A.6.3.4 River otter 358
A.6.3.5 Harbor seal 364
A.6.3.6 Summary of risk conclusions for wildlife 367

A.7.0 Selection of Primary Ecological Risk Drivers/Indicator Hazardous Substances 368
A.7.1 CONSIDERATIONS FOR IDENTIFICATION OF RISK DRIVERS 369
A.7.2 IDENTIFICATION OF RISK DRIVER CHEMICALS 369

A.8.0 Conclusions 377

A.9.0 References 383

Attachment 1 Benthic Invertebrate Species Identified in the Lower Duwamish Waterway

Attachment 2 Fish Species Identified in the Lower Duwamish Waterway

Attachment 3 Data Management Rules

Attachment 4 Results of Chemicals of Interest Screen

Attachment 5 All Acceptable Decapod Studies for Chemicals of Interest for Crab

Attachment 6 Sediment Toxicity Reference Values for Chlordane and DDTs

Attachment 7 Summary of Toxicity Data for Volatile Organic Compounds in Porewater

Attachment 8 Acceptable Toxicity Studies for the Selection of Fish Toxicity Reference Values

Attachment 9 Acceptable Toxicity Studies for the Selection of Bird Dietary Toxicity Reference Values

Attachment 10 Acceptable Toxicity Studies for the Selection of Mammal Dietary Toxicity Reference Values

Attachment 11 Exposure Concentration Calculation Methods

Attachment 12 Wildlife Exposure Dose Calculations

List of Tables

Table ES-1.	*COPCs and ROCs with HQs ≥ 1.0*	*7*
Table ES-2.	*Benthic invertebrate community COPCs with detected concentrations in LDW surface sediments greater than SMS criteria, DMMP guidelines, or TRVs*	*8*
Table A.1-1.	*Document organization for primary ERA components*	*2*
Table A.2-1.	*Habitat types represented in the LDW*	*5*
Table A.2-2.	*Threatened and candidate species listed under ESA or by Washington State Department of Fish and Wildlife*	*8*
Table A.2-3.	*Phase 2 intertidal benthic invertebrate community survey results*	*11*
Table A.2-4.	*2003 intertidal clam survey results*	*12*
Table A.2-5.	*Numbers of clams collected for chemical analysis in 2004*	*13*
Table A.2-6.	*Phase 2 subtidal benthic invertebrate community survey results*	*15*
Table A.2-7.	*Numbers of individual invertebrate species caught using trawls and traps in the LDW during the 2004 and 2005 surveys*	*17*
Table A.2-8.	*Summary of studies assessing the fish community in the LDW*	*19*
Table A.2-9.	*Bird species using the LDW*	*27*
Table A.2-10.	*ROCs selected for the LDW and a summary of the considerations for selection*	*45*
Table A.2-11.	*Summary of studies included in the baseline surface sediment dataset*	*47*
Table A.2-12.	*LDW tissue data used in this baseline ERA*	*51*
Table A.2-13.	*Summary of COPCs retained for benthic invertebrates based on surface sediment chemistry data*	*63*
Table A.2-14.	*Identification of COPCs for benthic invertebrates in porewater*	*67*
Table A.2-15.	*Chemicals identified as COIs for crabs*	*68*
Table A.2-16.	*Results of TRV search for chemicals identified as crab COIs through the initial screening process and detected in crab tissue*	*71*
Table A.2-17.	*Selected critical tissue-residue TRVs for crab COIs*	*72*
Table A.2-18.	*Maximum concentrations of COIs in crab tissue samples compared with NOAEL TRVs*	*73*
Table A.2-19.	*COPCs evaluated in the risk characterization for the benthic invertebrate community and for crabs*	*74*
Table A.2-20.	*Chemicals identified as COIs for fish ROCs*	*76*
Table A.2-21.	*Results of TRV search for fish COIs*	*78*
Table A.2-22.	*TRVs selected for fish COIs evaluated using a dietary approach*	*80*
Table A.2-23.	*TRVs selected for fish COIs evaluated using the critical tissue-residue approach*	*81*

Table A.2-24.	*Maximum COI concentrations in any fish ROC tissue compared to NOAEL TRVs*	*84*
Table A.2-25.	*Maximum tissue concentrations in each fish ROC compared to NOAEL TRVs for COIs analyzed using a critical tissue-residue approach*	*85*
Table A.2-26.	*Maximum dietary exposure concentrations compared to NOAEL TRVs for fish COIs analyzed using a dietary approach*	*87*
Table A.2-27.	*Maximum dietary PAH exposure concentrations (including all alkylated and non-alkylated PAHs) compared to total PAH and benzo(a)pyrene NOAEL TRVs for fish*	*88*
Table A.2-28.	*COPCs selected for fish ROCs*	*89*
Table A.2-29.	*Chemicals identified as COIs for wildlife ROCs*	*90*
Table A.2-30.	*Results of TRV search for COIs for birds*	*93*
Table A.2-31.	*Results of TRV search for COIs for mammals*	*94*
Table A.2-32.	*TRVs selected for bird COIs*	*95*
Table A.2-33.	*TRVs selected for mammal COIs*	*97*
Table A.2-34.	*Results of COPC screen for spotted sandpiper*	*100*
Table A.2-35.	*Results of COPC screen for great blue heron and osprey*	*101*
Table A.2-36.	*Results of COPC screen for river otter and harbor seal*	*102*
Table A.2-37.	*COPCs evaluated in the risk characterization for birds and mammals*	*104*
Table A.2-38.	*Assessment endpoints for ROCs and measures of effect and exposure*	*110*
Table A.3-1.	*Chemical concentrations and detection frequencies in LDW surface sediments for COPCs identified for the benthic invertebrate community*	*115*
Table A.3-2.	*Concentrations of cis-1,2-dichloroethene in porewater samples from GWI*	*117*
Table A.3-3.	*Concentrations of cis-1,2-dichloroethene in porewater samples from Boeing Plant 2/Jorgensen Forge*	*118*
Table A.3-4.	*Tributyltin concentrations in benthic invertebrate tissue samples and co-located sediment samples*	*119*
Table A.3-5.	*Biological effect endpoints used to determine the SQS and CSL for COPCs*	*122*
Table A.3-6.	*Available toxicity studies for selection of chlordane and total DDTs TRVs*	*124*
Table A.3-7.	*Biological effect endpoints for DMMP guidelines and selected TRVs*	*125*
Table A.3-8.	*SMS biological effects criteria for marine sediment toxicity tests*	*126*
Table A.3-9.	*Results of Phase 2 site-specific toxicity testing of surface sediment samples from the LDW*	*127*
Table A.3-10.	*Sediment toxicity datasets meeting project data quality objectives*	*131*
Table A.3-11.	*Summary of site-specific sediment toxicity test results for surface sediment samples collected by King County at Duwamish/Diagonal CSO/SD site*	*132*
Table A.3-12.	*Imposex stage criteria*	*134*
Table A.3-13.	*TBT concentrations in sediment and Nassarius mendicus RPS indices*	*136*
Table A.3-14.	*TBT critical tissue-residue toxicity studies for benthic invertebrates*	*137*

Table A.3-15.	*Estimated COPC concentrations in whole-body crab tissue*	*138*
Table A.3-16.	*Zinc and PCB critical tissue-residue toxicity studies for crabs and other decapods*	*140*
Table A.3-17.	*Selected critical tissue-residue TRVs for crabs*	*142*
Table A.4-1.	*ROC/COPC pairs evaluated for fish*	*143*
Table A.4-2.	*TBT and total PCB exposure concentrations in whole-body fish tissue*	*145*
Table A.4-3.	*Proportions of dietary items in dietary exposure estimates for each fish ROC*	*148*
Table A.4-4.	*Dietary exposure calculations and resulting COPC concentrations in fish ROC diets*	*150*
Table A.4-5.	*Juvenile chinook salmon stomach contents data for dietary COPCs*	*152*
Table A.4-6.	*PCB critical tissue-residue toxicity studies for fish*	*154*
Table A.4-7.	*TBT critical whole-body tissue-residue toxicity studies for fish*	*159*
Table A.4-8.	*Arsenic dietary toxicity studies for fish*	*160*
Table A.4-9.	*Cadmium dietary toxicity studies for fish*	*161*
Table A.4-10.	*Copper dietary toxicity studies for fish*	*164*
Table A.4-11.	*Vanadium dietary toxicity studies for fish*	*166*
Table A.4-12.	*TRVs selected for COPCs evaluated using the critical tissue-residue approach*	*167*
Table A.4-13.	*TRVs selected for COPCs calculated using the dietary approach*	*168*
Table A.5-1.	*ROC/COPC pairs evaluated for wildlife receptors*	*169*
Table A.5-2.	*Dietary fractions of prey items used in wildlife exposure calculations*	*173*
Table A.5-3.	*Exposure factor values for each ROC*	*174*
Table A.5-4.	*COPC concentrations in tissues of prey species ingested by great blue heron, osprey, river otter, and harbor seal*	*185*
Table A.5-5.	*COPC concentrations in benthic invertebrate tissues used to estimate spotted sandpiper exposure*	*187*
Table A.5-6.	*COPC concentrations in LDW sediment used to estimate exposure of wildlife ROCs, except spotted sandpiper*	*188*
Table A.5-7.	*COPC concentrations in sediment used to estimate spotted sandpiper exposure*	*189*
Table A.5-8.	*Concentrations of metals and BEHP in surface water used to estimate exposure of great blue heron, osprey, river otter, and harbor seal*	*191*
Table A.5-9.	*Concentrations of metals and BEHP in surface water used to estimate exposure of spotted sandpiper*	*191*
Table A.5-10.	*Total PCB concentrations in surface water used to estimate exposure of wildlife ROCs*	*192*
Table A.5-11.	*Estimated dietary doses of COPCs for spotted sandpiper*	*193*
Table A.5-12.	*Dietary exposure doses of COPCs for great blue heron, osprey, river otter, and harbor seal*	*194*
Table A.5-13.	*PCB dietary toxicity studies for birds*	*196*

Table A.5-14. 2,3,7,8-TCDD toxicity studies for birds for use in PCB TEQ risk analysis 199
Table A.5-15. Arsenic dietary toxicity studies for birds 201
Table A.5-16. Cadmium dietary toxicity studies for birds 202
Table A.5-17. Chromium dietary toxicity studies for birds 204
Table A.5-18. Cobalt dietary toxicity studies for birds 204
Table A.5-19. Copper dietary toxicity studies for birds 206
Table A.5-20. Lead dietary toxicity studies for birds 207
Table A.5-21. Mercury dietary toxicity studies for birds 209
Table A.5-22. Nickel dietary toxicity studies for birds 211
Table A.5-23. Selenium dietary toxicity studies for birds 213
Table A.5-24. Vanadium dietary toxicity studies for birds 214
Table A.5-25. Zinc dietary toxicity studies for birds 216
Table A.5-26. PCB dietary toxicity studies for mammals 219
Table A.5-27. 2,3,7,8-TCDD toxicity studies for mammals 222
Table A.5-28. Arsenic dietary toxicity studies for mammals 224
Table A.5-29. Cobalt dietary toxicity studies for mammals 225
Table A.5-30. Mercury dietary toxicity studies for mammals 227
Table A.5-31. Selenium dietary toxicity studies for mammals 228
Table A.5-32. TRVs for ROC/COPC pairs 234
Table A.6-1. Detection frequencies and frequencies of detected concentrations greater than SQS and CSL for all SMS COPCs 237
Table A.6-2. Sediment chemistry and toxicity test results for samples from the LDW 239
Table A.6-3. Detection frequencies and frequencies of detected concentrations above the SL/NOAEL and ML/LOAEL for COPCs without SMS chemical criteria 243
Table A.6-4. Number of detected concentrations for each COPC greater than SQS and CSL chemical criteria based on sediment chemistry data 244
Table A.6-5. Summary of chemical data for COPCs with at least one reporting limit greater than SQS/CSL chemical criteria 249
Table A.6-6. Summary of detected results and RLs for total DDTs and total chlordane relative to NOAELs and LOAELs 253
Table A.6-7. HQs for and benthic invertebrates exposed to cis-1,2-dichloroethene in porewater 260
Table A.6-8. TBT risk estimates for benthic invertebrates using the critical tissue-residue approach 263
Table A.6-9. HQs for crabs using whole-body exposure and effects data 264
Table A.6-10. Level of uncertainty associated with crab TRVs 266
Table A.6-11. Estimated total DDT and methoxychlor concentrations in whole-body crab tissue 267
Table A.6-12. Total DDT and methoxychlor critical tissue-residue toxicity studies for crabs and other decapods 268
Table A.6-13. Crab HQs using whole-body exposure and effects data 270

Table A.6-14.	*Summary of risk characterization for crab*	*270*
Table A.6-15.	*HQ calculations for juvenile chinook salmon*	*273*
Table A.6-16.	*Chemicals detected in ≥ 5% of baseline surface sediment samples that were not analyzed in LDW tissue samples*	*274*
Table A.6-17.	*Metals and PAHs with maximum total concentrations in water in any cell exceeding TRVs following the Tier 1 analysis*	*277*
Table A.6-18.	*Comparison of juvenile chinook salmon HQs assuming that they forage only in intertidal areas versus throughout the LDW*	*279*
Table A.6-19.	*COPC concentrations in sediment samples co-located with benthic invertebrate tissue samples relative to the site-wide baseline surface sediment dataset for dietary COPCs*	*280*
Table A.6-20.	*Toxicity studies for total PAHs and fish, including studies with a field component*	*283*
Table A.6-21.	*Cadmium dietary toxicity studies for fish*	*285*
Table A.6-22.	*LDW-specific studies of juvenile chinook salmon*	*286*
Table A.6-23.	*Critical tissue-residue toxicity studies of endrin in fish*	*290*
Table A.6-24.	*HQs for juvenile chinook salmon and endrin based on a critical tissue-residue approach*	*291*
Table A.6-25.	*Summary of risk characterization for juvenile chinook salmon*	*293*
Table A.6-26.	*HQ calculations for English sole*	*294*
Table A.6-27.	*HQ calculations for shiner surfperch and English sole for total PCBs*	*295*
Table A.6-28.	*HQs for BEHP, dimethyl phthalate, and di-n-butyl phthalate*	*296*
Table A.6-29.	*English sole dietary exposure estimates based on three assumed incidental sediment ingestion scenarios*	*297*
Table A.6-30.	*Dietary HQs for English sole as a function of sediment consumption*	*297*
Table A.6-31.	*English sole NOAEL-based HQs for chemicals without LOAEL TRVs*	*303*
Table A.6-32.	*HQs for English sole total PCBs and PCB TEQs*	*305*
Table A.6-33.	*Growth, reproductive effects, and survival studies using English sole collected from the LDW*	*305*
Table A.6-34.	*Critical tissue-residue toxicity studies of endosulfan in fish*	*307*
Table A.6-35.	*HQs for English sole and endosulfan and endrin*	*308*
Table A.6-36.	*Summary of risk characterization for English sole*	*310*
Table A.6-37.	*HQ calculations for Pacific staghorn sculpin*	*312*
Table A.6-38.	*Pacific staghorn sculpin HQs for BEHP, dimethyl phthalate, and di-n-butyl phthalate using reporting limits as exposure concentrations*	*314*
Table A.6-39.	*Shiner surfperch and Pacific staghorn sculpin UCLs and HQs for TBT*	*315*
Table A.6-40.	*Comparison of crab and benthic invertebrate tissue concentrations for Pacific staghorn sculpin dietary COPCs*	*316*
Table A.6-41.	*HQs for Pacific staghorn sculpin for arsenic, cadmium, copper, and vanadium with and without the use of larger crab tissue data*	*317*
Table A.6-42.	*Pacific staghorn sculpin NOAEL-based HQs for chemicals for which no LOAEL TRVs were identified*	*319*

Table A.6-43.	*HQs for Pacific staghorn sculpin and endosulfan and endrin*	*320*
Table A.6-44.	*Summary of risk characterization for Pacific staghorn sculpin*	*322*
Table A.6-45.	*HQ calculations for spotted sandpiper in the six exposure scenarios evaluated*	*326*
Table A.6-46.	*COPC concentrations in sediment samples co-located with benthic invertebrate tissue samples relative to the site-wide baseline surface sediment dataset for COPCs without a significant sediment/tissue relationship*	*335*
Table A.6-47.	*Level of uncertainty associated with TRVs for birds*	*337*
Table A.6-48.	*Exposure concentrations of total DDTs in benthic invertebrate tissue and sediment for spotted sandpiper*	*338*
Table A.6-49.	*Ingested doses of total DDTs for spotted sandpiper*	*339*
Table A.6-50.	*DDT, DDD, and DDE dietary toxicity studies for birds*	*340*
Table A.6-51.	*DDT HQs for spotted sandpiper*	*342*
Table A.6-52.	*Summary of risk characterization for spotted sandpiper*	*344*
Table A.6-53.	*HQ calculations for great blue heron*	*346*
Table A.6-54.	*Exposure concentrations of total DDTs*	*348*
Table A.6-55.	*Summary of risk characterization for great blue heron*	*350*
Table A.6-56.	*HQ calculations for osprey*	*351*
Table A.6-57.	*Exposure concentrations of total DDTs*	*353*
Table A.6-58.	*Estimated PCB concentration in osprey eggs*	*354*
Table A.6-59.	*Toxicity studies for PCBs in bird eggs*	*355*
Table A.6-60.	*HQs for PCBs estimated in osprey eggs*	*355*
Table A.6-61.	*Summary of risk characterization for osprey*	*357*
Table A.6-62.	*HQ calculations for river otter*	*358*
Table A.6-63.	*Contribution of each prey category to the ingested PCB dose for river otter*	*359*
Table A.6-64.	*Level of uncertainty associated with TRVs for mammals*	*362*
Table A.6-65.	*Summary of risk characterization for river otter*	*363*
Table A.6-66.	*HQ calculations for harbor seal*	*364*
Table A.6-67.	*Summary of risk characterization for harbor seal*	*366*
Table A.7-1.	*COPCs, COPCs that exceed relevant risk thresholds, and risk drivers*	*368*
Table A.7-2.	*Rationale for risk driver designation*	*370*
Table A.7-3.	*Chemicals with RLs greater than SQS that were not identified as risk drivers*	*374*
Table A.8-1.	*COPCs and ROCs with HQs ≥ 1.0*	*380*

List of Figures

Figure A.2-1.	*Screening process for benthic invertebrate COPCs using sediment chemistry data*	*61*
Figure A.2-2.	*Conceptual site model for fish and the benthic invertebrate community*	*106*
Figure A.2-3.	*Conceptual site model for wildlife*	*107*
Figure A.2-4.	*Generalized food web diagram for the LDW*	*108*
Figure A.3-1.	*Frequency distribution of TBT concentrations in LDW sediment samples*	*120*
Figure A.6-1.	*Copper concentrations in sediment samples co-located with benthic invertebrate tissue samples relative to the LDW-wide baseline surface sediment dataset*	*333*
Figure A.6-2.	*Nickel concentrations in surface sediment samples co-located with benthic invertebrate tissue samples relative to the LDW-wide baseline surface sediment dataset*	*334*

List of Maps

Map A.1-1.	*Lower Duwamish Waterway study area*
Map A.2-1.	*Locations of various LDW benthic invertebrate community surveys conducted from 1977 to 2004*
Map A.2-2.	*Tissue sampling locations in the Lower Duwamish Waterway*
Map A.2-3.	*Baseline surface sediment sampling locations and porewater sampling areas*
Map A.3-1.	*Porewater sampling locations at GWI*
Map A.3-2.	*Porewater sampling locations at Boeing Plant 2/Jorgensen Forge*
Map A.3-3.	*Gastropod sampling locations and TBT concentrations in baseline surface sediment samples*
Map A.3-4.	*TBT concentrations in baseline surface sediment samples and benthic invertebrate tissue and co-located sediment sampling locations*
Map A.3-5.	*Results of baseline toxicity testing conducted with LDW surface sediment samples*
Map A.5-1.	*Exposure areas and sampling locations of data used in the spotted sandpiper exposure assessment*
Map A.6-1a.	*Phase 2 sediment toxicity test results compared to SMS biological effects criteria and detected chemical concentrations at baseline surface sediment locations compared to SMS chemical criteria (RM 0.0-1.0)*
Map A.6-1b.	*Phase 2 sediment toxicity test results compared to SMS biological effects criteria and detected chemical concentrations at baseline surface sediment locations compared to SMS chemical criteria (RM 1.0-2.0)*
Map A.6-1c.	*Phase 2 sediment toxicity test results compared to SMS biological effects criteria and detected chemical concentrations at baseline surface sediment locations compared to SMS chemical criteria (RM 2.0-3.0)*

Map A.6-1d. *Phase 2 sediment toxicity test results compared to SMS biological effects criteria and detected chemical concentrations at baseline surface sediment locations compared to SMS chemical criteria (RM 3.0-4.0)*

Map A.6-1e. *Phase 2 sediment toxicity test results compared to SMS biological effects criteria and detected chemical concentrations at baseline surface sediment locations compared to SMS chemical criteria (RM 3.5-3.8)*

Map A.6-1f. *Phase 2 sediment toxicity test results compared to SMS biological effects criteria and detected chemical concentrations at baseline surface sediment locations compared to SMS chemical criteria (RM 4.0-5.0)*

Map A.6-2. *SL/ML categories for nickel and NOAEL/LOAEL categories for total DDTs and total chlordane at LDW baseline surface sediment sampling locations*

Map A.6-3. *Exceedances of SQS and CSL (chemical criteria and toxicity combined) using Thiessen polygons for the LDW baseline surface sediment dataset*

Map A.6-4. *Exceedances of SQS and CSL chemical criteria using Thiessen polygons for PCBs in the LDW baseline surface sediment dataset*

Map A.6-5. *Exceedances of SQS and CSL chemical criteria using Thiessen polygons for bis(2-ethylhexyl) phthalate in the LDW baseline surface sediment dataset*

Map A.6-6. *1,2,4-Trichlorobenzene detected concentrations and RLs compared to SQS and CSL chemical criteria at baseline surface sediment sampling locations*

Map A.6-7. *2,4-Dimethylphenol detected concentrations and RLs compared to SQS and CSL chemical criteria at baseline surface sediment sampling locations*

Map A.6-8. *Hexachlorobenzene detected concentrations and RLs compared to SQS and CSL chemical criteria at baseline surface sediment sampling locations*

Map A.6-9. *Baseline surface sediment locations with only non-detect RLs (no detected chemicals) greater than SQS or CSL chemical criteria*

Map A.6-10. *Areas with total PCB concentration greater than SQS and CSL, as represented by Thiessen polygons versus IDW interpolation*

Acronyms

Acronym	Definition
AET	apparent effects threshold
Ah	aryl hydrocarbon
ANOVA	analysis of variance
AOC	Administrative Order on Consent
ATSDR	Agency for Toxic Substance and Disease Registry
AWQC	ambient water quality criteria
BCA	benthic community analysis
BEHP	bis(2-ethylhexyl) phthalate
BMF	biomagnification factor
bw	body weight
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act

Acronym	Definition
COC	chemical of concern
COI	chemical of interest
COPC	chemical of potential concern
CSL	cleanup screening level of SMS
CSM	conceptual site model
CSO	combined sewer overflow
CSO/SD	combined sewer overflow/storm drain
DDTs	DDT and its metabolites
DMMP	Dredged Material Management Program
DNA	deoxyribonucleic acid
dw	dry weight
EC50	concentration that causes a non-lethal effect in 50% of an exposed population
Ecology	Washington State Department of Ecology
EPA	US Environmental Protection Agency
ERA	ecological risk assessment
ERED	Environmental Residue Effects Database
EROD	ethoxyresorufin-O-deethylase
ESA	Endangered Species Act
FAV	final acute value
FCV	final chronic value
FIR	food ingestion rate
FMR	free-living metabolic rate
FS	feasibility study
GWI	Great Western International
HHRA	human health risk assessment
HPAH	high-molecular-weight polycyclic aromatic hydrocarbon
HQ	hazard quotient
IDW	inverse distance weighting
IP	intraperitoneal
IR	ingestion rate
kcal	kilocalories
LAET	lowest apparent effects threshold
2LAET	second lowest apparent effects threshold
LC50	concentration that causes the death of 50% of a group of test animals

Acronym	Definition
LC100	concentration that causes the death of 100% of a group of test animals
LD50	dose that causes the death of 50% of a group of test animals
LDW	Lower Duwamish Waterway
LDWG	Lower Duwamish Waterway Group
LOAEL	lowest-observed-adverse-effect level
LOEC	lowest-observed-effect concentration
LPAH	low-molecular-weight polycyclic aromatic hydrocarbon
MHHW	mean higher high water
ML	maximum level of DMMP
MLLW	mean lower low water
MTCA	Model Toxics Control Act
NMFS	National Marine Fisheries Service
NOAEL	no-observed-adverse-effect level
NOEC	no-observed-effect concentration
NPL	National Priorities List
OC	organic carbon
PAH	polycyclic aromatic hydrocarbon
PCB	polychlorinated biphenyl
ppt	parts per thousand
PSAMP	Puget Sound Ambient Monitoring Program
PSDDA	Puget Sound Dredged Disposal Analysis
PSEP	Puget Sound Estuary Program
QAPP	quality assurance project plan
QA/QC	quality assurance/quality control
QSAR	quantitative structure-activity relationship
RCRA	Resource Conservation and Recovery Act
RGS	reporter gene system
RFI	RCRA facility investigation
RI	remedial investigation
RI/FS	remedial investigation/feasibility study
RL	reporting limit
RM	river mile
ROC	receptor of concern
RPF	relative potency factor

Acronym	Definition
RPS	relative penis size
SD	standard deviation
SL	screening level of DMMP
SMS	Washington State Sediment Management Standards
SQG	sediment quality guideline
SQS	sediment quality standards of SMS
SUF	site use factor
SVOC	semivolatile organic compound
SWAC	spatially weighted average concentration
T&E	threatened and endangered
TBT	tributyltin
TEC	toxic equivalence concentration
TEF	toxic equivalency factor
TCDD	tetrachlorodibenzo-*p*-dioxin
TEQ	toxic equivalent
TOC	total organic carbon
TRV	toxicity reference value
UCL	upper confidence limit
USACE	US Army Corps of Engineers
USGS	US Geological Survey
VOC	volatile organic compound
WAC	Washington Administrative Code
WDFW	Washington State Department of Fish and Wildlife
WERF	Water Environment Research Foundation
WHO	World Health Organization
WIR	water ingestion rate
WQA	water quality assessment
ww	wet weight

Executive Summary

This document presents the Phase 2 (baseline) Ecological Risk Assessment (ERA) for the Lower Duwamish Waterway (LDW), as outlined in the work plan for the Phase 2 Remedial Investigation (RI) (Windward 2004e). Baseline risk assessments, as defined in EPA (1988) guidance, "provide an evaluation of the potential threat to human health and the environment in the absence of any remedial action. They provide the basis for determining whether or not remedial action is necessary and the justification for performing remedial actions." The Phase 2 RI is being conducted by the Lower Duwamish Waterway Group (LDWG) under the oversight of the US Environmental Protection Agency (EPA) and the Washington State Department of Ecology (Ecology). These parties agreed in an Administrative Order on Consent (AOC) to conduct the RI in accordance with applicable Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) and Model Toxics Control Act (MTCA) guidance, policies, and procedures. Compliance with MTCA includes compliance with the Washington State Sediment Management Standards (SMS) (Washington Administrative Code [WAC] 173-204).

The baseline ERA presents risk estimates for benthic invertebrate, fish, and wildlife species that may be exposed to chemicals of potential concern (COPCs) found in sediment, water, and aquatic biota from the LDW. The dataset used in the baseline ERA consists of historical data and sediment and tissue chemistry data collected from the LDW during Phase 2 to supplement the historical data that were used in the Phase 1 ERA (Windward 2003a). The baseline ERA consists of separate sections on problem formulation, exposure assessment, effects assessment, risk characterization, and uncertainty analysis, each of which is briefly summarized below.

ES.1 PROBLEM FORMULATION

The problem formulation of the ERA establishes the overall scope of the assessment. Because it is impractical to evaluate risks to every potentially-exposed species, it is standard ERA practice and consistent with the SMS to focus on representative receptor species that typify groups of organisms with specific exposure pathways. One objective of selecting representative receptors is to choose species for which the risk conclusions will be protective or representative of other species that were not explicitly evaluated. For example, an assessment of risks to great blue herons would be assumed to be protective of all wading birds that eat fish because of the higher exposure potential of great blue herons than other wading birds. In addition, risks to some species were analyzed because those species are highly valued by society, such as endangered or threatened species.

Representative receptors of concern (ROCs) selected for this Phase 2 ERA were the benthic invertebrate community, crabs, English sole, Pacific staghorn sculpin, great blue heron, spotted sandpiper, osprey, river otter, and harbor seal. In addition,

juvenile chinook salmon was selected as an ROC because they are a federally protected species that use the LDW during outmigration to Puget Sound.

The problem formulation includes a description of the data available for conducting the ERA, the suitability of the data for risk assessment purposes, and a risk-based screening process that allows the risk assessment to focus on COPCs and eliminate chemicals that do not pose risks to ecological receptors.

Data used in the ERA consisted largely of:

- Surface sediment (uppermost 0 to 15 cm) chemistry data
- Site-specific sediment toxicity test results
- Sediment porewater chemistry data
- Tissue chemistry data for benthic invertebrates (including benthic infauna and epifauna, crabs, clams, and mussels), English sole, Pacific staghorn sculpin, shiner surfperch, and juvenile chinook salmon

For each ROC selected, COPCs were identified through a conservative risk-based screening process. COPCs identified included: 45 chemicals (including tributyltin [TBT], metals, polychlorinated biphenyls [PCBs] and other organic compounds) for benthic invertebrate communities; 2 chemicals (total PCBs and zinc) for crabs; 6 chemicals (arsenic, cadmium, copper, total PCBs, TBT, and vanadium) for at least one fish ROC, and 12 chemicals (arsenic, cadmium, chromium, cobalt, copper, lead, mercury, nickel, selenium, total PCBs, zinc, and vanadium) for at least one wildlife ROC. COPCs that were never detected in tissue because reporting limits (RLs) were above the screening criteria and COPCs without effect-level toxicity information were evaluated in the uncertainty analysis. In addition, organochlorine pesticides in some sediment samples and all tissue samples collected in 2004 were qualified as estimates (JN-qualified) because of probable analytical interference from PCBs, resulting in a high bias in concentration and a tentative identification (Windward 2005b, c). Therefore, risks from organochlorine pesticides were also assessed in the uncertainty analysis.

The problem formulation also presents the conceptual site models for the ROCs. Conceptual site models identify and describe pathways in which ROCs may be exposed to COPCs associated with sediment within the LDW. The pathways evaluated in the ERA included both direct sediment exposure and indirect exposure through the ingestion of prey or water from the LDW. Exposures associated with direct contact with water in the LDW were based on water data and risk estimates from the King County water quality assessment ERA (King County 1999b) and more recent water sampling for PCBs.

Finally, assessment and measurement endpoints were identified in the problem formulation. Survival, growth, and reproduction were the key endpoints under review for most ROCs in this assessment. Biomarker, behavioral, and histological endpoints

were not included as assessment endpoints. Typically, ERAs focus on ecological effects at the individual level or higher (i.e., population level). In this way, the emphasis is placed on endpoints that integrate an overall response by an organism, rather than indicators of a biochemical response that may or may not result in an ecologically relevant effect.

The representative ROCs, COPCs, pathways, and endpoints formed the scope for the Phase 2 ERA. Uncertainties associated with these analyses are discussed in the uncertainty analysis.

ES.2 EXPOSURE ASSESSMENT

The exposure assessment estimates the potential exposure of each ROC to the sediment-associated COPCs identified in the problem formulation. Exposure of benthic invertebrates to COPCs was assessed primarily by evaluating the distribution, concentration, and co-occurrence of COPCs in surface sediment, with the exception of risks to crabs and risks from sediment-associated TBT, which were both assessed using a critical tissue-residue approach. Risks to gastropods from exposure to TBT were also evaluated in two field studies of the most sensitive endpoint for these benthic invertebrates. In addition, risks to benthic invertebrates from exposure to volatile organic compounds (VOCs) were assessed using sediment porewater data.

Exposure of fish to COPCs was characterized based either on COPC concentrations in fish tissue or on COPC concentrations in likely fish prey. For wildlife ROCs, the exposure assessment identified equations and parameters to quantify the ingested dose of COPCs. Dietary doses for wildlife were estimated using available information on ROC biology and life histories, including body weight, feeding behavior, site usage, and diet.

ES.3 EFFECTS ASSESSMENT

Potential adverse effects (i.e., reduced survival, reduced growth, or impaired reproduction) were identified in the effects assessment. For the benthic invertebrate community, direct measures of sediment toxicity provided by site-specific sediment toxicity tests were given primary consideration over comparisons of sediment chemistry to chemical criteria. For locations without site-specific toxicity information, chemical criteria provided by the SMS were used to set benthic invertebrate effects levels for most of the COPCs. For COPCs without chemical-specific sediment criteria, toxicologically based guidelines from the Dredged Material Management Program (DMMP) or toxicity information from the literature was used. For gastropods, an invertebrate group highly sensitive to TBT, a direct, site-specific assessment of effects was also given primary consideration. For assessment of the effects on benthic invertebrates of VOCs in porewater, toxicity data from the literature were used.

For crabs, fish, and wildlife, a detailed evaluation was conducted of studies in the scientific literature documenting effects of COPCs on the ROCs or similar species. This

literature review identified COPC concentrations in receptor tissue or doses associated with no effects (i.e., safe concentrations or doses), in addition to concentrations or doses indicating a threshold of adverse effects. Both sets (i.e., no-observed-adverse-effect level [NOAEL] and lowest-observed-adverse-effect level [LOAEL]) of toxicity reference values (TRVs) are summarized in tables, and the rationale for TRV selection is provided.

ES.4 RISK CHARACTERIZATION AND UNCERTAINTY ANALYSIS

The exposure and effects data were compared in the risk characterization to assess the potential for sediment-associated COPCs to cause adverse effects to the ROCs. This analysis identified the following conclusions.

- **Benthic Invertebrate Community —** The goal of the SMS is to reduce and ultimately eliminate adverse effects on biological resources (WAC 173-204-100). Sediment chemistry and site-specific toxicity test results indicate that no adverse effects to benthic invertebrates living in intertidal and subtidal sediments are predicted for 75% of the LDW area (i.e., the area in which chemical concentrations were less than or equal to chemical sediment quality standard [SQS] criteria and where sediments were nontoxic according to biological SQS criteria). There is a higher likelihood for adverse effects in approximately 7% of the LDW area, which was designated as having chemical concentrations or biological effects in excess of cleanup screening level (CSL) criteria. The remaining 18% of the LDW area had chemical concentrations or biological effects between the SQS and CSL criteria, indicating that risks to benthic invertebrate communities are less certain in these areas than in areas with concentrations greater than one or more CSL values. Some uncertainty is associated with these area estimates because areas were estimated by interpolating from individual points at which sediments were sampled. The SQS and CSL criteria were exceeded by 39 chemicals; 2 additional chemicals exceeded only the SQS criteria.[1] Risks to the benthic invertebrate community from all VOCs detected in sediment porewater were very low, except for cis-1,2-dichloroethene, which had concentrations 21 times the no-effects concentration in a small area at RM 2.4; all concentrations of cis-1,2-dichloroethene were less than the concentration associated with adverse effects. Therefore, there is uncertainty whether exposure to cis-1,2-dichloroethene within the LDW is sufficiently high to result in adverse effects in this small area. Risks to benthic invertebrates from TBT were very low based on NOAEL-based hazard quotients (HQs) less than 1.0 and the absence

[1] Total DDTs, nickel, and total chlordane also exceeded their DMMP guidelines or literature-based TRVs at one or more locations.

of imposex in all gastropods, except one neogastropod with imposex characterized as Stage 2, a stage that is not expected to impact reproduction.

- **Crabs —** Exposure concentrations of total PCBs in tissue were equal to concentrations associated with adverse effects in crabs, indicating the potential for adverse effects. Exposure concentrations of zinc in tissue were greater than concentrations associated with no effects but less than those associated with adverse effects, indicating there is uncertainty whether exposure within the LDW is sufficiently high to result in adverse effects.
- **Fish —** Exposure concentrations for three of the six COPCs for fish (PCBs, cadmium, and vanadium) were greater than concentrations associated with adverse effects for English sole. LOAEL-based hazard quotients (HQs[2]) for cadmium and vanadium were both 1.2, and LOAEL-based HQs for PCBs ranged from 0.98 to 5.0 based on effects concentrations in the study reporting the lowest TRVs. Therefore, there is a potential for adverse effects from PCBs, but risk estimates are uncertain because the exposure concentration was in between the concentrations selected as LOAELs. Estimated exposures of English sole to two additional COPCs (arsenic and copper) were greater than their respective no-effects levels but were lower than the adverse effect levels associated with survival, growth, or reproduction, indicating that there is uncertainty whether exposure within the LDW is sufficiently high to result in adverse effects. Site-specific studies of English sole indicate the potential for reproductive effects that correlate with exposure to chemical mixtures in the field. However, the relationship of such effects to specific chemicals has not been established. Exposure concentrations of PCBs, cadmium, and vanadium for Pacific staghorn sculpin were equal to or greater than the concentrations associated with adverse effects in at least one area within the LDW. LOAEL-based HQs up to 1.0 and 1.2 were estimated for cadmium and vanadium, respectively, indicating a potential for adverse effects. LOAEL-based HQs for PCBs ranged from 0.30 to 3.8 based on effects concentrations in the study reporting the lowest TRVs. Therefore, there is a potential for adverse effects from PCBs, but risk estimates are uncertain because the exposure concentrations were in between the concentrations selected as the LOAEL range. The exposure concentrations of TBT and copper were greater than their respective no-effects concentrations for Pacific staghorn sculpin in at least one area within the LDW but less than those associated with adverse effects. Thus, the potential for adverse effects is uncertain. For juvenile chinook salmon, exposure concentrations of cadmium were greater than concentrations associated with adverse effects in any fish species but lower than concentrations

[2] The HQ is the ratio of the exposure concentration (or dose) to a concentration (or dose) associated with adverse effects.

associated with adverse effects in salmonids. Exposure concentrations of arsenic, copper, and vanadium in the diet of juvenile chinook salmon were greater than their respective no-effect concentrations but less than concentrations associated with adverse effects.

- **Birds —** Estimated exposures of spotted sandpiper to six of the 12 COPCs (copper, chromium, lead, mercury, PCB toxic equivalent (TEQ),[3] and vanadium) for spotted sandpiper were greater than the dietary doses associated with adverse effects on survival, growth, or reproduction in at least one area within the LDW (LOAEL-based HQs of up to 1.1, 1.8, 5.5, 1.0, 1.5, and 1.4, respectively). Therefore, there is a potential for adverse effects from these COPCs. Estimated doses to great blue heron of all four COPCs (chromium, lead, mercury, and total PCBs) were less than no-effects levels, indicating very low risk. For osprey, estimated doses of PCBs were greater than no-effect levels for osprey using a TEQ approach but less than those levels using a total PCBs approach; the latter risk estimate is less uncertain. Therefore, the potential for adverse effects from PCBs is uncertain for osprey. Estimated doses of the remaining three COPCs to osprey (chromium, lead, and mercury) were less than the doses associated with no-effects, indicating very low risk.
- **Mammals —** Estimated dietary doses of total PCBs were greater than doses associated with adverse effects for river otters, with a LOAEL-based HQ of 2.9. Estimated exposure of river otters to mercury was greater than a no-effects level but was less than adverse effects levels associated with survival, growth, or reproduction, indicating that the potential for effects is uncertain. Exposures of otter to the remaining three COPCs (arsenic, cobalt, and selenium) and exposures of harbor seals to both COPCs (mercury and total PCBs) were less than their respective no-effects levels, indicating very low risk.

Table ES-1 provides a summary of COPCs for crabs, fish, or wildlife for which either the NOAEL-based HQ was greater than 1.0 or the LOAEL-based HQ was greater than or equal to 1.0. Table ES-2 lists the COPCs for benthic invertebrates that exceeded SMS criteria, DMMP guidelines, or TRVs. In summary, risk estimates for PCB exposure indicated a potential for adverse effects to the benthic invertebrate community, crabs, English sole, Pacific staghorn sculpin, spotted sandpiper, and river otter. There is also a potential for adverse effects to osprey from PCB exposure, but risk estimates for this ROC are more uncertain because exposures were greater than no-effect levels but less than levels associated with adverse effects. Other COPCs with exposures greater than levels associated with adverse effects were cadmium, chromium, copper, lead, mercury, and vanadium. Numerous additional chemicals may pose a risk to the benthic invertebrate community as shown in Table ES-2.

[3] EPA refers to this as the toxic equivalence concentration or TEC.

Table ES-1. COPCs and ROCs with HQs ≥ 1.0

COPC	ROC	NOAEL HQ	LOAEL HQ
COPCs with LOAEL-Based HQs ≥ 1.0[a]			
Total PCBs	crab	**10**	**1.0**
	English sole	**4.9 – 25**	0.98 – **5.0**
	Pacific staghorn sculpin	**1.5 – 19**	0.30 – **3.8**
	river otter	**5.8**	**2.9**
PCB TEQs	spotted sandpiper	**1.9 – 15**	0.18 – **1.5**
Cadmium	juvenile chinook salmon	**5.0**	**1.0**
	English sole	**6.1**	**1.2**
	Pacific staghorn sculpin	**3.0 – 5.2**	0.60 – **1.0**
Chromium	spotted sandpiper	**1.3 – 8.8**	0.26 – **1.8**
Copper	spotted sandpiper	0.62 – **1.5**	0.45 – **1.1**
Lead	spotted sandpiper	0.58 – **19**	0.17 – **5.5**
Mercury	spotted sandpiper	**1.1 – 5.3**	0.21 – **1.0**
Vanadium	English sole	**5.9**	**1.2**
	Pacific staghorn sculpin	**3.2 – 5.9**	0.65 **– 1.2**
	spotted sandpiper	**2.0 – 2.7**	**1.0 – 1.4**
COPCs with NOAEL-Based HQs ≥ 1.0 and LOAEL-Based HQs < 1.0[b]			
Total PCBs	spotted sandpiper	0.51 – **2.0**	0.18 – 0.71
PCB TEQs	osprey	**1.6**	0.16
	river otter	**4.5**	0.59
Arsenic	juvenile chinook salmon	**1.1**	0.73
	English sole	**1.2**	0.80
	crab	**3.9**	na
Benzoic acid	English sole	**1.5**	na
	Pacific staghorn sculpin	**2.1**	na
Cadmium	Pacific staghorn sculpin	**3.0 – 4.9**	0.60 – 0.98
Chromium	juvenile chinook salmon	**2.1**	na
	English sole	**1.1**	na
Copper	Juvenile chinook salmon	**1.9**	0.93
	English sole	**1.9**	0.93
	Pacific staghorn sculpin	0.9 – **1.5**	0.45 – 0.77
Mercury	river otter	**2.8**	0.57
TBT	Pacific staghorn sculpin	**1.6** – **2.9**	0.18 – 0.33
Vanadium	juvenile chinook salmon	**4.0**	0.79
Zinc	crab	**2.5**	0.91

Note: HQs for fish are the highest HQs in cases where more than one approach was used.

[a] The LOAEL-based HQs for endrin were 1.2 and 3.1 for English sole and Pacific staghorn sculpin, respectively, based on risk calculations discussed in the uncertainty section. These calculations were presented only in the uncertainty section because of analytical interferences from PCB Aroclors in the pesticide analyses of LDW tissue samples, resulting in uncertainties in pesticide identification and a high bias in pesticide concentrations.

[b] The NOAEL-based HQs were ≥1 for the following COPC/ROC pairs based on risk calculations discussed in the uncertainty section: 1) total DDTs and spotted sandpiper (2.6 to 4.3), 2) endrin and juvenile chinook salmon (3.6), 3) alpha-endosulfan and English sole (6.8) and Pacific staghorn sculpin (2.3), 4) beta-endosulfan and English sole (29) and Pacific staghorn sculpin (6.6), 5) endrin and juvenile chinook salmon (3.6), and 6) methoxychlor and crab (3.6). These calculations were presented in the uncertainty section because of analytical interferences from PCB Aroclors in the pesticide analyses, resulting in uncertainties in pesticide identification and a high bias in pesticide concentrations.

COPC – chemical of potential concern
HQ – hazard quotient
LOAEL – low-observed-adverse-effect level
na – not available
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
ROC – receptor of concern
TBT – tributyltin
TEQ – toxic equivalent

Bold identifies NOAEL-based HQs greater than 1.0 or LOAEL-based HQs greater than or equal to 1.0.

Table ES-2. Benthic invertebrate community COPCs with detected concentrations in LDW surface sediments greater than SMS criteria, DMMP guidelines, or TRVs

COPC	Number of Detected Concentrations > SQS and ≤ CSL	Number of Detected Concentrations > CSL
Total PCBs	301	173
Bis(2-ethylhexyl) phthalate	48	58
Mercury	14	23
Lead	2	19
Zinc	26	16
Total chlordane[a]	19	14
Copper	0	12
Cadmium	2	11
Silver	0	10
Fluoranthene	31	8
Butyl benzyl phthalate	69	8
Indeno(1,2,3-cd)pyrene	15	8
Chromium	1	8
Arsenic	5	8
Phenol	18	7
Benzo(g,h,i)perylene	9	7
Benzoic acid	0	7
Dibenzo(a,h)anthracene	15	4
Nickel[b]	9	4
Total benzofluoranthenes	5	4
4-Methylphenol	0	4
Phenanthrene	24	3
Total HPAH	21	3
Acenaphthene	16	3
Fluorene	11	3

COPC	Number of Detected Concentrations > SQS and ≤ CSL	Number of Detected Concentrations > CSL
Benzo(a)anthracene	9	3
Dibenzofuran	7	3
Benzo(a)pyrene	5	3
Total LPAH	3	3
Pyrene	1	3
1,4-Dichlorobenzene	0	3
1,2-Dichlorobenzene	0	3
2-Methylnaphthalene	0	3
Dimethyl phthalate	0	2
Naphthalene	0	2
n-Nitrosodiphenylamine	0	2
Hexachlorobenzene	4	2
Benzyl alcohol	2	2
Chrysene	23	1
Total DDTs[a]	1	1
1,2,4-Trichlorobenzene	0	1
2,4-Dimethylphenol	0	1
Anthracene	2	0
Pentachlorophenol	1	0

[a] SMS criteria do not exist for these chemicals; number of exceedances was based on a comparison of sediment chemical concentrations to a TRV.

[b] SMS criteria do not exist for nickel. The DMMP SL and ML values were used for the comparison.

COPC – chemical of potential concern

CSL – cleanup screening level

HPAH – high-molecular-weight polycyclic aromatic hydrocarbon

LPAH – low-molecular-weight polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

SQS – sediment quality standards

Based on the risk estimates, uncertainties discussed in this ERA, natural background concentrations, and residual risks following early actions in the LDW, chemicals were identified as risk drivers for ecological receptors in accordance with EPA (1998) and MTCA (WAC 173-340-703) guidance. The risk drivers from both this ERA and the human health risk assessment (HHRA) will be the focus of remedial analyses in the feasibility study (FS).

In consultation with EPA and Ecology, PCBs were identified as a risk driver for river otter because estimated exposure concentrations for river otter were greater than the LOAEL by a factor of 2.9, and uncertainties in the risk estimate were relatively low. In addition, 41 chemicals were selected as risk drivers for the benthic invertebrate community because concentrations of these 41 chemicals exceeded Washington SMS in one or more locations.

Other COCs, chemicals that exceeded risk thresholds (LOAEL-based HQ ≥ 1.0) but were not selected as risk drivers may be addressed through focused evaluation in the FS, as part of the 5-year review, or included in the post-remedial monitoring program, as appropriate.

No quantitative ecological risk estimates were calculated for dioxins and furans and thus the level of ecological risk from dioxins and furans is unknown. Ecological risks associated with exposure to dioxins and furans within the LDW were not assessed for several reasons. Primarily, human health risks from dioxins and furans through seafood consumption were assumed to be unacceptable, and therefore, neither tissues from the LDW nor from background areas were analyzed for dioxins and furans. Dioxins and furans were determined to be a risk driver based on human health risks from both seafood consumption and direct contact pathways. Risk management decisions to address dioxin and furan contamination in LDW sediment will be based on MTCA and CERCLA regulations and guidance. Remedial decisions to address dioxin and furan contamination in sediment will be made by EPA and Ecology as part of the FS process and will be documented in the Record of Decision. Additional detail on dioxins and furans is provided in Section B.5.5.2 of the HHRA.

This ERA is based on the baseline surface sediment dataset, which includes sediment data collected prior to early actions in the LDW. Since these data were collected, early actions in the LDW have been conducted at two locations (Duwamish/Diagonal and the Boeing Developmental Center south storm drain in the Norfolk area). Therefore, the risks discussed in this ERA may represent an overestimate of current risks for areas where remediation has already occurred.

A.1.0 Introduction

This document presents the baseline ecological risk assessment (ERA) as part of the remedial investigation and feasibility study (RI/FS) for the Lower Duwamish Waterway (LDW). The LDW was added to the US Environmental Protection Agency's (EPA's) National Priorities List (NPL) under the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA), also known as Superfund, on September 13, 2001. The LDW was added to Ecology's Hazardous Sites List on February 26, 2002.

Under Superfund regulations, EPA requires that an RI/FS be conducted for all listed sites. An RI evaluates the nature and extent of chemical contamination, estimates baseline human health and ecological risks, and is used by risk managers to identify areas that should be remediated because they pose an unacceptable risk to human health or the environment. An FS proposes a number of alternative approaches to remediating the areas with unacceptable risk, and analyzes and compares these alternatives. The study area is shown in Map A.1-1.

The key parties involved in the LDW RI/FS are the City of Seattle, King County, the Port of Seattle, and The Boeing Company, working together for this project as the Lower Duwamish Waterway Group (LDWG), under the oversight of EPA and the Washington State Department of Ecology (Ecology). These parties agreed (in an Administrative Order on Consent, or AOC) to conduct the RI for the LDW in two phases. The first phase is complete, including a Phase 1 ERA based on data that existed at that time (Windward 2003c). The AOC requires that the RI be conducted in accordance with applicable CERCLA and MTCA guidance, policies, and procedures. Compliance with MTCA includes compliance with the Washington State Sediment Management Standards (SMS).

This document presents the Phase 2 (baseline) ERA as outlined in the Phase 2 RI work plan (Windward 2004e). Baseline risk assessments, as defined by EPA (1988) guidance for conducting an RI/FS, "provide an evaluation of the potential threat to human health and the environment in the absence of any remedial action. They provide the basis for determining whether or not remedial action is necessary and the justification for performing remedial actions." This baseline ERA presents risk estimates for ecological receptors that may come in contact with sediment–associated chemicals of potential concern (COPCs) through exposure to or ingestion of sediment, water, fish, and invertebrates (e.g., polychaete worms, clams, crabs) in the LDW.

This baseline ERA is based on data previously summarized in the Phase 1 ERA (Windward 2003b) and data collected since the Phase 1 RI was completed. It was developed in accordance with both national and regional EPA guidance (EPA 1992, 1997a, b, 1998). This baseline ERA includes the following sections:

- Section A.2.0 – Problem Formulation

- Section A.3.0 – Exposure and Effects Assessment: Benthic Invertebrates
- Section A.4.0 – Exposure and Effects Assessment: Fish
- Section A.5.0 – Exposure and Effects Assessment: Wildlife
- Section A.6.0 – Risk Characterization and Uncertainty Analysis
- Section A.7.0 – Selection of Primary Ecological Risk Drivers/Indicator Hazardous Substances
- Section A.8.0 - Conclusions
- Section A.9.0 – References

Details on site background, previous investigations, and environmental setting are provided in the Phase 2 RI and referenced accordingly. Table A.1-1 presents the sections in which the primary components of the risk assessment process are discussed for each receptor type.

Table A.1-1. Document organization for primary ERA components

ERA Component	ERA Section Number		
	Benthic Invertebrates	Fish	Wildlife
ROC selection	A.2.3.1	A.2.3.2	A.2.3.3
COPC selection	A.2.5.1	A.2.5.2	A.2.5.3
Conceptual site model	A.2.6	A.2.6	A.2.6
Exposure assessment	A.3.1 (benthic invertebrate community) A.3.3 (crabs)	A.4.1	A.5.1
Effects assessment	A.3.2 (benthic invertebrate community) A.3.4 (crabs)	A.4.2	A.5.2
Risk characterization and uncertainty analysis	A.6.1	A.6.2	A.6.3

[a] Available data used in the ERA are discussed in Section A.2.4.1, and exposure concentration calculations are presented in Attachment 11.

A.2.0 Problem Formulation

This section presents the problem formulation for this baseline ERA. Through the use of a risk-based screening approach, the problem formulation establishes which receptor of concern (ROC) and COPC pairs are further evaluated in the exposure and effects assessment, the risk characterization, and the uncertainty analysis. This section includes information regarding the environmental setting, ecological resources that use the site, selection of ROCs, a summary of relevant available data collected from the LDW, a COPC screen for ROCs, and the conceptual site model (CSM) for the LDW. Together, these elements establish the scope for this ERA.

A.2.1 ENVIRONMENTAL SETTING

This section presents an overview of the site's setting, hydrology, and habitat. Additional information regarding the setting, history, hydrology, and sediment regime of the LDW is presented in the Phase 2 RI.

A.2.1.1 Site description

The confluence of the Black and Green Rivers forms the Duwamish River 10.5 miles upstream from the southern end of Harbor Island. The LDW consists of the downstream portion of the Duwamish River, excluding the East and West Waterways around Harbor Island, and extends from river mile (RM) 0.0 near the southern tip of Harbor Island to upstream of the Upper Turning Basin, which is located near RM 4.8. The LDW is tidally influenced over its entire length, with the degree of tidal influence varying depending on stream flow and on tide stage at the mouth of the LDW. The mean tide range is 7.5 ft, and the mean diurnal range is 11 ft. Recorded tides have ranged from -4.6 to 14.7 ft mean lower low water (MLLW). Annual river discharge ranges from 43 to 51 m^3/sec (2,300 to 2,350 cfs) (NOAA 1998).

The US Army Corps of Engineers (USACE) maintains the LDW as a navigable waterway through dredging (Dexter et al. 1981). The typical cross section of the LDW includes a deeper, maintained navigation channel at the middle of the waterway, with intermittent shallow benches along the margins of the channel. The navigation channel is maintained throughout the study area, with typical depths ranging from greater than -30 ft MLLW downstream of RM 2.0 to less than -15 ft MLLW near the Upper Turning Basin. Shallower bench areas exist in the nearshore intertidal zones outside of the navigation channel, with variable dimensions and elevations. The width of the LDW is relatively uniform, ranging between about 500 and 700 ft. The navigation channel is approximately 200 ft wide.

The LDW is a highly modified estuary that has been straightened and dredged in the lower 4.8 miles to facilitate navigation and industrial development. The only remnant of the river's natural meanders exists west of Kellogg Island. The banks of the LDW are dominated by structures, including piers and buildings. Where they are not

occupied by structures, the banks are typically armored with a combination of riprap, concrete debris, and other structures for stabilization of banks, especially in mid to upper tidal elevations of the bank. Industrial land use dominates on the east bank in the immediate vicinity of the LDW. The west bank includes industrial, commercial, and mixed residential land uses in the vicinity of the LDW.

Exceptions to the industrialized condition of the LDW include the area around Kellogg Island, which is partially formed by a remnant meander of the natural Duwamish River channel, and some areas of mixed commercial, recreational, and residential uses within both the upstream and downstream areas of the LDW.

The LDW is a stratified saltwater wedge estuary (Stoner 1972). Circulation within a stratified estuary results from a net upstream movement of water within a bottom-layer saltwater wedge and a net downstream movement of fresher water in the layer that overrides the wedge. Recent average annual discharges from the Duwamish River have ranged between 43 and 51 m^3/s (2,300 to 2,350 cfs), as measured at the US Geological Survey (USGS) Tukwila gaging station, located at USGS-designated RM 12.4. Flow rates at the Auburn gaging station, located at USGS-designated RM 32.0, ranged from 4.3 to 329 m^3/s (200 to 15,200 cfs, the record high) between 1962 and 1994 (NOAA 1998). The saltwater wedge, which has its source in Elliott Bay, oscillates upstream and downstream with the tide and stream flow. At freshwater inflow greater than 1,000 cfs, the saltwater wedge does not extend upstream beyond the East Marginal Way Bridge (RM 6.3), regardless of the tide height (Stoner 1967). During periods of low freshwater inflow and high-tide stage, the saltwater wedge has extended as far upstream as the Foster Bridge at RM 8.7 (Stoner 1967).

Because of their circulation, estuaries naturally act as sediment traps for incoming sediment. Sediment from freshwater sources is transported into the estuary at the upstream end while sediment from coastal waters is transported into the estuary via the saltwater wedge. The presence of the man-made channel increases the cross-sectional area of the estuary relative to natural conditions, which results in a decrease in current velocities and an increase in deposition. Additional information related to LDW sediment stability and erosion potential is presented in the Phase 2 RI.

A.2.1.2 Habitat features

Benthic habitats within the LDW include intertidal habitat (exposed by low tides) and subtidal habitat (never exposed by low tides) (Table A.2-1). A typical cross section of the LDW includes intertidal habitats, subtidal transition areas (often with steep slopes), and a deeper navigation channel in the center of the LDW. Sediment characteristics (i.e., grain size and organic carbon [OC] content) vary throughout the LDW, depending on several physical parameters such as currents, quiescent areas (e.g., slips), and sediment sources (i.e., creeks and outfalls).

Table A.2-1. Habitat types represented in the LDW

Habitat Type	Description	Influential Physical Parameters	Extent and Condition in LDW
Upland	area outside the immediate influence of wetted area of the LDW (below +14 ft MLLW), many uplands consist of fill material	temperature, precipitation, soil type, groundwater elevation	dominated by industrial uses; some mixed-use residential, commercial, and recreational areas
Intertidal marshes	Intertidal area between -4 and +14 ft MLLW; exposed at low tides; marsh soils generally fine-textured and nutrient-rich, supporting grasses, sedges, rushes, and various other plants	salinity gradients, tidal variation, freshwater stream flow, wave action, water temperature, sediment characteristics and oxygen content	limited extent; various marsh habitat classifications: emergent marsh (e.g., Herring's House, Hamm Creek, and Upper Turning Basin restoration areas), tidal marsh (5 acres in LDW), high and low marsh on Kellogg Island (Blomberg et al. 1988)
Intertidal mudflats	Intertidal area between -4 and +14 ft MLLW, remnant mudflats isolated from upland riparian vegetation, exposed at low tides, sometimes shaded by overwater structures	salinity gradients, wave action, water temperature, sediment characteristics, and oxygen content	flats and shallows (approximately 30 to 50 acres in LDW) (Blomberg et al. 1988), approximately 8.6 miles of exposed sand/mud substrate (Battelle et al. 2001), largest remnant on Kellogg Island
Intertidal riprap	armored shoreline consisting of large rocks and rubble (riprap) or vertical wood or metal structures (sheet pile)	salinity gradients, wave action, water temperature, oxygen content	riprap-covered areas comprise approximately 17% of intertidal areas in LDW (USFWS 2000b)
Subtidal	area deeper than -4 ft MLLW, never exposed by low tide, includes navigation channel and transition areas, sediment composition ranges from sand to mud	salinity, sediment composition, grain size, and OC content, water depth, temperature	throughout LDW, including navigation channel

LDW – Lower Duwamish Waterway

MLLW – mean lower low water

OC – organic carbon

The majority of the LDW shoreline consists of riprap, pier aprons, or sheet piling (Tanner 1991). Shoreline armoring is usually present at the top of the intertidal zone; but areas of sloping mud, mudflats, and hard surfaces exist in the lower intertidal zone (Battelle et al. 2001). These hard surfaces support populations of encrusting organisms, such as barnacles, and burrowing organisms, such as shipworms (Leon 1980). However, because of the shoreline armoring, these intertidal flats are partially isolated from inputs of sediment, nutrients, and organic matter (i.e., woody debris) from upland riparian vegetation zones; this isolation degrades the habitat quality of these flats (Battelle et al. 2001). In addition, overwater structures, which are common throughout the LDW, often shade shallow and intertidal habitats, alter microclimates, and inhibit growth of aquatic plant communities, further degrading the value of nearshore habitats for native fauna (Battelle et al. 2001).

Sections of natural shoreline in the LDW occur only upstream of the Upper Turning Basin (Tanner 1991). Most (98%) of approximately 510 hectares (ha) (1,270 ac) of tidal marsh and 590 ha (1,450 ac) of flats and shallows and all of about 500 ha (1,230 ac) of tidal wetland have been either filled or dredged (Blomberg et al. 1988) or altered by the hydrologic changes resulting from the channelization of the estuary and

maintenance of the navigation channel. Remnant tidal marshes account for only 2 ha (5 ac), and mudflats account for 12 to 22 ha (30 to 54 ac) (Leon 1980).

Remnants of natural intertidal habitat are present in occasional patches throughout the LDW; the largest remnant of intertidal habitat surrounds Kellogg Island, located south of Harbor Island. Kellogg Island has been highly altered from its historical shape and function. In the 1950s and 1960s, it was filled with hydraulic dredge spoils by USACE. Later, in 1974, an upland component of Kellogg Island was created when the Port of Seattle deposited 1,700 m^3 (2,200 yd^3) of dredged materials on the south end of the island (Sato 1997). A mixture of introduced and native plant and tree species rapidly colonized the 7-ha island. Current habitat associated with the island includes high and low marsh, intertidal mudflats, and filled uplands (Canning et al. 1979). The island, including intertidal area near Kellogg Island, has been designated as a wildlife refuge (Hotchkiss 2006).

Subtidal habitat is variable throughout the LDW. Subtidal sediment composition ranges from sand to mud, depending on the sediment source and current speed (Windward 2003b). The sediments in the upstream portion of the LDW, near the head of the main channel at the Upper Turning Basin, are predominantly sand; whereas the sediments in the subtidal habitat further downstream (e.g., near Kellogg Island) are characterized as brown or brown-gray sandy mud overlying darker, more clayey mud.

Few surveys have investigated the vascular plant communities present in the LDW (Tanner 1991; Canning et al. 1979; USFWS 2000a; Cordell et al. 2001). The methods used to assess plant communities have ranged from analysis of aerial photos to field surveys. Many of these surveys were conducted to investigate habitat availability in the LDW and mainly addressed the plant communities of tidal marsh areas.

In estuaries in general, tidal elevation and salinity gradients determine the potential distribution of estuarine plants. Intertidal elevation gradients between MLLW and mean higher high water (MHHW) create habitats such as low-, mid-, and high-elevation tidal marshes. Salinity gradients range from saline to brackish to fresh tidal waters. The most productive areas for estuarine plant communities are found in tidal marshes. Marsh soils are generally fine-textured and nutrient-rich and support grasses, sedges, rushes, and various other types of plants associated with marine and estuarine habitats. In the LDW, there are 1.75 ha of habitat for vascular plants, primarily limited to portions of Kellogg Island and other small areas (USFWS 2000a).

Carex sp. and *Scirpus* sp. are the predominant type of vegetation between the Upper Turning Basin and Kellogg Island. Downstream from Kellogg Island are more marine plants, such as *Salicornia* sp., *Distichli* sp., and *Atriplex* sp. The interior high-marsh plant community of Kellogg Island, which is flooded only by higher spring tides, includes *Carex lyngbyei*, *Distichlis spicata*, *Juncus balticus* (Baltic rush), and *Phragmites* sp., a non-native species (Battelle et al. 2001). No eelgrass is found in the LDW, and

habitats with aquatic vegetation are rare (Battelle et al. 2001) Thus, plants present in the LDW are seldom present under water.

In recent years, there have been several restoration efforts in the LDW, including the Terminal 105 (T-105) channel along the western shoreline at RM 0.1, Herring's House off-channel marsh along the western shoreline at RM 0.5, Duwamish Diagonal south along the eastern shoreline at RM 0.65, Terminal 107 (T-107) at RM 0.6, the General Services Administration marsh along the eastern shoreline at RM 1.2, the Hamm Creek off-channel marsh on the western shoreline at RM 4.3, and two separate restoration sites in the Upper Turning Basin at RM 4.7. Restoration activities include removal of riprap and overwater structures, placement of log booms to decrease debris deposition and boat wakes, creation of intertidal benches to promote use by juvenile salmon, creation of new off-channel habitat (including emergent salt marshes), and creation of freshwater wetlands (including excavation and planting with native species) (Cordell et al. 2001).

A.2.2 RESOURCES POTENTIALLY AT RISK

This section provides an overview of the ecological resources that use the LDW, including threatened, endangered, and sensitive species. These resources are considered in three groups, which include species that could be directly or indirectly exposed to contaminated sediments: benthic invertebrates, fish, birds, and mammals. Representative species from these groups were selected as ROCs (Section A.2.3) and further evaluated to determine whether they may be at risk from contaminated sediments. Reptiles and amphibians are not likely to be exposed to sediment contamination in the LDW because habitat for these species is limited, and their presence has not been reported in any wildlife surveys conducted in the area[4] (Canning et al. 1979; Cordell et al. 1996; 1997; 1999). Therefore, they are not evaluated further in this ERA. In addition, risks to vascular plants and algae will not be evaluated in the Phase 2 ERA because they were evaluated as part of the Phase 1 ERA (Windward 2003b), and risk estimates were highly uncertain. No new information is available to resolve the uncertainties.

A.2.2.1 State and federal threatened, endangered, and sensitive species in the LDW

Fifteen species that are reported to occur either as residents or during migration in the LDW are listed under either the federal Endangered Species Act (ESA) or by the Washington State Department of Fish and Wildlife (WDFW) as candidate species, or threatened species (Table A.2-2). No species reported to occur in the LDW are listed under the ESA or by WDFW as endangered.

[4] A large tadpole was observed once in Slip 4.

Table A.2-2. Threatened and candidate species listed under ESA or by Washington State Department of Fish and Wildlife

COMMON NAME	SCIENTIFIC NAME	STATUS
Chinook salmon	*Oncorhynchus tshawytscha*	federal threatened species, state candidate species
Coho salmon	*Oncorhynchus kisutch*	federal species of concern
Puget Sound steelhead	*Oncorhynchus mykiss*	federal threatened species
River lamprey	*Lampetra ayresi*	federal species of concern, state candidate species
Bull trout	*Salvelinus confluentes*	federal threatened species, state candidate species
Pacific herring	*Clupea herengus pallasi*	federal species of concern, state candidate species
Pacific cod	*Gadus macrocephalus*	federal species of concern, state candidate species
Walleye pollock	*Theragra chalcogrammus*	state candidate species
Rockfish species	*Sebastes* spp.	state candidate species
Bald eagle	*Haliaeetus leucocephalus*	federal threatened species,[a] state threatened species
Peregrine falcon	*Falco peregrinus*	federal species of concern, state sensitive species[b]
Merlin	*Falco columbarius*	state candidate species
Common murre	*Uria aalge*	state candidate species
Common loon	*Gavia immer*	state sensitive species
Western grebe	*Aechmophorus occidentalis*	state candidate species

Source: WDFW (2007)

[a] Listing currently under review for removal.

[b] Downlisted from state endangered to state sensitive April 2002 (WDFW 2002)

Nine of these fifteen listed species are fish and six are birds. With the exception of chinook salmon, coho salmon, Puget Sound steelhead, bald eagle, western grebe, peregrine falcon, and Pacific herring, use of the LDW by these species is rare or incidental, and thus they are not likely to have frequent exposure to sediment-associated chemicals from the LDW. Reports of these rare or incidental species in the LDW include: loons (Canning et al. 1979, rare), merlin (Cordell et al. 1997, rare), common murre (believed to be rare), rockfish (Matsuda et al. 1968, rare; Malins et al. 1980, present; Windward 2005c, 2006b), river lamprey (Warner and Fritz 1995, rare; Matsuda et al. 1968, rare), walleye pollock (Matsuda et al. 1968, rare; Miller et al. 1975, rare), bull trout (Shannon 2006, rare; Warner and Fritz 1995, rare), and Pacific cod (Miller et al. 1975; 1977b; Weitkamp and Campbell 1980). Use of the LDW by chinook salmon, coho salmon, Puget Sound steelhead, bull trout, and herring is described in Section A.2.2.3. Use of the LDW by bald eagle, western grebe, and peregrine falcon is described in Section A.2.2.4.

The Puget Sound Southern Resident Orca whale distinct population segment was added to the federal endangered species list in February 2006 (50 CFR 224) and Washington State's endangered species list in 2004 (WDFW 2006). Orca whales do not use the LDW but are occasionally present in Elliott Bay and the Seattle area in the fall and early winter when chum salmon are running (Nelson 2006). Records from the Whale Museum's sightings database indicate that Orca whales may be seen in Elliott

Bay two or three times during the chum salmon season, although this may be an underestimate because it does not account for unreported sightings (Traxler 2006). Orca whales may be exposed to chemicals from the LDW through the consumption of prey that spend part of their time in the LDW.

A.2.2.2 Benthic invertebrate community

Benthic invertebrate communities (including species such as polychaetes and amphipods) are an important component of the LDW ecosystem because they serve as a major food resource for fish and wildlife, and because they are active in detrital processing and nutrient cycling.

In general, key physical factors that may influence the distribution and abundance of benthic invertebrates in the LDW are salinity, water depth (i.e., intertidal and subtidal), sediment grain size, and OC content. The LDW is a stratified salt-wedge estuary influenced by river flow and tidal effects (Section A.2.1.1). The daily salinity fluctuations select for species that are tolerant of such variability. A salinity range of approximately 5 to 8 parts per thousand (ppt) has been identified as a critical transition range that corresponds to a pronounced minimum of benthic invertebrate species richness (Levinton 1982). Benthic species richness, in general, diminishes steadily in an estuary until it reaches a minimum at the critical salinity (Levinton 1982). In general, a more diverse benthic invertebrate community exists in the downstream, more saline part of the waterway (RM 0.0 to RM 2.0) (Windward 2005f). Similarly, the benthic invertebrate community in the subtidal habitat is, in general, more diverse than the community in the intertidal habitat (Windward 2005f). The benthic invertebrate communities in the intertidal and subtidal habitats of the LDW are discussed in more detail in Section A.2.2.2.1, while larger epibenthic invertebrates are discussed in Section A.2.2.2.2. A table of benthic taxa identified in the LDW is presented in Attachment 1.

A.2.2.2.1 Epibenthos and infauna

Benthic invertebrates present in the LDW are characterized as either infaunal or epibenthic. The infaunal community is typified by burrowing polychaetes and bivalves. This community is dominated by deposit- and filter-feeding organisms. Many of the polychaetes, such as *Hobsonia florida* and *Capitella capitata* complex, are surface detrital- and deposit-feeding organisms. Bivalves obtain their food either from the water column (filter feeders) or the sediment surface (surface deposit feeders). Other infauna, such as oligochaetes, feed on bacteria, diatoms, and other microorganisms (King County 1999b). Small crustaceans (including copepods) feed on diatoms, detritus, and larvae. Epibenthic invertebrate communities include larger crustaceans, mussels, anemones, and echinoderms. Many of these invertebrates, such as echinoderms and anemones, are surface detrital- or filter-feeding organisms. The larger crustaceans are both carnivores and scavengers.

LDWG conducted a survey of the benthic communities (including infauna and smaller epibenthic species) at 12 intertidal locations throughout the LDW in 2004 (Map A.2-1) (Windward 2005f). Descriptions of the benthic communities were based primarily on samples collected with a 0.0024-m^2 core sampler to a depth of 10 cm. Five replicate core samples from each location were composited into one sample for community analysis. To augment the core data by enumerating larger benthic invertebrates from a larger area, samples were also collected within a 0.1-m^2 transect frame and sieved on a 2-mm sieve. Five replicate frame samples from each location were composited into one sample for these additional community analyses.

The numerical abundance of organisms in core samples ranged from 500 to 16,233/0.1-m^2 (Table A.2-3). A total of 61 invertebrate taxa were identified in the core and transect frame samples. The most abundant organisms at the 12 intertidal locations were annelids and crustaceans. The most abundant annelids were *Capitella capitata* complex, *Hobsonia florida, Manayunkia aestuarina*, and *Pygospio elegans*. The crustaceans were dominated by two amphipod species of the genus *Americorophium*. Entognathous hexapods (a subphylum of insects that includes springtails, *Collembola*) and mollusks, including *Macoma baltica,* were also identified at some locations. Echinoderms were not observed at any of the locations (Windward 2005f).

Swartz's dominance index was calculated for each intertidal location. Because replicate samples were not analyzed at each location, these indices should be viewed as qualitative indicators of community structure. Swartz's dominance index is defined as the minimum number of taxa that comprises 75% of the total abundance (Swartz et al. 1985; as cited in PTI 1993). As indicated by the index (Table A.2-3), the intertidal benthic invertebrate community consisted of relatively few species. The index ranged from 2 at location BCA-3 to 8 at location B2a. Numerous physical factors such as salinity, temperature fluctuations, desiccation, and wave action present physiological challenges to benthic invertebrates (Levinton 1982), and are therefore contributing factors to the relatively low number of dominant species in the intertidal areas. For example, salinity in the intertidal areas frequently reaches the critical range of 5 to 8 ppt from approximately Slip 1 (RM 1.0) to upstream sections of the LDW (King County 1999b). In these areas, the salinity is below 5 ppt more than 30% of the time (King County 1999a). Prior to the Phase 2 survey, Cordell et al. (2001) conducted epibenthic and infaunal surveys at seven restoration and reference sites throughout the LDW from 1993 through 1999. They found diversity and abundance of intertidal organisms varied seasonally and among locations in the LDW. The greatest diversity of organisms (i.e., species richness) occurred in the lower (higher salinity) LDW; diversity was comparatively lower in the Upper Turning Basin. Seasonally, species diversity and abundance increased from winter through summer as primary productivity increased. In spring, community composition was generally dominated by two to three species. By summer, the species composition was generally more evenly distributed among a greater number of species. At all sites sampled, the macrofauna were generally numerically dominated by oligochaetes, polychaete

Table A.2-3. Phase 2 intertidal benthic invertebrate community survey results

River Mile	Location ID[a]	Total Abundance[b]	Taxa Richness[c]	Swartz's Dominance Index	Taxa Richness by Major Taxonomic Group[d]					Abundance by Major Taxonomic Group[f]				
					Annelida	Crustacea	Insecta	Mollusca	Misc. Taxa[e]	Annelida	Crustacea	Insecta	Mollusca	Misc. Taxa[e]
0.2	B1a	4,842	13	3	4	6	1	0	2	3,950	667	8	0	217
0.6	BCA-1	11,667	22	6	7	9	2	1	3	6,083	5,417	17	8	142
0.8	B3a	11,958	25	4	9	13	0	2	1	9,175	2,767	0	17	0
0.9	B2a	16,233	30	8	8	12	2	3	5	7,725	8,183	25	42	258
1.4	B4a	8,858	23	4	6	13	2	0	2	6,933	1,833	25	0	67
2.1	B6a	1,475	19	5	8	5	0	2	4	1,217	100	0	17	142
2.3	B5a-2	500	10	3	6	2	0	0	2	475	17	0	0	8
2.9	BCA-3	9,050	16	2	8	6	0	0	2	7,658	1,267	0	0	125
3.1	B7a	8,600	20	4	6	11	1	0	2	5,983	2,592	8	0	17
3.5	B8a	14,100	21	6	7	11	0	0	3	7,383	6,583	0	0	133
4.5	B9a	5,875	24	5	8	13	1	0	2	3,992	1,867	8	0	8
4.8	B10a	6,600	20	6	7	10	0	0	3	3,292	2,767	0	0	542

[a] Sampling locations are shown on Map A.2-1.

[b] Total number of individual organisms in a standard 0.1-m^2 area determined by extrapolating the number in the composite sample representing a total area of 0.012 m^2 to the number expected in the larger area (0.1 m^2) by multiplying by 8.33.

[c] Total number of taxa in a composite of five core samples, representing a total area of 0.012 m^2, at each location.

[d] Total number of taxa in each major taxonomic group in a composite of five core samples, representing a total area of 0.012 m^2, at each location.

[e] Miscellaneous taxa include Nemertea, Nematoda, Cnidaria, and Platyhelminthes.

[f] Number of individual organisms in each major taxonomic group at each location in a standard 0.1-m^2 area determined by extrapolating what was enumerated in the composite sample representing a total area of 0.012 m^2 to the larger area (0.1 m^2) by multiplying by 8.33.

ID – identification

worms of the genus *Manayunkia*, and gammarid amphipods of the genus *Corophium*. The meiofauna, defined as organisms passing through a 0.5-mm sieve but retained on a 0.153-mm sieve, at all sites sampled were generally dominated by nematodes and harpacticoid copepods.

Williams (1990) and Leon (1980) also conducted benthic invertebrate surveys in the LDW (Map A.2-1). Williams (1990) identified 80 invertebrate taxa inhabiting intertidal habitats at Kellogg Island. Nematodes, oligochaetes, small harpacticoid copepods, ostracods, and sabellid polychaetes were the dominant invertebrates. Leon (1980) found 43 different benthic taxa in sediment cores from the intertidal mudflats at Kellogg Island. Most organisms occurred infrequently; nine taxa accounted for 97% of all individuals. Small marine worms of the genus *Manayunkia*, oligochaetes, and harpacticoid copepods made up nearly 80% of all individuals (Leon 1980). In comparison, there were very few organisms at a mudflat site with anoxic sediments near the Duwamish Shipyards, and there was a greater degree of seasonal variability in the benthic invertebrate community at a mudflat site in the marina near Kellogg Island (Leon 1980).

In 2003, LDWG conducted a clam survey at 11 intertidal locations between RM 0.0 and RM 4.0 (Map A.2-1) (Windward 2004b). A random sampling design, based on WDFW guidance (Campbell 1996), was employed to survey each of the 11 locations for clam abundance. The sediment was excavated to a depth of 30 cm and all clams were sorted from the substrate. The mean number of clams per 1 ft^2 ranged from 0.18 to 1.0 (Table A.2-4). The majority of clams collected were identified as *Macoma baltica* (60%), followed by *Mya arenaria* (20%) and *Macoma nasuta* (18%). Other less common *Macoma* species included *Macoma inquinata* and *Macoma secta* (both < 1%). The potential catch rates were also assessed at four areas with the highest abundances of clams found during the intertidal survey. *Mya arenaria* was the most common clam collected during the assessment of potential catch rates comprising more than 98% of the total biomass.

Table A.2-4. 2003 intertidal clam survey results

Beach	River Mile	Mean Number of Clams per 1-ft^2 Sample	Total Number of Individuals per Species					
			Macoma baltica	*Macoma inquinata*	*Macoma nasuta*	*Macoma secta*	*Macoma* spp.	*Mya arenaria*
1a	0.1	0.28	6	0	0	0	1	4
2a-island	0.5	1.0	75	0	12	1	1	14
2a-mainland	0.5	0.67	17	1	32	0	1	12
2b-mainland	0.5	0.70	2	0	4	0	0	1
2c-mainland	0.5	0.17	1	0	0	0	0	1
7	1.8	0.46	4	0	0	0	0	9
8	2.1	0.94	39	0	0	1	0	6
11	2.6	0.30	2	0	0	0	0	1

Beach	River Mile	Mean Number of Clams per 1-ft^2 Sample	Total Number of Individuals per Species					
			Macoma baltica	*Macoma inquinata*	*Macoma nasuta*	*Macoma secta*	*Macoma* spp.	*Mya arenaria*
1a	0.1	0.28	6	0	0	0	1	4
12	2.8	0.71	6	0	0	0	0	4
13a	2.9	0.47	2	0	3	0	0	3
16	3.5	0.18	19	0	0	0	1	2
All beaches	na	0.53	173	1	51	2	4	57

na – not applicable

In 2004, LDWG collected clams for chemical analysis at 10 intertidal locations throughout the LDW (Windward 2005b) (Map A.2-2). Two composite clam tissue samples were collected at 4 of these 10 locations. The collection method involved three field crew members actively searching and collecting clams from locations within the intertidal area with the highest clam abundance, as determined by evidence of siphon holes. The majority of clams collected for chemical analysis were *Mya arenaria,* with only a few *Macoma nasuta* collected at two of the locations (Table A.2-5).

Table A.2-5. Numbers of clams collected for chemical analysis in 2004

Beach[a]	Total Number of Individuals per Species	
	Mya arenaria	*Macoma nasuta*
C1-T	25	0
C2-T1	32	0
C2-T2	52	0
C3-T1	26	0
C3-T2	22	0
C4-T	22	0
C5-T	28	0
C6-T	22	0
C7-T1	17	3
C7-T2	22	0
C8-T	23	0
C9-T	22	0
C10-T1	19	2
C10-T2	17	1

[a] Replicate samples were collected at three beaches along two separate transects (T1 and T2).

C – clam sampling location

T – sampling transect

LDWG conducted a qualitative survey of the benthic communities (including infauna and smaller epibenthic species) at 14 subtidal locations throughout the LDW in 2004 (Map A.2-1) (Windward 2005f). Subtidal samples were collected with a 0.1-m^2 van

Veen grab sampler and organisms were retained using a 1.0-mm mesh sieve. Three replicate samples were composited into one sample at each location. The total number of organisms ranged from 72 to 2,300 per 0.1 m^2. A total of 246 invertebrate taxa were identified in the van Veen grab samples. In general, annelids, crustaceans, and mollusks were the most abundant organisms at the subtidal locations. The most abundant annelids were *Aphelochaeta* cf *glandaria, Capitella capitata* complex, *Hobsonia florida, Polydora cornuta*, and *Scoletoma luti*. The crustaceans were dominated by two amphipods of the genus *Americorophium,* especially at locations between RM 3.9 and RM 5.0. *Eogammarus confervicolus* and *Grandidierella japonica* were also common crustaceans. The most abundant mollusks were the bivalves *Axinopsida serricata, Parvilucina tenuisculpta*, and *Macoma* sp., and the most common gastropod was *Alvania compacta*. Echinoderms were also present at six locations with abundances ranging from 0.1 to 1.3% of all identified organisms.

As a way to summarize the subtidal benthic invertebrate community data, Swartz's dominance index was calculated for each subtidal location. Because replicate samples were not collected at each location, these indices should be viewed as qualitative indicators of community structure. The highest values of Swartz's dominance index were, in general, calculated for the more saline locations in the lower part of the waterway (RM 0 to RM 1.5), and the lowest value was calculated for a mostly brackish location in the Upper Turning Basin at RM 4.3. In the Upper Turning Basin, the salinity is less than 5 ppt 70 to 84% of the time (King County 1999a). This pattern is in accordance with general descriptions of benthic communities in estuaries, in which diversity diminishes steadily up an estuary along a salinity gradient (Levinton 1982). Table A.2-6 summarizes the results of the subtidal benthic invertebrate community survey.

Prior to the Phase 2 survey, Ecology (2000) evaluated the benthic invertebrate community at three subtidal locations in the LDW as part of the sediment quality reconnaissance study for central Puget Sound. The benthic invertebrate community at the three locations was dominated by annelids. Mollusks were also common, whereas crustaceans and echinoderms were present in low abundances.

King County (1999b) evaluated risks to benthic infauna and epibenthos as a component of their water quality assessment (WQA) of combined sewer overflow (CSO) discharges to the Duwamish River and Elliott Bay. Subtidal samples were collected with a 0.1-m^2 van Veen grab sampler and organisms were retained using a 1.0-mm mesh sieve. Sampling sites included transects located at Kellogg Island and downgradient from the Duwamish/Diagonal CSO (Map A.2-1). Polychaetes were abundant and were the dominant organisms at all subtidal locations, except at two stations downstream of the Duwamish/Diagonal CSO, where oligochaetes and mollusks were dominant. A Kellogg Island station also had relatively high abundance of mollusks. Arthropods were more abundant in deeper waters.

Table A.2-6. Phase 2 subtidal benthic invertebrate community survey results

River Mile	Location ID[a]	Total Abundance[b]	Taxa Richness[c]	Swartz's Dominance Index	Taxa Richness by Major Taxonomic Group[d]					Major Taxonomic Group Abundance[f]				
					Annelida	Crustacea	Echino-dermata	Mollusca	Misc. Taxa[e]	Annelida	Crustacea	Echino-dermata	Mollusca	Misc. Taxa[e]
0.1	B1b	326	107	15	43	22	4	28	10	176	77	3	66	3
0.6	BCA-4	937	107	13	62	12	2	25	6	499	60	<1	367	11
0.9	B2b	537	92	7	43	17	1	23	8	124	85	<1	323	4
1.0	B3b	559	93	12	54	14	2	22	1	397	42	<1	85	35
1.4	B4b	521	78	8	39	11	3	19	6	208	28	7	276	2
1.5	B5b	643	60	8	34	11	0	11	4	468	105	0	44	27
1.5	BCA-5	72	50	15	27	7 (1)[g]	0	12	3	52	6	0	13	2
1.7	BCA-2	328	54	6	29	12	1	11	1	282	17	<1	27	<1
2.2	B6b	1,137	83	3	41	17	0	21	4	1,041	17	0	77	2
2.7	B7b	497	75	5	38	16	1	18	2	391	11	<1	94	<1
3.9	B9b	935	36	6	17	12	0	6	1	435	493	0	8	<1
4.2	B8b	2,300	27	4	14	12	0	1	0	502	1,793	0	5	0
4.3	B10b	1,541	16	2	9	7	0	0	0	195	1,347	0	0	0
4.6	BCA-6	1,689	14	3	7	7	0	0	0	405	1,284	0	0	0

[a] Sampling locations are shown on Map A.2-1.

[b] Total number of individual organisms retained on a 1-mm sieve in a standard 0.1-m^2 area determined by extrapolating what was enumerated in three composite van Veen grab samples representing a total area of 0.3 m^2 to the smaller area (0.1 m^2) by dividing by 3.

[c] Total number of taxa in a composite of three van Veen grab samples, representing a total area of 0.3 m^2, at each location.

[d] Total number of taxa in each major taxonomic group in a composite of three van Veen grab samples, representing a total area of 0.3 m^2, at each location.

[e] Miscellaneous taxa include Nemertea, Nematoda, Cnidaria, and Platyhelminthes.

[f] Number of individual organisms in each major taxonomic group retained on a 1-mm sieve in a standard 0.1-m^2 area determined by extrapolating what was enumerated in three composite van Veen grab samples representing a total area of 0.3 m^2 to the smaller area (0.1 m^2) by dividing by 3.

[g] One insect specimen was collected at this location.

BCA – benthic community analysis (BCA locations were sampled only for the benthic invertebrate community analysis)

ID – identification

Williams (1990) sampled epibenthic sediment biota near Kellogg Island and found that nematodes, oligochaetes, small harpacticoids, and cumaceans dominated the subtidal epibenthos. As with the intertidal benthos, stations with finer sediments generally had a greater abundance of epibenthic biota.

Leon (1980) used van Veen grab samplers to characterize the epibenthic and infaunal sediment biota from subtidal locations near Kellogg Island. More than 60 different taxa were identified, which was greater than the number found in the intertidal habitat from the same survey. The most abundant taxon was deposit-feeding cirratulid polychaete worms. While some of these invertebrates were also found in intertidal habitats (oligochaetes, *Capitella* sp., *Pygospio* sp., ostracods), most subtidal species were deposit-feeding polychaete worms that are characteristic of the deeper, turbid waters of the LDW. Small deposit-feeding clams (*Macoma* sp., *Axinopsida* sp., and *Psephidia* sp.) and the amphipod *Anisogammarus* sp., which feeds on diatoms and green algae, were also present.

A.2.2.2.2 Larger epibenthic invertebrates

Larger epibenthic invertebrates identified in the LDW include crabs, shrimp, sea stars, anemones, and mussels. Mussels, anemones, and echinoderms are surface detrital- or filter-feeding organisms, whereas crabs and shrimps couple predaceous feeding with scavenging. Numerous larger epibenthic invertebrate species were caught during two fish and crab surveys conducted in late summer (August and September 2004 and 2005) throughout the LDW by LDWG (Windward 2005c, 2006b) (Maps A.2-1 and A.2-2; Table A.2-7). The invertebrates were collected in high-rise otter trawls, beach seines, crab traps, and shrimp traps. The most abundant epibenthic invertebrates were slender crabs, crangon shrimps, and coonstripe shrimps. Dungeness crabs were also caught in both surveys. The distribution of Dungeness crabs and other crabs was generally limited to the downstream portion of the LDW where the salinity is greater; however, a few adult Dungeness and slender crabs were caught between RM 4.2 and 4.5. A pilot survey and three quarterly surveys were performed in 2003 and 2004 to estimate the abundance of crabs and shrimps in the LDW (Windward 2004a). Three crab species and one shrimp species were caught in the surveys. Slender crabs and Dungeness crabs were the most abundant species. The majority of these crabs were caught in the downstream, more saline part of the LDW, with a few adults caught between RM 4.2 and 4.6. Red rock crabs and dock shrimps were less abundant and they were also primarily caught in the downstream portion of the LDW, with a few adults caught between RM 1.6 and 2.2. In October 1998, adult Dungeness and red rock crabs were collected at multiple locations near Kellogg Island but were not caught farther upstream (ESG 1999).

Table A.2-7. Numbers of individual invertebrate species caught using trawls and traps in the LDW during the 2004 and 2005 surveys

Common Name	Scientific Name	Number of Specimens Caught	
		2004	2005
Anemone, plumose	*Metridium senile*	nc	49
Anemone	unknown	81	nc
Ascidian[a]	unknown	1	nc
Crab, black-clawed	*Lophopanopeus bellus*	4	nc
Crab, decorator	*Loxorhynchus crispatus*	32	19
Crab, Dungeness	*Cancer magister*	62	33
Crab, hermit	*Pagurus* sp.	11	3
Crab, kelp	*Pugettia producta*	8	11
Crab, red rock	*Cancer productus*	16	19
Crab, slender	*Cancer gracilis*	942	483
Frilled dogwinkle	*Nucella lamellosa*	nc	2
Mussel, blue	*Mytilus edulis*	6	nc
Moon snail	*Polinices lewisii*	nc	5
Nudibranch, striped	*Armina californica*	39	118
Nudibranch	unknown	41	nc
Sea star, mottled	*Evasterias troschelii*	11	7
Sea star, sunflower	*Pycnopodia helianthoides*	23	18
Sea star, sand	*Luidia*	nc	2
Sea star	*Pisaster* sp.	50	27
Sea star	unknown	11	nc
Sea pen	unknown	38	8
Shrimp, coonstripe	*Pandalus danae*	314	231
Shrimp, crangon	*Crangon* sp.	538	172
Shrimp	unknown	8	nc
Solaster	*Solaster stimpsoni*	nc	1
Tunicate[a]	unknown	nc	2
Urchin	unknown	4	12

[a] Ascidian is a class of tunicates, and tunicate (or Urochordata) is the subphylum. However, because the two other classes of tunicates are pelagic, the two common names refer to the same group of invertebrates.

nc – not collected

During the 1989-1999 Puget Sound Ambient Monitoring Program (PSAMP), invertebrates were collected throughout the Puget Sound, including locations in the LDW (West et al. 2001). Epibenthic invertebrate species, similar to those caught in the LDWG surveys, were collected with otter trawls. The most common invertebrates were slender crabs and crangon shrimps. Other species caught during the PSAMP survey, but not in the LDWG surveys, included porcelain crabs (family Porcellanidae), chitons (class Polyplacophora), and several sea stars.

Commonly observed mollusks include epibenthic mussels which have been observed in large numbers on pilings and other structures in the downstream, more saline portion of the LDW with fewer mussels reported up to and slightly above the Upper Turning Basin (Windward 2000).

A.2.2.2.3 Summary

In summary, benthic invertebrates in the LDW consist of infauna and epibenthic organisms in both intertidal and subtidal habitats. Invertebrates surveyed in the LDW include more than 670 taxa, representing 178 families in 13 phyla (Attachment 1). Typical of most estuaries, the invertebrate community is dominated by annelids, mollusks, and crustaceans. Crustaceans are the most diverse of these three groups in the LDW, representing more than 250 taxa. These taxa included numerous macrofauna species from the orders Amphipoda, Isopoda, Cumacea, Tanaidacea, and Decapoda and numerous meiofauna species from the orders of Harpacticoida and Calanoida. The mollusks are represented by various bivalves and to a lesser extent by gastropods. The most abundant large epibenthic invertebrates include slender crabs, crangon, and coonstripe shrimps.

The taxonomic survey performed by LDWG, which was designed as a qualitative study of the benthic invertebrate community throughout the LDW, did not evaluate the distribution of benthic invertebrates in the LDW or their use of the LDW. Because of the qualitative nature of the study, no conclusions can be drawn about potential adverse effects of sediment chemicals on the benthic invertebrate community or about benthic invertebrate distribution among habitat types.

A.2.2.3 Fish

Diverse fish communities inhabit the LDW. Data are available from 14 studies conducted in the LDW investigating site usage by fish (Table A.2-8). Fifty-three resident and non-resident fish species were captured in the LDW during Phase 2 sampling (Windward 2004c, 2005c, 2006b). During historical sampling, Warner and Fritz (1995) recorded 33 resident and seasonal fish species, Miller et al. (1975; 1977b) observed a total of 29 species, and Matsuda et al. (1968) recorded a total of 28 species. In these studies, shiner surfperch, snake prickleback, Pacific sandlance, Pacific staghorn sculpin, longfin smelt, English sole, and starry flounder were particularly abundant, as were juvenile chinook, chum, and coho salmon. Fish numerical abundance reaches its maximum in late summer to early fall and is generally lowest in winter (Miller et al. 1977b; Dexter et al. 1981). Based on otter trawl data, species richness was shown to follow a similar trend but did not vary greatly with season (Miller et al. 1977b). Fish species reported to occur in LDW studies are listed in Attachment 2 with habitat, diet, and abundance information for each.

This section presents a summary of studies of LDW fish as well as a brief summary of the life history characteristics and dietary preferences of some fish species found in the LDW.

Table A.2-8. Summary of studies assessing the fish community in the LDW

Study	Year Completed	Location	Sampling Period	Equipment Type	No. of Locations Sampled
Phase 2 fish and crab tissue collection and chemical analyses (Windward 2005c, 2006b)	2005	four areas throughout the LDW	August and September, 2004 August and September, 2005	otter trawl, beach seine, shrimp traps, crab traps	24
Habitat utilization, migration timing, growth, and diet of juvenile chinook salmon in the Duwamish River and Estuary (Ruggerone et al. 2006)	2005	throughout LDW	February to July, 2005	beach seine	14
Fish assemblages and patterns of juvenile chinook salmon abundance, diet, and growth at restored sites in the Duwamish River (Cordell J et al. 2006)	2005	restoration and reference sites throughout LDW	February to July 2005	enclosure net	6
Phase 2 juvenile chinook salmon collection and chemical analyses (Windward 2004c)	2003	lower waterway (RM 0.1 to RM 0.9), and mid waterway (RM 1.4 to RM 2.9)	May (2 days) and June (3 days) 2003	beach seine	8
East Waterway channel deepening project, juvenile salmonid and epibenthic prey assessment (Shannon 2006)	2003	Kellogg Island and Harbor Island area	biweekly April – August 1998, 2000, 2003	beach seine	7 (2 in LDW)
East Waterway juvenile chinook salmon (Windward 2002)	2002	Kellogg Island	June 2002	beach seine	1
Waterway sediment operable unit, Harbor Island Superfund site. Assessing human health risks from ingestion of seafood (Robertson 2004)	1998	Harbor Island to south side of 1st Ave S. Bridge	single visit to each site	SCUBA	8 (6 in LDW)
PSAMP (West 2001)	1997	Kellogg Island	May 1992 – 1997	otter trawl	1
Distribution and growth of Green River chinook salmon and chum salmon outmigrants in the Duwamish estuary (Warner and Fritz 1995)	1994	Kellogg Island to above rapids	February – April (biweekly); April – May (weekly); May – September (biweekly) 1994	beach seine	9
Distribution and food habits of juvenile salmonids in the Duwamish Estuary (Meyer et al. 1981)	1980	Kellogg Island and at S Kenyon Street (RM 3.0)	April to June (weekly); July (biweekly) 1980	purse seine	2
				beach seine	2

Study	Year Completed	Location	Sampling Period	Equipment Type	No. of Locations Sampled
Port of Seattle Terminal 107 fisheries study (Weitkamp and Campbell 1980)	1978	Kellogg Island and adjacent channel	monthly October 1977 to February, July, and August 1978; more frequently from March to June 1978	purse seine	5
				beach seine	5
		South end of Kellogg Island	October 1977 – August 1978 (quarterly)	gill net (surface and bottom)	1
Chemical contaminants and biological abnormalities in central and southern Puget Sound (Malins et al. 1980)	1979	South end of Harbor Island	quarterly	7.5-m otter trawl	1 in LDW
Ecological survey of demersal fishes in the Duwamish River and at West Point or near Metro sewage treatment plants (Miller et al. 1975; 1977a; 1977b)	1974, 1975, and 1976	West Waterway to the Upper Turning Basin	1974 and 1975 (monthly) and January 1976	5-m otter trawl	8 (7 in LDW)
Fishes of the Green-Duwamish River (Matsuda et al. 1968)	1966	upper and lower LDW (exact locations unknown)	1964 – 1966 (weekly)	beach seine	2

Note: The majority of these studies used active capture techniques such as beach seining and otter trawls. These techniques preferentially capture less mobile species and are not effective for rough substrates or near structures. However, passive techniques employed in LDW sampling that included gill nets (Weitkamp and Campbell 1980), shrimp traps, and crab traps (Windward 2005c, 2006b) yielded no additional fish species beyond those observed using beach seines or otter trawls, indicating that the trawl data are generally reflective of the LDW fish community. Five of the 11 studies were conducted prior to 1986 when the Renton Wastewater Treatment Plant outfall was diverted from the Green River to Central Puget Sound. Because the diversion of the wastewater treatment plant effluent decreased summer flows by as much as 25% (~1.6 m3/s [56 cfs]), the diversity and abundance of fish in the LDW may have changed since these studies were conducted.

LDW – Lower Duwamish Waterway

PSAMP – Puget Sound Ambient Monitoring Program

RM – river mile

A.2.2.3.1 Pacific salmon

Five species of Pacific salmon (coho, chinook, chum, sockeye, and pink) occur in the LDW (Kerwin and Nelson 2000). These anadromous fish use the estuary for rearing and as a migration corridor for adults and juveniles. Among numerous beneficial uses identified for the LDW, habitat for outmigrating juvenile salmonids is one of the most important (Harper-Owes 1983).

The time spent by salmonids in the LDW is limited for all species. The amount of time spent each year depends on the specific life history of the species. Salmon found in the LDW spawn mainly in the middle reaches of the Green River and its tributaries (Grette and Salo 1986).

Adult salmon generally do not feed to any significant extent once they enter the estuary on their upstream spawning migrations. The peak timing of outmigration for juveniles of all species generally corresponds with March-to-June high flows. Outmigration usually lasts through mid-July to early August for most species (Warner and Fritz 1995; Nelson et al. 2004). During this time, juveniles use the estuary to feed and begin their physiological adaptation to higher salinity.

Chinook Salmon

Historically, the Green/Duwamish River supported spring and fall runs of chinook salmon. Fall-run chinook, a sub-population of the Puget Sound chinook population, are the only naturally sustaining run that still uses the Green/Duwamish River corridor. This run was among those listed as threatened under ESA in March 1999. These fish use the LDW for migration to and from spawning grounds in the mainstem Green River and larger tributary streams. Production is from hatcheries, naturally spawning hatchery-reared fish, and naturally spawning native fish (Grette and Salo 1986; WDFW 1993).

Returning fall chinook salmon enter the LDW from late June through mid November, with peak upstream migration in mid August (Grette and Salo 1986). In the mid-1970s, WDFW established an escapement goal of 5,800 naturally produced fall chinook using average escapement of natural origin fish and hatchery strays from 1965-1976 (Ames and Phinney 1977; as cited in Weitkamp and Ruggerone 2000). For the period from 1989-99, spawning escapements have been relatively high, averaging 8,578 fish, exceeding the WDFW goal for eight of the 10 years (WDFW unpublished data, as cited in King County 1999c). The contribution of Green River chinook salmon to the total chinook run entering Puget Sound and the Strait of Juan de Fuca ranged from 1.9 to 7.0% for the period 1979 to 1984 (Grette and Salo 1986).

In a 2003 tagging study of natural origin fry released approximately 30 miles upstream in February and March, fry were recovered at RM 5.0 to 6.0 of the LDW within one to 31 days after release (53% of the fry were found within 1 to 4 days) (Nelson et al. 2004). In 2002 and 2003, several year classes of outmigrant chinook (fry, yearlings, and

possibly 2-year-old fish) were found in the LDW between January and September (Warner and Fritz 1995; Nelson et al. 2004). Two peaks in abundance occurred: the first peak (of fry) was observed from late February to early March, and the second peak (of fingerlings) occurred between mid-May and mid-June. Peak catch data in 2002 and 2003 suggest that most naturally spawned fingerlings arrive in the estuary in May and reside in the LDW for approximately 2 weeks before departing to marine waters (Nelson et al. 2004). However, some fry may stay in the LDW from January until outmigration in June. Subyearlings have been consistently captured at RM 5.0 and 6.0,[5] suggesting that this is an important zone where juvenile chinook salmon transition between fresh and salt water (Ecology 2000; Nelson et al. 2004; Warner and Fritz 1995).

Ruggerone and Volk (2004) estimated residence time in the LDW prior to capture using incremental uptake of strontium in otoliths. Residence time of naturally spawned[6] chinook salmon collected from throughout the LDW averaged from a low of 16 ± 4 days in early July to a high of 58 ± 13 days in early September. The average residence time of individual hatchery chinook salmon increased from 16.6 days during late May through June (ranging from 6 to 25 days) to 45.6 days in mid-September.

Gut content analyses have shown that juvenile chinook in the LDW prey on a wide variety of benthic organisms such as *Corophium* spp. (amphipods) and *Cumella vulgaris*, drifting organisms such as adult dipterans, and zooplankton such as barnacle nauplius larvae (Cordell et al. 1997; 1999). Seasonal sampling suggests juvenile chinook shift their diet as different prey become available (Cordell et al. 1997; 1999).

Coho Salmon

Green River coho constituted from 0.9 to 1.4% of the total coho run entering Puget Sound and the Strait of Juan de Fuca for the period 1979-1984 (Grette and Salo 1986). Production was from hatcheries, naturally spawning hatchery-reared fish, and naturally spawning native fish (Grette and Salo 1986; WDFW 1993).

Adult coho return to the LDW between August and January, move through the LDW in a few days, and spawn and rear in all accessible reaches of the Green River drainage (Williams et al. 1975; as cited in Grette and Salo 1986). Juvenile coho rear in the Green River and move quickly through the LDW estuary as smolts (Warner and Fritz 1995; Weitkamp and Schadt 1982). The timing of outmigration is dependent on releases from Green River hatcheries (Warner and Fritz 1995; Weitkamp and Schadt 1982).

---

[5] Reported as RM 5.5 and 6.5 in original report; mapped locations were consistent with RM 5.0 and 6.0 as defined in this document (relative to the south end of Harbor Island).

[6] Fish that have not been fin-clipped and may be native fish or progeny of hatchery adults spawning in the wild.

Chum Salmon

The current status of the native chum population of the Green River watershed is unknown, but this population is suspected to have declined dramatically (Grette and Salo 1986; WDFW 1993). WDFW (1993) reported the state of the Green/Duwamish chum stock as unknown, whereas Nehlsen et al. (1991) reported the stock as at risk of extinction. Chum from the Green River constitute an insignificant portion of the total south Puget Sound run, and are not specifically addressed with harvest strategies for south Puget Sound stocks (Grette and Salo 1986).

Outmigrating chum salmon spend from several days to two months rearing in the Duwamish River estuary prior to moving offshore (Grette and Salo 1986). Warner and Fritz (1995) captured juvenile chum in beach seines throughout the LDW from February through September 1994 and showed a continuous increase in size of fish captured, suggesting a relatively extended residence time in the LDW. However, in the same study, catch rates declined rapidly following peak catches, suggesting that fish were likely moving through the estuary within a few days. Adult chum salmon return to the LDW between September and December.

Gut content analyses showed that in the LDW, juvenile chum preyed on both epibenthic species and drift insects during outmigration, with a large temporal variation in prey composition (Cordell et al. 1997; 1999).

Pink Salmon

Pink salmon appear infrequently and in low numbers. A run of odd-year pink salmon existed in the Green River in the 1930s (Williams et al. 1975; as cited in Grette and Salo 1986), though this run is believed to be currently extinct (Grette and Salo 1986). Warner and Fritz (1995) captured a total of 14 juvenile pink salmon in beach seines from nine stations throughout the LDW sampled approximately every two weeks from February through September 1994. Adult pink salmon have also been observed spawning in the mainstem Green River (Kerwin and Nelson 2000). Grette and Salo (1986) suggest that pink salmon have a high incidence of straying and that the few pink salmon captured in Green/Duwamish River are likely strays from other systems.

Sockeye Salmon

There is limited evidence that sockeye salmon spawn and rear in the Green River watershed (Jeanes and Hilgert 2000). Juvenile sockeye appear to have the shortest residence time in the nearshore of all salmon species (Kerwin and Nelson 2000).

A.2.2.3.2 Other salmonids

The Coastal-Puget Sound population of bull trout was proposed for listing under the federal ESA in June 1998 and was formally listed as threatened on November 1, 1999. The decline of bull trout has been primarily attributed to habitat degradation and fragmentation, blockage of migratory corridors, poor water quality, past fisheries management practices, and the introduction of non-native species (64FR 210: 58910-

58933). Bull trout were historically found in the LDW, but current stock status is unknown (WDFW 2000).

Muckleshoot tribal biologists captured one adult bull trout during beach seining in the LDW on May 24, 1994 during the period of peak juvenile salmon outmigration. However, it is unknown whether the fish reared in the Green River or was an opportunistic resident (Warner and Fritz 1995). Eight subadult bull trout ranging in length from 271 to 373 mm were captured in beach seines in the Upper Turning Basin during two sampling events in August and September 2000 (Shannon 2001). Peak numbers of juvenile shiner surfperch were captured at the same site the previous week, and near-peak numbers of shiner surfperch were captured in the same sampling in which the bull trout were caught (Shannon 2001). The co-occurrence of bull trout with high abundance of potential prey suggest that they may be opportunistically occupying the LDW to prey on these small fish. There is no evidence that bull trout are spawned or reared within the LDW. Bull trout juveniles typically remain in the upper tributaries for a period of two to three years prior to migrating to saltwater during spring. Adults typically return to their native streams in summer and fall (Grette and Salo 1986).

Summer steelhead (*Oncorhynchus mykiss*) is a non-native stock sustained by wild spawning of hatchery-reared fish (WDFW 1993). The run size is unknown, but approximated at a few hundred fish (WDFW 1993). The winter steelhead run consists of wild and hatchery fish with annual returns of 944 to 2,378 fish (WDFW 1993). Spawning of winter steelhead, which return to the Green River from December through May, generally begins about mid-March and continues to early June, with a peak in mid-May (Cropp 1985, as cited in Grette and Salo [1986]). Grette and Salo (1986) report that repeat spawners make up approximately 19% or less of returning wild adult steelhead in the Green River (1976/77 to 1983/84). Summer steelhead outmigrate from the Green River after rearing for two years as smolts, and do not have an extensive residence time in the LDW. Winter steelhead outmigrate from the Green River as subyearling adults and also do not rear extensively in the LDW.

Sea-run cutthroat trout may occur in the LDW, but little is known about this population. A total of 11 adult cutthroat trout were captured in beach seines at nine stations sampled approximately 30 times each throughout the LDW from February through June 1994 (Warner and Fritz 1995). In Washington, adult cutthroat return to their home stream from July to January, with the peak occurring in October and November (Wydoski and Whitney 2003). Smolt outmigration occurs from April through May (Wydoski and Whitney 2003).

A.2.2.3.3 Non-salmonid fishes

The most abundant non-salmonid fishes in the LDW are shiner surfperch, starry flounder, three spine stickleback, English sole, Pacific staghorn sculpin, Pacific herring, Pacific sandlance, surf smelt, and longfin smelt (Warner and Fritz 1995; Windward 2004c, 2005c, 2006b; Shannon 2006; West 2001).

Of these fish, shiner surfperch, longfin smelt, and Pacific herring are seasonally abundant in the LDW. Pacific herring, Pacific sandlance, surf smelt, and longfin smelt were encountered infrequently in recent beach seine and trawling attempts, but occasionally occurred in large numbers (Shannon 2006; Windward 2005c, 2006b). Three spine stickleback were abundant in monthly beach seine samples at both the Upper Turning Basin and Kellogg Island sampling locations June through September but were uncommon in February through May samples (Shannon 2006). Historical otter trawl data show peaks in longfin smelt abundance in summer, fall, and early winter (Miller et al. 1977b). Miller et al. (1977b) suggest that the fall-winter peak (80- to 115-mm fish) may represent part of a spawning run and that the late summer peak (30- to 50-mm fish) may represent downstream migrant young of the year. Pacific herring were reported in purse seine samples throughout the year (Weitkamp and Campbell 1980), were present in trawl samples in August and September (Windward 2005c, 2006b), and were reported in beach seine samples in May, June, July, November, and December (Weitkamp and Campbell 1980; Shannon 2006). In Puget Sound, threespine stickleback and surf smelt feed on both epibenthic and pelagic invertebrates. Epibenthic invertebrates constitute a slight majority of their diet (Miller et al. 1977c; Fresh et al. 1979). Pacific herring and longfin smelt generally feed on pelagic invertebrates but also ingest epibenthic invertebrates to a lesser extent (Miller et al. 1977c; Fresh et al. 1979).

In all studies, Pacific staghorn sculpin was consistently one of the most abundant fish captured in the LDW (Attachment 2). Taylor and Associates (Shannon 2006) captured Pacific staghorn sculpin during all months from February through October. Their abundance was highest from May through July. Miller et al. (1977b) reported that Pacific staghorn sculpin were abundant in otter trawls during all seasons, but were particularly abundant in the fall. Few Pacific staghorn sculpin larger than 150 mm were collected in beach seines by Weitcamp and Campbell (1980). During otter trawl sampling conducted by LDWG in August and September 2004 and 2005, juvenile (< 120 mm) and adult Pacific staghorn sculpin were commonly encountered throughout the LDW (Windward 2005c, 2006b).

Pacific staghorn sculpin are opportunistic feeders that feed at a higher trophic level than other common resident fish species. As discussed in a meeting of local fish experts (Windward 2004j), other upper-trophic-level fish that occur in the LDW, such as brown rockfish and sand sole, occur primarily as juveniles that feed at a similar trophic level as Pacific staghorn sculpin (e.g., brown rockfish) or have foraging ranges that likely extend well beyond the LDW (e.g., sand sole). Pacific staghorn sculpin feed mostly on crabs, shrimps, and benthic invertebrates, but also ingest larval, juvenile, and adult fish (Fresh et al. 1979; Wingert et al. 1979; Miller et al. 1977c). Larger sculpin are more likely to eat at a higher trophic level than smaller sculpin.

Shiner surfperch abundance peaks in summer during the bearing of young (Miller et al. 1975). Taylor and Associates recorded abundant shiner surfperch May through October with peak abundance in July (Shannon 2006). Shiner surfperch are

opportunistic omnivores, feeding on zooplankton, small crustaceans, algae, and detritus (Gordon 1965; Bane and Robinson 1970), as well as polychaetes, mollusks, and benthic organisms (Fresh et al. 1979; Wingert et al. 1979; Miller et al. 1977c).

English sole are common in the LDW over all seasons, with peak abundance in spring (Miller et al. 1977b). English sole were abundant in recent trawl samples but were absent from beach seine samples (Windward 2006b; Shannon 2006; West 2001; Windward 2005c). In Puget Sound, English sole are typically found on soft sand or mud bottoms at depths of 25 to 50 m (Smith 1936). Juvenile English sole (those less than 110 mm) ingest annelids (Smith 1936), copepods, amphipods, and mollusks (Holland 1954). Adult English sole studied in Puget Sound ingest clams, clam siphons, small mollusks, marine worms, small crabs, and small shrimps (Wingert et al. 1979; Fresh et al. 1979). It has been suggested that English sole exist in discrete populations with some site fidelity (Day 1976). Day (1976) conducted a tagging study in Puget Sound that suggested that fish captured and released at the same location remained within an area approximately equal to 5 to 10 km^2. In addition, catch rates for fish captured and released dozens of miles from their original capture site were higher at their original capture site than at the release site or other sites sampled.

English sole migrate seasonally to their spawning grounds in Puget Sound in winter (Forrester 1969) and typically spawn in Puget Sound during February and March (Smith 1936). In central Puget Sound, adult populations of English sole spawn in Elliott Bay and Port Gardner but disperse after spawning (Pallson 2001). Angell et al. (1975; as cited in King County 1999c) reported off-shore migration in winter and spring of all age groups of central Puget Sound English sole from Meadow Point to Carkeek Park (northwest Seattle) at depths of 3 to 30 m. Juveniles (10 to 25 mm standard length), not all completely metamorphosed, migrated from spawning areas to nursery grounds as pelagic fish and moved to benthic habitats in December or May and June (King County 1999b). Data from Malins et al. (1982) show that during the winter and spring, greater than 50% of the English sole in the LDW are juveniles (< 150 mm).

Starry flounder are also noted to migrate seasonally between very shallow water and in estuaries during the summer, moving into deeper water in the winter (Morrow 1980). Young and adult starry flounder are tolerant of fresh water and move up rivers as much as 120 km (Morrow 1980). Because they have a larger mouth, starry flounder are capable of consuming somewhat larger organisms than English sole ingest, although their diets greatly overlap. Starry flounder in Puget Sound were found to ingest primarily benthic invertebrates, with bivalves, amphipods, and shrimp serving as important prey items (Fresh et al. 1979).

A.2.2.3.4 Summary

In summary, the LDW provides habitat for many migrating and resident fish. Anadromous salmon are present during rearing and migration to and from spawning sites in the upper watershed. In the LDW, juvenile salmonids are an important part of

the food web, preying on various epibenthic, water column, and drift organisms and serving as prey for larger fish and wildlife. Of the non-salmonid fish, shiner surfperch, English sole, Pacific staghorn sculpin, snake prickleback, longfin smelt, and starry flounder are among the most abundant species in the LDW. Seasonal abundance of fish in the LDW varies, peaking in the summer and early fall. Fish in the LDW are primarily carnivorous and are likely to rely extensively on epibenthic invertebrates as prey.

A.2.2.4 Wildlife

The intertidal and subtidal habitats of the LDW support a diversity of wildlife species. Formal studies, field observations, and anecdotal reports indicate that up to 87 species of birds and 6 species of mammals utilize the LDW during at least part of the year to feed, rest, or reproduce. This section provides an overview of these bird and mammal species.

A.2.2.4.1 Birds

The bird species associated with the LDW are presented in Table A.2-9. The birds using the site can be grouped as follows:

- Passerine/upland species
- Raptors
- Shorebirds/waders
- Waterfowl
- Seabirds

Table A.2-9. Bird species using the LDW

COMMON NAME	SCIENTIFIC NAME	COMMON NAME	SCIENTIFIC NAME
Passerine/Upland Species			
Blackbird, red-winged	*Agelaius phoeniceus*	Sparrow, English (house)	*Passer domesticus*
Bushtit, common	*Psaltriparus minimus*	Sparrow, fox	*Passerella iliaca*
Chickadee, black-capped	*Poecile atricapillus*	Sparrow, golden-crowned	*Zonotrichia atricapilla*
Cowbird, brown-headed	*Molothrus ater*	Sparrow, savannah	*Passerculus sandwichensis*
Crow, northwestern	*Corvus corrinus*	Sparrow, song	*Melospiza melodia*
Dove, rock	*Columba livia*	Sparrow, white-crowned	*Zonotrichia leucophrys*
Finch, house	*Carpodacus mexicanus*	Starling, European	*Sturnus vulgaris*
Flicker, northern	*Colaptes auratus*	Swallow, barn	*Hirundo rustica*
Goldfinch, American	*Spinus tristis*	Swallow, cliff	*Petrochelidon pyrronota*
Hummingbird, Anna's	*Calypte anna*	Swallow, tree	*Iridoprocne bicolor*
Junco, dark-eyed	*Junco hyemalis.*	Swallow, violet-green	*Tachycineta thalassina*
Kingfisher, belted	*Ceryle alcyon*	Thrush, Swainson's	*Hylocichla ustulata*
Kinglet, ruby-crowned	*Regulus calendula*	Towhee, rufous-sided	*Pipilo erythrophthlamus*
Purple martin	*Progne subis*	Warbler, orange-crowned	*Vermivora celata*

Common Name	Scientific Name	Common Name	Scientific Name
Quail, California	*Lophortyx californicus*	Wren, Bewick's	*Thryomanes bewickii*
Robin, American	*Turdus migratorius*	Wren, house	*Troglodytes aedon*
Siskin, pine	*Carduelis pinus*		
Raptors			
Eagle, bald	*Haliaeetus leucocephalus*	Hawk, sharp-shinned	*Accipiter striatus*
Falcon, peregrine	*Falco peregrinus*	Hawk, Swainson's	*Buteo swainsoni*
Hawk, Cooper's	*Accipter cooperii*	Merlin	*Falco columbarius*
Hawk, red-tailed	*Buteo jamaicensis*	Osprey	*Pandion haliaetus*
Shorebirds/Waders			
Dowitcher	*Limnodromus sp.*	Sanderling	*Crocethia alba*
Dunlin	*Erolia alpina*	Sandpiper, least	*Calidris minutilla*
Heron, great blue	*Ardea herodias*	Sandpiper, spotted	*Actitis macularia*
Heron, green	*Butorides virescens*	Sandpiper, western	*Calidris mauri*
Killdeer	*Charadrius vociferus*	Yellowlegs, lesser	*Totanus flavipes*
Waterfowl			
Bufflehead	*Bucephala albeola*	Goose, domestic	*Branta domesticus*
Canvasback	*Aythya valisineria*	Mallard	*Anas platyrhynchos*
Coot, American	*Fulica americana*	Merganser, common	*Mergus merganser*
Duck, domestic	*Anas domesticus*	Merganser, hooded	*Lophodytes cucullatus*
Gadwall	*Anas strepera*	Merganser, red-breasted	*Mergus serrator*
Goldeneye, Barrow's	*Bucephala islandica*	Scoter, surf	*Melanitta perspicillata*
Goldeneye, common	*Bucephala clangula*	Teal, greenwinged	*Anas carolinensis*
Goose, cackling Canada	*Branta canadensis minima*	Wigeon, American	*Mareca americana*
Goose, Aleutian	*Branta canadensis*		
Seabirds			
Cormorant, double-crested	*Phalacrocorax auritus*	Gull, glaucous-winged	*Larus glaucescens*
Cormorant, pelagic	*Phalacrocorax pelagicus*	Gull, mew	*Larus canus*
Grebe, eared	*Podiceps capsicus*	Gull, ring-billed	*Larus delawarensis*
Grebe, horned	*Podiceps auritus*	Loon, common	Gavia immer
Grebe, pied-billed	*Podilymbus podiceps*	Loon, Pacific	Gavia Pacifica
Grebe, red-necked	*Podiceps grisegena*	Loon, red-throated	Gavia stellata
Grebe, western	*Aechmophorus occidentalis*	Murre, common	*Uria aalge*
Guillemot, pigeon	*Cepphus columba*	Tern, Caspian	*Hydroprogne caspia*

Canning et al. (1979) conducted extensive surveys of the avifauna of Kellogg Island, as well as occasional surveys of the entire LDW from September 1977 to July 1978. They recorded a total of 70 species: 26 passerines/upland birds, 3 raptors, 11 shorebirds/ waders, 17 waterfowl, and 13 seabirds. Kellogg Island had a much higher diversity of birds than the rest of the LDW because of its seclusion and greater variety of habitats.

Cordell et al. (1999) monitored bird populations monthly from 1995 to 1997 at four sites: two sites in the Upper Turning Basin, one on Kellogg Island, and one at Terminal 105. They recorded 75 species of birds: 32 passerine/upland birds, 7 raptors, 8 shorebirds/waders, 16 waterfowl, and 12 seabirds. Diversity and abundance were highest at the Kellogg Island site, but other areas of the LDW were also consistently

used by a wide variety of birds. Birds were most abundant in the spring and least abundant in the summer.

LDWG conducted a 4-day survey in June 2004 to identify the presence and quality of spotted sandpiper habitat along the LDW (Windward 2004h), as discussed below in Section A.2.2.4.4. This survey noted the presence of both spotted sandpiper and killdeer in specific locations along the LDW shoreline. Great blue herons, ospreys, and bald eagles were also observed along the LDW during this survey.

The following sections provide a brief summary of site usage by the various types of bird species in the LDW.

A.2.2.4.2 Passerines/upland birds

Thirty-two species of passerine/upland birds have been documented along the LDW (Canning et al. 1979; Cordell et al. 1999). Though generally associated with upland habitats, these birds occasionally forage in the exposed mudflats or use freshwater habitats along the river for bathing (Canning et al. 1979). Because they primarily use upland habitat, passerine birds likely experience very limited exposure to sediments in the LDW.

A.2.2.4.3 Raptors

Eight raptor species have been reported to use the LDW (Cordell et al. 1999). The bald eagle is listed under ESA as a threatened species but is currently under review for delisting. In Washington, it is also listed as a state-threatened species (WDFW 2006). Five nests were occupied within 5 miles of the LDW in 1999 (King County 1999c). The closest known nest is located in West Seattle within 1.6 km (1 mi) of the LDW. One or two pairs of resident eagles may be found in the LDW vicinity during the summer (King County 1999b). Overwintering migrant eagles are routinely observed in the vicinity of the LDW from the beginning of October through late March.

The bald eagle is an opportunistic forager with site-specific food habits based on available prey species (Buehler 2000). Bald eagles ingest dead and live fish, birds, and mammals. In most regions, bald eagles seek out aquatic habitats for foraging and prefer to ingest fish (Buehler 2000). Spawned-out salmon are a particularly important food item for eagles in the Pacific Northwest, though not in the LDW because salmon spawn farther upstream. Of the 45 individual fish identified in a study of prey remains at the base of eagle nest trees throughout Puget Sound, 8 were rockfish, 10 were starry flounder, and the remainder included cod, pollock, hake, cabezon, red Irish lord, sculpin, surfperch, salmon, plainfin midshipman, and channel catfish (Knight et al. 1990). Although eagles feed primarily on fish, birds, such as grebes, gulls, and waterfowl, make up a portion of their diet during winter months. Eagles have been reported to kill western grebe in the Duwamish River during winter (Strand 1999; as cited in King County 1999a). Eagles also have been reported to prey on great blue heron chicks (Norman 1999; as cited in King County 1999c).

Cooper's and sharp-shinned hawks have been observed to overwinter in the LDW. These relatively small raptors generally feed on birds up to the size of quail. They rarely feed on aquatic birds (Canning et al. 1979; Cordell et al. 1999). Red-tailed hawks, a resident species commonly observed along grassland/woodland margins along the LDW, feed primarily on rodents but have been observed pursuing ducklings in the study area. Swainson's hawks and merlin are rare in the LDW and not likely to prey on aquatic associated species (Canning et al. 1979; Cordell et al. 1999).

The US Geological Survey (USGS) observed five osprey nests along the LDW between RM 0.0 and RM 5.0 in 2003, as well as one nest on Harbor Island and three nests along the Duwamish and Green Rivers within about 4 miles of the Upper Turning Basin. Ospreys feed opportunistically and almost exclusively on live fish from fresh or salt water. Ospreys can penetrate about 1 m below the water surface. Therefore, they generally catch pelagic fish or those that frequent shallow flats and shorelines. A west-central Idaho osprey study reported 89% of fish ingested by osprey were 11 to 30 cm long, suggesting a preference for medium-sized fish (Van Daele and Van Daele 1982). A USGS study conducted in the LDW noted that approximately 73% of the prey delivered by osprey to their nests along the LDW were LDW fish species; salmon, perch, and sole/flounder accounted for approximately 33, 25, and 15% of total prey remains, respectively (Henny 2005). Twenty-five percent of the total prey remains were peamouth, a freshwater fish, and two percent were not reported. This dietary information was collected during only a portion of the nesting season. Additional dietary data have been collected and are currently being analyzed, but those data are not yet available (Kaiser 2006).

There are currently two known nesting pairs of peregrine falcons along the LDW; one in a nest box on the West Seattle Bridge, and one in a natural nest on the First Avenue South Bridge (Anderson 2006). Peregrine falcons in western Washington feed primarily on rock pigeons and European starlings, although they may also ingest some waterfowl (Anderson 2006).The peregrine falcon is listed as a federal species of concern and was downlisted from a state endangered species to a state sensitive species in April 2002 (WDFW 2006).

A.2.2.4.4 Shorebirds/waders

Eight species of shorebirds and wading birds have been documented in the LDW (Cordell et al. 1999), including green heron and great blue heron. Of these species, great blue heron make up the only sizeable or consistent population.

The great blue heron is a wading bird that has a range from the coasts of southeast Alaska and Northern British Columbia, through Canada and the US, and south to Belize and Guatemala. The great blue heron is found primarily in natural wetlands and along riverbanks, but can also be found in brackish marshes, lagoons, lakes, and along ocean shores. They were the most abundant shore/wading bird recorded by Cordell et al. (1996) in the LDW, and are a year-round resident. Great blue herons nest in colonies of up to several hundred pairs, preferably on islands or wooded swamps

(Butler 1992). A heron colony of up to 37 active nests was located in West Seattle a few hundred meters from Kellogg Island until 1999, but no successful nesting has occurred there since 2000 (Norman 2006). This colony was likely abandoned because of either eagle predation or disturbance from construction at the nearby Herring's House Park (Norman 2006). Great blue herons are sensitive to interferences during their nesting period, and disturbance by eagles and humans is a common reason for nesting failures (Heron Working Group 2006a). Other heron colonies in the vicinity of the LDW are located about 12 km (7.5 mi) south in Renton and 11 km (6.8 mi) northwest near Salmon Bay.

Great blue herons feed in shallow water primarily on small fish, such as juvenile salmonids, but they also take crustaceans, insects, amphibians, reptiles, and occasionally small mammals (Butler 1992; Kushlan 1978). Great blue herons hunt by sight and stalk or ambush their prey. They also feed by probing, quickly moving their bills in and out of the water and substrate. Great blue herons feed on small fish that range in size from 8 to 33 cm (Kirkpatrick 1940; Alexander 1977; Hoffman 1978). Butler (1992) reports that shiner surfperch is a major food source for female and hatchling great blue herons and may be important for juvenile survival.

The two most common shorebirds observed in the LDW are sandpipers and killdeer. The spotted, least, and western sandpipers are commonly observed in the LDW. These birds feed primarily on insects, small crustaceans and mollusks, worms, and other invertebrates, and rarely on seeds and berries. Least and western sandpipers occur in mixed flocks and are difficult to distinguish. These species nest primarily in northern Canada and Alaska in the summer months (Paulson 1993), but are reported to frequent Kellogg Island from September through May (Canning et al. 1979). Most are thought to be migrants, though some may reside in the LDW throughout the winter.

Spotted sandpipers are a common bird in western Washington, and are known to nest along the LDW. They have been observed in the LDW from late June through September (Cordell et al. 1996), but have also been known to overwinter locally (Paulson 1993). Spotted sandpipers breed in open habitats along the margins of water bodies (Oring and Lank 1986) and mostly forage in open areas on the ground within 200 m of the shoreline (Oring et al. 1997). Canning et al. (1979) recorded seven spotted sandpiper nests located on Kellogg Island, and at least three additional nest sites were suspected based on the behavior of adult or juvenile birds. LDWG conducted a survey in June 2004 to identify the presence and quality of spotted sandpiper habitat along the LDW (Windward 2004h). The results of this survey indicate that sandpipers may be nesting along Kellogg Island, at the Herring's House restored marsh area (western shoreline RM 0.5), near the channel beneath the 1st Avenue South Bridge (western shoreline RM 2.1), and along the western shoreline of the Upper Turning Basin (RM 4.6 to RM 4.9).

Killdeer, another common shorebird, feed in intertidal mudflats. Their diet includes small invertebrates, insects, and some vegetative matter. Killdeer are a common bird

that uses the LDW year-round. In a year-long survey conducted in 1977-78, 20 to 60 birds were reported to use the Kellogg Island area in the winter and approximately 10 birds in the fall and spring (Canning et al. 1979).[7] Two killdeer nests were observed on Kellogg Island (Canning et al. 1979). During the spotted sandpiper survey conducted in June 2004, killdeer were most frequently observed in the vicinity of Kellogg Island from RM 0.0 to RM 1.0, but were also observed at RM 2.1 to 2.2, RM 4.0 to 4.2, and RM 4.7 to 4.9 (the Upper Turning Basin) (Windward 2004h). Killdeer were confirmed to nest on Kellogg Island based on an observation of a killdeer leaving its nest during the sandpiper survey (Windward 2004h).

A.2.2.4.5 Waterfowl

Cordell et al. (1999) reported 16 species of waterfowl utilizing the LDW, including nine species of dabbling ducks. All species are generally migratory, though some non-migratory populations exist. In general, these birds overwinter in the Puget Sound area (and farther south) and migrate north in the summer. The dabbling ducks feed on aquatic plants, seeds, and grasses and to some extent small aquatic animals and insects. Feeding occurs primarily in shallow water and over intertidal mudflats. A resident population of approximately 25 mallards lives year-round in the LDW, and an additional population of approximately 15 mallards overwinters in the LDW. As many as 290 migratory mallards have been reported to move through the LDW (Canning et al. 1979). The other dabbling duck species use the LDW seasonally for nesting and forage during migration. The most significant of these are gadwalls. Approximately 10 gadwall nests were observed along the LDW in the vicinity of Kellogg Island during a survey conducted in 1977-78 (Canning et al. 1979).

Canvasback, greater scaup, bufflehead, and common and Barrow's goldeneye are reported to use the LDW. These birds dive for small aquatic animals and plants. Canvasback feed primarily on plants, scaup on equal portions of plants and animals, and bufflehead and goldeneyes exclusively on aquatic animals and insects. During a survey conducted in 1977-78, approximately 60 canvasbacks arrived in the LDW in November and departed in late February, using Kellogg Island as their primary feeding area (Canning et al. 1979). During the same survey, greater scaup and common and Barrow's goldeneyes arrived in the LDW in late November and departed by early May, and a small population of approximately eight buffleheads overwintered in the LDW from December to May (Canning et al. 1979). In general, Canning et al. (1979) found that feeding by all diving duck species occurred primarily in the vicinity of Kellogg Island in 1977-78.

All three species of North American mergansers have been recorded to use the LDW, two substantively. Common mergansers were reported to use the LDW from September to March in 1977-78; these birds did not winter on the LDW but were likely

[7] Killdeer were present in the summer, but the number observed was not reported.

migrating through the area (Canning et al. 1979). Approximately 30 red-breasted mergansers were reported to overwinter in the LDW from December 1977 to March 1978 (Canning et al. 1979). These birds feed primarily on small fish and are reported to feed in the deeper water of the channel (Canning et al. 1979).

A resident population of approximately 1,000 Canada geese resides in the vicinity of Lake Washington. The geese foraging in the LDW are thought to be a part of the Lake Washington population. Migratory Canada geese arrive in the LDW in January and February and remain until the end of July as a spring nesting population. Canada geese swim in the LDW and feed in intertidal habitats. They feed primarily on grass and terrestrial vegetation (Canning et al. 1979). In the LDW, 40 to 50 birds overwinter from September to April along Kellogg Island and the west bank of the waterway along the South Park district and in the Upper Turning Basin (Canning et al. 1979).

A.2.2.4.6 Seabirds

Thirteen species of seabird have been recorded in the LDW (Canning et al. 1979; Cordell et al. 1999), including two species of cormorants (pelagic and double-crested). Cormorants feed primarily on small fish and occasionally crustaceans. Wintering cormorants use the LDW from November to May, with large numbers present from December to April (Canning et al. 1979; Cordell et al. 1996).

Several species of gulls are reported to use the LDW. Gulls feed on fish and shellfish and are omnivorous scavengers. Glaucous-winged gulls and mew gulls are the only species reported to use the area in large numbers. Glaucous-winged gulls are reported to use the area throughout the year. Mew gulls frequent the area, occasionally in large numbers, from September through May (Canning et al. 1979).

Caspian terns have been observed near Kellogg Island (Luxon 2004). Pigeon guillemots and common murres have been reported in the LDW; however, their use of the LDW is infrequent. These birds feed primarily on pelagic fish, though bottomfish and crustaceans may also be taken.

Common loons are listed as a state sensitive species (WDFW 2006). They are present in Puget Sound in the winter and use local waters for resting during migrations to and from wintering areas farther south. Their diet consists primarily of small fish and other aquatic animals. Ten to 30 birds have been observed in the Seattle area during annual winter counts, although they are reported to be rare visitors to the LDW (Canning et al. 1979).

Three species of grebe are reported in the LDW. Of these, only western grebes are found in substantial numbers. Grebes and other marine bird species have been declining in recent years (Nysewander et al. 2001). Feeding behavior varies with species. In marine waters, the eared grebe primarily ingests crustaceans while the western grebe favors fish. The most common fish species ingested by western grebes are Pacific herring, pilchard, stickleback, sculpin, sea perch, and smelt. Western grebes occasionally feed on juvenile salmonids. The LDW population was estimated to

include about 90 birds in the 1970s (Canning et al. 1979). Grebes arrive in the LDW from October to November and depart by early May.

In summary, the LDW is a corridor frequented by a diverse avian group. It is utilized mostly by shore birds, waders, seabirds, and waterfowl, which feed in areas of mudflats and other shallow-water habitats. Raptors also use the LDW for foraging.

A.2.2.4.7 Mammals

Three marine mammal species may occasionally enter the LDW: harbor seal, California sea lion, and harbor porpoise (Dexter et al. 1981). Harbor seals and California sea lions have been documented in the LDW (WDFW 1999), but recent information on harbor porpoise usage was not available. The harbor seal can be found along both North American coasts (Hoover 1988; Payne and Selzer 1989). Along the Pacific coast they can be found from Alaska to Baja California and mainland Mexico, and are the most commonly observed pinniped species (Hoover 1988). They can be seen in protected harbors year-round (Boulva and McLaren 1979). Harbor seals are commonly seen in Elliott Bay and occasionally enter the LDW (Kenney 1982).

Harbor seals are opportunistic feeders, selecting prey based on availability and ease of capture (Pitcher 1980; Pitcher and Calkins 1979; Schaffer 1989). Their diet can vary seasonally and includes bottom-dwelling fishes, invertebrates, and species that congregate for spawning (Pitcher and Calkins 1979; Everitt et al. 1981; Lowry and Frost 1981; Roffe and Mate 1984). In Washington, the most important prey include Pacific whiting, tomcod, walleye pollock, flatfishes, Pacific herring, shiner surfperch, plainfin midshipman, and sculpins (NMFS 1997). Fish ingested are generally between 40 and 280 mm (Brown and Mate 1983). Harbor seals have been shown to forage over large areas ranging from 5 km (Stewart et al. 1989) to 55 km (Beach et al. 1985).

California sea lions and harbor porpoises are also opportunistic feeders, consuming various fish species depending on availability (Marine Mammal Center 2006). California sea lions and harbor porpoises will, like harbor seals, also feed on non-fish species such as squid and octopus (Yates 1998).

A survey of sea lions and harbor seals was conducted in the LDW from December 1998 to June 1999 (WDFW 1999). This survey monitored the presence of sea lions and harbor seals in the East and West Waterways and in the LDW up to the 16th Avenue South Bridge for a total of 307 hours on 52 days. In the LDW, sea lions were observed on 16 occasions and seals on 17 occasions, with most observations for both species occurring below the 1st Avenue South Bridge. In the East and West Waterways, sea lions were observed 69 times and seals 6 times; both species used the West Waterway most frequently.

Three species of semi-aquatic terrestrial mammals use the LDW: river otters, raccoons, and muskrats. Anecdotal information indicates that a river otter family lives year-round on Kellogg Island in the LDW, although otters were not observed by Cordell during wildlife surveys (Cordell 2001). River otters are almost exclusively aquatic and

prefer food-rich habitats such as the lower portions of streams and rivers, estuaries, and lakes and tributaries that feed rivers (Tabor and Wight 1977; Mowbray et al. 1979). Local river otters feed primarily on fish but will also feed on crabs and sometimes mussels and clams (Strand 1999; as cited in King County 1999a). River otters range over an area sufficiently large enough for foraging and reproduction (Melquist and Dronkert 1987); however, they are typically found in a limited number of activity centers within their overall range. In streams, the river otter's home range can average 30 km (Melquist and Hornocker 1983).

Raccoons are reported to be common along the forested ridge slopes to the west of the LDW. Raccoons are scavengers that feed on carrion and occasionally on fish. Muskrat populations are reported to exist at Terminal 107 and at the Upper Turning Basin (Canning et al. 1979). Muskrats are herbivores, feeding primarily on aquatic and semi-aquatic plants.

In summary, the LDW corridor provides habitat for a limited number of mammal species. It may serve as a significant part of the home range of a river otter family, but is used only occasionally as a foraging site by marine mammals.

A.2.3 RECEPTOR OF CONCERN SELECTION

This section presents the ROCs selected to represent benthic invertebrate, fish, and wildlife species based on a set of ROC selection criteria. Inherent to the ROC selection process is the realization that not all species in the LDW can be evaluated individually because of the large number and variety of species present. Instead, representative species are chosen to include species that are most likely to be exposed to contaminated sediment. In this way, species not selected should also be protected. Sensitive species are also preferred, although the relative sensitivity of most species is not known.

A systematic process was followed to select representative species as ROCs based on the available information for the resources presented in Section A.2.2. This process is consistent with SMS, available EPA guidance, and the process commonly used in Superfund risk assessments.

Key considerations in the selection of ROCs included:

- Potential for direct or indirect (e.g., ingestion of fish or invertebrates) exposure to sediment-associated chemicals
- Human and ecological significance
- Site usage
- Sensitivity to COPCs at the site
- Susceptibility to biomagnification of COPCs (i.e., higher-trophic-level species)
- Data availability

To ensure that ROCs were selected to represent all potential exposure pathways for sediment-associated COPCs, key direct and indirect exposure routes from sediment were identified (e.g., direct exposure to sediment or ingestion of prey associated with sediment either directly or through prey). Groups of organisms that may be exposed via these pathways were then identified, and representative species that are thought to be most exposed were selected from these groups representing the greatest potential for exposure. Next, human or ecological significance was considered (i.e., species valued by society, have special regulatory status [i.e., threatened or endangered], or serve a unique ecological function).

Site usage, sensitivity to COPCs at the site, and data availability were also evaluated to determine the final list of ROCs. Site usage is an important criterion because it determines the exposure of a species; species that occupy the LDW during a significant part of the year or during sensitive periods, such as gestation and rearing of young, were preferred. Sensitivity to COPCs was evaluated based on available toxicological data, although in many cases the availability of data specific to LDW resident species is low. Therefore, where necessary, toxicological information from surrogate species, or a wide range of species, was used because species-specific data were not available. Finally, data availability regarding both site-specific exposure and effects was assessed, and species for which there are related site-specific data (such as COPC concentrations in food, site usage, and feeding) and toxicological data (such as sediment toxicity tests) were preferred. The following sections provide additional rationale for each of the ROCs selected; a summary of the selected ROCs is provided in Section A.2.3.4.

A.2.3.1 Benthic invertebrate community

The benthic invertebrate community as a whole is evaluated in this Phase 2 ERA as an ROC. As discussed in Section A.2.2.2, a wide variety of benthic invertebrates inhabit the LDW; most of these species are in direct contact with sediment year round and have a limited home range. Benthic invertebrates use various techniques to nourish themselves, and thus may be exposed to sediment through several different pathways (e.g., filter feeder vs. detritus feeder). Benthic invertebrates include sediment dwellers (benthic infauna) and organisms closely associated with the sediment surface (epibenthos).

Benthic invertebrates are an important food source for other invertebrates, fish, birds, and mammals, and perform essential functions such as nutrient cycling. Thus, the diversity and abundance of benthic invertebrates is an important component of the ecosystem. In addition, benthic organisms have been shown to be sensitive to sediment-associated chemicals, and data are available to assess their exposure and predict or measure effects, including sediment criteria from the SMS.

Among the benthic invertebrates, gastropods were considered by EPA to be potentially at risk from tributyltin (TBT).[8] LDWG performed two focused investigations to address this risk (Windward 2004g; 2005j). The scope and findings of these investigations are summarized in Sections A.3.1.3, A.3.2.4, A.6.1.1.3, and A 6.1.3.

SMS, which were used to evaluate risks to the benthic invertebrate community ROC, were not developed to explicitly address issues associated with bioaccumulation of COPCs by benthic invertebrates. Although this issue is implicitly addressed through the incorporation of benthic invertebrate community structure into the overall development of the SMS, the sediment quality standards (SQS) and cleanup screening levels (CSLs) do not specifically address risks to higher-trophic-level benthic invertebrate species, such as crabs. Therefore, crabs were selected as an ROC to represent higher-trophic-level benthic invertebrate species present in the LDW.

Crabs have a relatively larger home range than most of the benthic invertebrate species covered by the SMS. Dungeness crabs are the largest crab species observed in the LDW. Mating usually occurs in offshore locations, but may occasionally occur in estuaries (Pauley et al. 1988). Spawning takes place offshore. Juvenile crabs are found in shallow coastal waters and estuaries, like the LDW, and large numbers can live among eelgrass or other aquatic vegetation.[9]

The diet of crabs is dependent on their life stage (Pauley et al. 1986). Larvae ingest both zooplankton and phytoplankton. The diet of juvenile crabs consists largely of fish, mollusks, and crustaceans. Adult crabs prey on clams, crustaceans, and fish. Crabs eat bivalves their first year, shrimp their second year, and teleost fish their third year. Megalopae[10] are preyed upon by many fish, including juvenile salmon. Juvenile crabs are preyed upon by various demersal fish in the nearshore area. Flatfish, such as starry flounder and English sole, are the most important predators. Adult and juvenile crabs are preyed upon by sea otters, fish, and octopuses. Cannibalism is also common among crabs.

A.2.3.2 Fish

The potential fish ROCs were grouped into the following three broad categories based on potential sediment exposure at the site:

- **Anadromous juvenile salmonids**—representing species such as juvenile chinook, chum, and coho salmon

[8] Risks to gastropods from exposure to all other chemicals evaluated in this baseline ERA were addressed by the assessment of risks to the benthic invertebrate community.

[9] No eelgrass is found in the LDW, and habitats with aquatic vegetation are rare (Battelle et al. 2001).

[10] Crab larvae progress through five zoeal states before molting into megalopae. Megalopae first appear in April in Washington waters, with abundance peaking in May through June, after which they molt into juveniles.

- **Benthivorous fish—** representing species such as English sole, rock sole, and starry flounder. This category is also protective of fish, such as Pacific herring and pile perch, which prey on pelagic and encrusting organisms.
- **Upper-trophic-level fish —**representing species such as bull trout, sand sole, and Pacific staghorn sculpin

Using the criteria discussed in Section A.2.3, the following fish species were selected as ROCs in the LDW:

- Wild juvenile chinook salmon – anadromous juvenile salmonids
- English sole – benthivorous fish
- Pacific staghorn sculpin – upper-trophic-level fish

The remainder of this section discusses the rationale for selecting each fish ROC and how these species serve as surrogates for protection of other similar and important species within the LDW.

A.2.3.2.1 Wild juvenile chinook salmon

Wild juvenile chinook salmon (*Oncorhynchus tshawytscha*) were selected primarily because the Puget Sound evolutionary significant unit of chinook salmon (to which the Green River belongs) is a federally threatened species under ESA. In addition, they serve as a surrogate for other juvenile anadromous salmon.

Juvenile chinook salmon are representative of other salmonids because they have estuarine residence times similar to or greater than those of other juvenile salmonids in the LDW. During their spring outmigration, juvenile chinook salmon are among the most abundant fish in the LDW and are an important prey item for birds and piscivorous fish in the LDW (Warner and Fritz 1995). Juvenile chum salmon are also present in large numbers in the LDW from April through June and may rear extensively in the LDW (Warner and Fritz 1995). Residence times of all species of juvenile salmonids in the LDW are uncertain; however, juvenile chinook salmon are generally regarded as the most estuarine-dependent juvenile salmonid and their exposure to sediment-associated chemicals is likely equal to or greater than that of other juvenile salmonids.

Juvenile chinook are exposed to sediment-associated chemicals, primarily through their ingestion of benthic invertebrates, which are an important prey item in their early estuarine residence (Cordell et al. 1999). Juvenile chinook salmon have been studied in the LDW, and data are available on their exposure within the LDW, as well as potential effects associated with this exposure. Although toxicity data are available for several salmonid species, there are insufficient data to suggest that any one juvenile salmon species is more sensitive than another; therefore, available toxicity data did not affect selection of juvenile chinook as an ROC over other salmon.

Furthermore, chinook salmon are an icon of the Pacific Northwest. They have been used for centuries by indigenous people as a primary food source and are an economic resource of the region as a commercial fishery species. It is likely that some yearling juvenile chinook salmon (i.e., fish that have reared for one year in fresh water) outmigrate through the LDW (Warner and Fritz 1995; Shannon 2001). Yearling chinook tend to occupy deeper water than subyearling chinook and prey mainly on pelagic organisms, including small fish (Healy 1991). Risks to piscivorous yearling juvenile chinook salmon are assumed to be addressed by the Pacific staghorn sculpin ROC, as discussed below.

A.2.3.2.2 English sole

English sole (*Pleuronectes vetulus*) were selected to represent benthic carnivorous fish in the LDW. English sole live in close proximity to sediment, thus giving them a high potential for direct exposure to sediment-associated chemicals. In addition, English sole feed extensively on sediment-associated invertebrates and thus are subject to exposure to sediment-associated chemicals through their diet.

A number of studies have examined potential effects of sediment-associated chemicals (e.g., polycyclic aromatic hydrocarbons [PAHs]) on flatfish in the LDW, particularly English sole (e.g., Johnson et al. 1997b). Several toxicological studies have used data from English sole collected in the LDW (Casillas et al. 1991b; Johnson and Landahl 1994; Johnson et al. 1988; 1997b; 1998; 1999; Kubin 1997; Malins et al. 1984; 1985a; 1985b; Schiewe et al. 1989). National Marine Fisheries Service (NMFS) data suggest English sole are as sensitive to the effects of PAHs as other flatfish species tested (Myers et al. 1998). English sole are caught recreationally in the LDW and have some value as a commercial fishery species (though not in the LDW). Available toxicity data are not sufficient to suggest that English sole are more or less sensitive than other LDW species represented by English sole. Therefore, except for regionally specific studies conducted with English sole, no preference is given to toxicological data conducted with fish closely related to English sole.

English sole is a surrogate for other benthopelagic, demersal, and benthic fish species. In general, benthic organisms preyed on by other fish in the LDW are similar to those preyed on by English sole (see Section A.2.2.3.3), so their primary exposure route to sediment is similar. Therefore, because of the direct sediment contact, exposure of benthic fish such as English sole to sediment-associated chemicals is assumed to be greater than fish with equivalent prey preferences residing in other LDW habitats.

English sole is one of the most abundant fish in the LDW and is closely related to starry flounder, another of the most abundant fish in the LDW. Exposure and effects studies with English sole should, therefore, be relevant to starry flounder.

Other LDW fish, such as pile perch, ingest organisms that encrust pilings and other vertical structures. However, because these prey organisms do not have direct contact with sediment, this exposure route is not likely to result in greater exposure to

sediment-associated chemicals than the ingestion of benthic invertebrates. Similarly, fish species such as herring, surf smelt, longfin smelt, and threespine stickleback that ingest significant quantities of pelagic prey, are likely to have less exposure to sediment-associated chemicals than English sole that consume benthic prey exclusively. The available information thus indicates that the assessment of risks for English sole adequately addresses risks from sediment-associated chemicals to fish with benthopelagic, demersal, and benthic habitat preferences.

A.2.3.2.3 Pacific staghorn sculpin

Pacific staghorn sculpin, a benthic carnivorous fish, were selected to represent upper-trophic-level fish in the LDW. Like English sole, Pacific staghorn sculpin are benthic fish that feed on benthic invertebrates, giving them a high potential for direct and indirect exposure to sediment-associated chemicals.

Pacific staghorn sculpin feed on larger invertebrates than English sole and also feed on fish. Because other upper-trophic-level fish species have foraging ranges that extend beyond the LDW, Pacific staghorn sculpin were selected to represent upper-trophic-level fish in the LDW, thus accounting for exposures to bioaccumulative and biomagnifying chemicals, such as mercury and polychlorinated biphenyls (PCBs). This distinction was made because upper-trophic-level fish may have higher body burdens of biomagnifying chemicals than other fish, such as English sole, that ingest primarily invertebrates at a lower trophic level.

Pacific staghorn sculpin are one of the most abundant fish species in the LDW and likely serve as a food resource to piscivorous wildlife such as bald eagles (Knight et al. 1990), harbor seals (NMFS 1997), and river otters (Strand 1999; as cited in King County 1999a). No toxicity data were identified for Pacific staghorn; therefore, it is unknown whether Pacific staghorn sculpin are more or less sensitive than other higher-trophic fish species.

In the Phase 1 ERA (Windward 2003b), piscivorous bull trout were selected as the ROC to represent upper-trophic-level fish, largely because of their ESA status. Bull trout were replaced as an ROC in Phase 2 because they only occasionally use the LDW,[11] and insufficient tissue could be collected for chemical analysis. Other primarily piscivorous fish, including rockfish and sand sole, were also considered as representative of upper-trophic-level species because adults of these species eat at a high trophic level and rockfish are longer lived than Pacific staghorn sculpin. Available data (presented in Section A.2.2.3 and Attachment 2) and expert opinion (Windward 2004j) suggest that sandsole and rockfish are not abundant in the LDW and their foraging ranges likely extend beyond the LDW.

[11] Bull trout are believed to forage significantly in the LDW only during times of large abundances of small fish, such as just after the live birthing period for perch (Shannon 2001).

Also, subadult rockfish present in the LDW are at a trophic level similar to or lower than Pacific staghorn sculpin. In a study from British Columbia, Murie (1995) found that 80% of subadult copper rockfish (less than 15 cm in length) ingested demersal crustaceans and about 10% of subadult copper rockfish ingested fish. Similarly, about 60% of subadult quillback rockfish (less than 20 cm in length) ingested demersal crustaceans and less than 10% ingested fish. Pacific staghorn sculpin stomach contents analyses from throughout Puget Sound showed that fish constitute from 7 to 48% of the biomass in their diets (Fresh et al. 1979; Wingert et al. 1979; Miller et al. 1977c). Because diets of subadult rockfish are similar to Pacific staghorn sculpin diets, rockfish are assumed to be represented by Pacific staghorn sculpin. Because adult rockfish occur only near Harbor Island (the northern boundary of the LDW) (Windward 2004i) and other piscivorous species (such as bull trout and sand sole) forage extensively outside the LDW, their exposure to LDW-associated chemicals is likely to be less than Pacific staghorn sculpin exposure.

A.2.3.3 Wildlife

The potential wildlife ROCs were grouped into the following three broad categories based on potential sediment exposure at the site:

- **Piscivorous/carnivorous birds**—including species such as great blue heron, western grebe, cormorant, osprey, and bald eagle
- **Benthivorous birds**—including species such as spotted sandpiper, killdeer, and dabbling ducks
- **Piscivorous mammals**—including species such as river otter and harbor seal

Other broad categories of wildlife receptors, such as herbivorous birds, passerine birds, or omnivorous mammals are assumed to be less exposed to COPCs in the LDW than the three categories listed above because of their foraging behavior and diet. Primarily herbivorous birds, such as geese and some diving ducks, may also feed on benthic invertebrates and may incidentally ingest sediment while foraging, but this exposure is assumed to be less than that of benthivorous birds such as shorebirds, which may ingest significant amounts of sediment while probing intertidal sediment for benthic invertebrates. Ingestion of algae is also a potential exposure pathway. The significance of this pathway depends upon the extent to which sediment-associated chemicals migrate through the water column, are taken up by algae, and then ingested in significant quantities. Exposure from ingestion of algae is assumed to be insignificant compared to other more direct pathways examined (e.g., incidental ingestion of sediment or contaminated benthic species). Passerine birds are also likely to experience limited exposure to contaminated sediments in the LDW because they primarily use upland habitat. Other mammals, such as raccoons, are expected to have less exposure to sediment-associated chemicals because their food is largely terrestrial in origin, especially when compared to the primarily piscivorous river otter and harbor seal.

Using the criteria discussed in Section A.2.3, the following wildlife species were selected as ROCs in the LDW:

- Great blue heron—piscivorous birds
- Osprey—piscivorous/carnivorous birds
- Spotted sandpiper—benthivorous birds
- River otter—piscivorous mammals
- Harbor seal—piscivorous mammals

Evaluation of these representative receptors should be protective of other exposed species in the LDW.

The remainder of this section discusses the rationale for selecting each ROC and how these species will serve as representative species for protection of other species within the LDW. Species-specific toxicological data were not available for any of the LDW wildlife species to determine which species might be most sensitive to COPCs, although data for some COPCs are available for mink, which are in the same family as river otter. Therefore, toxicological sensitivity was not considered in the wildlife ROC selection process.

A.2.3.3.1 Great blue heron

The great blue heron (*Ardea herodias*) was selected to represent the piscivorous bird category because they are year-round residents, known to feed in and around the LDW, and nest nearby. Additionally, they are susceptible to biomagnification of certain chemicals because of their trophic position and feeding habits. Site-specific data for chemicals in heron food resources are available. It is assumed that the great blue heron serves as a representative species for waterfowl and seabirds that also feed primarily on fish and aquatic invertebrates (i.e., loons, western grebe, mergansers, double-crested cormorant, pigeon guillemot, Caspian tern, common murre). Great blue herons also attract a high level of societal interest.

A.2.3.3.2 Osprey

The osprey (*Pandion haliaetus*) was selected to represent piscivorous birds in addition to the great blue heron, as well as carnivorous birds such as peregrine falcon and bald eagles. Ospreys were selected because of the high number of nesting pairs along the LDW, their relatively small home range during nesting and fledgling development, and their diet, which is composed almost exclusively of fish. All of these factors can result in higher exposures to sediment-associated chemicals in the LDW when compared to other raptors, such as bald eagles. In addition, site-specific information on osprey feeding preferences is available because ospreys are being studied by USGS in the LDW.

Bald eagle was evaluated in the Phase 1 ERA (Windward 2003b) as the piscivorous bird ROC. Bald eagle has been replaced by osprey as an ROC because of the higher

incidence of osprey in the LDW, localized foraging range while nesting in the LDW, diet preferences,[12] and higher ingestion rates (IRs) normalized for body weight. Risk predictions for osprey should be similar or higher than would be estimated for bald eagles because of these factors. In addition, both species have been shown to be sensitive to some chemicals (e.g., DDT and PCBs) (Anthony et al. 1999; Buck 1999; Bowerman et al. 1995; Wiemeyer et al. 1988). While bald eagles are listed under ESA as a federally threatened species, all raptors tend to have high human interest and ecological significance.

A.2.3.3.3 Spotted sandpiper

The spotted sandpiper (*Actitis macularia*) was chosen to represent the benthivorous bird category because it feeds in the intertidal areas of the LDW from June through September and survey data indicate that sandpiper nest near Kellogg Island and perhaps other areas along the LDW. Sandpipers feed on invertebrates by probing the sediment, therefore, potentially ingesting significant quantities of sediment in addition to benthic invertebrates. Sandpipers have a higher incidental rate of sediment ingestion (some sandpiper species can have up to 30% in the diet) than other LDW bird species, including ducks and geese (EPA 1993). It is assumed that because of the high potential exposure through direct ingestion of sediment, protection of the spotted sandpiper will also be protective of other benthivorous birds such as scaup and scoters (i.e., diving ducks), as well as geese and dabbling ducks. Based on available information presented in the *Wildlife Exposure Factors Handbook* (EPA 1993) for birds that ingest aquatic invertebrates, spotted sandpipers have a higher food ingestion rate (FIR) than lesser scaups and mallards, contributing to higher exposure of spotted sandpipers. In addition, spotted sandpiper exposure should be higher than exposure of herbivorous birds such as American coot, American widgeon, mallard, and geese because COPC exposure is likely higher through ingestion of invertebrates than through ingestion of plants.

A.2.3.3.4 River otter

The river otter (*Lutra canadensis*) was chosen from the three semi-aquatic mammals using the LDW (including raccoon and muskrat) to represent the piscivorous mammal group because otters are suspected to be year-round residents that reproduce and feed in and around the LDW. The river otter is susceptible to biomagnification of chemicals because of its high trophic position and feeding habits and is more likely to feed on fish from the LDW than are raccoons or muskrats. Mustelids are also known to be highly sensitive to PCBs and other chlorinated organic compounds. Site-specific data for chemicals in otter food resources are available, as are relevant toxicological data for mustelids. Otters also attract a high level of societal interest.

---

[12] Eagles may consume birds and mammals as part of their diet; these dietary components may be less exposed to contaminated sediment in the LDW food web than are fish.

A.2.3.3.5 Harbor seal

The harbor seal (*Phoca vituluna*) was chosen from the three marine mammals using the LDW (including sea lion and harbor porpoise) to represent piscivorous mammals. The harbor seal, like the river otter, is susceptible to biomagnification of chemicals because of its trophic position and feeding habits. Pinnipeds are suspected to be sensitive to PCBs and other chlorinated organic compounds. Site-specific data for chemicals in harbor seal food resources are available. Seals are protected under the Marine Mammal Protection Act, and attract a high level of societal interest. It is assumed that the harbor seal will act as a representative species for other marine mammals, such as sea lions or harbor porpoise, which may infrequently use the LDW.

A.2.3.4 Summary of ROC selection

In summary, the following species were selected as ROCs to represent the range of organisms exposed to sediment-associated chemicals in the LDW:

- Benthic invertebrate community
- Crabs—higher-trophic-level benthic invertebrate
- Juvenile chinook salmon—anadromous juvenile salmon
- English sole—benthivorous fish
- Pacific staghorn sculpin—upper-trophic-level fish
- Great blue heron—piscivorous birds
- Osprey—piscivorous/carnivorous birds
- Spotted sandpiper—benthivorous birds
- River otter—piscivorous mammals
- Harbor seal—piscivorous mammals

The selection criteria for each of the above receptors are presented in Table A.2-10 to summarize the rationale for ROC selection.

Table A.2-10. ROCs selected for the LDW and a summary of the considerations for selection

ROC	Exposure Route	Ecological Significance	Societal Significance	Site Use	Exposure Data Availability	Sensitivity
Benthic invertebrate community	direct contact, diet, sediment ingestion	food source for other invertebrates, fish, and mammals; nutrient cycling	target community for protection in the development of numerical sediment quality criteria	present year-round; multiple life stages	abundant surface sediment data available	because of the diversity of organisms in this ROC group, the range of sensitivities is represented
Crabs	direct contact, diet, incidental sediment ingestion	higher trophic level benthic invertebrate	some recreational and commercial value	present year-round; multiple life stages	site-specific tissue data available	effects data available for decapods
Pacific staghorn sculpin	incidental sediment ingestion; preys on both invertebrates and other fish thus potential for elevated exposure via bioaccumulation because of trophic position	serves as prey for higher trophic level species such as birds and mammals	considered to be a nuisance by anglers	adults and juveniles present year-round; may spawn in the LDW	site-specific tissue data and prey tissue data available	effects data available for other fish species; unknown relative sensitivity of sculpin
English sole	direct contact, diet, incidental sediment ingestion	important prey item for birds and fish; key benthic predator	some recreational and commercial value	juveniles present year-round; adults present except during spawning migrations to Puget Sound	site-specific tissue and prey tissue data available	NMFS data suggest that they are as sensitive as other flatfish species (Myers et al. 1998)
Juvenile chinook salmon	diet	important prey item for birds/fish; seasonally one of the most abundant juvenile salmonids in the LDW	T&E species; returning adults important to commercial, sport, and tribal fisheries	generally present April-July; most estuary-dependent juvenile salmonid	site-specific tissue and prey tissue data available	believed to be sensitive to a wide range of COPCs
Great blue heron	preys on fish thus potential for elevated exposure via bioaccumulation because of trophic position	high on food chain	valued by society	present year-round; feed in LDW	site-specific data available for some food resources	effects data available for other bird species; unknown relative sensitivity of heron
Osprey	preys on fish thus potential for elevated exposure via bioaccumulation because of trophic position	top of food chain	valued by society; protected under migratory bird treaty	nests along the LDW and forages in LDW	site-specific data available for food resources	effects data available for other bird species; unknown relative sensitivity of osprey

ROC	Exposure Route	Ecological Significance	Societal Significance	Site Use	Exposure Data Availability	Sensitivity
Spotted sandpiper	diet, incidental sediment ingestion	important role as an intermediate predator	protected under migratory bird treaty	present June to September; nests along LDW	site-specific data available for food resources	effects data available for other bird species; unknown relative sensitivity of sandpiper
River otter	preys on fish thus potential for elevated exposure via bioaccumulation because of trophic position	top of food chain	valued by society	present year-round	site-specific data available for food resources	some mustelids shown to be highly sensitive to some chemicals, e.g., PCBs
Harbor seal	preys on fish thus potential for elevated exposure via bioaccumulation because of trophic position	top of food chain	protected under Marine Mammal Act	infrequent	site-specific data available for food resources	pinnipeds suspected to be sensitive to some chemicals, e.g., PCBs

COPC – chemical of potential concern

LDW – Lower Duwamish Waterway

NMFS – National Marine Fisheries Service

PCB – polychlorinated biphenyl

ROC – receptor of concern

T&E – threatened and endangered (species listed under the Endangered Species Act)

A.2.4 DATA SELECTION, REDUCTION, AND SUITABILITY

This section presents the chemical data available for the LDW and provides an evaluation of the relevance of these data to assess exposure of ROCs to sediment-associated chemicals.

A.2.4.1 Data selection and reduction

Many environmental investigations conducted in the LDW have included the collection of chemistry data from samples of surface sediment, tissue, or water. This section presents the datasets selected for use in the ERA for surface sediments, tissue samples, surface water samples, and porewater samples. The data management rules are presented in Attachment 3.

A.2.4.1.1 Surface sediment chemistry

The baseline surface sediment chemistry memorandum (Windward 2006d) summarizes, in tabular format, all the surface sediment samples that have been collected in the LDW from 1990 to 2005 and identifies those that are included in the baseline surface sediment dataset and those that were excluded, along with the rationale for their exclusion (Windward 2006d). The surface sediment chemistry data used in this baseline ERA are hereafter referred to as the baseline surface sediment chemistry dataset. The baseline surface sediment sampling locations are shown on Map A.2-3. Surface sediment samples (i.e., 15 cm or less) were collected from approximately 570 intertidal locations; the remainder (samples from approximately 750 sampling locations) were collected from subtidal areas.

Surface sediment samples from approximately 1,300 sampling locations were analyzed for PCBs (as Aroclors), 800 for semivolatile organic compounds (SVOCs) (including PAHs and phthalates), and 830 for metals and trace elements. Organochlorine pesticides were analyzed in surface sediment samples from approximately 200 locations. Dioxins and furans were analyzed in sediment samples from 43 locations. A summary of baseline surface sediment chemistry data is provided in Table A.2-11.

Table A.2-11. Summary of studies included in the baseline surface sediment dataset

SAMPLING EVENT	EVENT CODE	YEAR	CHEMICALS	NUMBER OF SAMPLING LOCATIONS[a]	SOURCE
LDW RI. Chemical analyses of benthic invertebrate and clam tissue samples and co-located sediment samples.	LDWRI-Benthic	2005	metals, SVOCs, PCB Aroclors, selected PCB congeners on subset of samples, butyltins	35	Windward (2005b)
LDW RI. Data report: Surface sediment sampling for chemical analyses and toxicity testing.	LDWRI-SurfaceSedimentRound1 & Round2	2005	metals, SVOCs, PCB Aroclors, selected PCB congeners and dioxins and furans on subset of samples, butyltins	160	Windward (2005d; 2005e)

Sampling Event	Event Code	Year	Chemicals	Number of Sampling Locations[a]	Source
Boyer Towing dock replacement	Boyer Towing	2004	metals, SVOCs, PCB Aroclors, TBT	3	WR Consulting (2004)
Slip 4 early action area site characterization	Slip4-EarlyAction	2004	PCB Aroclors, mercury	30	Integral (2004)
Rhône-Poulenc surface/subsurface sampling	RhônePoulenc 2004	2004	metals, organochlorine pesticides, SVOCs	21	EPA (2005b)
Norfolk CSO sediment remediation project five-year monitoring program: Annual monitoring report - year 5, April 2004.	Norfolk-monit7	2004	metals, PCB Aroclors, SVOCs	4	unpublished data from King County
Triad Approach to Characterize PCB in a Washington Riverine Sediment Site	Jorgensen August 2004	2004	metals, PCB Aroclors, SVOCs	43	USACE (2004)
Duwamish/Diagonal Perimeter monitoring – pre-dredge	DuwDiag-October2003	2003	metals, PCB Aroclors, organochlorine pesticides, SVOCs	12	unpublished data from King County
Terminal 117 early action area site characterization	T117 Boundary Definition	2003-2004	PCB Aroclors; metals, TBT and SVOCs on selected samples	54	Windward et al. (2004a; 2004b)
Boeing Plant 2 transformer investigation – Phase 1	Plant 2-Trans-former Phase1	2003	PCB Aroclors	6[b]	Floyd Snider McCarthy (2004)
Norfolk CSO (Duwamish River) sediment cap recontamination. Phase I investigation.	Ecology-Norfolk	2002	PCB Aroclors	17	Ecology (2003a)
Norfolk CSO sediment remediation project five-year monitoring program: Annual monitoring report - year 3, April 2002.	Norfolk-monit5	2002	metals, PCB Aroclors, SVOCs	1	King County (2002)
Norfolk CSO five-year monitoring program, Year Two, April 2001	Norfolk-monit4	2001	metals, PCB Aroclors, SVOCs	1	King County (2001b)
Norfolk CSO five-year monitoring program – Twelve-month post construction	Norfolk-monit3	2000	metals, PCB Aroclors, SVOCs	1	King County (2000c)
Norfolk CSO five-year monitoring program – Supplemental nearshore sampling	Norfolk-monit2b	2000	metals, PCB Aroclors, SVOC	3	King County (2000b)
Outfall and nearshore sediment sampling report, Duwamish Facility	James Hardie Outfall	2000	metals, PCB Aroclors, SVOCs	9	Weston (2000)
Norfolk CSO five-year monitoring program – Six-month post construction	Norfolk-monit2a	1999	metals, PCB Aroclors, SVOCs	2	King County (2000d)
Norfolk CSO five-year monitoring program – Post backfill	Norfolk-monit1	1999	metals, PCB Aroclors, SVOCs	2	King County (1999e)

Sampling Event	Event Code	Year	Chemicals	Number of Sampling Locations[a]	Source
EPA Site Inspection: Lower Duwamish River	EPA SI	1998	metals, PCB Aroclors, selected PCB congeners, and SVOCs; organochlorine pesticides, dioxins and furans, TBT, and VOCs on subset of samples	251	Weston (1999a)
King County CSO water quality assessment for the Duwamish River and Elliott Bay	KC WQA	1997	metals, PCB Aroclors, SVOCs, tetrabutyltin	14	King County (1999d)
Duwamish Waterway Phase 1 site characterization	Boeing SiteChar	1997	metals, PCB Aroclors, SVOCs	79[c]	Exponent (1998)
Duwamish Waterway sediment characterization study	NOAA SiteChar	1997	total PCBs, selected PCB congeners, total polychlorinated terphenyls	299	NOAA (1997; 1998)
Seaboard Lumber site, Phase 2 site investigation	Seaboard-Ph2	1996	metals, PCB Aroclors, SVOCs	20	Herrera (1997)
Rhône-Poulenc seep sampling	Rhône-Poulenc RFI-3	1996	metals, phenols	14	Rhône-Poulenc (1996)
RCRA Facility Investigation Duwamish Waterway sediment investigation, Plant 2 – Phase 2b	Plant 2 RFI-2b	1996	metals, PCB Aroclors, phthalates	36	Weston (1998)
Duwamish/Diagonal cleanup Study – Phase 2	Duw/Diag-2	1996	metals, PCB Aroclors, SVOCs	10	King County (2000a)
Duwamish/Diagonal cleanup Study – Phase 1.5	Duw/Diag-1.5	1995	metals, PCB Aroclors, SVOCs	9	King County (2000a)
Norfolk CSO sediment cleanup study – Phase 3	Norfolk-cleanup3	1995	PCB Aroclors, SVOCs	12	King County (1996)
Norfolk CSO sediment cleanup study – Phase 2	Norfolk-cleanup2	1995	metals, organochlorine pesticides, PCB Aroclors, selected PCB congeners, SVOCs	2	King County (1996)
RCRA Facility Investigation Duwamish Waterway sediment investigation, Plant 2 – Phase 2a	Plant 2 RFI-2a	1995	metals, PCB Aroclors SVOCs	54	Weston (1998)
RCRA Facility Investigation Duwamish Waterway sediment investigation, Plant 2 – Phase 1	Plant 2 RFI-1	1995	metals, PCB Aroclors, TPH, SVOCs, VOCs	66	Weston (1998)
Duwamish/Diagonal cleanup Study – Phase 1	Duw/Diag-1	1994	metals, organochlorine pesticides, PCB Aroclors, SVOCs	31	King County (2001a)
Norfolk CSO sediment cleanup study – Phase 1	Norfolk-cleanup1	1994	metals, organochlorine pesticides, SVOCs, PCB Aroclors	13	King County (1996)
Rhône-Poulenc RCRA Facility Investigation for the Marginal Way facility – Round 2	Rhône-Poulenc RFI-2	1994	SVOCs	6	Rhône- Poulenc (1995)
Results of sampling and analysis, sediment monitoring plan, Duwamish Shipyard, Inc.	Duwamish Shipyard	1993	metals, SVOCs, TBT	1	Hart Crowser (1993)

Sampling Event	Event Code	Year	Chemicals	Number of Sampling Locations[a]	Source
Harbor Island Remedial Investigation	Harbor Island RI	1991	metals, organochlorine pesticides, PCB Aroclors, SVOCs, VOCs, TPH, TBT	9	Weston (1993)

[a] Samples are surface sediment grab samples from 0 to 15 cm unless otherwise noted.

[b] Five samples were collected from 0 to 5 cm. The top interval from 0 to 15 cm from a subsurface sediment core was also included in the baseline surface sediment chemistry dataset.

[c] Sample total does not include three reference samples that were collected upstream of the study area.

CSO – combined sewer overflow
EPA – US Environmental Protection Agency
KC – King County
NOAA – National Oceanic and Atmospheric Administration
PCB – polychlorinated biphenyl
RCRA – Resource Conservation and Recovery Act
RFI – RCRA facility investigation
RI – remedial investigation
SI – site inspection
SVOC – semivolatile organic compound
TBT – tributyltin
USACE – US Army Corps of Engineers
VOC – volatile organic compound
WQA – water quality assessment

Since the RI/FS began in December 2000, there have been two sediment removal actions at early action areas within the LDW (Duwamish/Diagonal and Norfolk areas). There is no EPA policy or guidance about whether baseline risk assessments should include or exclude risk reduction achieved by removal actions that occur during the RI/FS. LDWG plans to evaluate the risk reduction achieved at these two areas in the FS as part of the residual risk assessment. Therefore, data characterizing areas prior to early actions are included to represent baseline conditions; post-removal action data collected from within and adjacent to early actions were excluded from the baseline dataset but will be included in the FS.

Both intertidal and subtidal sediment chemistry data were used in this baseline ERA. The elevation dividing intertidal and subtidal locations was approximately –2 ft MLLW, which corresponds to the shoreline (i.e., land/water interface) elevation defined by the aerial photos taken by the US Fish and Wildlife Service (USFWS) in 1999 (2000c).

A.2.4.1.2 Tissue chemistry

Tissue chemistry data for the study area are available for several different tissue types from several sampling events conducted since 1995. Site-specific tissue chemistry data were used in the ERA for the following species: juvenile chinook salmon, English sole, starry flounder, shiner surfperch, Pacific staghorn sculpin, Dungeness crab, slender crab, mussels, clams, and benthic invertebrates (Table A.2-12; Map A.2-2).

Over 200 composite samples of crab, English sole, starry flounder, Pacific staghorn sculpin, shiner surfperch, and benthic invertebrate tissue were used in this ERA (Table A.2-12). PCBs, as Aroclors, were analyzed in almost all samples. Organochlorine pesticides and SVOCs, metals including mercury, and TBT, were also analyzed frequently in approximately 150 to 180 samples. Unless otherwise noted, conversions between wet weight and dry weight were based on sample-specific

moisture content. Three studies were conducted with limited analyte lists (King County 1999d; Windward 2002, 2006b). The goals of these studies did not require the analysis of all SMS chemicals. Methylmercury and chromium VI were not analyzed in any of the tissue samples. Measurements of total mercury and chromium were determined to be sufficient in each study. Chemicals that have not been analyzed in LDW tissue samples but were analyzed in surface sediments samples will be discussed in the uncertainty analysis (Section A.6.0). Collection locations for LDW tissue samples listed in Table A.2-12 are presented on Map A.2-2.

Table A.2-12. LDW tissue data used in this baseline ERA

Study	Sample Collection Year	Species	Number of Samples	Individuals per Sample	Sample Type	Chemicals
Chemical analyses of fish and crab tissue samples collected in 2005 (Windward 2006b)	2005	English sole	21[a]	5	whole body	PCB Aroclors
		Dungeness crab	3	5	edible meat[b]	
			3	5	hepatopancreas[b]	
		slender crab	1	5	edible meat[b]	
			1	10	hepatopancreas[b]	
		shiner surfperch	22	10	whole body	
		Pacific staghorn sculpin	4	10	whole body	
Chemical analyses of fish and crab tissue samples collected in 2004 (Windward 2005c)	2004	English sole	21	5	whole body	metals, SVOCs, organochlorine pesticides, PCB Aroclors (PCB congeners in subset of samples), TBT
		starry flounder	3	5	whole body	
		Dungeness crab	7	5	edible meat[b]	
			3	6 – 15	hepatopancreas[b]	
		slender crab	12	5	edible meat[b]	
			4	15 – 18	hepatopancreas[b]	
		shiner surfperch	24	9 – 10	whole body	
		Pacific staghorn sculpin	24	7 – 10	whole body	
Chemical analyses of benthic invertebrate and clam tissue samples and co-located sediment samples (Windward 2005b)	2004	benthic invertebrates	20	nd	whole body	metals, SVOCs, alkylated PAHs, organochlorine pesticides, PCB Aroclors (PCB congeners in subset of samples), TBT
		clams	14	19 – 52	whole body	
Lower Duwamish Waterway remedial investigation. Juvenile chinook salmon data report (Windward 2004c)	2003	chinook salmon	28	9 – 10	whole body	metals, SVOCs, alkylated PAHs, organochlorine pesticides, PCB Aroclors, TBT
			1	74[d]	stomach contents	
Tissue chemistry results for juvenile chinook salmon collected from Kellogg Island and East Waterway (Windward 2002)	2002	chinook salmon	6	6 – 7	whole body	mercury, PCB Aroclors

Study	Sample Collection Year	Species	Number of Samples	Individuals per Sample	Sample Type	Chemicals
King County Combined Sewer Overflow Water Quality Assessment for the Duwamish River and Elliott Bay (King County 1999d) [c]	1996-1997	Dungeness crab	2	3	edible meat[b]	metals, SVOCs, PCB Aroclors, TBT
			1	3	hepatopancreas[b]	
		shiner surfperch	3	10	whole body	
		mussels	22	50 – 100	whole body	
		amphipods	4	nd	whole body	

[a] Whole-body samples include 11 composite samples analyzed as whole bodies and 10 estimated composite samples. Estimated English sole whole-body concentrations were based on the relative weights and total PCB concentrations in skin-on fillet and remainder tissues collected in 2005.

[b] Data from hepatopancreas composite samples were mathematically combined with data from composite samples of edible meat to estimate concentrations in composite samples of edible meat plus hepatopancreas. Whole-body (i.e., edible meat plus hepatopancreas) crab concentrations were calculated for each edible meat sample assuming 69% (by weight) edible meat and 31% hepatopancreas, based on the relative weight of these tissues in a 16.6-cm Dungeness crab dissected by Windward in 2004.

[c] Approximately 30 additional mussel samples, beyond those indicated in the table, were analyzed as part of the caged mussel deployment designed to assess effects from the combined sewer overflows (King County 1999d). These data are not included in this ERA because they are not representative of concentrations in mussels that would be ingested by wildlife.

[d] An unknown fraction of these stomachs were empty.

nd – no data

PAH – polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

SVOC – semivolatile organic compound

TBT – tributyltin

Not all of the available tissue chemistry data were appropriate for use in this baseline ERA. Juvenile chinook salmon whole-body and stomach content data from Varanasi et al. (1993) and from NMFS (2002) were not used in the ERA because the quality assurance/quality control (QA/QC) data were not readily available for EPA review. The allocation of resources to further review these data was determined to not be warranted because both total PCB and total DDT concentrations in the NMFS dataset were similar to or lower than concentrations in LDWG data (Windward 2005h). However, total PAH concentrations in LDW juvenile chinook salmon stomach contents from three separate NMFS studies (Arkoosh et al. 1998; McCain et al. 1990; Stein et al. 1995) were higher than those in juvenile chinook salmon stomach contents collected in support of the Phase 2 RI from similar locations in the LDW (Windward 2004c).

Data associated with two individual shiner surfperch collected in Slip 4 (NMFS 2002) and 10 individual shiner surfperch collected and analyzed by LDWG in subarea T2E (RM 2.1 to 2.4) (Section 4 of the RI) were also excluded. These data were excluded because only data from composite samples were included in the ERA, as per EPA (2000c) guidance. Total PCB concentrations in the two individual shiner surfperch from Slip 4 were 940 and 2,100 µg/kg ww, and the total PCB concentrations in the 10 individual shiner surfperch from subarea T2E ranged from 172 to 1,140 µg/kg ww. These individual shiner surfperch concentrations were lower than the T2E composite sample concentrations. However, the individual sample concentrations were within the range of concentrations in shiner surfperch composite samples (530 to 18,400 µg/kg). The inclusion of these samples would not have affected the results of the risk assessment. The available data for individual shiner surfperch samples were also

deemed to be insufficient to estimate the variability in individuals in composite samples (EPA 2005a).

In addition, some site-specific tissue chemistry data were also excluded from the ERA because they are unrelated to site-specific contamination (e.g., data for adult salmon) or because the samples were not appropriate for ecological receptors (e.g., fillet samples).

The only available bird egg data were heron egg data collected by USFWS (Krausmann 2002a). There were insufficient QA/QC data available to include the data in the risk characterization.

A summary of LDW tissue chemistry data used in this baseline ERA is provided in the Phase 2 RI. The RI report also contains a compact disc with all raw data used in the RI and risk assessments.

A.2.4.1.3 Water chemistry

Water data are available from sampling events conducted by King County in 1996 and 1997 during their WQA (King County 1999d) and during an additional sampling event in 2005 in which water samples were collected and analyzed only for PCBs and conventional parameters (Mickelson and Williston 2006).

Grab samples were collected along transects at three locations in the LDW in 1996 and 1997: RM 1.1 (in the vicinity of the Brandon CSO outfall), RM 1.9 (in the vicinity of the Southwest Michigan CSO outfall), and RM 4.9 (in the vicinity of the Norfolk CSO outfall). At the Brandon and Southwest Michigan areas, there were three sampling locations, corresponding to the east and west banks and the center of the channel. Two depths were sampled at each location: 1 m below the surface and 1 m above the bottom. There were only two sampling locations at Norfolk, corresponding to the east and west banks, because of the narrow width and depth of the LDW at this location. Samples were collected weekly from October 1996 to June 1997, except during storm events, in which case sampling was conducted on three successive days following the storm. Chemicals relevant to the risk assessment that were analyzed in these samples are metals and trace elements (total and dissolved concentrations of antimony, arsenic, cadmium, cobalt, copper, lead, mercury, nickel, selenium, silver, thallium, vanadium, and zinc) and SVOCs.

In 2005, King County collected water column samples for PCB congener analysis during two dry-weather sampling events in August and September, and two wet-weather sampling events in November and December. Two locations were sampled in the LDW: one location at RM 0.0 at the southern end of Harbor Island, and one at RM 3.3 at the 16th Avenue South Bridge. All 209 PCB congeners were analyzed using high resolution methods, and the total PCB concentration for the risk assessment was calculated as the total of all detected PCB congener concentrations (data were blank-qualified, and RLs for undetected PCB congeners were not included).

King County also deployed semipermeable membrane devices and analyzed samples for organochlorine pesticides and PAHs. Although predicted water concentrations are available from these data, they were not used in the risk assessment.[13]

A.2.4.1.4 Porewater chemistry

The concentrations of volatile organic compounds (VOCs) in porewater were available from a Phase 2 porewater collection effort conducted by LDWG (Windward 2006a). Groundwater data for VOCs from 11 sites along the LDW were reviewed to determine which sites had the highest potential for VOC concentrations in porewater. As documented in the porewater QAPP (Windward 2005i), two sites (Great Western International [GWI] and Boeing Plant 2/Jorgensen Forge) were selected for porewater sampling and analysis (Windward 2006a). Porewater was sampled at multiple locations within each site using two methods. Piezometers were deployed at 12 locations and peepers were deployed at 16 locations (Windward 2006a). Samples were analyzed for 71 VOCs. No VOCs were detected in the piezometer samples, and 16 VOCs were detected in at least one of the peeper samples.

In 1997, porewater samples were collected at 15 locations throughout LDW subtidal areas as part of the EPA Site Inspection for the LDW (Weston 1999b). The samples were analyzed for a suite of metals and TBT. These data were not used in the risk assessment because benthic invertebrate exposure from metals in porewater was addressed in the sediment exposure evaluation by using SMS, toxicologically based DMMP guidelines, and TRVs. TBT porewater data were not used for the assessment of gastropods because TBT porewater data were not available from the imposex locations. Exposure of other invertebrates to TBT was addressed by the critical tissue-residue approach.

A.2.4.2 Suitability of data for risk assessment

There are several factors to consider in assessing the suitability and sufficiency of environmental data for risk assessments (EPA 1989, 1990). Of primary importance is the degree to which the data adequately represent site-related contamination, and the expected exposure of ecological receptors at the site. Other important considerations are data quality criteria goals, documentation, analytical methods, reporting limits (RLs), and level of review associated with the data. Because data from many different investigations were available for the LDW, these factors were evaluated for each data set to determine whether it was reasonable to combine these data for use in this ERA.

---

[13] Water data were only used for wildlife exposure estimates. Predicted PAH water concentrations from SPMDs were not used because PAHs did not screen in as COPCs for wildlife. Predicted pesticide water concentrations (i.e., total DDTs) from SPMDs were not used because of uncertainties in calculating the predicted concentration, and because the contribution of COPCs from water to the total exposure dose was very small and was not expected to affect risk calculations (see Section 5.1.3.3).

A.2.4.2.1 Representativeness to site-related contamination and receptor exposure

This section provides an overview of the representativeness of the available sediment, tissue, and water data.

Sediment

Many historical environmental sampling events included collection of sediment from the LDW (Table A.2-11). The studies have been designed for both reconnaissance (e.g., Boeing SiteChar, EPA SI, and NOAA SiteChar) and focused investigation of suspected areas of contamination (e.g., Boeing Resource Conservation and Recovery Act [RCRA] facility investigation [RFI], Rhône-Poulenc RFI). The extensive coverage of the reconnaissance surveys, and the focused intensity of facility investigations, indicate available sediment chemistry data are sufficiently representative of the general range of environmental conditions within the LDW. Because samples of environmental media have often been collected to characterize particularly contaminated areas, the dataset as a whole may have a relatively higher proportion of elevated concentrations than is representative of the LDW as a whole.

Existing data used in Phase 1 were supplemented with Phase 2 surface sediment data. Phase 2 surface sediment samples were collected from more than 165 stations according to the following criteria:

- Low historical spatial coverage, particularly at sites where single SQS or CSL exceedances were observed with few nearby sampling locations
- Special use areas (e.g., intertidal areas with public access or used by wildlife) that had previously been incompletely characterized
- Proximity to potential historical or current chemical sources, including seeps identified as being of concern
- Historical sediment chemical concentrations above the SQS or CSL
- Analyte considerations including chemicals with relatively low numbers of historical samples or historical locations that did not have sufficiently low RLs for certain chemicals

Tissue

To be representative, tissue data must provide a reasonable indication of COPC exposure by ROCs at a site. Key considerations in the representativeness of site data are:

- Representativeness of the tissue data with respect to capture location, timing, and home range of the species relative to the site
- Availability of tissue data for COPCs at the site
- Representativeness of tissue data with respect to ROCs at the site and their primary prey items

The home range of fish collected in the LDW may be greater or smaller than the area of the LDW Superfund site. For English sole, for example, considerable uncertainty exists regarding preferred foraging habitat and home range; no site-specific home range estimates have been published for English sole in the LDW. A few home range estimates have been developed using best professional judgment, such as the 9 km^2 home range of English sole, as reported in the Puget Sound Dredged Disposal Analysis (PSDDA 1988a) and 2 km^2 based on a literature review (Stern et al. 2003). One tagging study (Day 1976) suggests that English sole may have some site fidelity, although the "sites" defined in that study are relatively large compared to the LDW. Also, the extent of migration was not established.

When the home range of a particular species does not match the LDW site boundaries, measured body-burdens may over- or underestimate contamination associated with the site. For example, in winter, English sole migrate to Elliott Bay to spawn (Forrester 1969) and, therefore, could be exposed to some chemicals outside of the LDW. Also, juvenile chinook salmon pass through the LDW in their migration from either upstream spawning locations or hatcheries. Juvenile chinook salmon released from hatcheries have a small contaminant load before entering the LDW, which is generally attributed to the low concentrations of various chemicals (e.g., PCBs) present in hatchery feed (Easton et al. 2002). As such, a portion of their overall contaminant body burden is not associated with LDW exposure (Meador 2000).

Also, the age of the fish captured can influence the body burden. Older fish tend to have higher concentrations of biomagnifying COPCs in their tissues, and thus ingestion of these individuals could result in higher exposures to piscivorous receptors. The available English sole, Pacific staghorn sculpin, and perch data represent adult fish.

Uncertainty in exposure assumptions associated with tissue data used in the ERA is discussed further in the uncertainty analysis (Section A.6.2).

Water

The only use of the surface water data in this baseline ERA was to estimate a relatively small percentage of the exposure of wildlife receptors to COPCs through the ingestion of water. Water data for all COPCs, except PCBs, were based on a large dataset of samples collected on a weekly basis from October 1996 to June 1997 along transects at three areas in the LDW. The PCB data were collected from a downstream location and a location midway through the LDW (RM 3.3), and were collected four times in 2005 to represent a seasonal range of flow conditions. The water data are assumed to represent conditions to which wildlife receptors can be exposed in the LDW because the LDW is a well-mixed system.

Porewater

Porewater data were collected from two areas of the LDW to investigate potential worst case exposure conditions for VOCs to assess benthic invertebrate risk. Porewater

samples were collected using peepers deployed 10 cm beneath the sediment surface and left to equilibrate for two weeks. Because of their placement within the biologically active zone and the equilibration time allowed, these samples should reasonably represent benthic invertebrate exposure conditions within these areas.

A.2.4.2.2 QA/QC results

All datasets used in this ERA have been validated by the original study authors or by outside third parties. No additional data validation was performed for this ERA. Some results were qualified as unusable[14] by the data validators. Data qualified as unusable were not used in this ERA and were not included in summary information provided in Tables A.2-11 and A.2-12.

Analytical interference with the quantification of organochlorine pesticides from the presence of PCB congeners occurred during the pesticide analysis of benthic invertebrate tissue (Windward 2005b) and fish and crab tissue (Windward 2005c). This issue was identified by both the analytical laboratory and the data validators. The organochlorine pesticides were analyzed using EPA Method 8081 (gas chromatography with electron capture detection [GC/ECD]), which is a standard method used in many environmental investigations for organochlorine pesticides. The detected results for both the benthic invertebrates and clams (Windward 2005b) and fish and crab samples collected in 2004 (Windward 2005c) were qualified JN, which indicates "the presence of an analyte that has been 'tentatively identified' and the associated numerical value represents its approximate concentration" (EPA 1999b). These data were qualified based on the probable interference in the analysis from PCB congeners.

The JN-qualified results are highly uncertain and biased high. The high bias for DDTs was confirmed by reanalyzing six sediment samples co-located with benthic invertebrate tissue samples and eight fish and crab tissue samples that had high PCB and DDT concentrations using a gas chromatography/mass spectrometry (GC/MS) method that is not susceptible to analytical interference by PCBs for organochlorine pesticides. The GC/MS method is less sensitive than EPA Method 8081, and therefore, could not be used for the original analyses and could only be used for confirmation in the high concentration samples. The only DDT isomers that were detected in the confirmation analyses were 4,4'-DDE and 4,4'-DDD. The four isomers most commonly detected in the GC/ECD analyses were 4,4'-DDT, 2,4'DDT, 4,4'-DDE, and 4,4'-DDD; the highest concentrations were reported for the two DDT isomers.

The confirmation analysis results confirmed the JN-qualification of the original sample results. Specifically, all the results from the confirmation analyses were lower than the original results. The total DDT concentrations in the confirmation analyses ranged from 4% to 60% of the original sediment results (Windward 2005g). The two highest

[14] Approximately 1,000 results were qualified as unusable out of more than 140,000 analytical results.

total DDT concentrations were not detected in the confirmation analyses with RLs that were 3% and 8% of the original results. Eight fish and crab tissue samples (three crab hepatopancreas, four shiner surfperch, and one English sole whole body) were reanalyzed and the reanalysis confirmed the presence of DDTs in six of the samples with total DDT concentrations that were 5 to 34% of the original results (Windward 2005g). Thus, the original reported concentrations of DDT compounds appear to reflect the presence of both PCB congeners and DDT isomers in the sample, and were elevated because of analytical interference.

The DDT confirmation analyses were run using subsamples of the original sample extracts that had been archived frozen. The confirmation analyses were conducted six months after extraction, which greatly exceeds the maximum 40-day extract holding time. Therefore, the results of these analyses were treated as qualitative and useful as an estimate of the DDT isomer concentrations, but were not incorporated into the project database, or used in the risk calculations presented in this ERA.

The JN-qualified organochlorine pesticide results were used in the problem formulation screens to identify COPCs for each receptor (Section A.2.5). However, because of the high uncertainty inherent in the organochlorine pesticide concentrations in tissue samples, the characterization of risks associated with tissue organochlorine pesticide results is presented in the relevant uncertainty analysis sections rather than the risk characterization sections to highlight the uncertainty in the data.

A.2.5 SELECTION OF CHEMICALS OF POTENTIAL CONCERN

This section presents a risk-based screen conducted to identify which chemicals should be identified as COPCs for each of the ROCs. Through this COPC screening process, a clear distinction can be made between chemical/receptor pairs that should be evaluated in more detail using more realistic assumptions (Sections A.3.0 through A.6.0) and pairs for which no additional analysis is warranted. The COPC selection process is presented separately in this section for benthic invertebrates, fish, and wildlife ROCs. A brief summary of the ecotoxicology of potential COPCs was presented in the Phase 1 ERA (Windward 2003b).

Quantitative ecological risk analyses were not conducted for dioxins and furans in this baseline ERA, and dioxins and furans were not selected as COPCs for any ecological receptor. Because dioxin and furan congeners were not analyzed in tissue samples collected from the LDW, risks to ecological receptors associated with exposure to dioxins and furans are unknown.

Risks resulting from exposure to dioxins and furans were not assessed because dioxins and furans were not analyzed in tissue samples collected from the LDW for the following reasons:

- Risks from dioxins and furans were assumed to be unacceptable in the LDW, particularly from a human health seafood-consumption perspective. Dioxins and furans are commonly detected in all media regardless of whether there are any local sources of dioxins and furans, and dioxins and furans are toxic to humans, fish, and wildlife species.
- Because of the ubiquity of dioxins and furans in the environment, it is important to consider the degree of contamination of sediments with dioxins and furans in a regional context. In accordance with MTCA and CERCLA regulations and guidance regarding use of information from background areas in contaminated site investigations, comparisons with background are necessary to understand exposures that may have resulted from site-specific releases relative to exposures associated with the general urban environment. Few tissue data are available for dioxins and furans within the greater Seattle metropolitan area and Puget Sound. Therefore, the collection of a large background tissue dataset, in addition to extensive tissue collection within the LDW, would have been required to conduct a robust comparison between the LDW and the general urban environment.

Sediment data for dioxins and furans are available for both the LDW site and background areas. However, sediment data were not considered sufficient for evaluating risks associated with seafood consumption to either human or ecological receptors because of uncertainties associated with predicting tissue concentrations when only sediment data are available. Although the concentrations of dioxins and furans in most sediment samples from the LDW were comparable to those found throughout the Seattle area (Section 4 of the RI), dioxins and furans were detected in sediment in a few samples at concentrations that were much higher than the concentrations detected at upstream or other locations in the greater Seattle metropolitan area.

Risk management decisions to address the dioxin and furan contamination in LDW sediments will be based on MTCA and CERCLA regulations and guidance. Remedial decisions will be made by EPA and Ecology as part of the FS process and will be documented in the Record of Decision. Additional information regarding the nature and extent of dioxins and furans in LDW sediments is presented in the Phase 2 RI and in the baseline HHRA (Appendix B).

A.2.5.1 Benthic invertebrates

This section presents the screening approach that was used to identify COPCs for the benthic invertebrate community and for crabs. The benthic invertebrate community screen was conducted separately for two media: sediment and porewater. The sediment screen was conducted to identify COPCs in sediment for which SMS or other sediment-based guidelines are available. VOCs were screened using porewater data. VOCs may be present in sediment porewater in areas associated with groundwater

discharge to the LDW if there is a continuing source of VOC-contaminated groundwater (Church et al. 2002; Chadwick et al. 1999). VOCs do not have a high affinity for sediment because of their generally low OC/water partition coefficients (K_{OC}) (Mabey et al. 1982). Therefore, the exposure of sediment-dwelling organisms (i.e., benthic invertebrates) to VOCs is most appropriately assessed through analysis of sediment porewater rather than bulk sediment. COPCs for crabs, also presented in this section, were identified based on a critical tissue-residue approach.

A.2.5.1.1 Screening methods and results for benthic communities based on surface sediment chemistry data

The Phase 2 work plan established the use of sediment numerical chemical standards promulgated under the Washington SMS to evaluate whether individual sediment-associated chemicals should be retained as COPCs for benthic invertebrates in this ERA. Thus, surface sediment baseline data described in Section A.2.4.1.1 were compared to SQS for all chemicals listed in the SMS. For chemicals with no SQS and detected in at least 5% of LDW surface sediment samples, maximum concentrations were compared to DMMP guidelines (USACE et al. 2000) that were determined to be toxicologically based (Figure A.2-1). In cases where no DMMP value was available or the available DMMP value was not toxicologically based (i.e., total DDTs), the scientific literature was searched for toxicologically based toxicity reference values (TRVs) for these chemicals. Chemicals exceeding the DMMP guidelines or toxicologically based TRVs were identified as COPCs for benthic invertebrates.

Chemicals detected in greater than 5% of the sediment samples, but without SMS or toxicologically based guidelines or TRVs, are discussed in the uncertainty analysis (Section A.6.1.1.1). Chemicals with RLs greater than the SQS are discussed in the uncertainty analysis and in the Phase 2 RI.



Figure A.2-1. Screening process for benthic invertebrate COPCs using sediment chemistry data

As previously discussed, the SQS were promulgated to address risks to benthic invertebrate communities as a whole, except for higher-trophic-level invertebrates, such as crabs, that may be at greater risk of exposure through bioaccumulation.[15] SQS values are based on apparent effects thresholds (AETs), which are defined as the highest "no effect" chemical concentration above which a significant adverse biological effect always occurred among the several hundred samples used for its derivation. Biological endpoints included in derivation of the SQS chemical standard were field measures of benthic infaunal abundance, laboratory toxicity tests with marine benthic invertebrate organisms (i.e., amphipods [survival] and oysters [percent abnormal development of oyster larvae]), and laboratory toxicity tests with bacteria (Microtox [decrease luminescence from the bacteria *Vibrio fisheri*]). Representatives of these groups are found throughout the LDW. Under the provisions of the SMS, surface sediments with chemical concentrations equal to or less than all the SQS are designated as having no adverse effects on biological resources (Washington Administrative Code [WAC] 173-204-310(1)(a)).[16]

The baseline surface sediment data described in Section A.2.4.1.1 were compared to SQS for all chemicals listed in Ecology's SMS. Many SQS values are expressed as concentrations normalized to total organic carbon (TOC). At very low or high TOC concentrations, normalization is not appropriate (Michelsen and Bragdon-Cook 1993). Concentrations of organic chemicals were not normalized to TOC for samples with TOC concentrations less than or equal to 0.5% or greater than or equal to 4.0%. In these cases, dry weight chemical concentrations were compared to the lowest AET (LAET), which is functionally equivalent to the SQS, and the second lowest AET (2LAET), which is functionally equivalent to the CSL.

Table A.2-13 presents the chemicals identified as COPCs for benthic invertebrates and retained for further evaluation in this baseline ERA. A total of 44 chemicals were retained as COPCs for benthic invertebrates.[17] In addition, TBT was also included as a COPC for benthic invertebrates in this baseline ERA based on Phase 1 ERA (Windward 2003b) results and the focused investigations regarding imposex in gastropods conducted for this baseline ERA.

---

[15] Risks to crabs were evaluated using a tissue approach in Sections A.3.3, A.3.4, and A.6.1.2. Also, SMS are not intended to be protective of other receptors (such as fish and wildlife) exposed to sediment-associated COPCs through bioaccumulation. Risks to these receptors are presented in Sections A.6.2 and A.6.3.

[16] Because of the SQS derivation process, there is some uncertainty regarding the prediction of effects based solely on comparison with the SQS, even though this is designated under the provisions of the SMS.

[17] Six SMS chemicals were screened out: 2-methylphenol, acenaphthylene, diethyl phthalate, di-n-butyl phthalate, di-n-octyl phthalate, and hexachlorobutadiene.

Table A.2-13. Summary of COPCs retained for benthic invertebrates based on surface sediment chemistry data

COPC[a]	Maximum Concentration	Unit	Maximum Concentration in Comparable Units	Unit	Screening Criteria	
					SQS, LAET, SL, or Toxicologically Based TRV	Unit
Metals						
Arsenic	1,100	mg/kg dw	1,100	mg/kg dw	57	mg/kg dw
Cadmium	120	mg/kg dw	120	mg/kg dw	5.1	mg/kg dw
Chromium	1,100 J	mg/kg dw	1,100 J	mg/kg dw	260	mg/kg dw
Copper	12,000 J	mg/kg dw	12,000 J	mg/kg dw	390	mg/kg dw
Lead	23,000	mg/kg dw	23,000	mg/kg dw	450	mg/kg dw
Mercury	4.6 J	mg/kg dw	4.6 J	mg/kg dw	0.41	mg/kg dw
Nickel	910	mg/kg dw	910	mg/kg dw	140[b]	mg/kg dw
Silver	270	mg/kg dw	270	mg/kg dw	6.1	mg/kg dw
Zinc	9,700	mg/kg dw	9,700	mg/kg dw	410	mg/kg dw
Organometals						
TBT	3,000	µg/kg dw	na[f]	na[f]	na[f]	na[f]
PAHs						
Acenaphthene	5,200	µg/kg dw	260	mg/kg OC	16	mg/kg OC
Anthracene	10,000	µg/kg dw	380	mg/kg OC	220	mg/kg OC
Benz(a)anthracene	8,400	µg/kg dw	440	mg/kg OC	110	mg/kg OC
Benzo(a)pyrene	7,900	µg/kg dw	420	mg/kg OC	99	mg/kg OC
Benzo(g,h,i)perylene	3,800	µg/kg dw	180	mg/kg OC	31	mg/kg OC
Chrysene	7,700	µg/kg dw	410	mg/kg OC	110	mg/kg OC
Dibenzo (a,h)anthracene	1,500	µg/kg dw	71	mg/kg OC	12	mg/kg OC
Fluoranthene	24,000	µg/kg dw	1,300	mg/kg OC	160	mg/kg OC
Fluorene	6,800	µg/kg dw	260	mg/kg OC	23	mg/kg OC
Indeno (1,2,3,-c,d)pyrene	4,300	µg/kg dw	200	mg/kg OC	34	mg/kg OC
Naphthalene	5,300	µg/kg dw	260	mg/kg OC	99	mg/kg OC

COPC[a]	Maximum Concentration	Unit	Maximum Concentration in Comparable Units	Unit	Screening Criteria	
					SQS, LAET, SL, or Toxicologically Based TRV	Unit
Phenanthrene	28,000	µg/kg dw	1,500	mg/kg OC	100	mg/kg OC
Pyrene	4,400	µg/kg dw	4,400	µg/kg dw	2,600[c]	µg/kg dw
Total benzofluoranthenes	17,000	µg/kg dw	890	mg/kg OC	230	mg/kg OC
HPAH	85,000	µg/kg dw	4,500	mg/kg OC	960	mg/kg OC
LPAH	44,000	µg/kg dw	1,700	mg/kg OC	370	mg/kg OC
Phthalates						
Bis(2-ethylhexyl) phthalate	14,000	µg/kg dw	14,000	µg/kg dw	1,300[c]	µg/kg dw
Butyl benzyl phthalate	7,100	µg/kg dw	340	mg/kg OC	4.9	mg/kg OC
Dimethyl phthalate	1,400 J	µg/kg dw	140 J	mg/kg OC	53	mg/kg OC
Other SVOCs						
1,2-Dichlorobenzene	520 J	µg/kg dw	520 J	µg/kg dw	35[c]	µg/kg dw
1,4-Dichlorobenzene	1,600 J	µg/kg dw	1,600 J	µg/kg dw	110[c]	µg/kg dw
1,2,4-Trichlorobenzene	72 J	µg/kg dw	72 J	µg/kg dw	31[c]	µg/kg dw
2-Methylnaphthalene	3,300	µg/kg dw	160	mg/kg OC	38	mg/kg OC
4-Methylphenol	4,600 J	µg/kg dw	4,600 J	µg/kg dw	670	µg/kg dw
2,4-Dimethylphenol	290 J	µg/kg dw	290 J	µg/kg dw	29	µg/kg dw
Benzoic acid	4,500	µg/kg dw	4,500	µg/kg dw	650	µg/kg dw
Benzyl alcohol	670	µg/kg dw	670	µg/kg dw	57	µg/kg dw
Dibenzofuran	4,200	µg/kg dw	220	mg/kg OC	15	mg/kg OC
Hexachlorobenzene	95 J	µg/kg dw	3.7 J	mg/kg OC	0.38	mg/kg OC
N-Nitrosodiphenylamine	110	µg/kg dw	110	µg/kg dw	28[c]	µg/kg dw
Pentachlorophenol	410	µg/kg dw	410	µg/kg dw	360	µg/kg dw
Phenol	2,800	µg/kg dw	2,800	µg/kg dw	420	µg/kg dw
PCBs						
Total PCBs	220,000	µg/kg dw	10,000	mg/kg OC	12	mg/kg OC

COPC[a]	Maximum Concentration	Unit	Maximum Concentration in Comparable Units	Unit	Screening Criteria	
					SQS, LAET, SL, or Toxicologically Based TRV	Unit
Organochlorine Pesticides						
Total DDTs	2,900 J	µg/kg dw	2,900 J	µg/kg dw	567[d]	µg/kg dw
Total chlordane	230	µg/kg dw	230	µg/kg dw	2.8[e]	µg/kg dw

[a] COPCs identified based on: 1) a comparison of maximum surface sediment concentrations to SMS SQS, or 2) for chemicals with no SQS, detected in at least 5% of LDW surface sediment samples and a comparison between maximum surface sediment concentrations and DMMP guidelines or toxicologically based TRVs.

[b] DMMP SL guideline.

[c] Comparison is based on LAET because TOC normalization was not appropriate for the sample.

[d] Literature based TRV – (Lotufo et al. 2001b).

[e] Literature based TRV – PSDDA AET evaluation 1994 (Gries and Waldow 1996).

[f] TBT was included as a COPC based on the results of the Phase 1 ERA (Windward 2003b), and was evaluated using benthic invertebrate tissue data consistent with EPA (1999a) and Meador et al. (2002). Tissue-residue data integrate multiple exposure pathways and provide a direct measure of exposure (EPA 1999a); therefore, tissue-residue-based TRVs were selected over the porewater-based DMMP SL guideline for TBT.

COPC – chemical of potential concern

DMMP – Dredged Material Management Program

dw – dry weight

HPAH – high-molecular-weight polycyclic aromatic hydrocarbon

J – estimated concentration

LAET – lowest apparent effects threshold

LPAH – low-molecular-weight polycyclic aromatic hydrocarbon

na – not applicable

OC – organic carbon

SL – screening level (DMMP)

SMS – Washington State Sediment Management Standards

SQS – sediment quality standards (SMS)

TBT – tributyltin

TRV – toxicity reference value

A.2.5.1.2 Screening methods and results for benthic communities based on porewater chemistry data

COPCs for benthic invertebrates potentially exposed to VOCs in porewater were also identified. The Phase 2 RI work plan (Windward 2004e) identified the need to assess risks to benthic invertebrates from porewater collected in areas where VOCs have been historically detected in groundwater at upland properties immediately adjacent to the LDW. Available data for VOCs in groundwater at 11 sites along the LDW were reviewed and two sites, GWI and Boeing Plant 2/Jorgensen Forge, were selected for porewater characterization because they had the highest potential for VOC concentrations in porewater based on their proximity to the LDW and existing groundwater data. The two locations are shown on Map A.2-3.

VOCs were detected in samples collected using peepers, but were not detected in deeper samples collected with piezometers. Four VOCs were detected in porewater samples collected at Boeing Plant 2/Jorgensen Forge. In the GWI area, 16 VOCs were detected in the porewater samples. COPCs in porewater for benthic invertebrates were identified by comparing maximum VOC concentrations detected in porewater to literature-based no-observed-effect concentration (NOEC) TRVs.

A literature search was conducted for relevant aquatic toxicity studies on growth, mortality (including immobilization), or reproductive (including developmental) endpoints using two databases, ECOTOX and BIOSIS. Studies with aquatic invertebrate species were preferred because the purpose of the porewater study was to evaluate risks to benthic invertebrates. In many cases, the available toxicity data were uncertain because lowest-observed-effect concentrations (LOECs) were rarely reported, NOECs were dependent on the selected test dilution series, and very few studies reported both effect and no-effect concentrations for a single species and endpoint. Most of the LOECs were based on LC50s (concentrations that result in the death of 50% of a test population). No uncertainty factors were assigned to the toxicity data. In addition, some of the exposure scenarios in the studies were of relatively short duration (48 to 96 hr). Use of these toxicity data may underestimate risks to benthic invertebrates in the LDW. Table A.2-14 presents the TRVs that were selected for 15 of the 16 VOCs. No toxicity data were available for isopropylbenzene. A discussion of all TRVs selected and all toxicological studies evaluated during the TRV search is presented in Attachment 7.

All maximum VOC concentrations in porewater were less than NOECs, except for cis-1,2-dichloroethene (Table A.2-14). The NOEC for cis-1,2-dichloroethene was derived by dividing the LOEC (an LC50 [concentration that causes the death of 50% of a group of test animals]) by 50 because a NOEC was not reported in the literature. Because the maximum concentration of cis-1,2-dichloroethene was greater than the calculated NOEC, it was retained as a COPC.

Table A.2-14. Identification of COPCs for benthic invertebrates in porewater

COPC	Concentration in Porewater		
	Maximum Detected Concentration (µg/L)[a]	NOEC (µg/L)	LOEC (µg/L)[b]
1,1-Dichloroethane	16	7,800[c]	39,600[d]
1,1-Dichloroethene	4.9	2,400	11,600
1,2-Dichlorobenzene	1.2	11[e]	550[f]
1,2-Dichloroethane	15	139[e]	6,927
1,2-Dichloropropane	2.5	840[e]	42,000
1,4-Dichlorobenzene	0.3	3[e]	147
Benzene	9.4	180	1,100[g]
Carbon disulfide	0.7	38[e]	1,900
Chlorobenzene	1.4	1,400	2,500[g]
cis-1,2-Dichloroethene	**2,900**	136[e]	6,785
Isopropylbenzene	0.3	na	na
Tetrachloroethene	1.1	331	332
Toluene	3.5	737	14,700
trans-1,2-Dichloroethene	21 J	136[e]	6,785
Trichloroethene	2.5	2,200	14,000
Vinyl chloride	2,500	12,800[c]	65,300[d]

[a] The maximum detected concentration in the 20 porewater samples analyzed as part of the Phase 2 RI.

[b] Concentration is based on a LC50 (concentration that causes the death of 50% of a group of test animals) unless otherwise noted.

[c] Concentration is a final chronic value (FCV) based on the narcosis model.

[d] Concentration is a final acute value (FAV) based on the narcosis model.

[e] A literature-based NOEC was not available; the NOEC was derived by dividing the LC50 by 50.

[f] Concentration is an EC50 (concentration causing a non-lethal effect in 50% of an exposed population).

[g] Concentration is an LC100 (concentration that causes the death of 100% of a group of test animals).

COPC – chemical of potential concern

HQ – hazard quotient

LOEC – lowest-observed-effect concentration

na – not available

NOEC – no-observed-effect concentration

Bold identifies maximum porewater concentrations greater than the NOEC.

A.2.5.1.3 Crabs

COPCs were identified for crabs using a critical tissue-residue approach and a two-step process. The first step was to identify a list of chemicals of interest (COIs). Chemicals were identified as COIs for crabs if they met at least two of the following three criteria:

- Detection in at least 5% of LDW surface sediment samples
- Identification as a bioaccumulative chemical in EPA (2000)
- Detection in any LDW-collected crab tissue sample

Table A.2-15 presents a summary of the metals, organometals, and organic compounds identified as COIs for crabs. Detailed results of the COI screening step for crabs are presented in Attachment 4.

Table A.2-15. Chemicals identified as COIs for crabs

COIs		
Metals		
Antimony	Copper	Selenium
Arsenic	Lead	Silver
Cadmium	Mercury	Thallium
Chromium	Methylmercury[a]	Vanadium
Chromium VI[a]	Molybdenum	Zinc
Cobalt	Nickel	
Organometals		
Monobutyltin as ion	Dibutyltin as ion	TBT as ion
PAHs		
2-Methylnaphthalene	Benzo(b)fluoranthene	Fluoranthene
Acenaphthene	Benzo(g,h,i)perylene	Fluorene
Acenaphthylene	Benzo(k)fluoranthene	Indeno(1,2,3-cd)pyrene
Anthracene	Chrysene	Naphthalene
Benzo(a)anthracene	Dibenzo(a,h)anthracene	Phenanthrene
Benzo(a)pyrene	Dibenzofuran	Pyrene
Phthalates		
Bis(2-ethylhexyl) phthalate	Diethyl phthalate	Di-n-butyl phthalate
Butyl benzyl phthalate	Dimethyl phthalate	
Other SVOCs		
Benzyl alcohol	Phenol	
Hexachlorobenzene		
PCBs		
Total PCBs		
Dioxins and Furans		
Organochlorine Pesticides		
Total DDTs[b]	gamma-BHC	Heptachlor epoxide
Dieldrin	alpha-Endosulfan	Methoxychlor
alpha-BHC	Endrin aldehyde	Total chlordane[c]

[a] Chromium VI and methylmercury (both bioaccumulative chemicals and detected in at least 5% of surface sediment samples) were not analyzed in any of the tissue samples; these chemicals are assumed to be incorporated in total mercury and total chromium risk analyses.

[b] Includes 2,4-DDE, 2,4-DDD, 2,4-DDT, 4,4-DDD, 4,4-DDE, and 4,4-DDT.

[c] Includes alpha-chlordane and gamma-chlordane.

COI – chemical of interest

PCB – polychlorinated biphenyl

SVOC – semivolatile organic compound

TBT – tributyltin

In the second step of the screening process, the maximum exposure concentration of each COI was compared to a no-observed-adverse-effect level (NOAEL) for that chemical. If the maximum exposure concentration was greater than the NOAEL, the chemical was identified as a COPC for crabs. This step was conducted for all COIs, except dioxins and furans. Dioxins and furans were identified as COIs for crabs because they were detected in at least 5% of LDW surface sediment samples and are bioaccumulative (Attachment 4). However, quantitative ecological risk analysis was not conducted for dioxins and furans in this baseline ERA, as described in Section A.2.5.

For the remaining COIs, the scientific literature was searched[18] to identify TRVs for COIs. The literature search included BIOSIS, EPA's ECOTOX database, aquatic life sciences database, USACE's Environmental Residue Effects Database (ERED), and Jarvinen and Ankley (1999). Original sources of toxicity data were obtained and reviewed to verify effects data summarized in the databases as well as the suitability of the studies. The databases were searched for studies that evaluated effects on survival, growth, and reproduction (including developmental effects).

TRVs were based on the risk evaluation method used. For crabs, the TRV search focused on chemical tissue-residue data associated with effects on decapods to support the critical tissue-residue approach (see Section A.3.4). For critical tissue-residue studies to be acceptable, the concentration in tissue had to be analyzed as part of the study. Acceptable toxicological data that met the following criteria were compiled for crabs.

- All selected TRVs were based on laboratory toxicological studies. Studies using field-collected data (i.e., field-collected crabs) were not considered acceptable. Field studies were not used to derive TRVs because adverse effects observed in organisms from field studies may be attributed to the presence of multiple chemicals and/or other uncontrolled environmental factors, rather than to a single test chemical.
- Selected TRVs were based preferentially on dietary, sediment, or water exposure studies.

After the literature search was conducted, all acceptable studies for TRV derivation were compiled, and presented in Attachment 5.

For each COI, a TRV was selected for both the NOAEL and the lowest-observed-adverse-effect level (LOAEL). The NOAEL represents the level below which adverse effects would not be expected. The NOAEL was compared to the maximum exposure concentration for each COI to identify COPCs. The LOAEL represents the level above

[18] For COIs that were evaluated in the Phase 1 ERA (Windward 2003b), literature published after January 2001 was searched for new toxicity data. If any COIs were not evaluated in the Phase 1 ERA, the literature was searched for appropriate toxicity data and was not limited by date.

which an effect would be expected. LOAELs are presented in this section for informational purposes only; they were not used in the screening process.

The LOAEL was selected from among the list of possible TRVs presented in Attachments 5 through 10 if it was the lowest dose at which an effect was observed for any of the three endpoints evaluated and a clear dose-response relationship was observed. The NOAEL was selected as the highest level below the selected LOAEL with the same endpoint. If no NOAEL with the same endpoint as the selected LOAEL was available, the NOAEL was selected as the highest NOAEL below the selected LOAEL based on another endpoint (survival, growth, or reproduction).

For COIs without NOAELs lower than the selected LOAEL, the NOAEL was determined using the following uncertainty factors following EPA Region 10 guidance (EPA 1997b):

- Acute or subchronic LOAEL/10
- Chronic or critical lifestage[19] LOAEL/5
- LC50 (or similar)/50

Based on the available literature, crab critical tissue-residue TRVs were developed for chemicals identified as COIs and detected in crab tissue according to the guidelines outlined above. TRVs were available for 14 of the 55 COIs (Table A.2-16). No studies with reproductive endpoints were identified; all acceptable studies addressed either growth or survival. TRVs selected from the acceptable studies are presented in Table A.2-17. Chemicals with no TRVs are discussed in the uncertainty analysis (Section A.6.1.2.2).

To identify COPCs for crabs, critical tissue-residue NOAELs were compared to maximum chemical concentrations detected in the corresponding LDW crab tissue samples (e.g., hepatopancreas NOAELs were compared with the maximum concentration detected in hepatopancreas tissue; whole-body NOAELs were compared to the estimated maximum whole-body concentration, as calculated from hepatopancreas and edible muscle tissue concentrations[20]). NOAELs were available in the literature for 10 COIs. NOAELs were estimated for naphthalene, PCBs, and methoxychlor as the LOAEL (based on survival data) divided by 10.

---

[19] Chronic exposure is defined as >15% of an organism's lifespan (Calabrese and Baldwin 1993). Exposure is assumed to be chronic if the duration is greater than 10 weeks for birds and greater than one year for mammals (Sample et al. 1996). For fish, chronic exposure duration was assumed to be 28 days or greater. A critical lifestage is one that occurs during reproduction, gestation, or development (Sample et al. 1996).

[20] Chemical concentrations in hepatopancreas tissue are likely to provide a conservative estimate of internal organs in general because the hepatopancreas constitutes the great majority of organ mass and has a relatively high lipid content relative to other organs.

Table A.2-16. Results of TRV search for chemicals identified as crab COIs through the initial screening process and detected in crab tissue

COIs for Crabs		
Chemicals with TRVs		
Arsenic	Vanadium	Methoxychlor
Cadmium	TBT	Total chlordane[a]
Chromium	Zinc	Total DDTs[b]
Copper	Naphthalene	Total PCBs[c]
Mercury	Heptachlor epoxide	
Chemicals without TRVs		
Antimony	Di-n-butyl phthalate	Dibenzo(a,h)anthracene
Cobalt	2-Methylnaphthalene	Dibenzofuran
Lead	Benzyl alcohol	Fluoranthene
Molybdenum	Hexachlorobenzene	Fluorene
Nickel	Phenol	Indeno(1,2,3-cd)pyrene
Monobutyltin	Acenaphthene	Phenanthrene
Dibutyltin	Acenaphthylene	Pyrene
Selenium	Anthracene	alpha-BHC
Silver	Benzo(a)anthracene	gamma-BHC
Thallium	Benzo(a)pyrene	Dieldrin
Bis(2-ethylhexyl) phthalate	Benzo(b)fluoranthene	alpha-Endosulfan
Butyl benzyl phthalate	Benzo(g,h,i)perylene	Endrin aldehyde
Diethyl phthalate	Benzo(k)fluoranthene	
Dimethyl phthalate	Chrysene	

[a] Alpha-chlordane and gamma-chlordane were detected in crab tissue. The two chemicals were assessed as total chlordane.

[b] 2,4-DDD, 2,4-DDE, 2,4-DDT, 4,4-DDD, 4,4-DDE, and 4,4-DDT were detected in crab tissue. The five chemicals were assessed as total DDTs.

[c] Aroclor-1248, Aroclor-1254, and Aroclor-1260 were detected in crab tissue, and assessed as total PCBs.

COI – chemical of interest

PCB – polychlorinated biphenyl

TBT – tributyltin

TRV – toxicity reference value

Table A.2-17. Selected critical tissue-residue TRVs for crab COIs

COI	Test Species	Tissue Type	NOAEL	LOAEL	Units	Endpoint	Source
Arsenic	juvenile grass shrimp *(Palaemonetes pugio)*	whole body	1.28 [a]	na	mg/kg ww	growth	Lindsay and Sanders (1990)
Cadmium	grass shrimp *(Palaemonetes pugio)*	whole body	0.6 [a]	na	mg/kg ww	survival	Rule and Alden (1996)
	grass shrimp *(Palaemonetes pugio)*	whole body	na	2.6[a]	mg/kg ww	survival	Vernberg et al. (1977)
Chromium	juvenile (2nd instar) sand crab (*Portunus pelagicus*)	whole body	1	3.2	mg/kg ww	growth	Mortimer and Miller (1994)
Copper	adult crayfish (*Orconectes rusticus*)	whole body	50[a]	na	mg/kg ww	survival	Evans (1980)
Mercury	adult Norway lobster (*Nephrops norvegicus*)	hepatopancreas	0.99[a,b,]	na	mg/kg ww	survival	Canli and Furness (1995)
	adult male shore crab (*Eriocheir sinensis*)	hepatopancreas	na	1[c,d]	mg/kg ww	survival	Bianchini and Giles (1996)
Vanadium	shrimp (*Lysmata seticaudata*)	whole body	0.6	na	mg/kg ww	survival	Miramand et al. (1981)
Zinc	crayfish (*Orconectes virilis*)	whole body	12.7[a]	35.2[a]	mg/kg ww	survival	Mirenda (1986a)
TBT	juvenile blue crab (*Callinectes sapidus*)	whole body	120	na	µg/kg dw	growth	Rice et al. (1989)
Naphthalene	spot shrimp (*Pandalus platyceros*)	whole body	5.0[e]	50	µg/kg ww	survival	Sandborn and Malins (1977)
PCBs (Aroclor 1016)	grass shrimp (*Palaemonetes pugio*)	whole body	110[e]	1,100[f]	µg/kg ww	survival	Hansen et al. (1974b)
Total chlordane	pink shrimp (*Penaeus duorarum*)	whole body	710	1,700	µg/kg ww	survival	Parrish et al. (1976)
Total DDTs	pink shrimp (*Penaeus duorarum*)	whole body	na	60	µg/kg ww	survival	Nimmo et al. (1970)
	crayfish (*Orconectes nais*)	whole body	46	na	µg/kg ww	survival	Johnson et al. (1971)
Heptachlor epoxide	pink shrimp (*Penaeus duorarum*)	whole body	54	180	µg/kg ww	survival	Schimmel et al. (1976)
Methoxychlor	juvenile Dungeness crab (*Cancer magister*)	whole body	15[e]	150	µg/kg ww	survival	Armstrong et al. (1976)

[a] Converted from dry weight to wet weight using a moisture content of 80% (Jarvinen and Ankley 1999) (80% was also the average moisture content of two crab samples collected by King County in 1997 (King County 1999b); the mean % moisture in LDW crabs was 82.7% (Windward 2005c)).

[b] Dietary exposure route

[c] Full equilibrium between water and tissue may not have been reached because of a short exposure time (≤ 48 hrs).

[d] Concentration is lowest of three crab species tested (*Carcinus maenas*, *Eriocheir sinensis*, and *Cancer pagurus*).

[e] Calculated from LOAEL by dividing by 10.

[f] Survival was reduced by 33%.

dw – dry weight
PCB – polychlorinated biphenyl
ww – wet weight
na – not available
TBT – tributyltin

Two chemicals, zinc and PCBs, were identified as COPCs for crabs (Table A.2-18) and retained for further evaluation in this baseline ERA. The maximum concentrations of arsenic, total DDTs, and methoxychlor also exceeded NOAELs for these COIs, but were not identified as COPCs for the following reasons. No LOAEL TRV was identified for arsenic. Because the arsenic NOAEL is not bounded by a LOAEL, this assessment does not indicate a potential threshold above which adverse effects may occur. The two organochlorine pesticides were not identified as COPCs because of the uncertainty associated with JN-qualified organochlorine pesticide tissue data (see Section A.2.4.2.2). Therefore, arsenic, total DDTs, and methoxychlor will be discussed further in the uncertainty analysis (Section A.6.1.2.2). A summary of the COPCs that will be evaluated for benthic invertebrates and crabs in the risk characterization is presented in Table A.2-19.

Table A.2-18. Maximum concentrations of COIs in crab tissue samples compared with NOAEL TRVs

COI	Unit	Maximum Concentration in Crab Tissue[a]	NOAEL TRV
Arsenic	mg/kg ww	**11 M**	1.28
Cadmium	mg/kg ww	0.2951 M	0.6
Chromium	mg/kg ww	0.14 M	1
Copper	mg/kg ww	24 M	50
Mercury	mg/kg ww	0.067[b]	0.99
Vanadium	mg/kg ww	0.2 JM	0.6
Zinc	mg/kg ww	**37.3 M**	12.7
TBT	µg/kg ww	75 M	120
Naphthalene	µg/kg ww	3.2 M	5.0[c]
Total PCBs	µg/kg ww	**1,900 JM**	110[c]
Total chlordane	µg/kg ww	26 JNM	710
Total DDTs	µg/kg ww	**150 JNM**	46
Heptachlor epoxide	µg/kg ww	5.5 JNM	54
Methoxychlor	µg/kg ww	**90 JNM**	15[c]

[a] All whole-body concentrations except as noted.

[b] Hepatopancreas

[c] Calculated from LOAEL by dividing by 10.

COI – chemical of interest

J – estimated concentration

M – estimated concentration based on a weighted mean of hepatopancreas and edible meat data

N – tentative identification

NOAEL – no-observed-adverse-effect level

PCB – polychlorinated biphenyl

TBT – tributyltin

TRV – toxicity reference value

ww – wet weight

Bold identifies maximum crab tissue concentrations greater than the NOAEL.

Table A.2-19. COPCs evaluated in the risk characterization for the benthic invertebrate community and for crabs

COPC	Benthic Invertebrate Community	Crabs
Metals		
Arsenic	X	
Cadmium	X	
Chromium	X	
Copper	X	
Lead	X	
Mercury	X	
Nickel	X	
Silver	X	
Zinc	X	X
Organometals		
TBT	X	
PAHs		
Acenaphthene	X	
Anthracene	X	
Benz(a)anthracene	X	
Benzo(a)pyrene	X	
Benzo(g,h,i)perylene	X	
Chrysene	X	
Dibenzo (a,h)anthracene	X	
Fluoranthene	X	
Fluorene	X	
Indeno (1,2,3,-c,d)pyrene	X	
Naphthalene	X	
Phenanthrene	X	
Pyrene	X	
Total benzofluoranthenes	X	
HPAH	X	
LPAH	X	
Phthalates		
Bis(2-ethylhexyl) phthalate	X	
Butyl benzyl phthalate	X	
Dimethyl phthalate	X	

COPC	Benthic Invertebrate Community	Crabs
Other SVOCs		
1,2-Dichlorobenzene	X	
1,4-Dichlorobenzene	X	
1,2,4-Trichlorobenzene	X	
2-Methylnaphthalene	X	
4-Methylphenol	X	
2,4-Dimethylphenol	X	
Benzoic acid	X	
Benzyl alcohol	X	
Dibenzofuran	X	
Hexachlorobenzene	X	
n-Nitrosodiphenylamine	X	
Pentachlorophenol	X	
Phenol	X	
VOCs		
cis-1,2-Dichloroethene	X	
PCBs		
Total PCBs	X	X
Organochlorine Pesticides		
Total DDTs	X	
Total chlordane	X	

COPC – chemical of potential concern
PCB – polychlorinated biphenyl
SVOC – semivolatile organic compound
TBT – tributyltin
VOC – volatile organic compound

A.2.5.2 Fish

COPCs were identified for fish using a two-step process. The first step was to identify a list of COIs. Chemicals were identified as COIs for fish if they met at least two of the following three criteria:

- Detection in at least 5% of LDW surface sediment samples
- Identification as a bioaccumulative chemical in EPA (2000)
- Detection in any LDW-collected tissue sample

Table A.2-20 presents a summary of chemicals identified as COIs as a result of this screen. Detailed results of the COI screening step for fish are presented in Attachment 4.

Table A.2-20. Chemicals identified as COIs for fish ROCs

COIs		
Metals		
Antimony	Copper	Selenium
Arsenic	Lead	Silver
Cadmium	Mercury	Thallium
Chromium	Methylmercury[a]	Vanadium
Chromium VI[a]	Molybdenum	Zinc
Cobalt	Nickel	
Organometals		
Monobutyltin as ion	TBT as ion	
Dibutyltin as ion	Tetrabutyltin as ion	
PAHs		
C1-Chrysenes	C3-Phenanthrenes/anthracenes	Benzo(g,h,i)perylene
C1-Dibenzothiophenes	C4-Naphthalenes	Benzo(k)fluoranthene
C1-Fluoranthene/pyrene	C4-Phenanthrenes/anthracenes	Chrysene
C1-Fluorenes	1-Methylnaphthalene	Dibenzo(a,h)anthracene
C1-Phenanthrenes/anthracenes	2-Methylnaphthalene	Dibenzofuran
C2-Chrysenes	Acenaphthene	Fluoranthene
C2-Dibenzothiophenes	Acenaphthylene	Fluorene
C2-Fluorenes	Anthracene	Indeno(1,2,3-cd)pyrene
C2-Naphthalenes	Benzo(a)anthracene	Naphthalene
C2-Phenanthrenes/anthracenes	Benzo(a)pyrene	Perylene
C3-Fluorenes	Benzo(b)fluoranthene	Phenanthrene
C3-Naphthalenes	Benzo(e)pyrene	Pyrene
Phthalates		
Bis(2-ethylhexyl) phthalate	Diethyl phthalate	Di-n-butyl phthalate
Butyl benzyl phthalate	Dimethyl phthalate	
Other SVOCs		
4-Chlorophenyl phenyl ether	Biphenyl	Hexachlorobenzene
4-Methylphenol	Carbazole	Phenol
Benzoic acid	Dibenzothiophene	
PCBs		
Total PCBs		
Dioxins and Furans		

COIs		
Organochlorine Pesticides		
Total DDTs[b]	delta-BHC	Heptachlor
Aldrin	gamma-BHC	Heptachlor epoxide
Dieldrin	alpha-Endosulfan	Methoxychlor
alpha-BHC	beta-Endosulfan	Total chlordane[c]
beta-BHC	Endrin	

[a] Chromium VI and methylmercury (both bioaccumulative chemicals and detected in at least 5% of surface sediment samples) were not analyzed in any of the tissue samples; these chemicals are assumed to be incorporated in total mercury and total chromium risk analyses.

[b] Includes 2,4'-DDD, 2,4'-DDE, 2,4'-DDT, 4,4'-DDD, 4,4'-DDE, and 4,4'-DDT.

[c] Includes alpha-chlordane, gamma-chlordane, oxychlordane, cis-nonachlor, and trans-nonachlor.

COI – chemical of interest

PCB – polychlorinated biphenyl

SVOC – semivolatile organic compound

TBT – tributyltin

In the second step of the screening process, the maximum exposure concentration of each COI was compared to a NOAEL for that chemical. If the maximum exposure concentration was greater than the NOAEL, the chemical was identified as a COPC for fish. This step was conducted for all COIs, except dioxins and furans, as discussed in Section A.2.5.

For COIs other than dioxins and furans, the scientific literature was searched[21] to identify TRVs for COIs. The literature search included BIOSIS, EPA's ECOTOX database, aquatic life sciences database, USACE's ERED, and Jarvinen and Ankley (1999). Original sources of toxicity data were obtained and reviewed to verify effects data summarized in the databases as well as the suitability of the studies. The databases were searched for studies that evaluated effects on survival, growth, and reproduction (including developmental effects).

TRVs were based on the risk evaluation method used. For fish, databases were searched for dietary studies for metals and PAHs and for critical tissue-residue data for other chemicals (see Section A.4.1 for further explanation of the two methods used). For critical tissue-residue studies to be acceptable, the concentration in tissue had to be analyzed as part of the study. Acceptable toxicological data that met the following criteria were compiled for fish.

- All selected TRVs were based on laboratory toxicological studies. Studies using field-collected data (i.e., field-collected fish or fish fed field-collected diets) were not considered acceptable. Field studies were not used to derive TRVs because

[21] For COIs that were evaluated in the Phase 1 ERA (Windward 2003b), toxicity literature published after January 2001 was searched for new toxicity data. If any COIs not evaluated in Phase 1 ERA, toxicity literature was searched for appropriate toxicity data and was not limited by date.

adverse effects observed in organisms from field studies may be attributed to the presence of multiple chemicals and/or other uncontrolled environmental factors, rather than to a single test chemical.[22]

- Selected TRVs were based preferentially on dietary, sediment, or water exposure studies. Studies conducted using intraperitoneal (IP) or egg injection or oral gavage as exposure routes were not considered representative of the ROC exposure conditions but were used if no other studies were available.
- All selected TRVs were based on whole-body tissue concentrations or egg concentrations that were converted to adult tissue concentrations using adult-to-egg conversion factors from the literature.

After the literature search was conducted, all acceptable studies for TRV derivation were compiled, as presented in Attachment 8. For each COI, TRVs were selected for both the NOAEL and the LOAEL. TRV selection rules and uncertainty factors discussed above for crab in Section A.2.5.1.3 were used for fish as well.

TRVs were available for 45 of the 86 fish COIs (Table A.2-21). Selected TRVs are presented in Tables A.2-22 and A.2-23 for COIs evaluated using the dietary approach and critical tissue-residue approach, respectively. Chemicals with no TRVs are discussed in the uncertainty analysis (Section A.6.2.1.2).

Table A.2-21. Results of TRV search for fish COIs

COIs for Fish		
Chemicals with TRVs		
Arsenic	Benzo(k)fluoranthene[a]	beta-Endosulfan
Cadmium	Chrysene[a]	Total DDTs
Chromium	Dibenzo(a,h)anthracene[a]	Dieldrin
Copper	Fluoranthene[a]	Endrin
Lead	Fluorene[a]	gamma-BHC (lindane)
Selenium	Indeno(1,2,3-cd)pyrene[a]	Heptachlor
Silver	Naphthalene[a]	Heptachlor epoxide
Vanadium	Phenanthrene[a]	Methoxychlor
Zinc	Pyrene[a]	Total chlordane
2-Methylnaphthalene[a]	Mercury	Bis(2-ethylhexyl) phthalate
Acenaphthene[a]	TBT as ion	Butyl benzyl phthalate
Anthracene[a]	4-Methylphenol	Diethyl phthalate
Benzo(a)anthracene[a]	Benzoic acid	Dimethyl phthalate
Benzo(a)pyrene[b]	Hexachlorobenzene	Di-n-butyl phthalate
Benzo(b)fluoranthene[a]	Phenol	Total PCBs

[22] The uncertainty associated with not including TRVs derived from field-collected data is explored in the uncertainty analysis subsections of the risk characterization (Section A.6.2).

COIs for Fish		
Benzo(g,h,i)perylene[a]	alpha-Endosulfan	
Chemicals without TRVs		
Antimony	C1-Fluoranthene/Pyrene	Dibutyltin as ion
Chromium VI	C1-Fluorenes	Monobutyltin as ion
Cobalt	C1-Phenanthrenes/anthracenes	Tetrabutyltin as ion
Molybdenum	C2-Chrysenes	4-Chlorophenyl phenyl ether
Nickel	C2-Dibenzothiophenes	Biphenyl
Thallium	C2-Fluorenes	Carbazole
1-Methylnaphthalene	C2-Naphthalenes	Dibenzothiophene
Acenaphthylene	C2-Phenanthrenes/Anthracenes	Aldrin
Benzo(e)pyrene	C3-Fluorenes	alpha-BHC
Dibenzofuran	C3-Naphthalenes	beta-BHC
Perylene	C3-Phenanthrenes/Anthracenes	delta-BHC
C1-Chrysenes	C4-Naphthalenes	
C1-Dibenzothiophenes	C4-Phenanthrenes/Anthracenes	

[a] PAHs included in the mixtures of PAHs evaluated by Palm et al. (2003) or Meador et al. (2006).

[b] Benzo(a)pyrene was included in the mixtures of PAHs evaluated by Palm et al. (2003) and Meador et al. (2006); TRVs for exposure to benzo(a)pyrene as a single chemical were also available.

COI – chemical of interest

PAH – polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

TBT – tributyltin

TRV – toxicity reference value

Table A.2-22. TRVs selected for fish COIs evaluated using a dietary approach

COI	Test Species	NOAEL (mg/kg dw)	LOAEL (mg/kg dw)	Endpoint	Source
Arsenic	rainbow trout	20	30	growth	Oladimeji et al. (1984)
Cadmium	rockfish	0.1[a]	0.5	growth	Kim et al. (2004); Kang et al. (2005)
Chromium	grey mullet	9.42	na	growth	Walsh et al. (1994)
Copper	rainbow trout	50	100	growth	Kang et al. (2005)
Lead	rainbow trout	7,040	na	growth	Goettl et al. (1976)
Silver	rainbow trout	3,000	na	growth	Galvez and Wood (1999)
Vanadium	rainbow trout	2.04[a]	10.2	growth	Hilton and Bettger (1988)
Zinc	rainbow trout	1,900	nr	growth	Mount et al. (1994)
	rainbow trout	nr	2,000	growth	Takeda and Shimma (1977)
Benzo(a)pyrene	rainbow trout	100	nr	growth	Hart and Heddle (1991)
	English sole	nr	116	growth	Rice et al. (2000)
Total PAHs[c]	chinook salmon	324	951	growth	Meador et al. (2006)

[a] NOAEL estimated using an uncertainty factor of 5 (chronic LOAEL to NOAEL).

[b] NOAEL estimated using an uncertainty factor of 10 (acute/subchronic LOAEL to NOAEL).

[c] Mixture comprises the following 21 PAHs included in Meador et al. (2006) diet: naphthalene, 2-methylnaphthalene, dimethylnaphthalene, dibenzothiophene, acenaphthene, fluorene, 1,8-dimethyl(9H)fluorene, phenanthrene, 9-ethylphenanthrene, 9-ethyl-10-methylphenanthrene, 1-methyl-7-isopropylphenanthrene, anthracene, fluoranthene, pyrene, methyl pyrene, benzo(a)anthracene, chrysene, benz(a)pyrene, benzo(k)fluoranthene, benzo(g,h,i)perylene, dibenzanthracene.

COI – chemical of interest

dw – dry weight

LOAEL – lowest-observed-adverse-effect level

na – not available; no LOAELs identified in the literature search; selected NOAEL is the highest unbounded NOAEL in the literature reviewed.

NOAEL – no-observed-adverse-effect level

nr – not relevant; NOAEL and LOAEL TRVs were derived from separate studies reporting the same endpoint.

PAH – polycyclic aromatic hydrocarbon

Table A.2-23. TRVs selected for fish COIs evaluated using the critical tissue-residue approach

COI	Test Species	NOAEL (µg/kg ww)	LOAEL (µg/kg ww)	Endpoint	Source
Mercury[a]	golden shiner	230	nr	survival	Webber and Haines (2003)
	mummichog	nr	470	survival	Matta et al. (2001)
Selenium	[national criterion]	1,200[b]	1,600[c]	adverse effects	EPA (2004b)
TBT	Japanese flounder	18	159	growth	Shimasaki et al. (2003)
Bis(2-ethylhexyl) phthalate	rainbow trout	390[d]	na	reproduction	Mehrle and Mayer (1976)
Butyl benzyl phthalate	bluegill	6,450	na	survival	Barrows et al. (1980)
Di(n)butyl phthalate	sheepshead minnow	1,170	na	survival	Wofford et al. (1981)
Dimethyl phthalate	bluegill	498	na	survival	Barrows et al. (1980)
Diethyl phthalate	bluegill	1,102	na	survival	Barrows et al. (1980)
4-methylphenol	rainbow trout	1,530[e]	76,500	survival	Kaiser et al. (1984)
Benzoic acid	mosquito fish	3,380	na	survival	Lu and Metcalf (1975)
Hexachlorobenzene	fathead minnow	468,000	na	survival	Schuytema et al. (1990)
Phenol	rainbow trout	1,470[e]	73,400	survival	McKim and Schmeider (1990)
PCBs	common barbel	104 – 528[f]	520 – 2,640[f]	reproduction	Hugla and Thome (1999)
alpha-Endosulfan	spot	0.62[e]	31	survival	Schimmel et al. (1977a)
beta-Endosulfan	spot	0.62[e]	31	survival	Schimmel et al. (1977a)
DDTs (total)	cutthroat trout	1,800[g]	1,800[g]	survival	Allison et al. (1964)
Dieldrin	rainbow trout	120	200	survival	Shubat and Curtis (1986)
Endrin	largemouth bass	1.15[h]	11.5	survival	Fabacher (1976)
gamma-BHC (lindane)	sheepshead minnow	1,580[e]	79,000	survival	Schimmel et al. (1977b)
Heptachlor	spot	30[e]	1,500	survival	Schimmel et al. (1976)
Heptachlor epoxide	bluegill	80[h]	800	growth	Andrews et al. (1966)
Methoxychlor	brook trout	50	300	growth	Oladimeji and Leduc (1975)
Total chlordane	goldfish	710	nr	survival	Moore et al. (1977)
	goldfish	nr	1,360	survival	Feroz and Khan (1979)

[a] TRVs were based on exposure to methylmercury. Methylmercury was not analyzed in any of the tissue samples from the LDW; the total mercury concentration in LDW tissue samples was compared to the methylmercury TRV.

[b] National criterion for selenium in summer-collected fish. Dry weight concentration converted to wet weight assuming 80% moisture content.

[c] National criterion for selenium in winter-collected fish. Dry weight concentration converted to wet weight assuming 80% moisture content.

[d] Fry concentration calculated using a bioconcentration factor and water concentration reported in Mehrle and Mayer (1976).

[e] NOAEL estimated using an uncertainty factor of 50 (LC50 to NOAEL).

[f] A LOAEL range was selected from this study because the specific LOAEL was unclear because of uncertainties associated with this study. The NOAEL range was estimated using an uncertainty factor of 5 (chronic LOAEL to NOAEL) .

[g] The LOAEL is tissue concentration at 111 days (3.7 months) in fish exposed to 0.1 mg/L DDT in water where significant mortality occurred after approximately 4 months (approximately 120 days). The NOAEL (1,800 µg/kg ww) is the highest tissue concentration (at 466 days) in fish exposed to 0.03 mg/L DDT in water at which significant mortality did not occur over the entire exposure duration.

[h] NOAEL estimated using an uncertainty factor of 10 (acute/subchronic LOAEL to NOAEL).

COI – chemical of interest

na – not available; no LOAELs identified in the literature search; selected NOAEL is the highest unbounded NOAEL in the literature reviewed.

nr – not relevant; NOAEL and LOAEL TRVs were derived from separate studies reporting the same endpoint.

PCB – polychlorinated biphenyl

TBT – tributyltin

TRV – toxicity reference value

ww – wet weight

COPCs were screened for fish ROCs using one of two approaches depending on the type of chemical. For chemicals evaluated using a critical tissue-residue approach, maximum detected concentrations of COIs in the whole-body tissue of fish ROCs were compared with effects data reported in the literature. Use of the critical tissue-residue approach integrates all exposure pathways and reduces uncertainty associated with the relative uptake and depuration rates of a chemical. For COPCs that are highly regulated or metabolized by fish, such as PAHs (Varanasi 1989) and most metals (Bury et al. 2003, as cited in Meyer et al. 2005), comparison of chemical concentrations in prey to suitable dietary TRVs is preferable. Therefore, a dietary approach was used to identify COPCs for PAHs and metals (except butyltins, mercury, and selenium). For PAHs and metals, the concentration in the diet more accurately reflects the toxic dose than whole-body tissue residues.

For chemicals evaluated using a dietary approach, maximum detected concentrations of COIs in fish diets (expressed as mg/kg dw) were compared with dietary exposure concentrations reported in the literature (expressed as mg/kg dw). An alternative way of expressing exposure that would have explicitly considered an ROC's rate of chemical uptake would have involved the calculation of "dietary dose" expressed as µg/g fish/day. Because fish prey consumption is variable, use of a dietary dose approach is becoming more prevalent as a way to normalize dietary exposure among species (e.g., Clearwater et al. 2002). This method could also be used to predict a total dose from both water and dietary exposure, although little progress has been made in this regard (e.g., Borgmann et al. 2005). Because use of a dose-based approach for the purpose of estimating effects from dietary exposure is in its infancy, components of dose (such as ration size, feeding frequency, and food wastage) are often not reported in toxicity papers. Therefore, it is difficult to estimate accurate doses from available effects data. In addition, daily food consumption rates are not standardized for fish species as they are for wildlife, making fish dietary dose exposure calculations uncertain. Therefore, a dietary concentration approach rather than a dietary dose approach was used as the dietary approach for fish exposure and effects calculations in this ERA.

To screen COIs using the critical tissue-residue approach, maximum COI concentrations in ROC fish species were compared to NOAELs from the literature. This screen was conducted in two parts. First, the maximum fish tissue concentration for each COI in any fish ROC species was compared to its respective NOAEL. COIs with maximum tissue concentrations greater than the NOAEL were identified as COPCs. Second, a species-specific screen was conducted to determine which ROC/COPC pairs would be evaluated in more detail in the exposure and effects assessment (Section A.5.0) and which pairs would not warrant additional analysis.

Maximum concentrations (or RLs for non-detected COIs) of nine COIs in LDW fish tissue were greater than their respective NOAEL TRVs (Table A.2-24). Of these chemicals, LOAEL TRVs were not identified for benzoic acid, dimethyl phthalate, or

di-n-butyl phthalate. Because the NOAELs for these chemicals are not bounded by LOAELs, they do not indicate potential thresholds above which adverse effects may occur. Therefore, risks associated with these chemicals are discussed in the uncertainty analysis (Section A.6.2).

Table A.2-24. Maximum COI concentrations in any fish ROC tissue compared to NOAEL TRVs

COI	Maximum Chemical Concentration in Fish ROC Tissue (µg/kg ww)	NOAEL TRV (µg/kg ww)
Mercury	39	230
Selenium	320	1,200
TBT as ion	**80**	18
Bis(2-ethylhexyl) phthalate	**5,000 U**[a]	390
Butyl benzyl phthalate	1,400	6,450
Diethyl phthalate	900 J	1,102
Dimethyl phthalate	**580 U**	498
Di-n-butyl phthalate	**2,300**	1,170
4-Methylphenol	380 J	1,530
Benzoic acid	**6,800 J**	3,380
Hexachlorobenzene	6.6 J	468,000
Phenol	200 J	1,470
Total PCBs	**4,700**	104 – 528
alpha-Endosulfan	**6.6 J**	0.62
beta-Endosulfan	**18 J**	0.62
Total DDTs	280 J	1,800
Dieldrin	5.7	120
Endrin	**36**	1.15
gamma-BHC (lindane)	5.6 J	1,580
Heptachlor	6.8 J	30
Heptachlor epoxide	45 J	80
Methoxychlor	0.49	50
Total chlordane	59 J	710

[a] RL is elevated because of analytical dilutions. A subset of 49 fish tissue samples with RLs of 7,700 µg/kg was reanalyzed, and BEHP was not detected with RLs ranging from 66 to 130 µg/kg ww.

COI – chemical of interest
J – estimated concentration
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
ROC – receptor of concern
SVOC – semivolatile organic compound
TRV – toxicity reference value
U – not detected at reporting limit shown
ww – wet weight

Bold identifies maximum fish tissue concentrations greater than the NOAEL.

Bis(2-ethylhexyl) phthalate (BEHP) and dimethyl phthalate were not detected in any fish ROC tissue sample but had RLs that were greater than the NOAELs. Uncertainty associated with RLs above NOAELs for these chemicals are described in the uncertainty analysis (Section A.6.2).

Based on the above COPC screen, TBT, alpha-endosulfan, beta-endosulfan, endrin, and total PCBs were identified as COPCs for further evaluation. A juvenile chinook-specific TRV was identified for PCBs because the study providing the selected PCB TRV (Hugla and Thome 1999) reported results following exposures of adults and documented reproductive endpoints. Because exposures of juvenile chinook salmon to chemicals in the LDW are limited to the period during which they migrate through the waterway, the selected TRV for adult fish (Hugla and Thome 1999) would not accurately reflect either the lifestage or exposure regime of the migratory juvenile chinook in the LDW. The lowest LOAEL for a non-reproductive endpoint was 46,000 µg/kg ww for survival of spot (Hansen et al. 1971). The highest NOAEL below this LOAEL (27,000 µg/kg ww) was selected as the screening NOAEL TRV for juvenile chinook salmon. This TRV is likely protective of juvenile chinook salmon growth because no adverse effects on growth of salmonid species have been observed at whole-body tissue concentrations as high as 31,000 µg/kg ww (Mauck et al. 1978).

Based on the ROC-specific screen, where maximum concentrations in each ROC tissue were compared to NOAEL TRVs, Pacific staghorn sculpin/TBT, English sole/total PCBs, and Pacific staghorn sculpin/total PCBs were identified as ROC/COPC pairs (Table A.2-25), and will be evaluated in this baseline ERA. Because of uncertainties in the JN-qualified organochlorine pesticides data (as discussed in Section A.2.4.2), risks associated with alpha-endosulfan and beta-endosulfan in English sole and Pacific staghorn sculpin, and endrin in all three fish ROCs are discussed in the uncertainty analysis (Section A.6.2).

Table A.2-25. Maximum tissue concentrations in each fish ROC compared to NOAEL TRVs for COIs analyzed using a critical tissue-residue approach

ROC	COI	Maximum Detected Chemical Concentration in Fish ROC Tissue (µg/kg ww)	NOAEL TRV (µg/kg ww)
Juvenile chinook salmon	TBT	14	18
	alpha-endosulfan	1.5	3.1
	beta-endosulfan	0.94	3.1
	endrin	**6.5**	1.15
	total PCBs	1,200	27,000

ROC	COI	Maximum Detected Chemical Concentration in Fish ROC Tissue (µg/kg ww)	NOAEL TRV (µg/kg ww)
English sole	TBT	9.9	18
	alpha-endosulfan	**6.6**	3.1
	beta-endosulfan	**18**	3.1
	endrin	**14**	1.15
	total PCBs	**4,700**	104 – 528
Pacific staghorn sculpin	TBT	**80**	18
	alpha-endosulfan	**3.6**	3.1
	beta-endosulfan	**6.4**	3.1
	endrin	**36**	1.15
	total PCBs	**2,800**	104 – 528

COI – chemical of interest

PCB – polychlorinated biphenyl

ROC – receptor of concern

TBT – tributyltin

TRV – toxicity reference value

ww – wet weight

Bold identifies maximum fish tissue concentrations greater than the NOAEL.

For COIs evaluated using a dietary approach, maximum dietary exposure concentrations were represented by a weighted average of 10% maximum sediment concentrations (to account for exposure via incidental sediment ingestion) and 90% maximum benthic invertebrate tissue concentrations (Equation 2-1).[23] Ten percent was selected as an upper-bound estimate of sediment ingestion based on discussions with fish experts (see Section A.4.1.2).

$$\text{Maximum [diet]} = \text{Maximum [sed]} \times 10\% + \text{Maximum [tissue]} \times 90\% \qquad \textbf{Equation 2-1}$$

Where:

diet = dietary concentration (mg/kg dw)
sed = sediment concentration (mg/kg dw)
tissue = tissue concentration (mg/kg dw)

Based on this COPC screen, five COIs evaluated using the dietary approach were selected as COPCs (TableA.2-26). Of these chemicals, a LOAEL TRV was not identified for chromium. Because the NOAEL for chromium is not bounded by a LOAEL, it does not indicate a potential threshold above which adverse effects may occur. Therefore, risks associated with chromium are discussed in the uncertainty analysis

[23] Note that maximum chemical concentrations in benthic invertebrate tissue samples exceeded those in shiner surfperch tissue samples for all dietary COIs. Therefore, this prey assumption conservatively screens COPCs for piscivorous fish, such as Pacific staghorn sculpin.

(Section A.6.2). The remaining four chemicals (arsenic, cadmium, copper, and vanadium) were identified as COPCs for all fish ROCs.

Table A.2-26. Maximum dietary exposure concentrations compared to NOAEL TRVs for fish COIs analyzed using a dietary approach

COI	Maximum LDW-Wide Sediment Concentration (mg/kg dw)	Maximum LDW-Wide Benthic Invertebrate Concentration (mg/kg dw)	Maximum Dietary Exposure Concentration (mg/kg dw)	NOAEL TRV (mg/kg dw)
Arsenic	1,100	116	**210**	20
Cadmium	120	1.3 J	**13**	0.1
Chromium	1,100 J	58.2	**160**	9.42
Copper	12,000 J	170	**1,400**	50
Lead	23,000	217.9	2,500	7,040
Silver	270	2.496	29	3,000
Vanadium	150	22.2	**35**	2.04
Zinc	9,700	384	1,300	1,900
Benzo(a)pyrene	7.9	1.3	2.0	100
Total PAHs[a]	119	21.7 J	34	324

[a] PAH mixture includes the following 21 PAHs: naphthalene, 2-methylnaphthalene, dimethylnaphthalene, dibenzothiophene, acenaphthene, fluorene, 1,8-dimethyl(9H)fluorene, phenanthrene, 9-ethylphenanthrene, 9-ethyl-10-methylphenanthrene, 1-methyl-7-isopropylphenanthrene, anthracene, fluoranthene, pyrene, methyl pyrene, benz(a)anthracene, chrysene, benzo(a)pyrene, benzo(k)fluoranthene, benzo(g,h,i)perylene, dibenzanthracene. These chemicals were evaluated as a PAH mixture by Meador et al. (2006).

COI – chemical of interest
dw – dry weight
J – estimated concentration
NOAEL – no-observed-adverse-effect level
PAH – polycyclic aromatic hydrocarbon
TRV – toxicity reference value

Bold identifies maximum dietary exposure concentrations greater than the NOAEL.

Based on the above COPC screen, PAHs were not selected as a COPC for any fish ROC because the maximum dietary exposure concentration was less than the NOAEL. However, of the 38 chemicals with no TRVs, 15 chemicals were alkylated PAHs, and 5 chemicals were non-alkylated PAHs. These alkylated and non-alkylated PAHs were not included in the calculation of total PAHs in the COPC screen above because they were not included in the PAH mixture used in the TRV study (Meador et al. 2006). To address this uncertainty, an additional PAH screen was conducted by comparing TRVs for benzo(a)pyrene (Hart and Heddle 1991; Rice et al. 2000) and a PAH mixture (Meador et al. 2006) to maximum concentrations of total PAHs (i.e., the sum of all alkylated and non-alkylated PAHs analyzed) calculated in fish diets using Equation 2.1 and detected in juvenile chinook stomach contents (Table A.2-27). Maximum dietary exposure concentrations of all PAHs were lower than the available TRVs (Table A.2-27), thus supporting the exclusion of PAHs as a COPC for all fish species.

Table A.2-27. Maximum dietary PAH exposure concentrations (including all alkylated and non-alkylated PAHs) compared to total PAH and benzo(a)pyrene NOAEL TRVs for fish

Chemical in TRV Study	TRVs		Maximum Exposure Concentration of Total PAHs			
	NOAEL (mg/kg dw)	LOAEL (mg/kg dw)	Juvenile Chinook Salmon Stomach Contents (mg/kg dw)	Sediment (mg/kg dw)	Benthic Invertebrate Prey (mg/kg dw)	Maximum Dietary Exposure Concentration (mg/kg dw)
Benzo(a)pyrene	100[a]	116[b]	14.4 JN	133[d]	36.1J[e]	46[f]
Total PAHs	324[c]	951[c]				

[a] No effects on growth of rainbow trout (Hart and Heddle 1991).

[b] Effects on growth of English sole (Rice et al. 2000).

[c] TRVs based on growth of juvenile chinook salmon exposed to a mixture of naphthalene, 2-methylnaphthalene, dimethylnaphthalene, dibenzothiophene, acenaphthene, fluorene, 1,8-dimethyl(9H)fluorene, phenanthrene, 9-ethylphenanthrene, 9-ethyl-10-methylphenanthrene, 1-methyl-7-isopropylphenanthrene, anthracene, fluoranthene, pyrene, methyl pyrene, benz(a)anthracene, chrysene, benzo(a)pyrene, benzo(k)fluoranthene, benzo(g,h,i)perylene, dibenzanthracene (Meador et al. 2006).

[d] Maximum concentration of total PAHs in LDW sediment.

[e] Maximum total PAH concentration in benthic invertebrate tissue collected from the LDW.

[f] Maximum concentration of total PAHs in diet calculated using Equation 2-1.

dw – dry weight
J – estimated concentration
N – tentative identification
na – not available
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PAH – polycyclic aromatic hydrocarbon
TRV – toxicity reference value

Although PAHs were not selected as a COPC for fish, regional field studies have suggested a correlation between PAH contamination of sediment and adverse effects on reproduction observed in English sole from Puget Sound, including English sole from the LDW (as discussed in Section A.6.2.2). Dietary effects data for PAHs are limited, however, and no laboratory studies investigating effects of ingested PAHs on fish reproduction are available.[24]

The LOAEL of 951 mg/kg dw in diet reported in Meador et al. (2006) was selected as the LOAEL TRV for fish in this section because it was the lowest LOAEL reported. The NOAEL of 324 mg/kg dw from this study was also selected. In the selected study, Meador et al. (2006) exposed juvenile chinook salmon to a mixture of 21 PAHs[25] in a

[24] Four studies that evaluated the dietary toxicity of benzo(a)pyrene to fish were identified (Wu et al. 2003; Hart and Heddle 1991; Hendricks et al. 1985; Rice et al. 2000). Effects investigated included growth or survival for English sole, rainbow trout, and aeriolated grouper. Two studies that evaluated the dietary toxicity of PAH mixtures to fish were identified (Palm et al. 2003; Meador et al. 2006) (see Attachment 8, Table 1). Both studies evaluated effects on growth of juvenile chinook salmon. The uncertainty section (Section A.6.2.1) discusses growth effects on English sole reported in Rice et al. (2000), in which fish were exposed to polychaetes that had previously been exposed to field-collected creosote-contaminated sediments from a Superfund site.

[25] The PAH mixture included the following chemicals: naphthalene, 2-methylnaphthalene, dimethylnaphthalene, dibenzothiophene, acenaphthene, fluorene, 1,8-dimethyl(9H)fluorene,

synthetic diet. Reduced body weight in exposed fish was observed at the LOAEL relative to the control. Palm et al. (2003) observed no adverse effects on juvenile chinook salmon growth following dietary exposure to a mixture of 14 PAHs for 7 weeks at total PAH concentrations of up to 280 mg/kg dw.

Because screening exposure assumptions would be similar for all fish ROCs, no ROC-specific screen was conducted for COPCs evaluated using a dietary approach. Therefore, the COIs that were identified in the dietary screen as COPCs were selected as COPCs for all three fish ROCs. The COPCs that will be evaluated for each fish ROC are presented in Table A.2-28.

Table A.2-28. COPCs selected for fish ROCs

COPC	Juvenile Chinook Salmon	English Sole	Pacific Staghorn Sculpin
Arsenic	X	X	X
Cadmium	X	X	X
Copper	X	X	X
Vanadium	X	X	X
TBT			X
Total PCBs		X	X

COPC – chemical of potential concern

PCB – polychlorinated biphenyl

TBT – tributyltin

A.2.5.3 Wildlife

COPCs were identified for wildlife using a two-step process. The first step was to identify a list of COIs. Chemicals were identified as COIs for wildlife (with the exception of spotted sandpiper) if they met at least two of the following three criteria:

- Detection in at least 5% of LDW surface sediment samples
- Identification as a bioaccumulative chemical in EPA (2000)
- Detection in any LDW-collected tissue sample

For spotted sandpiper, the COI screen was more focused because sandpipers ingest benthic invertebrates only from intertidal areas. Thus, chemicals were identified as COIs for spotted sandpiper if they met at least two of the following three criteria:

- Detection in at least 5% of LDW intertidal surface sediment samples
- Identification as a bioaccumulative chemical in EPA (2000)

---

phenanthrene, 9-ethylphenanthrene, 9-ethyl-10-methylphenanthrene, 1-methyl-7-isopropylphenanthrene, anthracene, fluoranthene, pyrene, methyl pyrene, benz(a)anthracene, chrysene, benzo(a)pyrene, benzo(k)fluoranthene, benzo(g,h,i)perylene, dibenzanthracene.

- Detection in any LDW-collected benthic invertebrate tissue sample[26]

Table A.2-29 presents a summary of chemicals identified as COIs as a result of this screen. Detailed results of the COI screening step for wildlife are presented in Attachment 4.

Table A.2-29. Chemicals identified as COIs for wildlife ROCs

COIs		
Metals		
Antimony	Copper	Selenium
Arsenic	Lead	Silver
Cadmium	Mercury	Thallium
Chromium	Methylmercury[a]	Vanadium
Chromium VI[a, b]	Molybdenum	Zinc
Cobalt	Nickel	
Organometals		
Monobutyltin as ion	TBT as ion	
Dibutyltin as ion	Tetrabutyltin as ion[b]	
PAHs		
C1-Chrysenes[b]	C3-Phenanthrenes/anthracenes	Benzo(g,h,i)perylene
C1-Dibenzothiophenes	C4-Naphthalenes	Benzo(k)fluoranthene
C1-Fluoranthene/pyrene	C4-Phenanthrenes/anthracenes	Chrysene
C1-Fluorenes	1-Methylnaphthalene	Dibenzo(a,h)anthracene
C1-Phenanthrenes/anthracenes	2-Methylnaphthalene	Dibenzofuran
C2-Chrysenes[b]	Acenaphthene	Fluoranthene
C2-Dibenzothiophenes	Acenaphthylene	Fluorene
C2-Fluorenes	Anthracene	Indeno(1,2,3-cd)pyrene
C2-Naphthalenes	Benzo(a)anthracene	Naphthalene
C2-Phenanthrenes/anthracenes	Benzo(a)pyrene	Perylene
C3-Fluorenes	Benzo(b)fluoranthene	Phenanthrene
C3-Naphthalenes	Benzo(e)pyrene	Pyrene
Phthalates		
Bis(2-ethylhexyl) phthalate	Diethyl phthalate	Di-n-butyl phthalate
Butyl benzyl phthalate[b]	Dimethyl phthalate[b]	
Other SVOCs		
4-Chlorophenyl phenyl ether[b]	Biphenyl	Hexachlorobenzene
4-Methylphenol	Carbazole[b]	Phenol
Benzoic acid	Dibenzothiophene	
PCBs		
Total PCBs		

[26] Includes amphipods, benthic invertebrates, crabs, and clams.

COIs		
Dioxins and Furans		
Organochlorine Pesticides		
Total DDTs[c]	delta-BHC	Endrin aldehyde[d]
Aldrin	gamma-BHC	Heptachlor
Dieldrin	alpha-Endosulfan	Heptachlor epoxide
alpha-BHC	beta-Endosulfan	Methoxychlor
beta-BHC	Endrin	Total chlordane[e]

[a] Chromium VI and methylmercury (both bioaccumulative chemicals and detected in at least 5% of surface sediment samples) were not analyzed in any of the tissue samples; these chemicals are assumed to be incorporated in total mercury and total chromium risk analyses.

[b] Not a COI for spotted sandpiper.

[c] Includes 2,4'-DDD, 2,4'-DDE, 2,4'-DDT, 4,4'-DDD, 4,4'-DDE, and 4,4'-DDT.

[d] COI for spotted sandpiper only.

[e] Includes alpha-chlordane, gamma-chlordane, oxychlordane, cis-nonachlor, and trans-nonachlor.

COI – chemical of interest

PCB – polychlorinated biphenyl

SVOC – semivolatile organic compound

TBT – tributyltin

In the second step of the screening process, the maximum exposure dose of each COI was compared to a NOAEL for that chemical. If the maximum exposure dose was greater than the NOAEL, the chemical was identified as a COPC for wildlife. This step was conducted for all COIs, except dioxins and furans, as discussed in Section A.2.5.

For COIs other than dioxins and furans, the scientific literature was searched[27] to identify toxicity reference values (TRVs) for COIs. The literature search included BIOSIS, EPA's ECOTOX database, the National Library of Medicine's TOXNET database, the US Fish and Wildlife Service's Contaminant Review series, the Oak Ridge National Laboratory's database, and EPA's IRIS database. Original sources of toxicity data were obtained and reviewed to verify effects data summarized in the databases as well as the suitability of the studies. The databases were searched for studies that evaluated effects on survival, growth, and reproduction (including developmental effects).

TRVs were based on the risk evaluation method used. For wildlife, dietary dose studies were identified. In many cases, the toxicity literature presented data only as a concentration in food, so these values were converted to a daily dose (mg/kg bw/day) using the animal's body weight and ingestion rate (IR). The following guidelines were considered in the selection of TRVs for wildlife.

[27] For COIs that were evaluated in the Phase 1 ERA (Windward 2003b), toxicity literature published after January 2001 was searched for new toxicity data. If any COIs were not evaluated in Phase 1 ERA, toxicity literature was searched for appropriate toxicity data and was not limited by date.

- Studies using field-collected data were not used to develop TRVs, but were considered if no other toxicity data were available for a COI.
- Studies conducted using IP injection, intramuscular injection, forced ingestion, or oral gavage as exposure routes were not considered for deriving TRVs unless no other toxicity data are available for a COI.
- Studies using drinking water as the exposure medium were not used to develop TRVs because bioavailability from water may be different from that of food. If no other toxicity data were available, then drinking water studies were considered.
- Studies with egg production endpoints for chicken or quail, such as Edens and Garlich (1983) and Edens et al. (1976) are considered highly uncertain and were only considered if data from other more appropriate studies were not available. These data are considered uncertain because chickens and quail have been bred to have high egg-laying rates. Even with a significant reduction in their baseline egg production, these egg production rates may be much higher than those of any wild avian species. These differences in reproductive physiology result in high uncertainty in extrapolating a reproductive effect threshold from egg production rates for chickens or quails.
- Toxicity studies conducted with chemical forms not likely found in the LDW, such as the fungicide methylmercury dicyandiamide, were not used to develop TRVs. Toxicity of these chemical forms is not comparable to the toxicity of forms of chemicals present in the LDW.

After the literature search was conducted, all acceptable studies for TRV derivation were compiled, as presented in Attachments 9 and 10. For each COI, TRVs were selected for both the NOAEL and the LOAEL. TRV selection rules and uncertainty factors discussed above for crab were used for wildlife as well.

Of the 86 COIs, TRVs were identified for 29 COIs for birds and 41 COIs for mammals (Tables A.2-30 and A.2-31, respectively). Selected TRVs are presented in Table A.2-32 for birds and in Table A.2-33 for mammals.

Table A.2-30. Results of TRV search for COIs for birds

COIs for Birds		
COIs with TRVs		
Arsenic	Thallium	Aldrin
Cadmium	Vanadium	Dieldrin
Chromium	Zinc	gamma-BHC
Cobalt	TBT as ion	alpha-Endosulfan
Copper	Benzo(a)pyrene	beta-Endosulfan
Lead	Total PAH	Endrin
Mercury	Bis(2-ethylhexyl) phthalate	Heptachlor
Molybdenum	Hexachlorobenzene	Methoxychlor
Nickel	Total PCBs	Total chlordane
Selenium	Total DDTs	
COIs without TRVs		
Antimony	Benzo(k)fluoranthene	Di-n-butyl phthalate
Silver	Chrysene	4-Chlorophenyl phenyl ether
Monobutyltin as ion	Dibenzo(a,h)anthracene	4-Methylphenol
Dibutyltin as ion	Dibenzofuran	Benzoic acid
Tetrabutyltin as ion	Fluoranthene	Biphenyl
1-Methylnaphthalene	Fluorene	Carbazole
2-Methylnaphthalene	Indeno(1,2,3-cd)pyrene	Dibenzothiophene
Acenaphthene	Naphthalene	Phenol
Acenaphthylene	Perylene	alpha-BHC
Anthracene	Phenanthrene	beta-BHC
Benzo(a)anthracene	Pyrene	delta-BHC
Benzo(b)fluoranthene	Butyl benzyl phthalate	Heptachlor epoxide
Benzo(e)pyrene	Diethyl phthalate	
Benzo(g,h,i)perylene	Dimethyl phthalate	

COI – chemical of interest

PAH – polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

TBT – tributyltin

TRV – toxicity reference value

Table A.2-31. Results of TRV search for COIs for mammals

COIs for Mammals		
Chemicals with TRVs		
Antimony	Dibutyltin as ion	Phenol
Arsenic	TBT as ion	Total PCBs
Cadmium	1-Methylnaphthalene	Total DDTs
Chromium	2-Methylnaphthalene	Aldrin
Cobalt	Benzo(a)pyrene	Dieldrin
Copper	Naphthalene	beta-BHC
Lead	Bis(2-ethylhexyl) phthalate	gamma-BHC
Mercury	Butyl benzyl phthalate	alpha-Endosulfan
Molybdenum	Diethyl phthalate	beta-Endosulfan
Nickel	Di-n-butyl phthalate	Endrin
Selenium	Benzoic acid	Heptachlor
Thallium	Biphenyl	Methoxychlor
Vanadium	Dibenzothiophene	Total chlordane
Zinc	Hexachlorobenzene	
Chemicals without TRVs		
Silver	Benzo(g,h,i)perylene	Phenanthrene
Monobutyltin as ion	Benzo(k)fluoranthene	Pyrene
Tetrabutyltin as ion	Chrysene	Dimethyl phthalate
Acenaphthene	Dibenzo(a,h)anthracene	4-Chlorophenyl phenyl ether
Acenaphthylene	Dibenzofuran	4-Methylphenol
Anthracene	Fluoranthene	Carbazole
Benzo(a)anthracene	Fluorene	alpha-BHC
Benzo(b)fluoranthene	Indeno(1,2,3-cd)pyrene	delta-BHC
Benzo(e)pyrene	Perylene	Heptachlor epoxide

COI – chemical of interest

PCB – polychlorinated biphenyl

TBT – tributyltin

TRV – toxicity reference value

Table A.2-32. TRVs selected for bird COIs

COI	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Endpoint	Source
Metals					
Arsenic	mallard	10	40	reproduction	Stanley et al. (1994)
Cadmium	chicken	1.5	na	growth	Cain et al. (1983)
	Japanese quail	na	4.0	growth	Richardson et al. (1974)
Chromium	black duck	1.0	5.0	reproduction	Haseltine et al. (unpublished), as cited in Sample et al. (1996)
Cobalt	chicken	2.31[a]	23.1	growth	Diaz et al. (1994)
Copper	chicken	ns	29	growth	Smith (1969)
	chicken	21	ns	growth	Poupoulis and Jensen (1976)
Lead	Japanese quail	ns	20	reproduction	Edens et al. (1976)
	American kestrel	5.82	na	reproduction	Pattee (1984)
Mercury	great egret	0.018[b]	0.091	growth	Spalding et al. (2000)
Molybdenum	chicken	6.0[b]	30	reproduction	Lepore and Miller (1965)
Nickel	mallard	77	107	growth	Cain and Pafford (1981)
Selenium	mallard	0.50	0.82	reproduction	Heinz et al. (1987)
Thallium	pheasant	2.4[a]	24	survival	Hudson et al. (1984)
Vanadium	chicken	1.2	2.3	growth	Ousterhout and Berg (1981)
Zinc	chicken	82	124	growth	Roberson and Schaible (1960)
Organometals					
TBT	Japanese quail	1.4	3.6	reproduction	Coenen et al. (1992)
PAHs					
Benzo(a)pyrene	pigeon	0.28[b]	1.4	reproduction	Hough el al. (1993)
Total PAHs	mallard	8.0	40	growth	Patton and Dieter (1980)
Phthalates					
BEHP	chicken	65.8[c]	329	reproduction	Ishida et al. (1982)
PCBs					
PCBs	screech owl	0.49	na	reproduction	McLane and Hughes (1980)
	ringed turtle dove	na	1.4	reproduction	Peakall et al. (1972); Peakall and Peakall (1973)
Organochlorine Pesticides					
Aldrin	quail	0.008[b]	0.04	survival	Dewitt (1956)
Total chlordane	bobwhite quail	0.6	na	growth, survival	Ludke (1976)
	bobwhite quail	na	2.0	survival	Hill et al. (1975); Heath et al. (1972)
Total DDTs	mallard	0.064[d]	0.32	reproduction	Davison and Sell (1974)
Dieldrin	quail	0.08	0.12	survival	Dewitt (1956)
Endosulfan	gray partridge	10	na	reproduction	Abiola (1992)

COI	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Endpoint	Source
Endrin	quail	0.070	0.20	survival	DeWitt (1956)
gamma-BHC	mallard	1.6	3.6	reproduction	Chakravarty and Lahiri (1986); Chakravarty et al. (1986)
Heptachlor	bobwhite quail	0.5[a]	5.0	survival	Hill et al. (1975); Heath et al. (1972)
Hexachloro-benzene	Japanese quail	na	1.2	reproduction	Schwetz et al. (1974)
	Japanese quail	1.1	na	reproduction	Vos et al. (1971)
Methoxychlor	zebra finch	34.6	346	reproduction	Gee et al. (2004)[e]
				survival	Millam et al. (2002)[e]

[a] NOAEL estimated from an acute or subchronic LOAEL using an uncertainty factor of 10.

[b] NOAEL estimated from a chronic LOAEL using an uncertainty factor of 5.

[c] There was a NOAEL of 1.45 mg/kg bw/day from a study that reported no effect on eggshell thinning, but this is an unbounded NOAEL at a substantially lower concentration than the study with observed effects. Therefore, the NOAEL was estimated from the reproductive LOAEL using an uncertainty factor of 5.

[d] There was a NOAEL of 0.19 mg/kg bw/day from a study that reported no effect on eggshell thinning from exposure of barn owls to DDT (Mendenhall et al. 1983). However, as discussed in Section A.6.3.1.2, there is evidence indicating that p,p'-DDE rather than DDT is the likely cause of eggshell thinning (Lundholm 1997). Therefore, the NOAEL was estimated from the DDE LOAEL for eggshell thinning using a factor of 5.

[e] Both studies had the same LOAEL and NOAEL.

BEHP – bis(2-ethylhexyl) phthalate

bw – body weight

COI – chemical of interest

LOAEL – lowest-observed-adverse-effect level

na – not available

NOAEL – no-observed-adverse-effect level

ns – NOAEL or LOAEL not selected from this study

PAH – polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

TBT – tributyltin

TRV – toxicity reference value

Table A.2-33. TRVs selected for mammal COIs

COI	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Endpoint	Source
Metals					
Antimony	rat	1,489	na	growth, survival	Hext et al. (1999)
Arsenic	rat	2.6	5.4	growth	Byron et al. (1967)
Cadmium	rat	3.5	13	growth	Machemer and Lorke (1981)
Chromium	rat	1,466	na	growth, survival	Ivankovic and Preussman (1975)
Cobalt	rat	0.1	1.0	growth	Chetty et al. (1979)
Copper	mink	18	26	reproduction	Aulerich et al. (1982)
Lead	rat	11	90	growth	Azar et al. (1973)
Mercury	rat	0.0017[a]	0.0084	growth	Verschuuren et al. (1976)
Molybdenum	mouse	0.258[b]	2.58	reproduction, survival	Schroeder and Mitchener (1971)
Nickel	rat	na	20	reproduction	Ambrose et al. (1976)
	rat	8.4	ns	growth	
Selenium	rat	0.055	0.08	growth	Halverson et al. (1966)
Thallium	rat	0.74	na	growth	Formigli et al. (1986)
Vanadium	mouse	1.05	na	growth	Schroeder and Balassa (1967)
	rat	na	2.7	growth	Adachi et al. (2000)
Zinc	rat	160	320	reproduction	Schlicker and Cox (1968)
Organometals					
TBT	rat	0.4	2.0	reproduction	Omura et al. (2001)
Dibutyltin	rat	na	7.6	reproduction, growth	Ema et al. (2003)
	rat	3.8	na	growth	Harazono and Ema (2003)
PAHs					
1-Methylnaphthalene	mouse	150	na	growth	Murata et al. (1993)
2-Methylnaphthalene	mouse	54	114	growth	Murata et al. (1997)
Benzo(a)pyrene	mouse	2.0[a]	10	reproduction	MacKenzie and Angevine (1981)
Naphthalene	mouse	133	na	growth, survival	Shopp et al. (1984)
Phthalates					
BEHP	mouse	44	91	reproduction	Tyl et al. (1988)
Butyl benzyl phthalate	rat	250	750	growth, reproduction	Tyl et al. (2004)
Diethyl phthalate	mouse	1,860	3,721	growth, reproduction	Lamb et al. (1987)
Di-n-butyl phthalate	rat	16[a]	80	reproduction	Wine et al. (1997)
Other SVOCs					
Benzoic acid	rat	80	na	growth, survival	Ignat'ev (1965), as cited in IRIS (EPA 2006)
	rat	na	750	growth	Marquardt 1980

COI	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Endpoint	Source
Biphenyl	rat	50	250	survival	Ambrose et al. (1960), as cited in IRIS (EPA 2006)
Dibenzothiophene	mouse	4.7[b]	47[c]	survival	Leighton (1989)
	mouse	na	300	reproduction, survival	Plasterer et al. (1985)
Phenol	rat	60	120	growth	Argus Research Laboratories (1997), as cited in IRIS (EPA 2006)[d]
	rat	60	120	reproduction	Charles River Laboratories (1988) and NTP (1983), as cited in IRIS (EPA 2006)[d]
PCBs					
PCBs	mink	0.045[e]	0.089	reproduction	Brunstrom et al. (2001)
Organochlorine Pesticides					
Aldrin	rat	0.8	4.1	survival	Reuber (1980)
beta-BHC	rat	5.7	31	growth, survival	Van Velsen et al. (1986)
Total chlordane	mouse	0.18	0.92	growth	Khasawinah and Grutsch (1989)
Total DDTs	rat	1.2	na	reproduction	Duby et al. (1971)
	mouse	na	1.3	reproduction	Ware and Good (1967)
Dieldrin	mouse	0.038[a]	0.19	reproduction	Treon and Cleveland (1955)
Endosulfan	mouse	0.84	2.5	growth, survival	Hack et al. (1995)
Endrin	rat	0.40	ns	survival, growth	Treon et al. (1955)
	mouse	na	0.92	survival, reproduction	Good and Ware (1969)
gamma-BHC	rat	64	na	growth	Srinivasan et al. (1991)
Heptachlor	mink	1.0	1.8	growth, survival, reproduction	Crum et al. (1993)
Hexachlorobenzene	mink and ferret	0.026[a]	0.13	reproduction	Bleavins et al. (1984)
Methoxychlor	rat	17	na	growth, reproduction	Masutomi et al. (2003)
	rat	na	56	growth, reproduction	You et al. (2002)

[a] NOAEL estimated from an chronic LOAEL using an uncertainty factor of 5.

[b] NOAEL estimated from an acute or subchronic LOAEL using an uncertainty factor of 10.

[c] LOAEL estimated from an LD50 using an uncertainty factor of 10.

[d] Both studies had the same LOAEL and NOAEL.

[e] NOAEL estimated from a chronic LOAEL using an uncertainty factor of 2; the rationale for using this uncertainty factor is discussed in Section A.5.2.2.1.

BEHP – bis(2-ethylhexyl) phthalate
bw – body weight
COI – chemical of interest
LOAEL – lowest-observed-adverse-effect level
na – not available
NOAEL – no-observed-adverse-effect level
ns – NOAEL or LOAEL not selected from this study
PAH – polycyclic aromatic hydrocarbon
PCB – polychlorinated biphenyl
SVOC – semivolatile organic compound
TBT – tributyltin
TRV – toxicity reference value

To identify COPCs for birds and mammals, NOAEL TRVs were compared to the estimated maximum dietary dose (expressed in mg/kg dw-day) for each of the five wildlife ROCs. The maximum dietary exposure dose was calculated using the following equation:

$$\text{Dose} = \frac{\text{FIR} \times C_{food}}{\text{BW}} \qquad \textbf{Equation 2-2}$$

Where:

Dose	=	COIs ingested per day via food or intertidal sediment (mg COI/kg body weight/day)
FIR	=	food ingestion rate (kg dw food/day)
C_{food} (sandpiper)	=	80% maximum [benthic invertebrate tissue] + 20% maximum [intertidal sediment] (mg COI/kg dw)
C_{food} (other ROCs)	=	maximum concentration in prey tissue (mg COI/kg dw)
BW	=	wildlife species body weight (kg ww)

The maximum daily ingested doses for great blue heron, osprey, river otter, and harbor seal were calculated using the maximum detected tissue concentration[28] in any of the appropriate prey species in the Phase 2 ERA dataset (see Table A.2-12). For spotted sandpiper, the maximum dose was calculated using a weighted average of 80% of the maximum concentration in any benthic invertebrate tissue sample plus 20% of the maximum concentration in any intertidal sediment sample.[29]

The body weights and FIRs used to calculate the maximum dietary exposure doses for each wildlife ROC are presented in Table A.5-3. The derivation of these values is described in detail in Section A.5.1.2.

Estimates of the maximum daily ingested doses of 11 metals (arsenic, cadmium, chromium, cobalt, copper, lead, mercury, nickel, selenium, vanadium, and zinc), two organic compounds (total PCBs and total DDTs) were greater than their respective NOAEL TRVs for spotted sandpiper (Table A.2-34). Total DDTs will be discussed in the uncertainty analysis because of the uncertainty associated with organochlorine pesticide tissue data (see Section A.2.4.2.2). Risks from the 11 metals and total PCBs are evaluated in the baseline ERA for spotted sandpiper.

[28] Incidental sediment ingestion was not considered in the screens for wildlife ROCs (except spotted sandpiper) because of the small amount of sediment assumed to be ingested (2% or less of diet).

[29] In risk calculations presented in Section 5.1, sediment ingestion rates are calculated separately from food ingestion rates. In those calculations, food is ingested at 100% of the food ingestion rate, with sediment comprising an additional component of the diet.

Table A.2-34. Results of COPC screen for spotted sandpiper

COI	Maximum Concentration		Calculated Maximum Dietary Exposure Dose (mg/kg bw/day)[a]	NOAEL TRV (mg/kg bw/day)
	Benthic Invertebrate Prey (mg/kg dw)	Intertidal Sediment (mg/kg dw)		
Metals				
Arsenic	43 J	1,100	**40**	10
Cadmium	0.91	120	**3.9**	1.5
Chromium	58	1,100	**43**	1.0
Cobalt	5.2	140	**5.2**	2.31
Copper	170	12,000	**400**	21
Lead	220	23,000	**750**	5.82
Mercury	0.44	4.6	**0.21**	0.018
Molybdenum	2.6	49	1.9	6.0
Nickel	6.7	910	**29**	77
Selenium	2.4	20	**0.92**	0.50
Thallium	0.026	30	0.94	2.4
Vanadium	164	150	**6.7**	1.2
Zinc	380	9,700	**350**	82
Organometals				
TBT as ion	4.2	0.099	0.53	1.4
PAHs				
Benzo(a)pyrene	1.3	2.9	0.25	0.28
Total PAH	24	130	7.0	8.0
Phthalates				
Bis(2-ethylhexyl) phthalate	14 J	14	2.2	65.8
PCBs				
Total PCBs	21	220	**9.5**	0.49
Organochlorine Pesticides				
Aldrin	0.0073 JN	0.00081	0.00094	0.008
alpha-Endosulfan	0.040 JN	0.071	0.0072	10
beta-Endosulfan	0.082 JN	0.010	0.011	10
Total chlordane	0.11 JN	0.23	0.021	0.6
Total DDTs	1.9 JN	2.9	**0.33**	0.064
Dieldrin	0.032 JN	0.28	0.013	0.08
Endrin	0.0099 JN	0.0091	0.0015	0.070
gamma-BHC	0.018 JN	0.0067	0.0025	1.6
Heptachlor	0.11 JN	0.00089	0.014	0.5
Methoxychlor	0.058 JN	0.099	0.010	34.6

[a] Calculated using Equation 2-2.

COI – chemical of interest
dw – dry weight
J – estimated concentration
N – tentative identification
PAH – polycyclic aromatic hydrocarbon
PCB – polychlorinated biphenyl
TBT – tributyltin
TRV – toxicity reference value

Bold identifies maximum dietary exposure concentrations greater than the NOAEL.

Maximum dietary exposures of great blue heron and osprey to chromium, lead, mercury, total PCBs, and total DDTs exceeded their respective NOAEL TRVs (Table A.2-35). Total DDT is discussed in the uncertainty assessment because of the uncertainty associated with JN-qualified organochlorine pesticide tissue data (see Section A.2.4.2.2). Risks from chromium, lead, mercury, and total PCBs are evaluated in the exposure and effects assessment and risk characterization for great blue heron and osprey (Sections A.6.3.2 and A.6.3.3, respectively).

Table A.2-35. Results of COPC screen for great blue heron and osprey

COI	Maximum Chemical Concentration in Fish (mg/kg dw)	Calculated Maximum Dietary Exposure Dose (mg/kg bw/day)[a]		NOAEL TRV (mg/kg bw/day)
		Great Blue Heron	Osprey	
Metals				
Arsenic	120	5.0	5.9	10
Cadmium	6.5	0.27	0.32	1.5
Chromium	58	**2.4**	**2.8**	1.0
Cobalt	5.2	0.22	0.25	2.31
Copper	380	16	19	21
Lead	220	**9.2**	**11**	5.82
Mercury	0.49 M	**0.020**	**0.024**	0.018
Molybdenum	2.7	0.11	0.13	6.0
Nickel	23	0.96	1.1	17
Selenium	3.6	0.15	0.18	0.50
Thallium	0.054	0.0023	0.0026	2.4
Vanadium	22	0.92	1.1	1.2
Zinc	380	16	19	82
Organometals				
TBT as ion	4.2	0.18	0.21	1.4
PAHs				
Benzo(a)pyrene	1.3	0.054	0.063	0.28
Total PAHs	24 J	1.0	1.2	8.0
Phthalates				
Bis(2-ethylhexyl) phthalate	14	0.58	0.68	65.8
PCBs				
Total PCBs	69	**2.9**	**3.4**	0.49
Organochlorine Pesticides				
Aldrin	0.025 JN	0.0010	0.0012	0.008
alpha-Endosulfan	0.21 JN	0.0088	0.010	10
beta-Endosulfan	0.18 JN	0.0075	0.0088	10
Total chlordane	1.2 JN	0.050	0.059	0.60
Total DDTs	3.8 JN	**0.16**	**0.19**	0.064
Dieldrin	0.032 JN	0.0013	0.0016	0.08
Endrin	0.17 JN	0.0071	0.0083	0.070

COI	Maximum Chemical Concentration in Fish (mg/kg dw)	Calculated Maximum Dietary Exposure Dose (mg/kg bw/day)[a]		NOAEL TRV (mg/kg bw/day)
		Great Blue Heron	Osprey	
gamma-BHC	0.033 JNM	0.033	0.0016	1.6
Heptachlor	0.11 JN	0.0046	0.0054	0.5
Hexachlorobenzene	0.032 JNM	0.0013	0.0016	1.1
Methoxychlor	0.70 JNM	0.029	0.034	34.6

[a] Calculated using Equation 2-2.

bw – body weight
COI – chemical of interest
COPC – chemical of potential concern
dw – dry weight
J – estimated concentration
M – mean concentration
N – tentative identification
NOAEL – no-observed-adverse-effect level
PAH – polycyclic aromatic hydrocarbon
PCB – polychlorinated biphenyl
TBT – tributyltin
TRV – toxicity reference value

Bold identifies maximum dietary exposure concentrations greater than the NOAEL.

For river otter, estimated maximum exposures to arsenic, cobalt, mercury, selenium, and PCBs were greater than their respective NOAEL TRVs (Table A.2-36). For harbor seal, estimated maximum exposures to mercury and PCBs exceeded their respective NOAEL TRVs (Table A.2-37). Risks associated with these COPCs are evaluated in this baseline ERA. A summary of the COPCs that will be evaluated for each wildlife ROC in the risk characterization is presented in Table A.2-29.

Table A.2-36. Results of COPC screen for river otter and harbor seal

COI	Maximum Chemical Concentration in Fish (mg/kg dw)	Calculated Maximum Dietary Exposure Dose (mg/kg bw/day)[a]		NOAEL TRV (mg/kg bw/day)
		River Otter	Harbor Seal	
Metals				
Antimony	1.9	0.062	0.014	1,489
Arsenic	120	**3.9**	0.089	2.6
Cadmium	6.5	0.21	0.048	3.5
Chromium	58	1.9	0.43	1,466
Cobalt	5.2	**0.17**	0.039	0.1
Copper	380	12	2.8	18
Lead	218	7.2	1.6	11
Mercury	0.49 M	**0.016**	**0.0036**	0.0017
Molybdenum	2.78	0.088	0.020	0.258
Nickel	238	0.75	0.17	8.4
Selenium	3.68	**0.12**	0.027	0.055
Thallium	0.054	0.0018	0.00040	0.74
Vanadium	228	0.72	0.16	1.05
Zinc	380	12	2.8	160

COI	Maximum Chemical Concentration in Fish (mg/kg dw)	Calculated Maximum Dietary Exposure Dose (mg/kg bw/day)[a]		NOAEL TRV (mg/kg bw/day)
		River Otter	Harbor Seal	
Organometals				
Dibutyltin	0.51	0.017	0.0038	3.8
TBT as ion	4.2	0.14	0.031	0.4
PAHs				
1-Methylnaphthalene	0.075	0.0024	0.00056	150
2-Methylnaphthalene	0.084	0.0027	0.00063	54
Benzo(a)pyrene	1.3	0.042	0.010	2.0
Naphthalene	0.13 J	0.0042	0.0010	133
Phthalates				
Bis(2-ethylhexyl) phthalate	14	0.46	0.10	44
Butyl benzyl phthalate	16	0.52	0.12	250
Diethyl phthalate	5.7 JM	0.19	0.042	1,860
Di-n-butyl phthalate	9.3	0.30	0.069	16
Other SVOCs				
Benzoic acid	220	7.2	1.6	80
Biphenyl	0.076 J	0.0025	0.00057	50
Dibenzothiophene	0.27 J	0.0088	0.0020	4.7
Phenol	12	0.39	0.089	60
PCBs				
Total PCBs	69	**2.2**	**0.51**	0.045
Organochlorine Pesticides				
Aldrin	0.025 JN	0.00081	0.00019	0.8
alpha-Endosulfan	0.21 JN	0.0068	0.0016	0.84
beta-BHC	0.13 JN	0.0042	0.0010	5.7
beta-Endosulfan	0.18 JN	0.0059	0.0013	0.84
Total chlordane	1.2 JN	0.039	0.0089	0.18
Total DDTs	3.8 JN	0.12	0.028	1.2
Dieldrin	0.032 JN	0.0010	0.00024	0.038
Endrin	0.17 JN	0.0055	0.0013	0.40
gamma-BHC	0.033 JNM	0.0011	0.00025	64
Heptachlor	0.11 JN	0.0036	0.00082	1.0
Hexachlorobenzene	0.032 JNM	0.0010	0.00024	0.026
Methoxychlor	0.70 JNM	0.023	0.0052	17

[a] Calculated using Equation 2-2.

bw – body weight
COI – chemical of interest
COPC – chemical of potential concern
dw – dry weight
J – estimated concentration
M – mean concentration
N – tentative identification
NOAEL – no-observed-adverse-effect level
PAH – polycyclic aromatic hydrocarbon
PCB – polychlorinated biphenyl
SVOC – semivolatile organic compound
TBT – tributyltin
TRV – toxicity reference value

Bold identifies maximum dietary exposure concentrations greater than the NOAEL.

Table A.2-37. COPCs evaluated in the risk characterization for birds and mammals

COPC	Spotted Sandpiper	Great Blue Heron and Osprey	River Otter	Harbor Seal
Arsenic	X		X	
Cadmium	X			
Chromium	X	X		
Cobalt	X		X	
Copper	X			
Lead	X	X		
Mercury	X	X	X	X
Nickel	X			
Selenium	X		X	
Vanadium	X			
Zinc	X			
Total PCBs	X	X	X	X

COPC – chemical of potential concern

PCB – polychlorinated biphenyl

This COPC screen for birds evaluated exposure and effects using a dietary approach. An alternative method to assess exposure of birds involves chemical analysis of bird eggs. In 1998, the USGS collected eggs from the great blue heron colony in West Seattle and analyzed them for PCBs (Krausmann 2002a). The results indicated that great blue herons were exposed to PCBs, although the source of the exposure was not established. Because the available QA/QC data were insufficient to use these data quantitatively in this baseline ERA, exposure of great blue herons to PCBs was evaluated using a dietary exposure approach. Osprey eggs collected from nests along the LDW have also been analyzed for PCBs by USGS. Because these data are not yet available from USGS, COPC concentrations in osprey eggs were estimated using biomagnification factors (BMFs) in the uncertainty analysis. In the uncertainty analysis, estimated PCB concentrations in osprey eggs are compared to egg TRVs from the literature.

A.2.6 Conceptual Site Model

A CSM is a graphical representation of chemical sources, transport mechanisms, exposure pathways, exposure routes, and potentially exposed receptors. This section presents the CSM that synthesizes pathways of exposure of ROCs to chemical stressors. Based on this model and assessment endpoints for this risk assessment, measures of exposure and effect are selected and discussed. These assessment endpoints determine which endpoints will be examined in detail in this baseline ERA for each ROC/COPC combination that was retained for further analysis based on the analyses in Section A.2.0.

Although chemical sources other than sediment exist in the LDW, the exposure assessment for each ROC focused on scenarios that include a direct (i.e., ingestion or direct contact) or indirect (i.e., ingestion of fish or benthic invertebrates) pathway for sediment-associated chemicals. Sources of chemical contamination to the sediments are discussed further in the Phase 2 RI.

Ecological risks from exposure to surface water were previously evaluated quantitatively by King County (1999b). The surface water risk estimates from that ERA have been incorporated in this ERA for fish. Additionally, surface water exposure data have been incorporated into bird and mammal exposure calculations in this ERA.

A.2.6.1 Exposure pathways

This section discusses the potential for ROCs in the LDW to be exposed significantly to COPCs. For COPCs to pose risk to ROCs, the exposure pathway must be complete. Identifying complete exposure pathways prior to a quantitative evaluation allows the assessment to focus on only those chemicals that can reach ecological receptors (EPA 1997a, b). An exposure pathway is considered complete if a chemical can travel from a source to ecological receptors and the receptor is exposed via one or more exposure routes (EPA 1997a, b). Complete pathways can be of varying importance, so key pathways that reflect maximum exposures to ecological receptors sensitive to that chemical (EPA 1997a, b) are identified as having more importance than pathways likely to provide a very low fraction of the total exposure of an ROC to a chemical.

Pathways for the exposure of ROCs to sediment-associated chemicals in the LDW were designated in one of four ways: complete and significant, complete and significance unknown, complete and insignificant, or incomplete. Each of the four designations is defined below, including whether it will be further evaluated in this ERA. This section also presents a brief rationale for each designation by receptor. The CSM is presented in Figures A.2-2 and A.2-3 for aquatic species and wildlife, respectively.

- **Complete and significant:** There is a direct link between the receptor and chemical via this pathway, and the specific pathway is considered to be potentially important. Pathways classified as complete and significant will be addressed in greater detail in the exposure and effects assessment (Sections A.3.0, A.4.0, and A.5.0).
- **Complete and significance unknown:** There is a direct link between the receptor and the chemical via this pathway; however, there is insufficient data available to quantify the significance of the pathway in the overall assessment of exposure. Pathways classified as complete and significance unknown will be discussed qualitatively in the uncertainty analysis (Section A.6.0).
- **Complete and insignificant:** There is a direct link between the receptor and the chemical via this pathway; however, the significance of this pathway in terms of

overall exposure is considered to be very low. Pathways classified as complete and insignificant will not be evaluated further in this baseline ERA.

- **Incomplete:** There is no direct pathway between the receptor and the chemical. Pathways classified as incomplete will not be evaluated further in this baseline ERA.



Figure A.2-2. Conceptual site model for fish and the benthic invertebrate community



Figure A.2-3. Conceptual site model for wildlife

A.2.6.2 Food web diagram

To understand the potential exposure pathways of a sediment-associated chemical to upper trophic-level ROCs, knowledge of food web relationships is important. The generalized food web diagram for the LDW shows the relationship between major trophic groups, and lists several representative species (Figure A.2-4).[30] The relationship among trophic groups illustrates the pathways for chemical transfer through the food web through the ingestion of prey; Figure A.2-4 provides additional detail for the prey ingestion pathways identified in Figures A.2-2 and A.2-3.

[30] Note that some organisms could have representatives in more than one box, depending on their life stage.



Figure A.2-4. Generalized food web diagram for the LDW

A.2.6.3 Assessment endpoints and measures of effect and exposure

An assessment endpoint is defined as an explicit expression of the ecological value that is to be protected (EPA 1992). Ecological values include those roles and processes vital to ecosystem function, those providing critical resources such as habitat and fisheries, and the perception of value by humans (e.g., threatened or endangered [T&E] species). An assessment endpoint must define both the valued entity and the characteristic of the entity to be protected. They provide direction for the risk assessment and are the basis for the analyses. Unless an ecological receptor is listed as a T&E species, assessment endpoints are selected that are relevant to population-level rather than individual effects. For T&E species, risks to individuals are important to evaluate (EPA 1998), although specific guidance regarding this approach is not available. At other EPA Region 10 Superfund sites, such as Coeur d'Alene and Blackbird Mine, greater emphasis has been placed on the NOAELs than on the LOAELs for the protection of T&E species.

Selection of assessment endpoints was based on available information regarding the ecological relevance of the endpoint and on societal values. In addition, assessment endpoints were evaluated to ensure that their protection would likely result in protection of other valued entities within the system. Finally, endpoints selected must be amenable to assessment either through previously existing data or data that were collected as part of the Phase 2 RI.

Assessment endpoints for each ROC are listed in Table A.2-38 along with the measures of exposure and effect used in the exposure and effects assessments (Sections A.3.0 through A.5.0). Survival, growth, and reproduction[31] are the key endpoints under review for most species in this assessment. Biomarker and histological endpoints are not included as assessment endpoints. Typically, ERAs focus on ecological effects at the individual level or higher (i.e., population level). In this way, the emphasis is placed on endpoints that integrate an overall response by an organism, rather than indicators of a biochemical response that may or may not result in an ecologically relevant effect. For biomarkers to be useful in determining sediment-associated risk, they must have clear dose-response data relating exposure to ecologically significant effects. Other responses, such as biliary fluorescent aromatic compounds and deoxyribonucleic acid (DNA) adducts, are categorized as a measure of exposure rather than as an assessment endpoint. Research is ongoing in the area of biomarkers to better understand their significance for potential use in ERAs.

[31] The fish reproductive endpoint includes early life stage (i.e., egg and embryo) survival and growth through the fry stage. The wildlife reproduction endpoint includes survival and growth of offspring after exposure of parents.

Table A.2-38. Assessment endpoints for ROCs and measures of effect and exposure

ROC	Assessment Endpoint	Assessment Scale	Measures of Effect	Measures of Exposure
Benthic				
Benthic invertebrate communities	survival, growth, reproduction	potential exposure area: small exposure areas for individuals assessment scale: small exposure areas throughout the LDW	SMS and toxicologically based sediment guidelines or TRVs[a]	chemical concentrations in sediment
			water based TRVs for VOCs	VOC concentrations in porewater
			site-specific toxicity tests	chemical concentrations in sediment samples co-located with toxicity test samples
			tissue-based TRVs for TBT (excluding imposex in gastropods)	TBT concentrations in sediment samples co-located with benthic invertebrate tissue collection
			assessment of imposex in field-collected gastropods	TBT concentrations in sediment samples co-located with gastropod collection
Crabs	survival, growth, reproduction	potential exposure area: crab may forage throughout the LDW assessment scale: LDW-wide	tissue-based TRVs for decapods	chemical concentrations in crab tissue collected from four tissue sampling areas located throughout the LDW
Fish				
Juvenile chinook salmon	survival and growth	potential exposure area: juvenile salmonids will migrate throughout the LDW and forage in shallow areas assessment scale: intertidal areas throughout the LDW	tissue-based TRVs for chemicals evaluated using a critical tissue-residue approach	chemical concentrations in juvenile chinook salmon tissue collected from middle and lower segments of the LDW
			dietary-based TRVs for chemicals evaluated using a dietary approach	chemical concentrations in juvenile chinook salmon prey collected from intertidal habitat throughout the LDW, stomach contents collected from juvenile chinook salmon captured throughout the LDW, and sediment collected from intertidal habitats throughout the LDW

ROC	Assessment Endpoint	Assessment Scale	Measures of Effect	Measures of Exposure
English sole	survival, growth, reproduction	potential exposure area: English sole may forage throughout the LDW assessment scale: LDW-wide	tissue-based TRVs for chemicals evaluated using a critical tissue-residue approach	chemical concentrations in English sole tissue collected from four tissue sampling areas located throughout the LDW
			dietary-based TRVs for chemicals evaluated using a dietary approach	chemical concentrations in English sole prey and sediment collected throughout the LDW
Pacific staghorn sculpin	survival, growth, reproduction	potential exposure area: sculpin may forage throughout the LDW or small segments of LDW assessment scale: LDW-wide and four modeling areas	tissue-based TRVs for chemicals evaluated using a critical tissue-residue approach	chemical concentrations in sculpin tissue collected from four tissue sampling areas located throughout the LDW
			dietary-based TRVs for chemicals evaluated using a dietary approach	chemical concentrations in sculpin prey and sediment collected throughout the LDW and divided into four modeling areas
Wildlife				
Great blue heron	survival, growth, reproduction	potential exposure area: herons may forage in areas of shallow water depths throughout the LDW assessment scale: LDW-wide intertidal	dietary-based TRVs for birds	chemical concentrations in heron prey collected throughout the LDW and in sediment collected from intertidal habitats throughout the LDW
Osprey	survival, growth, reproduction	potential exposure area: osprey may forage from the top meter of water throughout the LDW assessment scale: LDW-wide	dietary-based TRVs for birds	chemical concentrations in osprey prey collected throughout the LDW and in sediment collected from intertidal habitats throughout the LDW
Spotted sandpiper	survival, growth, reproduction	potential exposure area: sandpipers predominantly forage within small home range segments of the LDW assessment scale: three intertidal modeling areas	dietary-based TRVs for birds	chemical concentrations in sandpiper prey and sediment collected from intertidal habitats throughout the LDW

ROC	Assessment Endpoint	Assessment Scale	Measures of Effect	Measures of Exposure
River otter	survival, growth, reproduction	potential exposure area: river otters may forage throughout the LDW assessment scale: LDW-wide	dietary-based TRVs for mammals	chemical concentrations in river otter prey and sediment collected throughout the LDW
Harbor seal	survival, growth, reproduction	potential exposure area: harbor seals may forage throughout the LDW assessment scale: LDW-wide	dietary-based TRVs for mammals	chemical concentrations in harbor seal prey and sediment collected throughout the LDW

[a] A DMMP SL guideline is available for TBT; however, this guideline was not used in this ERA because it is based on an interstitial water concentration. TBT was included as a COPC based on the results of the Phase 1 ERA (Windward 2003b); the potential for adverse effects associated with exposure to TBT was evaluated using benthic invertebrate tissue data consistent with EPA (1999a) and Meador et al. (2002) and through a direct assessment of effects (i.e., imposex) on gastropods collected in the LDW.

DMMP – Dredged Material Management Program
LDW – Lower Duwamish Waterway
ROC – receptor of concern
SL – screening level
SMS – Washington State Sediment Management Standards
TBT – tributyltin
TRV – toxicity reference value
VOC – volatile organic compound

A.3.0 Exposure and Effects Assessment: Benthic Invertebrates

The benthic invertebrate community as a whole and crabs were selected as ROCs in the problem formulation to represent benthic invertebrates that may be exposed to sediment-associated chemicals in the LDW (Section A.2.3.1). In addition, site-specific measures of effect were evaluated for one benthic invertebrate group, meso- and neogastropods, because they are more sensitive to TBT exposure then other benthic invertebrates.

COPCs for the benthic invertebrate community were identified in the problem formulation (Section A.2.5.1). Forty-four chemicals based on the sediment chemistry data and one chemical based on porewater chemistry data were retained as COPCs for the benthic invertebrate community. TBT was also retained as a COPC based on the Phase 1 ERA (2003b). Two chemicals (total PCBs and zinc) were identified as COPCs for crabs in the problem formulation (Section A.2.5.1.3)

This section is divided into a benthic invertebrate community exposure assessment (Section A.3.1), benthic invertebrate community effects assessment (Section A.3.2), crab exposure assessment (Section A.3.3), and crab effects assessment (Section A.3.4). The exposure and effects data presented in this section are combined in the risk characterization (Section A.6.1).

A.3.1 Benthic Invertebrate Community Exposure Assessment

This section presents the exposure assessment for the benthic invertebrate community based on surface sediment chemistry data for all COPCs except two: cis-1,2-dichloroethene and TBT. Exposures to cis-1,2-dichloroethene were assessed using porewater data. Exposures to TBT were assessed using a critical tissue-residue approach. TBT effects were also evaluated using a direct measure of effect on meso- and neogastropods, as discussed further in Section A.3.2.4. The sediment exposure assessment for all COPCs, except TBT, is presented in Section A.3.1.1, the porewater exposure assessment is presented in Section A.3.1.2, and the TBT exposure assessment is presented in Section A.3.1.3.

A.3.1.1 Sediment exposure assessment

In this section, surface sediment data for COPCs are presented to characterize the exposure regime for the benthic invertebrate community. Concentrations and detection frequencies of COPCs in the baseline surface sediment dataset[32] are

[32] In addition to surface sediment grab samples, representing chemical concentrations within the upper 10 cm of sediment at a particular location, the baseline dataset also includes co-located sediment samples from the clam and benthic invertebrate sampling events (Windward 2005b, d, e). These co-

presented in Table A.3-1. Benthic invertebrates have small home ranges. Therefore, exposure is assessed based on the concentration of a COPC at a particular location. The spatial distribution of concentrations is also relevant from a risk perspective for the benthic invertebrate community as a whole as a food resource for other species in the LDW. Spatial scale is discussed in the risk characterization (Section A.6.1.1) and uncertainties associated with assessing spatial scale are presented in the uncertainty analysis (Section A.6.1.1). Chemicals with RLs greater than the SQS or CSL chemical criteria are discussed in the uncertainty analysis and are also discussed in the Phase 2 RI. The locations of surface sediment samples are shown on Map A.2-3.

Table A.3-1. Chemical concentrations and detection frequencies in LDW surface sediments for COPCs identified for the benthic invertebrate community

COPC	Unit	Detection Frequency (%)[a]	Detected Concentration		
			Minimum	Maximum	Median
Metals					
Arsenic	mg/kg dw	754 / 814 (93)	1.2	1,100	12
Cadmium	mg/kg dw	565 / 797 (71)	0.030 J	120	0.5
Chromium	mg/kg dw	811 / 811 (100)	4.8	1,100 J	29
Copper	mg/kg dw	814 / 814 (100)	5	12,000 J	52.3
Lead	mg/kg dw	814 / 814 (100)	2	23,000	36
Mercury	mg/kg dw	715 / 831 (86)	0.021	4.6 J	0.16
Nickel	mg/kg dw	771 / 773 (100)	5	910	22
Silver	mg/kg dw	481 / 782 (62)	0.020	270	0.42
Zinc	mg/kg dw	810 / 811 (100)	16	9,700	114
Organometals					
TBT[b]	µg/kg dw	143 /159 (90)	0.28J	3,000	29
PAHs					
2-Methylnaphthalene	µg/kg dw	139 / 780 (18)	1.0 J	3,300	27
Acenaphthene	µg/kg dw	301 / 790 (38)	1.0 J	5,200	40
Anthracene	µg/kg dw	552 / 790 (70)	2.0	10,000	82
Benzo(a)anthracene	µg/kg dw	717 / 790 (91)	3.6 J	8,400	195
Benzo(a)pyrene	µg/kg dw	716 / 784 (91)	5.8 J	7,900	200
Benzo(g,h,i)perylene	µg/kg dw	648 / 785 (83)	6.1	3,800	120
Total benzofluoranthenes	µg/kg dw	725 / 784 (92)	4.6	17,000	450
Chrysene	µg/kg dw	739 / 790 (94)	12	7,700	290
Dibenzo(a,h)anthracene	µg/kg dw	400 / 790 (51)	1.6 J	1,500	50
Dibenzofuran	µg/kg dw	246 / 789 (31)	1.0 J	4,200	33
Fluoranthene	µg/kg dw	759 / 790 (96)	18	24,000	430

located sediment samples consisted of surface sediment grab samples that were collected over a small area and composited.

COPC	Unit	Detection Frequency (%)[a]	Detected Concentration		
			Minimum	Maximum	Median
Fluorene	µg/kg dw	371 / 790 (47)	1.4 J	6,800	42
Indeno(1,2,3-cd)pyrene	µg/kg dw	692 / 785 (88)	6.5	4,300	130
Naphthalene	µg/kg dw	148 / 780 (19)	3.0 J	5,300	31
Phenanthrene	µg/kg dw	724 / 790 (92)	7.1	28,000	190
Pyrene	µg/kg dw	750 / 790 (95)	7 J	16,000	410
Total HPAH	µg/kg dw	767 / 790 (97)	20	85,000	2,100
Total LPAH	µg/kg dw	729 / 790 (92)	9.1	44,000	270
Phthalates					
Bis(2-ethylhexyl) phthalate	µg/kg dw	635 / 794 (80)	5.4	14,000	335
Butyl benzyl phthalate	µg/kg dw	390 / 784 (50)	2.0	7,100	40
Dimethyl phthalate	µg/kg dw	136 / 784 (17)	2.0 J	1,400 J	22
Other SVOCs					
1,2,4-Trichlorobenzene	µg/kg dw	5 / 778 (0.64)	1.6 J	72 J	2.9
1,2-Dichlorobenzene	µg/kg dw	18 / 778 (2.3)	1.3 J	520 J	2.2
1,4-Dichlorobenzene	µg/kg dw	35 / 778 (4.5)	0.74 J	1,600 J	6.4
2,4-Dimethylphenol	µg/kg dw	1 / 773 (0.13)	290 J	290 J	290
4-Methylphenol	µg/kg dw	78 / 793 (9.8)	4.8 J	4,600 J	33
Benzoic acid	µg/kg dw	69 / 781 (8.8)	54 J	4,500	220
Benzyl alcohol	µg/kg dw	14 / 771 (1.8)	8.2 J	670	25
Hexachlorobenzene	µg/kg dw	46 / 781 (5.9)	0.4 J	95 J	1.4
N-Nitrosodiphenylamine	µg/kg dw	23 / 780 (3.0)	6.5	230	8
Pentachlorophenol	µg/kg dw	12 / 747 (1.6)	14 J	410	97
Phenol	µg/kg dw	254 / 793 (32)	10 J	2,800	70
PCBs					
Total PCBs	µg/kg dw	1,203 / 1,288 (93)	1.6 J	220,000	150
Organochlorine Pesticides					
Total DDTs	µg/kg dw	78 / 197 (40)	0.72 J	2,900 J	8.6
Total chlordane[c]	µg/kg dw	33 / 197 (17)	0.20 J	230	3.7

[a] Number of detected concentrations per number of surface sediment samples analyzed for that chemical in the baseline dataset.

[b] TBT was evaluated using both a tissue-residue approach and imposex evaluation of gastropods. Sediment data are presented here for completeness.

[c] Total chlordane includes the calculated total chlordane for Phase 2 data and chlordane as reported in a subset of historical data (King County 1999b).

COPC – chemical of potential concern

dw – dry weight

HPAH – high-molecular-weight polycyclic aromatic hydrocarbon

J – estimated concentration

LPAH – low-molecular-weight polycyclic aromatic hydrocarbon

PAH – polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

SVOC – semivolatile organic compound

TBT – tributyltin

A.3.1.2 Porewater exposure assessment

As discussed in Section A.2.5.1.2 and in the *Quality Assurance Project Plan: Porewater Sampling of Lower Duwamish Waterway* (Windward 2005i), two sites, GWI and Boeing Plant 2/Jorgensen Forge, were selected as porewater sampling locations (Maps A.3-1 and A.3-2).

Cis-1,2-dichloroethene was the only VOC with a maximum detected concentration in porewater greater than a NOEC, and thus was the only COPC identified in porewater for the benthic invertebrate community. Tables A.3-2 and A.3-3 present the concentrations of cis-1,2-dichloroethene detected in porewater samples collected using peepers at GWI and Boeing Plant 2/Jorgensen Forge locations, respectively. Cis-1,2-dichloroethene was detected at all locations at GWI and at six locations at Boeing Plant 2/Jorgensen Forge. Porewater from GWI peeper PE-06 had the highest concentration of cis-1,2-dichloroethene (2,900 µg/L). This peeper was adjacent to a known seep (seep S-13), where cis-1,2-dichloroethene was detected at a concentration of 5,400 µg/L in an earlier sampling event (Windward 2005i). The other porewater samples collected at GWI had detected concentrations of cis-1,2-dichloroethene ranging from 0.5 µg/L to 630 µg/L. The concentrations of cis-1,2-dichloroethene in porewater samples collected at Boeing Plant 2/Jorgensen Forge were equal to or less than 1.7 µg/L. The evaluation is based on individual data points because benthic invertebrates have limited home ranges.

Table A.3-2. Concentrations of cis-1,2-dichloroethene in porewater samples from GWI

Sample ID	cis-1,2-Dichloroethene Concentration (µg/L)
LDW-PW-G-PE-01	6.1
LDW-PW-G-PE-02	46
LDW-PW-G-PE-03	0.5
LDW-PW-G-PE-04	2.4
LDW-PW-G-PE-05	630
LDW-PW-G-PE-06	2,900
LDW-PW-G-PE-07	18
LDW-PW-G-PE-08	20
LDW-PW-G-PE-203[a]	41
LDW-PW-G-PE-204[a]	27

[a] Field replicates from LDW-PW-G-PE-08.

ID – identification

Table A.3-3. Concentrations of cis-1,2-dichloroethene in porewater samples from Boeing Plant 2/Jorgensen Forge

Sample ID	cis-1,2-Dichloroethene Concentration (µg/L)
LDW-PW-B-PE-09	0.4
LDW-PW-B-PE-10	0.2 U
LDW-PW-B-PE-201[a]	0.4
LDW-PW-B-PE-202[a]	1.0
LDW-PW-B-PE-11	1.7
LDW-PW-B-PE-12	0.9
LDW-PW-B-PE-13	0.5
LDW-PW-B-PE-14	0.2
LDW-PW-B-PE-15	0.2 U
LDW-PW-B-PE-16	0.2 U

[a] Field replicates from LDW-PW-G-PE-10.

ID – identification

U – not detected at reporting limit shown

A.3.1.3 TBT exposure assessment

TBT was identified as a COPC for the benthic invertebrate community in Phase 1. The Phase 2 RI work plan (Windward 2004e) identified two approaches to evaluate risks to the benthic invertebrate community from TBT. The first approach included a direct measurement of an effect endpoint on meso- and neogastropods, which are the marine invertebrates that are most sensitive to TBT. The potential for effects was directly evaluated through site-specific assessments of imposex in gastropods collected from the LDW over a range of TBT concentrations in sediment (see Section A.3.2.4 for details on this approach). The second approach evaluated potential effects from TBT exposure on survival, growth, and reproduction of benthic invertebrates using a critical tissue-residue approach.

Gastropod sampling events, which were conducted in 2004 and 2005, are detailed in the gastropod pilot survey (Windward 2004f) and 2005 gastropod imposex study (Windward 2005j). Gastropod sampling locations are shown on Map A.3-3. Attempts were made to collect gastropods in both intertidal and subtidal habitats in the LDW. Gastropods were not found at the intertidal locations[33] surveyed but were found in most of the subtidal locations surveyed. Gastropod abundances were higher in the downstream, more saline portions of the LDW (Windward 2006c).

[33] A wide range of intertidal habitats with a range of chemical concentrations were surveyed for gastropods. It is not known why no gastropods were found at intertidal locations.

To assess exposure of the benthic invertebrate community to TBT, co-located benthic invertebrate tissue and sediment samples were collected and analyzed[34] as part of Phase 2 from 20 locations (Table A.3-4) (10 intertidal and 10 subtidal locations throughout the LDW [Map A.3-4]). All benthic invertebrates, including polychaetes, amphipods, and mollusks collected at each of the 20 locations, were combined into one composite tissue sample for each of the 20 locations to meet the biomass requirements for chemical analyses.

The cumulative frequency distribution of TBT concentrations in the co-located sediment samples is provided in Figure A.3-1. This figure demonstrates the range of TBT sediment concentrations sampled. A significant non-linear regression relationship was observed between the TBT concentrations in benthic invertebrate tissue and co-located sediment samples (Attachment 11). The regression equation was:

$$\text{tissue} = 145.4 \times \text{sediment}^{0.1801} \ (r^2 = 0.587) \qquad \textbf{Equation 3-1}$$

The maximum TBT concentration estimated for tissue (610 µg/kg dw) was estimated using the regression equation and the maximum TBT concentration in sediment (3,000 µg/kg dw). The estimated maximum tissue concentration was similar to the empirical maximum concentration (550 µg/kg dw) because of the distribution of the tissue sampling locations relative to the entire sediment dataset for TBT (Figure A.3-1).

Table A.3-4. Tributyltin concentrations in benthic invertebrate tissue samples and co-located sediment samples

Sample Location	Benthic Invertebrate Tissue TBT Concentration (µg/kg dw)	Sediment TBT Concentration (µg/kg dw)
Intertidal locations		
B1a	60 U	0.35 J
B2a	280	22
B3a	120	2.1
B4a	320	32
B5a-2	140	6.4
B6a	96	2.3
B7a	450	5.6
B8a	270	5.8
B9a	97	1.6 J
B10a	38 J	3.6
Subtidal locations		
B1b	550	2,300 J
B2b	500	63

[34] The benthic invertebrate tissue samples were also analyzed for a variety of metals, SVOCs, PCBs, and pesticides (Windward 2005b).

Sample Location	Benthic Invertebrate Tissue TBT Concentration (µg/kg dw)	Sediment TBT Concentration (µg/kg dw)
B3b	310	320
B4b	360	96
B5b	280	30
B6b	250	20
B7b	270	13
B8b[a]	140	1.7 J
B9b[a]	170	6.7
B10b[a]	74	2.3

[a] Originally designated subtidal in the quality assurance project plan (QAPP) for the collection effort (Windward 2004d) but designated intertidal in the baseline dataset used in the ERA and HHRA.

dw – dry weight

J – estimated concentration

TBT – tributyltin



Figure A.3-1. Frequency distribution of TBT concentrations in LDW sediment samples

A.3.2 BENTHIC INVERTEBRATE COMMUNITY EFFECTS ASSESSMENT

This section presents the effects assessment for the benthic invertebrate community. The potential effects of sediment-associated COPCs on the benthic invertebrate community were evaluated through:

- Comparisons of surface sediment chemical concentrations with SQS and CSL chemical criteria from the Washington State SMS (Section A.3.2.1)
- Site-specific sediment toxicity tests (Section A.3.2.2.)

Possible effects of the single COPC identified in porewater were evaluated using NOECs and LOECs selected from the literature (Section A.3.2.3).

Effects of exposure to TBT were evaluated using:

- Direct assessment of imposex on LDW-collected gastropods (Section A.3.2.4.1)
- Tissue-based toxicological data in the literature (for benthic invertebrates) (Section A.3.2.4.2)

Information on effects of COPCs presented in this section is combined with the exposure data in the risk characterization (Section A.6.1), and the uncertainties are discussed in the uncertainty analysis (Section A.6.1).

A.3.2.1 Sediment effects assessment

Effects on the benthic invertebrate community were assessed by comparing the COPC concentrations in LDW surface sediment to the SQS CSL chemical criteria derived in 1991 when Ecology adopted the SMS (WAC 173-204). These criteria are based on AETs (defined in Section A.2.5.1.1) developed for the Puget Sound Estuary Program (PSEP) (Barrick et al. 1988). An AET is the highest "no effect" chemical-specific sediment concentration above which a significant adverse biological effect always occurred among the several hundred samples used in its derivation. The methods used to calculate the AETs are described by Barrick et al. (1988) and Gries and Waldow (1996).

AETs were empirically derived using data from field-collected sediment samples that contained diverse chemical mixtures analyzed simultaneously for chemistry and toxicity. The data used to derive the 1988 AETs were collected from various locations in Puget Sound between March 1982 and September 1986. AETs for four endpoints[35] (i.e., amphipod mortality, abnormal development of oyster larvae, benthic invertebrate community abundance, and Microtox® bioluminescence) were developed for 47 chemicals. In general, the lowest AET for each chemical was identified as the SQS; the second lowest AET was identified as the CSL. The SQS corresponds to a sediment quality that will result in no adverse effects to biological resources; the CSL

[35] The specific tests associated with each of these endpoints are described in greater detail in the SMS rule (WAC 173-204).

corresponds to a sediment quality that will result in minor adverse effects (WAC 173-204). Table A.3-5 presents the biological effect endpoints that provide the basis for the SQS and CSL chemical criteria for COPCs.

Table A.3-5. Biological effect endpoints used to determine the SQS and CSL for COPCs

Chemical	SQS	CSL	Unit	Biological Endpoint Used to Establish SQS	Biological Endpoint Used to Establish CSL
Metals					
Arsenic	57	93	mg/kg dw	community abundance	amphipod mortality
Cadmium	5.1	6.7	mg/kg dw	community abundance	amphipod mortality
Chromium	260	270	mg/kg dw	community abundance	amphipod mortality
Copper	390	390	mg/kg dw	oyster abnormality	Microtox®
Lead	450	530	mg/kg dw	community abundance	Microtox®
Mercury	0.41	0.59	mg/kg dw	Microtox®	oyster abnormality
Silver	6.1	6.1	mg/kg dw	amphipod mortality	amphipod mortality
Zinc	410	960	mg/kg dw	community abundance	amphipod mortality
PAHs					
2-Methylnaphthalene	38	64	mg/kg OC	na	community abundance
Acenaphthene	16	57	mg/kg OC	oyster abnormality	community abundance
Anthracene	220	1,200	mg/kg OC	community abundance	amphipod mortality
Benzo(a)anthracene	110	270	mg/kg OC	oyster abnormality	amphipod mortality
Total benzofluoranthenes	230	450	mg/kg OC	oyster abnormality	amphipod mortality
Benzo(a)pyrene	99	210	mg/kg OC	oyster abnormality	amphipod mortality
Benzo(g,h,i)perylene	31	78	mg/kg OC	oyster abnormality	amphipod mortality
Chrysene	110	460	mg/kg OC	oyster abnormality	amphipod mortality
Dibenzo (a,h)anthracene	12	33	mg/kg OC	na	Microtox®
Fluoranthene	160	1,200	mg/kg OC	oyster abnormality	community abundance
Fluorene	23	79	mg/kg OC	oyster abnormality	community abundance
Indeno (1,2,3,-c,d)pyrene	34[a]	88	mg/kg OC	oyster abnormality	amphipod mortality
Naphthalene	99	170	mg/kg OC	oyster abnormality	community abundance
Phenanthrene	100[b]	480	mg/kg OC	oyster abnormality	community abundance
Pyrene	1,000	1,400	mg/kg OC	amphipod mortality	community abundance
HPAH	960	5,300	mg/kg OC	oyster abnormality	amphipod mortality
LPAH	370	780	mg/kg OC	oyster abnormality	community abundance
Phthalates					
Bis(2-ethylhexyl) phthalate	47	78	mg/kg OC	Microtox®	amphipod mortality
Butyl benzyl phthalate	4.9	64	mg/kg OC	Microtox®	community abundance
Dimethyl phthalate	53	53	mg/kg OC	amphipod mortality	community abundance
Other SVOCs					
1,2,4-Trichlorobenzene	0.81	1.8	mg/kg OC	Microtox®	amphipod mortality
1,2-Dichlorobenzene	2.3	2.3	mg/kg OC	oyster abnormality	community abundance and Microtox®
1,4-Dichlorobenzene	3.1	9	mg/kg OC	oyster abnormality	amphipod mortality
2,4-Dimethylphenol	29	29	µg/kg dw	oyster abnormality	Microtox®
4-Methylphenol	670	670	µg/kg dw	oyster abnormality	Microtox®

CHEMICAL	SQS	CSL	UNIT	BIOLOGICAL ENDPOINT USED TO ESTABLISH SQS	BIOLOGICAL ENDPOINT USED TO ESTABLISH CSL
Benzoic acid	650	650	µg/kg dw	oyster abnormality	community abundance, Microtox®
Benzyl alcohol	57	73	µg/kg dw	Microtox®	oyster abnormality
Dibenzofuran	15	58	mg/kg OC	oyster abnormality	community abundance
Hexachlorobenzene	0.38	2.3	mg/kg OC	community abundance	Microtox®
n-Nitrosodiphenylamine	11	11	mg/kg OC	community abundance	community abundance
Pentachlorophenol	360	690	µg/kg dw	amphipod mortality	community abundance
Phenol	420	1,200	µg/kg dw	oyster abnormality	amphipod mortality, community abundance, and Microtox®
PCBs					
Total PCBs	12	65	mg/kg OC	Microtox®	community abundance

Source: Washington State Sediment Management Standards (WAC 173-204); Barrick et al. (1988).

[a] The SQS for indeno(1,2,3,-c,d)pyrene is 34 mg/kg OC; the lowest AET, based on oyster abnormality, is 33 mg/kg OC.

[b] The SQS for phenanthrene is 100 mg/kg OC; the lowest AET, based on oyster abnormality, is 120 mg/kg OC.

CSL – cleanup screening level
dw – dry weight
HPAH – high-molecular-weight polycyclic aromatic hydrocarbon
LPAH – low-molecular-weight polycyclic aromatic hydrocarbon
na – not available
OC – organic carbon
PAH – polycyclic aromatic hydrocarbon
PCB – polychlorinated biphenyl
SQS – sediment quality standard
SVOC – semivolatile organic compound

Nickel, total DDTs, and total chlordane were identified as additional COPCs in the screening process. TRVs for nickel were based on toxicologically based DMMP guidelines for nickel. Because the DMMP guidelines for total DDTs and total chlordane were not toxicologically based,[36] TRVs were selected from the scientific literature. A search was conducted for relevant sediment toxicity studies for DDT and its metabolites and for chlordane using two databases, ECOTOX and BIOSIS. The databases were searched for relevant toxicity studies involving invertebrate species with growth, mortality, or reproductive endpoints. No toxicity studies were available in the scientific literature for chlordane. However, sediment quality guidelines (SQGs) and AETs other than those used to set SMS were available and were used to derive TRVs for chlordane (Table A.3-6). The lowest SQG (4.79 µg/kg dw) was selected as the LOAEL TRV. The highest NOAEL (2.8 µg/kg dw) below the LOAEL was selected as the NOAEL TRV. For DDTs, the lowest LOAEL (1,063 µg/kg dw) based on an individual laboratory toxicity study was selected as the LOAEL TRV. The NOAEL (567 µg/kg dw) for the same endpoint (survival) from the same study was selected as the NOAEL TRV. Table A.3-7 presents the basis for the screening level (SL) and maximum level (ML) guidelines for nickel and the selected TRVs for total DDTs and total chlordane. All reviewed sediment toxicity studies are presented in Attachment 6.

[36] The DMMP SLs for pesticides were selected to be approximately equal to the limit of quantification (i.e., 5 times the instrument detection limit). Insufficient data were available to establish the MLs for pesticides (PSDDA 1988b).

Table A.3-6. Available toxicity studies for selection of chlordane and total DDTs TRVs

Analyte	NOAEL (µg/kg dw)	LOAEL (µg/kg dw)	Biological Endpoint Determining NOAEL	Biological Endpoint Determining LOAEL	Source
Chlordane	2.8	na	amphipod AET	na	Gries and Waldow (1996)
Chlordane	2.26	4.79	All components of the aquatic ecosystem (e.g., bacteria, algae, macrophytes, invertebrates, fish) were considered, if data were available.	All components of the aquatic ecosystem (e.g., bacteria, algae, macrophytes, invertebrates, fish) were considered, if data were available.	Canadian environmental quality guidelines (CCME 2002)
p,p-DDE	9	na	benthic AET	na	Gries and Waldow (1996)
p,p-DDD	16	na	benthic AET	na	Gries and Waldow (1996)
p,p-DDT	12	na	echinoderm AET	na	Gries and Waldow (1996)
Total DDTs	24	na	amphipod AET	na	Gries and Waldow (1996)
Total DDT	8510	na	*Neanthes arenaceodentata*; full life-cycle toxicity test (survival, growth, fecundity, reproduction)	na	Chapman (1996)
DDTs	567	1,063	*Hyalella azteca* survival	*Hyalella azteca* survival	Lotufo et al. (2001b)
DDTs	na	1,985	na	*Leptocheirus plumulosus* survival	Lotufo et al. (2001a)
DDTs	na	2,910	na	*Diporeia* spp. survival	Lotufo et al. (2001b)
DDTs	na	3,510	na	*Hyalella azteca* survival	Lotufo et al. (2001b)
DDD	1,200	4,000[a]	*Hyalella azteca* survival	*Hyalella azteca* survival	Ingersoll et al. (2005)
DDD	1,200	4,000[a]	*Hyalella azteca* reproduction	*Hyalella azteca* reproduction	Ingersoll et al. (2005)
DDT	na	6,180	na	*Rhepoxynius abronius* survival	Murdoch et al. (1997)
Total DDT	na	7,500	na	significant reduced feeding rate in marine polychaete (*Heteromastus filiformis*)	Mulsow and Landrum (1995)
Total DDT	na	11,000	na	*Hyalella azteca* LC50	Nebeker et al. (1989)
DDTs	na	308,000	na	*Neanthes arenaceodentata* growth	Lotufo et al. (2000)

[a] The Ingersoll study also evaluated a growth endpoint. There was no dose response for the growth endpoint with the lowest effect value at 30 µg/kg dw.

AET – apparent effects threshold

CCME – Canadian Council of Ministers of the Environment

dw – dry weight

na – not available

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

TRV – toxicity reference value

Table A.3-7. Biological effect endpoints for DMMP guidelines and selected TRVs

Chemical	Unit	SL or NOAEL	ML or LOAEL	Biological Endpoint Determining SL or NOAEL	Biological Endpoint Determining ML or LOAEL
Nickel	mg/kg dw	140[a]	370[a]	amphipod mortality and community abundance	nr
Total DDTs	µg/kg dw	567[b]	1,063[b]	amphipod mortality	amphipod mortality
Total chlordane	µg/kg dw	2.8[c]	4.79[d]	amphipod mortality	CCME (2002)

[a] Source: USACE et al. (2000); Barrick et al. (1988).

[b] Literature-based TRV – (Lotufo et al. 2001b).

[c] Literature-based TRV – PSDDA AET evaluation 1994 (Gries and Waldow 1996).

[d] Literature-based TRV – probable effect levels for chlordane (CCME 2002).

CCME – Canadian Council of Ministers of the Environment

DMMP – Dredged Material Management Program

dw – dry weight

LC50 – concentration that causes the death of 50% of a group of test animals

LOAEL – lowest-observed-adverse-effect level

ML – maximum level (DMMP)

nr – not reported

NOAEL – no-observed-adverse-effect level

SL – screening level (DMMP)

A.3.2.2 Site-specific toxicity tests assessment

This section describes the results of site-specific toxicity tests conducted on LDW sediment samples to assess the potential effects of sediment-associated chemicals on benthic invertebrates. In general, toxicity testing was performed on sediment samples with at least one detected chemical exceeding SQS.[37] The SMS provide both chemical and biological effects criteria. Because AETs, which form the basis for the chemical criteria, are based on sediment samples with a mixture of chemicals from a large number of locations, toxicity tests either confirm or overrule the SMS designation based on sediment chemistry.

To generate more specific information about the nature and severity of effects on benthic invertebrates exposed to sediments with at least one chemical concentration exceeding the SQS, three toxicity tests were conducted with surface sediments (0 to 10 cm) collected at 48 locations (Map A.3-5) (Windward 2005d, e). The toxicity tests included:

- Acute 10-day amphipod (*Eohaustorius estuarius*) mortality test
- Acute 48-hr bivalve larvae (*Mytilus galloprovincialis*) normal survival test
- Chronic 20-day juvenile polychaete (*Neanthes arenaceodentata*) survival and growth test

[37] Two samples were also tested that did not have any SQS exceedances because they were near sources and were requested by Ecology. In addition, several samples were not tested if they were heavily contaminated; these samples were assumed to be toxic.

The results from the three sediment toxicity tests were evaluated using the SMS rules for marine toxicity tests (Ecology 2003b). The performance criteria and biological effects criteria (SQS and CSL of the SMS) are summarized in Table A.3-8.

Table A.3-8. SMS biological effects criteria for marine sediment toxicity tests

Toxicity Test	Biological Effects Criteria	
	SQS	CSL
Amphipod	mean mortality > 25% on an absolute basis, and statistically different from the reference sediment ($p \leq 0.05$)	mean mortality greater than the value in the reference sediment plus 30%, and statistically different from the reference sediment ($p \leq 0.05$)
Polychaete[a]	mean individual growth rate < 70% of that of the reference sediment and statistically different ($p \leq 0.05$)	mean individual growth rate < 50% of that of the reference sediment and statistically different ($p \leq 0.05$)
Bivalve larvae	mean normal survivorship < 85% of that of the reference sediment and statistically different ($p \leq 0.10$)	mean normal survivorship <70% of that of the reference sediment and statistically different ($p \leq 0.10$)

[a] The mortality endpoint for the polychaete toxicity test is not used for determination of SMS compliance.

CSL – cleanup screening level

SMS – Washington State Sediment Management Standards

SQS – sediment quality standard

For the amphipod mortality endpoint, 11 of the 48 samples failed the biological effects criteria of the SMS at the CSL level, and 5 of the 48 samples failed the biological effects criteria at the SQS level (Table A.3-9). For the polychaete growth endpoint, 8 of the 48 samples failed the biological effects criteria at the SQS level; no samples failed the polychaete criteria at the CSL level. For the bivalve survival/development endpoint, 8 of the 48 samples failed the biological effects criteria at the CSL level, and 12 of the 48 samples failed the biological effects criteria at the SQS level.

An exceedance of the SQS biological effects criteria in any two toxicity tests at one location is considered a CSL exceedance at that location (WAC 173-204-420(3)). Based on this guideline, of the 48 samples tested, 18 sediment samples did not exceed the biological effects criteria, 11 sediment samples exceeded the SQS biological effects criteria, and 19 sediment samples exceeded the CSL biological effects criteria (Table A.3-9; Map A.3-5).

Table A.3-9. Results of Phase 2 site-specific toxicity testing of surface sediment samples from the LDW

Sample ID	Amphipod Toxicity Test		Polychaete Toxicity Test			Bivalve Larval Toxicity Test		Overall SMS Exceedance
	Percent Mean Mortality ± SD	SMS Exceedance[a, b]	Mean Mortality ± SD	Mean Individual Growth Rate (mg/day) ± SD	SMS Exceedance[a, c, d]	Percent Mean Normal Survivorship ± SD	SMS Exceedance[a, e]	
LDW-SS2-010	39.0 ± 16.0	CSL	0.0 ± 0.0	0.76 ± 0.20	no exceedances	31.8 ± 15.1	CSL	CSL
LDW-SS6-010	47.0 ± 21.7	CSL	4.0 ± 8.9	0.81 ± 0.04	no exceedances	23.6 ± 16.3	CSL	CSL
LDW-SS15-010	28.0 ± 8.4	SQS	0.0 ± 0.0	0.73 ± 0.10	no exceedances	75.4 ± 8.7	no exceedances	SQS
LDW-SS16-010	16.0 ± 10.2	no exceedances	4.0 ± 8.9	0.96 ± 0.13	no exceedances	64.3 ± 3.2	SQS	SQS
LDW-SS17-010	35.0 ± 20.9	SQS	0.0 ± 0.0	0.92 ± 0.22	no exceedances	62.9 ± 7.6	no exceedances	SQS
LDW-SS21-010	37.0 ± 23.6	CSL	0.0 ± 0.0	0.99 ± 0.10	no exceedances	72.9 ± 3.6	no exceedances	CSL
LDW-SS22-010	32.0 ± 10.4	SQS	4.0 ± 8.9	0.70 ± 0.13	SQS	51.9 ± 1.7[f]	SQS	CSL
LDW-SS24-010	7.0 ± 4.5	no exceedances	0.0 ± 0.0	0.77 ± 0.17	SQS	18.3 ± 3.3	CSL	CSL
LDW-SS26-010	23.0 ± 11.0	no exceedances	0.0 ± 0.0	0.86 ± 0.17	no exceedances	76.7 ± 7.4	no exceedances	no exceedances
LDW-SS29-010	12.0 ± 7.6	no exceedances	0.0 ± 0.0	0.90 ± 0.08	no exceedances	64.7 ± 6.8	no exceedances	no exceedances
LDW-SS31-010	43.0 ± 5.7	CSL	4.0 ± 8.9	0.81 ± 0.10	no exceedances	62.9 ± 7.6	SQS	CSL
LDW-SS32-010	34.0 ± 11.9	SQS	0.0 ± 0.0	0.69 ± 0.10	no exceedances	78.9 ± 15.7[f]	no exceedances	SQS
LDW-SS37-010	45.0 ± 14.6	CSL	0.0 ± 0.0	0.78 ± 0.05	no exceedances	65.8 ± 19.2[g]	SQS	CSL
LDW-SS39-010	29.0 ± 11.4	SQS	0.0 ± 0.0	0.76 ± 0.12	SQS	83.4 ± 10.1	no exceedances	CSL
LDW-SS40-010	36.0 ± 12.9	CSL	0.0 ± 0.0	0.78 ± 0.14	no exceedances	79.7 ± 4.9	no exceedances	CSL
LDW-SS49-010	49.0 ± 19.5	CSL	0.0 ± 0.0	0.79 ± 0.22	no exceedances	55.1 ± 17.4	SQS	CSL
LDW-SS50-010	39.0 ± 10.8	CSL	0.0 ± 0.0	0.73 ± 0.11	no exceedances	70.2 ± 10.2	no exceedances	CSL
LDW-SS56-010	6.0 ± 4.2	no exceedances	0.0 ± 0.0	0.85 ± 0.14	no exceedances	67.6 ± 7.2	SQS	SQS
LDW-SS57-010	13.0 ± 12.5	no exceedances	0.0 ± 0.0	0.78 ± 0.13	no exceedances	55.3 ± 15.1	CSL	CSL
LDW-SS58-010	5.0 ± 5.0	no exceedances	0.0 ± 0.0	0.69 ± 0.07	SQS	61.0 ± 4.0	SQS	CSL
LDW-SS60-010	7.0 ± 5.7	no exceedances	4.0 ± 8.9	0.77 ± 0.17	no exceedances	84.7 ± 6.0	no exceedances	no exceedances
LDW-SS63-010	5.0 ± 6.1	no exceedances	0.0 ± 0.0	0.68 ± 0.19	no exceedances	80.0 ± 1.6	no exceedances	no exceedances

Sample ID	Amphipod Toxicity Test		Polychaete Toxicity Test			Bivalve Larval Toxicity Test		Overall SMS Exceedance
	Percent Mean Mortality ± SD	SMS Exceedance[a, b]	Mean Mortality ± SD	Mean Individual Growth Rate (mg/day) ± SD	SMS Exceedance[a, c, d]	Percent Mean Normal Survivorship ± SD	SMS Exceedance[a, e]	
LDW-SS68-010	12.0 ± 9.1	no exceedances	0.0 ± 0.0	0.85 ± 0.10	no exceedances	71.6 ± 12.5	no exceedances	no exceedances
LDW-SS69b-010	37.0 ± 15.7	CSL	0.0 ± 0.0	0.85 ± 0.29	no exceedances	59.0 ± 13.6	SQS	CSL
LDW-SS70-010	15.0 ± 7.1	no exceedances	4.0 ± 8.9	0.78 ± 0.12	no exceedances	60.7 ± 12.0	SQS	SQS
LDW-SS71-010	5.0 ± 6.1	no exceedances	0.0 ± 0.0	0.92 ± 0.16	no exceedances	61.8 ± 8.7	no exceedances	no exceedances
LDW-SS73-010	12.0 ± 9.1	no exceedances	0.0 ± 0.0	0.86 ± 0.11	no exceedances	56.8 ± 13.3	SQS	SQS
LDW-SS75-010	8.0 ± 7.6	no exceedances	4.0 ± 8.9	0.69 ± 0.16	no exceedances	76.5 ± 7.5	no exceedances	no exceedances
LDW-SS77-010	16.0 ± 9.6	no exceedances	4.0 ± 8.9	0.95 ± 0.12	no exceedances	10.1 ± 4.0	CSL	CSL
LDW-SS85-010	1.0 ± 2.2	no exceedances	0.0 ± 0.0	0.87 ± 0.18	no exceedances	86.8 ± 5.3	no exceedances	no exceedances
LDW-SS88-010	48.0 ± 25.9	CSL	0.0 ± 0.0	0.68 ± 0.14	no exceedances	11.9 ± 5.3	CSL	CSL
LDW-SS89-010	5.0 ± 3.5	no exceedances	0.0 ± 0.0	0.84 ± 0.26	no exceedances	86.9 ± 5.1[g]	no exceedances	no exceedances
LDW-SS92-010	1.0 ± 2.2	no exceedances	0.0 ± 0.0	0.79 ± 0.17	no exceedances	78.3 ± 14.3	no exceedances	no exceedances
LDW-SS106-010	6.0 ± 4.2	no exceedances	0.0 ± 0.0	0.91 ± 0.11	no exceedances	61.4 ± 8.8	no exceedances	no exceedances
LDW-SS112-010	4.0 ± 4.2	no exceedances	0.0 ± 0.0	0.82 ± 0.06	no exceedances	65.4 ± 9.6	no exceedances	no exceedances
LDW-SS114-010	85.0 ± 7.1	CSL	4.0 ± 8.9	0.77 ± 0.19	no exceedances	56.6 ± 8.5	SQS	CSL
LDW-SS115-010	9.0 ± 4.2	no exceedances	0.0 ± 0.0	0.76 ± 0.21	no exceedances	77.6 ± 11.0	no exceedances	no exceedances
LDW-SS119-010	3.0 ± 2.7	no exceedances	0.0 ± 0.0	0.79 ± 0.06	no exceedances	68.8 ± 7.6	no exceedances	no exceedances
LDW-SS120-010	3.0 ± 2.7	no exceedances	0.0 ± 0.0	0.71 ± 0.09	no exceedances	56.3 ± 11.2	SQS	SQS
LDW-SS121-010	4.0 ± 4.2	no exceedances	0.0 ± 0.0	0.90 ± 0.12	no exceedances	76.4 ± 8.0	no exceedances	no exceedances
LDW-SS122-010	7.0 ± 4.5	no exceedances	0.0 ± 0.0	0.83 ± 0.14	no exceedances	68.5 ± 14.9	no exceedances	no exceedances
LDW-SS143-010	6.0 ± 5.5	no exceedances	0.0 ± 0.0	0.75 ± 0.07	no exceedances	72.8 ± 3.2	no exceedances	no exceedances
LDW-SS144-010	1.0 ± 2.2	no exceedances	0.0 ± 0.0	0.72 ± 0.11	SQS	66.4 ± 12.1	no exceedances	SQS[h]
LDW-SS148-010	6.0 ± 6.5	no exceedances	0.0 ± 0.0	0.78 ± 0.08	SQS	29.9 ± 6.6	CSL	CSL
LDW-SS157-010	8.0 ± 7.6	no exceedances	4.0 ± 8.9	0.78 ± 0.14	SQS	71.6 ± 8.5	no exceedances	SQS[h]
LDW-SS158-010	12.0 ± 4.5	no exceedances	0.0 ± 0.0	0.81 ± 0.10	no exceedances	67.5 ± 6.5	no exceedances	no exceedances

Sample ID	Amphipod Toxicity Test		Polychaete Toxicity Test			Bivalve Larval Toxicity Test		Overall SMS Exceedance
	Percent Mean Mortality ± SD	SMS Exceedance[a, b]	Mean Mortality ± SD	Mean Individual Growth Rate (mg/day) ± SD	SMS Exceedance[a, c, d]	Percent Mean Normal Survivorship ± SD	SMS Exceedance[a, e]	
LDW-SSB2b-010	25.0 ± 12.2	no exceedances	0.0 ± 0.0	1.02 ± 0.10	no exceedances	42.1 ± 20.0	CSL	CSL
LDW-SSB6a-010	2.0 ± 4.5	no exceedances	0.0 ± 0.0	0.82 ± 0.14	SQS	60.1 ± 13.3	no exceedances	SQS[h]

[a] Statistical analyses in SedQual Release 5 included Wilk-Shapiro test for normality and Levene's test for equality of variances followed by the appropriate statistical test for significance (i.e., Student's t-test, approximate t-test, or Mann-Whitney).

[b] SQS – mean mortality > 25% on an absolute basis and statistically different from the reference sediment ($p \leq 0.05$); CSL – mean mortality greater than the value in the reference sediment plus 30% and statistically different from the reference sediment ($p \leq 0.05$). Reference sediment results are presented in the Round 1 and Round 2 surface sediment data reports (Windward 2005d, e).

[c] SQS – mean individual growth rate <70% of that of the reference sediment and statistically different ($p \leq 0.05$).

[d] No exceedance was reported for the polychaete growth endpoint for some of the sediment samples because of high variability in the reference and/or test samples.

[e] SQS – mean normal survivorship < 85% of that of the reference sediment and statistically different ($p \leq 0.10$); CSL – mean normal survivorship < 70% of that of the reference sediment and statistically different ($p \leq 0.10$).

[f] One of the five replicates from each of these tests was double-inoculated, so those replicates were not used in calculating mean normal survivorship and mean effective mortality for those test sediments.

[g] One of the five replicates from each of these tests was not inoculated at test initiation, so those replicates were not used in calculating mean normal survivorship and mean effective mortality for those test sediments.

[h] Exceeded the SQS criterion based on reduction in polychaete growth alone.

CSL – cleanup screening level

ID – identification

SMS – Washington State Sediment Management Standard

SD – standard deviation

SQS – sediment quality standard

Twelve other site-specific toxicity studies have been conducted in the LDW since 1990[38] (Table A.3-10). However, only 1 of these 12 studies (with a total of 7 samples, as shown in bold in Table A.3-10) was conducted on surface sediments (0 to 15 cm) that remained in place and are thus relevant to this study. One other study, *Sediment Quality in Puget Sound* (NOAA and Ecology 2000), was a monitoring program intended to characterize only the most recently deposited surface sediment (0 to 2 cm). The remainder of the studies were dredged material characterization studies that tested sediments from the 0- to 4-ft (or deeper) horizons, and those sediments were subsequently dredged and removed from the LDW.

Collection locations for samples analyzed for sediment toxicity in the one relevant historical study (King County 2000a) are presented in Map A.3-5. The seven locations sampled by King County were selected to help determine the removal boundary for the Duwamish/Diagonal cleanup. Part of the area for the Duwamish/Diagonal cleanup study (King County 2000a) was dredged and capped in 2004. Three of the seven locations sampled were within the removal area and were dredged. However, because the baseline dataset includes pre-removal conditions, a summary of all results from that study is included.

The seven sediment samples collected for the Duwamish/Diagonal cleanup study (King County 2000a) were tested for toxicity using three standard SMS confirmatory tests (i.e., amphipod [*Rhepoxynius abronius*] mortality, echinoderm embryo effective mortality,[39] and polychaete growth). One of seven samples (L9443-7) failed the biological effects criteria of the SMS at the SQS level for both the echinoderm embryo effective mortality and the polychaete growth endpoints (Table A.3-11). The results of the polychaete and echinoderm embryo tests for sample L9443-7 were similar to the results from several other samples tested concurrently. The control sample to which the results from sample L9443-7 were compared had a higher polychaete growth rate (0.77 vs. 0.60 mg/day) and lower effective echinoderm embryo mortality (15 vs. 29%) compared to the reference sample to which all other test sediments were compared.[40]

Thus, based on the available data, one of the seven sediment samples analyzed by King County (2000a) exhibited toxicity in exceedance of the state's biological effects criteria.

---

[38] At the beginning of the RI process, EPA and Ecology agreed that data collected within the past 10 years would be suitable to characterize the LDW; therefore, data collected in 1990 or later were acceptable. While some sampling events are now older than 10 years, they are included in Table A.3-9 to remain consistent with the list of suitable data sets developed at the beginning of the process.

[39] Combined mortality and abnormal development.

[40] This sample was not compared to the other reference samples because the grain size was not comparable. The percent fines for sample L9443-7 (7.9%) was much lower than in the remaining samples (49.5% to 91.3%) and in-batch reference sediment (54.5%). Therefore, the SMS comparison was performed on the West Beach control sediment, which had < 10 % fines.

Table A.3-10. Sediment toxicity datasets meeting project data quality objectives

Study	Year Conducted	Number of Samples[a]	Source
Sediment Quality in Puget Sound	1998	3	Ecology (2000)
Duwamish/Diagonal CSO/SD Site Assessment Report – Draft[b]	1996	7	King County (2000)
Sediment Sampling and Analysis, James Hardie Gypsum Inc., Duwamish Waterway, Seattle, Washington	1998 – 1999	7	Spearman (1999)
Dredge Material Characterization, Hurlen Construction Company and Boyer Alaska Barge Lines Berthing Areas, Duwamish Waterway, Seattle, Washington	1998	4	Hart Crowser (1998)
Proposed Dredging of Slip No. 4, Duwamish River, Seattle, Washington	1995	4	PTI (1996)
PSDDA Chemical Characterization of Duwamish Waterway and Upper Turning Basin. FY97 Operations and Maintenance Dredging, Seattle, Washington	1996	3	Striplin Environmental (1996)
Lone Star Northwest and James Hardie Gypsum Kaiser Dock Upgrade, Duwamish Waterway PSDDA Sampling and Analysis Results	1995	4	Hartman Associates (1995)
Lone Star Northwest West Terminal, Duwamish River PSDDA Sampling and Analysis Results	1992	1	Hartman Associates (1992)
Sediment Sampling and Analysis, Brown Morton Properties, Duwamish Waterway, Seattle, Washington	1991	1	Spearman (1991a)
Sediment Sampling and Analysis, South Park Marina, Duwamish Waterway, Seattle, Washington	1991	2	Spearman (1991b)
PSDDA Bioassays for Duwamish Channel Sediments	1991	14	SAIC (1992)
Duwamish Channel and Settling Basin Sediment Bioassays	1990	4	PTI (1990)

[a] With the exception of the *Sediment Quality in Puget Sound* report and the *Duwamish/Diagonal CSO/SD Site Assessment Report,* subsurface sediment samples from these studies were collected for dredged material characterization studies. The tested sediments have been removed.

[b] Surface sediment was characterized in this study; all other studies were dredged material characterizations and are not discussed further.

CSO/SD – combined sewer overflow/storm drain

PSDDA – Puget Sound Dredged Disposal Analysis

Table A.3-11. Summary of site-specific sediment toxicity test results for surface sediment samples collected by King County at Duwamish/Diagonal CSO/SD site

Sampling Location[a]	Sample ID	Echinoderm Embryo Effective Mortality (%)	SQS or CSL Exceedance[b]	Polychaete Growth Rate (mg/day)	SQS or CSL Exceedance[c]	Amphipod Mortality (%)	SQS or CSL Exceedance[d]	Overall SMS Exceedance
West Beach	control	15	na	0.77	na	1.0	na	na
Carr Inlet	reference	29	na	0.60	na	8.0	na	na
DUD200	L9443-1	32	no exceedances	0.60	no exceedances	13	no exceedances	no exceedances
DUD201	L9443-2	35	no exceedances	0.55	no exceedances	21	no exceedances	no exceedances
DUD202	L9443-3	35	no exceedances	0.62	no exceedances	18	no exceedances	no exceedances
DUD203	L9443-4	33	no exceedances	0.59	no exceedances	22	no exceedances	no exceedances
DUD204	L9443-5	17	no exceedances	0.51	no exceedances	26	no exceedances	no exceedances
DUD205	L9443-6	16	no exceedances	0.54	no exceedances	19	no exceedances	no exceedances
DUD206	L9443-7[e]	34	SQS	0.52	SQS	4.0	no exceedances	CSL

Source: King County (2000a)

[a] Sampling locations DUD200 through DUD206 are shown on Map A.3-5.

[b] SQS – mean normal survivorship < 85% of that of the reference sediment and statistically different ($p \leq 0.10$); CSL – mean normal survivorship < 70% of that of the reference sediment and statistically different ($p \leq 0.10$).

[c] SQS – mean individual growth rate <70% of that of the reference sediment and statistically different ($p \leq 0.05$).

[d] SQS – mean mortality > 25% on an absolute basis and statistically different from the reference sediment ($p \leq 0.05$); CSL – mean mortality greater than the value in the reference sediment plus 30% and statistically different from the reference sediment ($p \leq 0.05$).

[e] Test results for this sample were statistically compared to control sample results rather than the reference sample results because reference sample was not a suitable grain size match.

CSL – cleanup screening level

CSO/SD – combined sewer overflow/storm drain

ID – identification

na – not applicable

SMS – Washington State Sediment Management Standards

SQS – sediment quality standard

A.3.2.3 Porewater effects assessment

As discussed in Section A.2.5.1.2, a literature search was conducted for relevant aquatic toxicity studies for VOCs using two databases, ECOTOX and BIOSIS. The databases were searched for relevant toxicity studies involving invertebrate species with growth, mortality (including immobilization), or reproductive (including developmental) endpoints. Studies with invertebrates were preferred because the purpose of the porewater study was to evaluate risks to the benthic invertebrate community. However, if there were no invertebrate data, then fish studies with growth, survival, or reproductive endpoints were included in the search. Toxicity data for both freshwater and marine species were included because of the wide salinity range in the LDW and the paucity of toxicity data. Guidelines for the selection of NOECs and LOECs were:

- The study identified with the lowest effect level (preferably a LOEC) for each VOC was selected if the study was acceptable.
- The study with the lowest NOEC for each VOC was selected provided that more than one NOEC was not available for a given test species and endpoint. When there were multiple NOECs for the same test species/endpoint for a given VOC, the study with the highest NOEC for that test species/endpoint was selected if the study was acceptable.
- The study must include negative control tests. For no-effect results, the study must have used an exposure period of no less than 48 hrs for daphnids and no less than 96 hrs for fish. In addition, to represent conditions relevant to those found in the LDW, the salinity must be no greater than 35 parts per thousand in tests using *Artemia salina* (brine shrimp).

Two studies evaluated the toxicity of 1,2-dichloroethene. The lowest toxicity value for 1,2-dichloroethene was derived from a study testing toxicity to *Artemia salina* (Sanchez-Fortun et al. 1997). This study was reviewed in detail and found to be acceptable. In this study, *Artemia* shrimp larvae were added to synthetic seawater at a salinity of 35 parts per thousand (ppt) in plastic petri dishes containing the appropriate concentration of 1,2-dichloroethene. Controls containing untreated synthetic seawater were also tested. The petri dishes were incubated at 25°C. Larvae were evaluated for mortality after 72 hours. The reported LC50 was 6,785 µg/L. The only other toxicity data available for this chemical is an LC50 of 140,000 µg/L from a bluegill study. Therefore, the lower LC50 for *Artemia* (6,785 µg/L) was selected as the TRV for 1,2-dichloroethene. Because no NOAELs were available for this chemical, the LC50-based LOAEL was divided by 50 to derive a NOAEL TRV of 136 µg/L.

A.3.2.4 TBT effects assessment

Potential adverse effects to benthic invertebrates from TBT exposure were evaluated using a direct measure of imposex in LDW-collected gastropods (the benthic

invertebrate most sensitive to TBT based on the imposex endpoint) and a critical tissue-residue approach for benthic invertebrates. The imposex assessment is presented in Section A.3.2.4.1 and the critical tissue-residue assessment is presented in Section A.3.2.4.2.

A.3.2.4.1 Imposex assessment

Of the benthic invertebrates that have been identified in the LDW, meso- and neogastropods[41] have been reported to be particularly sensitive to toxic effects of TBT (Meador et al. 2002). Specifically, at sufficiently high tissue concentrations, TBT is known to cause the development of male sexual organs in females in some meso- and neogastropod species. This condition, known as imposex (Gibbs et al. 1988), may interfere with gastropod reproduction and potentially results in population-level effects (Meador et al. 2002). Therefore, a site-specific evaluation of imposex in gastropods was conducted in the LDW to directly assess the toxic endpoint. This direct approach is preferable because it reduces uncertainty in the risk analysis. Imposex evaluation was performed on as many different meso-and neogastropods collected in the LDW as possible, including the three most abundant neogastropod species, *Nassarius mendicus, Astyris gausapata*, and *Olivella baetica*.

Mature female gastropods were examined for imposex using the methods reported in Oehlmann et al. (1991) and Spence et al. (1990). Oehlmann et al. (1991) determined the imposex stage based on the presence of male reproductive organs in females, including vas deferens[42] and penis. Table A.3-12 presents the imposex stage criteria for this evaluation. Complete sterilization is associated with stages 5 and 6. Stage 4 is generally referred to as transitional, and stages 1 through 3 are described as early stages.

Table A.3-12. Imposex stage criteria

IMPOSEX STAGE	CHARACTERISTICS
1	development of small penis or small section of vas deferens
2	development of either: 1. larger penis with a penis duct, or 2. two sections of vas deferens, or 3. both a penis and a vas deferens section
3	development of either: 1. larger penis with vas deferens section, or 2. a complete vas deferens, or 3. a larger penis with a penis duct and a vas deferens section
4	development of a larger penis with penis duct and a complete vas deferens (last fertile imposex stage)

[41] Mesogastropods and neogastropods are snails in the taxonomic orders of Mesogastropoda and Neogastropoda, respectively.

[42] A sperm-carrying duct.

IMPOSEX STAGE	CHARACTERISTICS
5	development of a prostate gland or occlusion of the vulva (infertile stage)
6	infertile stage with aborted capsules

Source: Oehlmann et al. (1991)

In imposexed specimens where the vas deferens could not be seen, penis length was measured in both genders to assess the level of imposex using the relative penis size (RPS) approach (Gibbs et al. 1988). The RPS index was calculated as follows:

$$\text{RPS index} = (\text{mean length of female penis}^3/\text{mean length of male penis}^3) \times 100 \qquad \textbf{Equation 3-2}$$

Imposex evaluation was performed on as many different meso- and neogastropods as possible collected in LDW. Three neogastropods, *Olivella baetica*, *Astyris gausapata,* and *Nassarius mendicus,* were collected in large numbers and assessed for imposex. All other meso- and neogastropods were collected in low numbers (one to five); and because the majority of the specimens were immature, only two mature female mesogastropods (one of each *Polinices* sp. and *Lacuna vincta*) could be assessed. Imposex was not present in these mesogastropods.

No signs of imposex were found in 127 female *A. gausapata,* which was the most abundant gastropod species in the LDW, over a range of sediment TBT concentrations from 94 to 3,000 µg/kg dw.[43] Similarly, no signs of imposex were found in 19 female *O. baetica* collected at one location with a sediment TBT concentration of 2,300 µg/kg dw. Imposex was observed only in *N. mendicus* females collected at locations with sediment TBT concentrations ranging from 34 to 358 µg/kg dw. The degree of imposex in all female *N. mendicus* examined was found to be similar to Stage 2, except that in all cases penises were present but no vasa deferentia[44] were observed (Windward 2006c). Because the vas deferens could not be seen, the RPS index was calculated. The index ranged from 0.1% to 2.6% (Table A.3-13). A site-wide RPS index of 1.8% was also calculated for all of the locations sampled in 2005 by using the mean female penis length and the mean male penis length from all locations. A station-specific RPS index was also calculated for the two stations that had males and at least one female with imposex. The RPS indices for these two stations (G17b and G18b) were 2.0% and 3.4%, respectively. According to Spence et al. (1990), in general, sterile females are absent at RPS indices below 5%, between 5% and 40% the percentage of sterility increases, and at RPS indices exceeding 40%, most or all females are sterile.

[43] The sediment TBT concentrations were not analyzed synoptically with the imposex evaluation but were analyzed during previous sampling events.

[44] Plural of vas deferens.

Table A.3-13. TBT concentrations in sediment and *Nassarius mendicus* RPS indices

Location ID	TBT Concentration in Sediment (µg/kg dw)[a]	Mean Female Penis Length (mm) (±st dev) (number measured)	Mean Male Penis Length (mm) (number measured)[b]	RPS Index (%)
2004				
G1b	358	0.86 (1)	6.83 (7)	0.2
G2b	144	1.56 ± 0.49 (4)	6.83 (7)	1.2
G3b	94	1.8 (1)	6.83 (7)	1.8
G6b	34	0.70 ± 0.56 (3)	6.83 (7)	0.1
G8b	117	1.1 (1)	6.83 (7)	0.4
2005				
G17b	320	3.3 (1)	11.2 ± 1.5 (5)	2.6
G18b	350	3.1 ± 0.9 (2)	11.2 ± 1.5 (5)	2.1
G19b	250	2.6 ± 0.1 (2)	11.2 ± 1.5 (5)	1.2

[a] TBT concentration in sediment from a single grab or composite sample previously collected at target location, as reported in the Phase 1 RI (Windward 2003a) or in sediment data reports (Windward 2005b, d, e).

[b] Because penis length in males is not known to be affected by TBT exposures, the mean penis length was calculated for the two separate sampling events and used to calculate the RPS index for each location.

dw – dry weight

ID – identification

RPS – relative penis size

TBT – tributyltin

A.3.2.4.2 Critical tissue-residue assessment

Potential effects from TBT exposure on survival, growth, and reproduction of benthic invertebrates were also evaluated using a critical tissue-residue approach. Excluding studies involving the imposex endpoint for gastropods, which was addressed through direct measurement of the imposex endpoint as discussed in Section A.3.2.4.1 above, five studies were identified that reported tissue concentrations of TBT associated with adverse effects (Table A.3-14). The LOAELs for effects on growth and reproduction ranged from 2.36 to 5.44 mg/kg dw. The lowest LOAEL (2.36 mg/kg dw) was selected as the TRV because of the relevance of the sediment exposure as well as the polychaete growth endpoint. The juvenile polychaetes exhibited a reduction in growth of 25% relative to the control sediment at a TBT concentration of 101 ng/g dw in sediment, which resulted in a tissue concentration of 2.36 mg/kg dw (Meador and Rice 2001). The associated NOAEL of 0.97 mg/kg dw was the only NOAEL below the LOAEL.

Table A.3-14. TBT critical tissue-residue toxicity studies for benthic invertebrates

Test Species	NOAEL (mg/kg dw)	LOAEL (mg /kg dw)	Effect	Exposure Conditions	Source
Polychaete (*Armandia brevis*)	0.97	2.36	reduced growth	sediment 42 days	Meador and Rice (2001)
Polychaete (*Neanthes arenaceodentata*)	2.99	6.27	impaired reproduction	aqueous 10 weeks	Moore et al. (1991)
Blue mussel (*Mytilus edulis)*	3.96	5.44	reduced growth	aqueous 4 days	Widdows and Page (1993)
Amphipod (*Hyalella azteca*)	na	32	reduced survival (LC50)	aqueous 10 days	Borgmann et al. (1996)
Polychaete (*Armandia brevis*)	na	41	reduced survival (LC50)	aqueous 10 days	Meador (1997)
Amphipod (*Eohaustorius estuarius*)	na	59	reduced survival (LC50)	aqueous 10 days	Meador (1997)
Amphipod (*Rhepoxynius abronius*)	na	54	reduced survival (LC50)	aqueous 10 days	Meador (1997)

dw – dry weight

LC50 – concentration that causes the death of 50% of a group of test animals

LOAEL – lowest-observed-adverse-effect level

na – not available

NOAEL – no-observed-adverse-effect level

A.3.3 Crab Exposure Assessment

This section presents the assessment of crab exposure to COPCs identified in the problem formulation (Section A.2.5.1.3). Two chemicals, zinc and total PCBs, were identified as COPCs for crabs.

As part of Phase 2 investigations, 10 Dungeness crab and 13 slender crab edible meat samples were analyzed for butyltins, SVOCs (including PAHs), metals, total PCBs as Aroclors, and organochlorine pesticides. In addition, chemical analyses were performed on hepatopancreas samples from six Dungeness crabs and five slender crabs. The edible meat samples were analyzed as composite samples created by homogenizing edible meat from five[45] individual specimens together. Similarly, the hepatopancreas samples were created by homogenizing the hepatopancreas from 15[46] individual specimens together (Windward 2005c). The compositing plan took into consideration sampling area in LDW, specimen size, and gender.

[45] One composite sample included edible meat from six Dungeness crabs to provide enough tissue for analysis.

[46] One composite sample included hepatopancreas tissue from 16 slender crabs to provide enough tissue for analysis.

One other study, conducted by King County in 1997, was included in the exposure dataset. King County collected two composite Dungeness crab edible meat samples and one composite hepatopancreas sample (King County 1999b). The samples were analyzed for total PCBs as Aroclors, metals, and SVOCs. The crab samples were collected between Slip 1 and Harbor Island. Map A.2-2 presents the crab tissue sampling location for these samples.

The exposure data were evaluated as edible meat, hepatopancreas, and whole-body depending on the available effects data. Because effects data for several chemicals, including zinc and total PCBs, were available only for whole-body crabs, the whole-body concentrations for Dungeness and slender crabs were estimated using the following equation:

$$C_{wb} = (C_h \times F_h) + (C_{em} \times F_{em})$$ **Equation 3-3**

Where:

C = concentration
wb = whole-body
h = hepatopancreas
F = fraction
em = average edible meat

The hepatopancreas and edible meat fractions were estimated to be 0.31 and 0.69, respectively, based on the ratio of masses of these tissues in a 16.6 cm[47] Dungeness crab dissected at Windward[48] with 158 g edible meat and 49 g hepatopancreas tissue mass. Similar results were presented in Atar and Secer (2003). Based on width/mass relationship data presented in Atar and Secer (2003), 50 g of edible meat mass and 15 g hepatopancreas mass is predicted for a 9-cm crab, assuming a constant relationship among the weights of edible meat, hepatopancreas, and whole body. A summary of whole-body tissue-residue estimates for zinc and total PCBs is presented in Table A.3-15. Data for Dungeness and slender crabs were combined for the entire LDW to estimate risks to the crab community.

Table A.3-15. Estimated COPC concentrations in whole-body crab tissue

COPC	Number of Samples	Unit	Minimum Concentration	Maximum Concentration	Mean Concentration	UCL
Zinc	21	mg/kg ww	24.6	37.3	30.8	32
Total PCBs	25	µg/kg ww	250	1,900	888	1,100

COPC – chemical of potential concern
PCB – polychlorinated biphenyl
UCL – upper confidence limit on the mean
ww – wet weight

[47] Maximum width of the shell from tip of spine to tip of spine.
[48] A live Dungeness crab was purchased and dissected at Windward to determine the relative weights of edible meat and hepatopancreas.

A.3.4 CRAB EFFECTS ASSESSMENT

This section presents toxicological studies that reported critical tissue residues of zinc and total PCBs associated with potential effects on survival, growth, and reproduction in decapods.

The literature search for effects data was performed as described in the problem formulation (Section A.2.5.1.3). Guidelines for the selection of NOAELs and LOAELs were:

- Studies with decapod tissue-residue data were retained and preference was given to whole-body tissue data.
- Controlled laboratory studies of single chemical exposure with statistical significance were given preference.
- Chronic exposure duration (30-plus days) studies were preferred if available.

Table A.3-16 presents all of the toxicological studies for the two crab COPCs. All toxicological studies evaluated during the TRV search are presented in Attachment 5.

Only one toxicity study that evaluated the effects of zinc exposure on decapods was identified. Crayfish (*Orconectes virilis*) were exposed to five concentrations of zinc (5.2, 12.2, 26.8, 63.3, and 130 mg/L) for 2 weeks (Mirenda 1986b). Following exposure to zinc concentrations of 26.8 mg/L and 12.2 mg/L, crayfish mortality was 23% and 5.7%, respectively. These two concentrations were selected as the LOAEL and NOAEL because the mortality observed in the 12.2 mg/L exposure was not significantly different than mortality in the control. At the end of the exposure duration, the mean whole-body zinc concentration in crayfish was 35.2 mg/kg ww (LOAEL) and 12.7 mg/kg ww (NOAEL).

Four studies were identified that evaluated the effects of PCBs on decapod species (Duke et al. 1970; Hansen et al. 1974b; Nimmo et al. 1974; Sanders and Chandler 1972). Three shrimp species (brown, grass, and pink shrimp) were exposed to Aroclor 1254 and two shrimp species (grass and brown shrimp) were exposed to Aroclor 1016 in water for 2 to 20 days. Effects on survival were assessed in all studies. Reported NOAELs ranged from 1 µg/L to 10 µg/L and reported LOAELs ranged from 1 µg/L to 100 µg/L. Whole-body tissue concentrations associated with NOAEL exposures ranged from 1,300 µg/kg ww to 18,000 µg/kg ww, and ranged from 1,100 µg/kg ww to 42,000 µg/kg ww for LOAEL exposures. The effects of PCBs (Aroclor 1254) on the survival of two other decapod species, crayfish and blue crab, were also evaluated. No effects were observed in either species following water exposure for 20 to 21 days. NOAELs ranged from a whole-body tissue concentration of 1,220 µg/kg ww (crayfish) to 23,000 µg/kg ww (blue crab).

Table A.3-16. Zinc and PCB critical tissue-residue toxicity studies for crabs and other decapods

Chemical	Test Species	Tissue Type	NOAEL	LOAEL	Unit	Exposure Route and Duration	Effect	Source
Zinc	crayfish (*Orconectes virilis*)	whole body	**12.7**[a]	**35.2**[a]	mg/kg ww	12.2 and 26.8 mg/L in water for 2 weeks	reduced survival	Mirenda (1986a)
PCBs (Aroclor 1016)	grass shrimp (*Palaemonetes pugio*)	whole body	**110**[b]	**1,100**[c]	µg/kg ww	0.4 µg/L in water for 96 hours	reduced survival	Hansen et al. (1974b)
PCBs (Aroclor 1016)	brown shrimp (*Penaeus aztecus*)	whole body	3,800	42,000	µg/kg ww	1 and 10 µg/L in water for 96 hours	reduced survival	Hansen et al. (1974b)
PCBs (Aroclor 1254)	crayfish (*Orconectes nais*)	whole body	1,220[a, d]	na	µg/kg ww	1.2 µg/L in water for 96 hours	no effect on survival	Sanders and Chandler (1972)
PCBs (Aroclor 1254)	pink shrimp (*Panaeus duorarum*)	whole body	1,300	3,900	µg/kg ww	10 and 100 µg/L in water for 48 hours	reduced survival	Duke et al. (1970)
PCBs (Aroclor 1254)	pink shrimp (*Panaeus duorarum*)	whole body	na	16,000	µg/kg ww	5.0 µg/L in water for 20 days	reduced survival	Duke et al. (1970)
PCBs (Aroclor 1254)	blue crab (*Callinectes sapidus*)	whole body	23,000	na	µg/kg ww	5.0 µg/L in water for 20 days	no effect on survival	Duke et al. (1970)
PCBs (Aroclor 1254)	grass shrimp *(Palaemonetes pugio)*	whole body	18,000	27,000	µg/kg ww	1.3 and 4.0 µg/L in water for 16 days	reduced survival	Nimmo et al. (1974)

[a] Converted from dry weight to wet weight using a moisture content of 80% (Jarvinen and Ankley 1999); 80% is also the average moisture content of two crab samples collected by King County in 1997 (King County 1999b). Mean % moisture in LDW crabs was 82.7% (Windward 2005c).

[b] The LOAEL TRV was divided by an uncertainty factor of 10 to obtain the NOAEL TRV.

[c] Survival reduced 33%.

[d] In the study, the data from the 96-hour test were extrapolated to a 21-day exposure using a biomagnification factor.

LOAEL – lowest-observed-adverse-effect level

na – not available

NOAEL – no-observed-adverse-effect level

PCB – polychlorinated biphenyl

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

The lowest LOAEL of 1,100 µg/kg ww based on a study of grass shrimp and Aroclor 1016 was selected as the TRV (Hansen et al. 1974b). Because this LOAEL was lower than any of the available NOAELs, a NOAEL of 110 µg/kg ww was calculated from the same study by dividing this LOAEL by 10.

The selected NOAELs and LOAELs for zinc and total PCBs were based on acute exposure studies and no uncertainty factor was applied to derive chronic values. This uncertainty is discussed in the uncertainty analysis (Section A.6.1.2).

A.3.5 SUMMARY OF BENTHIC INVERTEBRATE COMMUNITY ASSESSMENT

A.3.5.1 Exposure assessment

Exposure of the benthic invertebrate community to COPCs was evaluated based on surface sediment, porewater, and benthic invertebrate tissue data, depending on the COPC. Exposure of the benthic invertebrate community to 44 COPCs in sediment was assessed based on the magnitude of detected concentrations at a particular location. Exposure of the single COPC in porewater (cis-1,2-dichloroethene) was evaluated based on the concentrations of this COPC in peeper samples at two sites in the LDW (Tables A.3-2 and A.3-3). To evaluate exposure to TBT, concentrations of TBT in benthic invertebrate tissue were estimated using a regression between benthic invertebrate tissue and co-located sediment to derive an estimated UCL concentration in tissue (Table A.3-4). For meso- and neogastropods, which are particularly sensitive to TBT, a direct measurement of the most sensitive endpoint, imposex, was used (Table A.3-13).

A.3.5.2 Effects assessment

In the effects assessment, the type of biological endpoints used to establish the SMS and DMMP guidelines for the COPCs were discussed. TRVs for most sediment COPCs were based on SMS chemical criteria. However, TRVs for three sediment COPCs were derived from either toxicologically based DMMP guidelines or the scientific literature because no chemical criteria were available. In addition, measures of site-specific toxicity were discussed. The results of site-specific toxicity testing were presented for 48 locations tested by LDWG. Of the 48 samples tested, 18 sediment samples did not exceed the biological effects criteria, 11 sediment samples exceeded the SQS biological effects criteria, and 19 sediment samples exceeded the CSL biological effects criteria. In addition, one of seven sediment samples from a historical surface sediment toxicity study (conducted by King County) was toxic.

The literature was searched for toxicity studies that evaluated effects associated with cis-1,2-dichloroethene in porewater. The lowest toxicity value for 1,2-dichloroethene was derived from a study testing toxicity to *Artemia salina* (Sanchez-Fortun et al. 1997). The LC50 for *Artemia* (6,785 µg/L) from this was selected as the LOAEL TRV for 1,2-dichloroethene. Because no NOAELs were available for this chemical, the LC50-based LOAEL was divided by 50 to derive a NOAEL of 136 µg/L (Sanchez-Fortun et al. 1997).

Possible effects of TBT were evaluated using two approaches: 1) a site-specific study providing a direct assessment of imposex in LDW-collected meso- and neogastropods, and 2) a critical tissue-residue approach for benthic invertebrates. Imposex analysis was performed on three neogastropod species, *Nassarius mendicus, Astyris gausapata*, and *Olivella baetica*, and on a limited number (2 to 5) of four mesogastropod species (*Natica* sp., *Polinices* sp., *L. vincta*, and *Melanella* sp.). Signs of imposex (stage 2 and maximum RPS index of 3.4) were observed only in *N. mendicus*.

To assess effects of TBT using the critical tissue-residue approach, the lowest LOAEL (2.36 mg/kg dw), which was associated with reduced growth in the polychaete, *Armandia brevis,* following exposure to TBT-spiked sediment (Meador and Rice 2001), was selected. The associated NOAEL of 0.97 mg/kg dw from the same study was selected as the NOAEL.

A.3.6 SUMMARY OF CRAB ASSESSMENT

A.3.6.1 Exposure assessment

Two chemicals, zinc and total PCBs, were identified as COPCs for crabs in the problem formulation. Exposure of crabs to these COPCs in sediment was assessed as the concentrations of these chemicals in crab tissue. The concentrations of zinc and total PCBs in whole-body tissue were estimated from LDW crab edible meat and hepatopancreas samples (Section A.3.3; Table A.3-15). UCL concentrations of 32 mg/kg ww and 1,100 µg/kg ww were calculated for zinc and total PCBs, respectively.

A.3.6.2 Effects assessment

The effects assessment discussed the effects on crabs from exposure to zinc and total PCBs based on critical tissue-residue TRVs for crabs and decapods. A summary of selected NOAEL and LOAEL TRVs is presented in Table A.3-17.

Table A.3-17. Selected critical tissue-residue TRVs for crabs

COPC	TEST SPECIES	NOAEL	LOAEL	UNIT	EFFECT	SOURCE
Zinc	crayfish (*Orconectes virilis*)	12.7[a]	35.2[a]	mg/kg ww	reduced survival	Mirenda (1986a)
Total PCBs	grass shrimp (*Palaemonetes pugio*)	110[b]	1,100[c]	µg/kg ww	reduced survival	Hansen et al. (1974b)

[a] Converted from dry weight to wet weight using a moisture content of 80%(Jarvinen and Ankley 1999); 80% is also the average moisture content of two crab samples collected by King County in 1997 (King County 1999b). Mean % moisture in LDW crabs was 82.7% (Windward 2005c).

[b] The LOAEL TRV was divided by an uncertainty factor of 10 to obtain the NOAEL TRV.

[c] Survival was reduced by 33%.

COPC – chemical of potential concern
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
ww – wet weight

A.4.0 Exposure and Effects Assessment: Fish

Three ROCs were selected in the problem formulation to represent fish that use the LDW (Section A.2.3) and may be exposed to sediment-associated chemicals:

- Juvenile chinook salmon
- Pacific staghorn sculpin
- English sole

ROC/COPC pairs identified in the problem formulation are summarized in Table A.4-1.

Table A.4-1. ROC/COPC pairs evaluated for fish

COPC	ROCs		
	Juvenile Chinook Salmon	English Sole	Pacific Staghorn Sculpin
Arsenic[a]	X	X	X
Cadmium[a]	X	X	X
Copper[a]	X	X	X
Vanadium[a]	X	X	X
TBT[b]			X
PCBs[b]		X	X

[a] Evaluated using the dietary approach.

[b] Evaluated using the critical tissue-residue approach.

COPC – chemical of potential concern

ROC – receptor of concern

PCB – polychlorinated biphenyl

TBT – tributyltin

In this ERA, risks to fish were evaluated using two approaches, depending on the COPC. Exposures to TBT and PCBs were evaluated based on concentrations of COPCs in fish tissue, referred to as a critical tissue-residue approach. Exposures to arsenic, cadmium, copper, and vanadium were evaluated based on concentrations in fish prey or stomach contents, referred to as a dietary approach. With this latter approach, the concentration of the relevant COPC in the fish diet was computed and compared to a LOAEL or NOAEL expressed as a concentration in food.

The application of these approaches is presented in this section, which is divided into an exposure assessment (Section A.4.1) and an effects assessment (Section A.4.2). Data presented in these sections are synthesized in the risk characterization (Section A.6.2) to assess risks to fish in the LDW, and uncertainties are also discussed.

A.4.1 EXPOSURE ASSESSMENT

In this section, relevant data were analyzed to determine representative exposure concentrations in the LDW for each ROC/COPC pair identified in Table A.4-1. The LDW dataset used to estimate exposure of fish to COPCs is summarized in Section A.2.4.1.

This section is divided into two subsections to assess exposure based on the approaches discussed above. The first subsection (Section A.4.1.1) presents whole-body concentrations of TBT and PCBs in ROC tissue for the critical tissue-residue approach. The second subsection (Section A.4.1.2) presents concentrations in dietary items and concentrations in juvenile chinook salmon stomach contents for the dietary approach.

A.4.1.1 Critical tissue-residue exposure assessment

The whole-body tissue concentration integrates exposure from all pathways (e.g., direct sediment and water contact and diet) to a fish over its foraging range. The following ROC/COPC pairs were evaluated using a critical tissue -residue approach:

- Pacific staghorn sculpin/TBT
- Pacific staghorn sculpin/total PCBs
- English sole/total PCBs

Foraging ranges for the fish ROCs are not precisely known. The English sole foraging range is assumed to be the entire LDW, whereas the Pacific staghorn sculpin foraging range is more uncertain. Thus, Pacific staghorn sculpin exposure was estimated assuming two different foraging ranges: 1) the entire LDW, and 2) four segments, each representing approximately one-quarter of the LDW. Therefore, for both species, exposures occurring across the entire LDW were estimated as the UCL on the mean concentration of all composite tissue samples from each ROC collected throughout the LDW.

To estimate exposures of Pacific staghorn sculpin in smaller areas, separate UCLs were also calculated for the PCB and TBT sculpin tissue samples from each of the four Phase 2 fish collection areas of the LDW (Map A.2-2). UCL calculation methods and results, along with minimum, mean, and maximum concentrations, are presented in Attachment 11. Resulting exposures for each ROC/COPC pair are presented in Table A.4-2.

Table A.4-2. TBT and total PCB exposure concentrations in whole-body fish tissue

ROC	COPC	LOCATION	UCL CONCENTRATION (µg/kg ww)
English sole[a]	total PCBs	LDW-wide	2,600
Pacific staghorn sculpin	TBT	LDW-wide	36
		T1	37
		T2	36
		T3	28
		T4	53
	total PCBs	LDW-wide	1,100
		T1	800
		T2	920
		T3	2,000
		T4	940

[a] Includes 42 English sole and 3 starry flounder whole-body/composite samples; starry flounder were used as a surrogate for English sole because insufficient numbers of English sole were collected in Area T4 during Phase 2 sampling (see Map A.2-2). Ten of the English sole whole-body sample concentrations were estimated based on the relative weights and total PCB concentrations detected in corresponding skin-on fillet and remainder samples.

COPC – chemical of potential concern
PCB – polychlorinated biphenyl
ROC – receptor of concern
TBT – tributyltin
UCL – upper confidence limit on the mean
ww – wet weight

Composite tissue samples were collected in all four Phase 2 tissue collection areas in order to maximize the number of individuals analyzed to provide the best estimate of the population mean for each species. It should be noted, however, that the use of composite samples may have resulted in a lower estimate of the variance of the population than if individual fish tissue samples had been analyzed. The UCL of the mean of composite samples may therefore underestimate the UCL that would be calculated from an analysis of the same number of individual samples. Because the actual variance at the individual level is unknown, the effect on the estimate of the UCL of analyzing composite tissue samples rather than individual tissue samples is unknown.

A.4.1.2 Dietary exposure

This section presents the approach used to estimate exposure through the diet of the three fish ROCs to arsenic, cadmium, copper, and vanadium. Comparison of chemical concentrations in prey to suitable dietary TRVs is preferable for COPCs that are highly regulated or metabolized by fish. Most aquatic organisms have specific mechanisms for uptake, internal transport, sequestration, and depuration of metals (Meyer et al. 2005). Essential metals are regulated because they are necessary for normal metabolic function, whereas other metals appear to be regulated because they mimic essential

elements and are transported by the same mechanisms (Bury et al. 2003 as cited in Meyer et al. 2005).

The primary exposure route of these COPCs was assumed to be ingestion of food. Dermal contact and incidental sediment ingestion were also considered complete exposure pathways for English sole and Pacific staghorn sculpin (see Figure A.2-2) but with unknown significance. Direct water contact and water ingestion were considered complete and significant pathways for all three fish ROCs. Risks associated with water exposure were evaluated in the King County WQA (King County 1999d); the results of that risk assessment are summarized in the risk characterization (Section A.6.2.1).

The dietary exposure approach requires an approximation of the COPC concentration in an ROC's diet. This section presents the dietary exposure assumptions for each fish ROC, the dietary exposure calculation methods, and the results of the exposure calculations. To approximate the dietary concentrations for each fish ROC, the feeding habits of each ROC were considered. ROC-specific exposure assumptions are described below.

Stomach contents analyses of juvenile chinook salmon from the LDW indicate that they typically ingest benthic invertebrates such as amphipods, worms, and clam siphons,[49] as well as drift organisms and zooplankton (Cordell et al. 1997, 1999, 2001). Because no zooplankton or drift organism tissue concentration data were available, juvenile chinook salmon were assumed to ingest only benthic invertebrates. Because benthic invertebrates live in close contact with sediments, they have greater potential for sediment exposure than do other juvenile chinook salmon prey items; therefore, these exposure assumptions are conservative (i.e., may overestimate exposures but unlikely to underestimate them). Juvenile chinook from the LDW were found to have no appreciable amounts of sediment in their stomachs (Cordell 2001); therefore, they were assumed to have no incidental sediment ingestion. Juvenile chinook salmon generally do not use deep-water habitats (Tabor et al. 2004); exposure was estimated assuming juvenile chinook salmon are exposed primarily in intertidal areas. Therefore, juvenile chinook salmon dietary exposures were based on intertidal benthic invertebrate tissue data. In addition to benthic invertebrate tissue data, exposure was also separately estimated based on chemical analysis of stomach contents of juvenile chinook salmon collected from the LDW.

Stomach contents analyses of English sole collected from Puget Sound show that English sole almost exclusively ingest benthic invertebrates such as marine worms, amphipods, bivalves, and mollusks (Fresh et al. 1979; Wingert et al. 1979). Based on

[49] Windward (2005b) clam data were not included in exposure calculations because benthic invertebrate samples included clams less than 2.0 cm, which are assumed to represent clam tissues consumed by fish ROCs. Mollusks constituted from 0 to 41% (median 17%) of tissue mass in benthic invertebrate samples (Windward 2005b).

these stomach contents analyses, all prey of English sole were assumed to be represented by the benthic invertebrate tissue chemistry data. In addition, incidental sediment ingestion of 1% was assumed based on anecdotal stomach contents observations of English sole and other bottom-feeding fish (Johnson 2006; Lange 2006).[50] Although English sole foraging ranges are uncertain, the available information (Day 1976; Stern et al. 2003) suggests that English sole forage in an area as large as or larger than the LDW. Therefore, English sole dietary exposure was calculated only on a site-wide basis.

Stomach contents analyses of Pacific staghorn sculpin collected from Puget Sound show that they ingest small fish and benthic invertebrates such as crabs, shrimp, marine worms, and amphipods (Wingert et al. 1979; Fresh et al. 1979; Miller et al. 1977c). Based on the relative proportions of biomass reported in stomach contents analyses, the average proportions of fish and benthic invertebrates in dietary exposure estimates were assumed to be 44 and 55%, respectively (Wingert et al. 1979; Fresh et al. 1979; Miller et al. 1977c). Shiner surfperch were selected as representative prey fish because they are numerically dominant in the fish community in the LDW (Windward 2005c, 2006b) and, thus, are likely to be ingested by Pacific staghorn sculpin. They also have a primarily benthic diet so they represent bioaccumulation of sediment-associated chemicals through the food chain (Wingert et al. 1979; Fresh et al. 1979; Miller et al. 1977c). All invertebrate prey were assumed to be represented by the benthic invertebrate tissue chemistry data. Incidental sediment ingestion of 1% was assumed based on Pacific staghorn sculpin's primarily epifaunal diet (Lange 2006). No data are available to determine Pacific staghorn sculpin foraging areas; however, based on opinions expressed at an LDW fish experts meeting with LDWG, EPA, and Ecology that took place on March 31, 2004, Pacific staghorn sculpin may have foraging areas as large as or smaller than the size of the LDW. Therefore, two Pacific staghorn sculpin dietary exposure scenarios were evaluated (an LDW-wide scenario and an area-specific scenario). For the area-specific scenario, exposures were estimated for four modeling areas (M1, M2, M3, and M4), defined as the four fish and crab tissue sampling areas, which extended out to the center points between each adjacent pair of tissue sampling areas (Map A.2-2). Modeling areas, rather than tissue sampling areas, were used in this exposure scenario in order to ensure that all available sediment chemistry and benthic invertebrate tissue concentrations were used in calculating dietary exposure for Pacific staghorn sculpin.

[50] Uncertainty in incidental sediment ingestion is explored in the uncertainty analysis (Section A.6.2.2.2) where exposure is also calculated assuming 10% sediment ingestion.

Using the ROC-specific assumptions described above, COPC concentrations in the diet of each fish ROC were calculated as the weighted average of COPC concentrations in sediment and prey tissue using Equation 4-1.

$$C_{diet} = \sum_{i=1}^{n} X_i C_i \qquad \textbf{Equation 4-1}$$

Where:

C_{diet} = COPC concentration in the diet (mg/kg dw)

X_i = proportion of a particular food item (or sediment) in the diet (unitless)

C_i = COPC concentration in the prey item (mg/kg dw)

n = number of dietary items

The relative proportions of each prey item in each fish ROC's diet are summarized in Table A.4-3.

Table A.4-3. Proportions of dietary items in dietary exposure estimates for each fish ROC

ROC	Prey Item	Proportion in Diet (unitless)	Source
Juvenile chinook salmon	benthic invertebrates (intertidal only)	1	Windward (2004c); Cordell et al. (1997)
English sole	benthic invertebrates (LDW-wide)	0.99	Fresh et al. (1979); Wingert et al. (1979)
	sediment (LDW-wide)	0.01	Johnson (2006); Lange (2006)
Pacific staghorn sculpin	shiner surfperch	0.44	Fresh et. al (1979); Wingert et al. (1979); Miller et al. (1977c)
	benthic invertebrates (LDW-wide)	0.55	
	sediment (LDW-wide)	0.01	(Lange 2006)

LDW – Lower Duwamish Waterway

ROC – receptor of concern

Concentrations in each prey item were estimated as the UCL on the mean concentration, which was calculated as described in Attachment 11. For benthic invertebrate prey, UCLs were calculated in two ways (Attachment 11). First, if the regression between co-located sediment and benthic invertebrate tissue data was significant for a given COPC, then the UCL in tissue was predicted from the regression equation using the arithmetic mean concentration in sediment. LDW-wide arithmetic mean concentrations in sediment were used to estimate UCL prey concentrations for English sole and for the Pacific staghorn sculpin LDW-wide exposure scenario. The mean sediment concentrations for the four modeling areas were used for the Pacific staghorn sculpin area-specific exposure scenarios, and the mean concentration of intertidal sediment samples was used for juvenile chinook salmon. If the regression

between co-located sediment and benthic invertebrate tissue data was not significant, then the UCL was calculated using the benthic invertebrate tissue dataset.[51] All LDW benthic invertebrate tissue data were included in the calculations for English sole and for the Pacific staghorn sculpin LDW-wide exposure scenario. Benthic invertebrate tissue data collected from within the individual modeling areas were used for the Pacific staghorn sculpin area-specific exposure scenarios, and intertidal benthic invertebrate tissue data were used for juvenile chinook salmon exposure calculations.

Resulting exposure concentrations for each ROC/COPC pair are presented in Table A.4-4. Dietary exposure concentrations for juvenile chinook salmon based on the single composite stomach contents sample from 72 fish[52] are presented in Table A.4-5.

[51] The regression was significant for arsenic but not for cadmium, copper, or vanadium (Attachment 11).
[52] Note that an unknown fraction of these stomachs were empty. Ruggerone et al. (2006) found that 4 to 6% of juvenile chinook from the LDW had empty stomachs.

Table A.4-4. Dietary exposure calculations and resulting COPC concentrations in fish ROC diets

Chemical	Exposure Area	Concentration in Dietary Component (UCL) (mg/kg dw)			Fraction of Dietary Component in Diet			Concentration in Diet (mg/kg dw)
		Sediment	Benthic Invertebrates	Shiner Surfperch	Sediment	Benthic Invertebrates	Shiner Surfperch	
Juvenile chinook salmon								
Arsenic	LDW-wide	na	22[a]	na	0	1	na	22
Cadmium		na	0.50[b]	na	0	1	na	0.50
Copper		na	93[b]	na	0	1	na	93
Vanadium		na	8.1[b]	na	0	1	na	8.1
English sole								
Arsenic	LDW-wide	30	24[c]	na	0.01	0.99	na	24
Cadmium		2.4	0.60	na	0.01	0.99	na	0.61
Copper		200	92	na	0.01	0.99	na	93
Vanadium		60	12	na	0.01	0.99	na	12
Pacific staghorn sculpin								
Arsenic	LDW-wide	30	24[c]	4.2	0.01	0.55	0.44	15
	M1	34	23	4.7	0.01	0.55	0.44	15
	M2	46	25	5.0	0.01	0.55	0.44	16
	M3	40	23	3.7	0.01	0.55	0.44	15
	M4	11	18	4.1	0.01	0.55	0.44	12
Cadmium	LDW-wide	2.4	0.60	0.066	0.01	0.55	0.44	0.38
	M1	1.1	0.81	0.079	0.01	0.55	0.44	0.49
	M2	0.55	0.48	0.071	0.01	0.55	0.44	0.30
	M3	6.8	0.77	0.068	0.01	0.55	0.44	0.52
	M4	0.33	0.54	0.056	0.01	0.55	0.44	0.32

Chemical	Exposure Area	Concentration in Dietary Component (UCL) (mg/kg dw)			Fraction of Dietary Component in Diet			Concentration in Diet (mg/kg dw)
		Sediment	Benthic Invertebrates	Shiner Surfperch	Sediment	Benthic Invertebrates	Shiner Surfperch	
Copper	LDW-wide	200	92	7.0	0.01	0.55	0.44	56
	M1	97	110	7.6	0.01	0.55	0.44	65
	M2	170	130	7.5	0.01	0.55	0.44	77
	M3	510	65	8.3	0.01	0.55	0.44	45
	M4	41	81	6.4	0.01	0.55	0.44	48
Vanadium	LDW-wide	60	12	2.0	0.01	0.55	0.44	8.1
	M1	63	12	3.6	0.01	0.55	0.44	8.8
	M2	60	18	3.7	0.01	0.55	0.44	12
	M3	60	20	2.0	0.01	0.55	0.44	12
	M4	60	10	1.2	0.01	0.55	0.44	6.6

Note: Calculated using Equation 4-1.

[a] UCL on the mean tissue concentration predicted from the intertidal arithmetic mean sediment concentration using a sediment:tissue regression (see Attachment 11).

[b] UCL on the arithmetic mean of intertidal tissue concentration data (see Attachment 11).

[c] UCL on the mean tissue concentration predicted from the LDW-wide arithmetic mean sediment concentration using a sediment:tissue regression (see Attachment 11).

COPC – chemical of potential concern

dw – dry weight

na – not applicable

ROC – receptor of concern

UCL – upper confidence limit on the mean

Table A.4-5. Juvenile chinook salmon stomach contents data for dietary COPCs

COPC	EXPOSURE CONCENTRATION (mg/kg dw)
Arsenic	3.9
Cadmium	0.46
Copper	42
Vanadium	na

COPC – chemical of potential concern

dw – dry weight

na – not analyzed

A.4.2 EFFECTS ASSESSMENT

This section presents the toxicological data and describes the selection of TRVs for the COPCs identified for fish ROCs. The literature search and guidelines for TRV selection for fish ROCs are described in detail in Section A.2.5.2. Toxicological studies reporting critical tissue residues of PCBs and TBT associated with potential adverse effects on survival, growth, or reproduction[53] were reviewed. Dietary studies involving these endpoints and arsenic, cadmium, copper, and vanadium were also reviewed. Growth and survival were the only endpoints evaluated for assessing risks to juvenile chinook salmon.[54] TRVs based on reproductive studies were not selected for juvenile chinook salmon because of their life stage at the time of exposure (i.e., migrating juveniles) and because their exposure to LDW sediments as adults is limited. Toxicological data presented in this section are assessed in combination with exposure data (presented in Section A.4.1) in the risk characterization (Section A.6.2.1).

The results of several site-specific and region-specific studies that have assessed potential toxicological effects on juvenile chinook salmon and English sole are presented in the uncertainty analysis (Sections A.6.2.1.2 and A.6.2.1.3, respectively). These studies were not used as a primary source of TRVs because they involved exposure to chemical mixtures; thus, chemical-specific NOAELs and LOAELs cannot be determined from these studies.

A.4.2.1 COPCs evaluated using the critical tissue-residue approach

The following ROC/COPC pairs were evaluated using a critical tissue-residue approach:

- Pacific staghorn sculpin/TBT

[53] The reproductive endpoint is inclusive of early life stage developmental effects (e.g., growth from egg through fry stage, embryo development/viability).

[54] PAH and PCB studies evaluating effects on immunocompetence are also discussed in the uncertainty analysis (Section A.6.2.2).

- Pacific staghorn sculpin/total PCBs
- English sole/total PCBs

The toxicological studies identified for these COPCs are summarized in the following subsections, and the selected NOAEL and LOAEL TRVs are presented. The guidelines for determining the acceptability of toxicity studies for TRV selection are described in detail in Section A.2.5.2. TRVs to evaluate risks to fish from dioxin-like PCB congeners are discussed in the uncertainty analysis (Section A.6.2) because of the uncertainty in the TRV and in fish toxic equivalency factors (TEFs).[55]

A.4.2.1.1 PCBs

Eighteen papers on the potential adverse effects of PCB mixtures on fish were reviewed (Fisher et al. 1994; Freeman and Idler 1975; Hansen et al. 1971; 1974a; 1974b; 1975; Hattula and Karlog 1972; Hendricks et al. 1981; Hugla and Thome 1999; Lieb et al. 1974; Matta et al. 2001; Mauck et al. 1978; Mayer et al. 1977; 1985; McCarthy et al. 2003; Nebeker et al. 1974; Powell et al. 2003; van Wezel et al. 1995). Concentrations in whole-body fish tissue were reported in 14 of these studies (Hansen et al. 1971; 1974a; 1974b; 1975; Hattula and Karlog 1972; Hugla and Thome 1999; Lieb et al. 1974; Matta et al. 2001; Mauck et al. 1978; Mayer et al. 1977; 1985; Nebeker et al. 1974; Powell et al. 2003; van Wezel et al. 1995), concentrations in fish eggs or embryos were reported in four studies (Fisher et al. 1994; Freeman and Idler 1975; Hendricks et al. 1981; McCarthy et al. 2003).

Critical tissue concentrations of PCBs were reported in the toxicological studies reviewed for selection of TRVs in 16 species (i.e., Atlantic croaker, Atlantic salmon, brook trout, channel catfish, coho salmon, common barbel, fathead minnow, goldfish, chinook salmon, pinfish, rainbow trout, mummichog, sheepshead minnow, and spot) . Adverse effects reported in the toxicological studies reviewed included reduced body weight, mortality, reduced early life stage or fry growth and survival, and reduced fecundity, hatchability and spawning success following exposure to PCBs via diet, water, or maternal transfer to eggs. Table A.4-6 presents a summary of the critical tissue-residue NOAELs and LOAELs reported for PCBs in these studies. NOAEL and LOAEL concentrations in eggs and embryos are presented separately from whole-body NOAELs and LOAELs because they are not directly comparable to the adult whole-body tissue concentrations from the LDW used to characterize exposure. Because egg and embryo NOAELs and LOAELs are not directly comparable, these studies were not selected as TRVs but are discussed below and in the uncertainty analysis (Section A.6.2)

[55] EPA refers to the TEF as a toxicity equivalence factor.

Table A.4-6. PCB critical tissue-residue toxicity studies for fish

Chemical	Test Species	Tissue Analyzed	NOAEL (µg/kg ww)	LOAEL (µg/kg ww)	Exposure Route and Duration	Effect	Source	Notes
Studies Reporting Whole-Body NOAELs and LOAELs								
Aroclor 1260	common barbel	whole body	na	**520**	maternal exposure for 50 days	reduced fecundity	Hugla and Thome (1999)	1
Aroclor 1254	juvenile chinook salmon	whole body	980	na	17 mg/kg ww in food for 4 weeks	no effect on growth or survival	Powell et al. (2003)	2
Aroclor 1260	common barbel	whole body	520	**2,640**	maternal exposure for 75 days	lack of spawning in first reproductive season; egg and larval mortality	Hugla and Thome (1999)	1
Aroclor 1254	rainbow trout (14 weeks)	whole body	8,000	na	15 mg/kg dw food for 32 weeks	no effect on growth or survival	Lieb et al. (1974)	
Aroclor 1254	Sheepshead minnow (adult)	whole body	1,900	9,300	maternal exposure to 0.32 µg/L in water for 28 days	decreased fry survival in the first week after hatch	Hansen et al. (1974a)	3
Aroclor 1268	Mummichog (adult)	whole body	15,000	na	15 µg/g in food for 6 weeks	no effect on fertilization, hatching, or larval survival	Matta et al. (2001)	4
Aroclor 1254	Spot	whole body	27,000	46,000	1 and 5 µg/L in water for 20 days	reduced survival	Hansen et al. (1971)	5
Aroclor 1254	brook trout embryos	whole body	31,000	71,000	0.69 and 1.5 µg/L water for 128 days (10 days prior to hatch and 118 days after)	reduced fry growth	Mauck et al. (1978)	6
Aroclor 1016	Sheepshead minnow	whole body	110,000	na	10 µg/L in water for 4 weeks	no effect on fertilization success, survival of embryos, or fry survival	Hansen et al. (1975)	7
Aroclor 1016	pinfish	whole body	na	106,000	21 µg/L in water for 33 days	50% mortality	Hansen et al. (1974b)	
Aroclor 1254: 1260 mixture	juvenile rainbow trout	whole body	120,000	na	2.9 µg/L in water for 90 days	no effect on survival	Mayer et al. (1985)	8
Aroclor 1254: 1260 mixture	juvenile rainbow trout	whole body	70,000	120,000	1.5 and 2.9 µg/L in water for 90 days	reduced growth	Mayer et al. (1985)	

Chemical	Test Species	Tissue Analyzed	NOAEL (µg/kg ww)	LOAEL (µg/kg ww)	Exposure Route and Duration	Effect	Source	Notes
Aroclor 1254	brook trout embryos	whole body	71,000	125,000	1.5 and 3.1 µg/L water for 128 days (10 days prior to hatch and 118 days after)	reduced fry survival	Mauck et al. (1978)	
Aroclor 1016	Sheepshead minnow fry	whole body	57,000	200,000	10 and 32 µg/L in water for 4 weeks	reduced fry survival	Hansen et al. (1975)	
Clophen A50	Goldfish	whole body	na	250,000	4,000 µg/L in water for 5 to 21 days	reduced survival	Hattula and Karlog (1972)	9
Aroclor 1254	fathead minnow	whole body	na	429,000 (female)	1.8 µg/L in water for 8 months	reduced spawning	Nebeker et al. (1974)	10
Aroclor 1242, 1254, or 1260	fathead minnow (6 months)	whole body	na	1,860 – 749,000	0.006 to 0.54 µmol/L in water for100 to 300 hours	range of lethal body burdens (concentration associated with mortality of individuals)	van Wezel et al. (1995)	11
Studies Reporting Only Egg and Embryo NOAELs and LOAELs								
1:1:1:1 Aroclor 1016, 1221, 1254, 1260 mixture	Atlantic salmon	embryo	na	857	embryos exposed to 625 µg/L PCB in water for 48 hours and observed through fry stage	reduced fry body weight	Fisher et al. (1994)	12
Aroclor 1254	rainbow trout	embryos	na	1,640	maternal exposure to 200 mg/kg in food for 60 days	reduced fry growth in offspring	Hendricks et al. (1981)	
Aroclor 1254	Atlantic croaker	egg	na	3,200	maternal transfer	reduced larval growth	McCarthy et al. (2003)	13
Aroclor 1254	brook trout	embryo	na	77,900	200 µg/L in water for 21 days	reduced hatchability (75%)	Freeman and Idler (1975)	

Notes:

1. Whole-body tissue residues were the weighted sum of 10 different tissues (i.e., blood, brain, muscle, skin, liver, gonads, adipose tissues, kidney, digestive tract, and skeleton) (Leroy 2007 [pers. comm.]). Tissue concentrations were converted from dry weight to wet weight assuming 20% solids; all endpoints except first reproductive season spawning were evaluated 1 year after exposure.
2. Whole-body tissue concentrations ranged from 740 to 980 µg/kg ww following the 28-day treatment.
3. Concentrations in maternal adults.
4. Two generations of progeny were observed.
5. Mortality did not appear to be directly related to PCB tissue concentration because tissue concentration increased with exposure duration.

6. At the LOAEL, growth was significantly less than controls at 48 days post hatch, but not at 118 days after hatching. At NOAEL and LOAEL concentrations, study provides tissue concentrations only after 7 days and 118 days of exposure. LOAEL and NOAEL are tissue concentrations in fry at 118 days post hatch. Tissue concentrations at 7 days post-hatch associated with no effects (1,800 µg/kg ww) and low effects (3,200 µg/kg ww) were lower than the concentration at 118 days.
7. For juvenile fish, the LOAEL and NOAEL for reduced survival were 220,000 and 57,000 µg/kg ww, respectively.
8. Survival was not significantly different from control; exposure dose was 1:2 ratio of Aroclor 1254:1260.
9. LOAEL is lethal tissue concentration.
10. LOAEL is average terminal tissue concentration of two female replicate groups. The number of eggs and number of spawnings per female were highly variable (up to 70-fold difference) between duplicate exposures and were not dose responsive at treatment levels below concentrations where no spawning occurred.
11. Tissue concentrations of individual fish that died in less than 20 hours ranged from 1,860 to 30,000 µg/kg ww; tissue concentrations of individual fish that died at 100 to 300 hours ranged from 120,000 to 749,000 µg/kg ww.
12. Growth data from 176 days post-exposure. Effects were not dose responsive; no effects were observed in fish from eggs with 1,534 µg/kg ww PCBs.
13. Only a single dose was evaluated.

LOAEL – lowest-observed-adverse-effect level

na – not available

NOAEL – no-observed-adverse-effect level

PCB – polychlorinated biphenyl

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as TRVs.

Whole-body effect-level concentrations ranged over three orders of magnitude across the fish species included in the toxicological studies reviewed. Whole-body tissue LOAELs ranged from 520µg/kg ww for reduced barbel fecundity (Hugla and Thome 1999) to 429,000 µg/kg ww for reduced spawning in fathead minnows (Nebeker et al. 1974).

In the study reporting the lowest LOAEL, Hugla and Thome (1999) exposed 3- to 5-year-old common barbel from the University of Liege hatchery to 2,500 µg/kg PCBs in food for 50 days or to 12,500 µg/kg PCBs in food for 75 days (nominal concentrations) and analyzed effects on reproduction. Fish were reared at elevated temperatures (Leroy 2007 [pers. comm.]). Treatments were not replicated;16 fish in each treatment were exposed in a single tank (Leroy 2007 [pers. comm.]). Spawning success was monitored in the first reproductive season, and fish were kept in PCB-free water for 1 year and evaluated for additional adverse effects. PCB concentrations in whole fish[56] were reported following 50 or 75 days of exposure. In the first reproductive season, no spawning was reported at the high exposure level. No adverse effects were reported for the lower exposure level associated with the first reproductive season. One year following exposure, significant reductions in fecundity were reported at both exposure levels corresponding to whole-fish concentrations of 520 and 2,640 µg/kg ww for the low and high exposure levels, respectively. Mortality of eggs from the high dietary exposure group was close to 100% and was significantly higher than controls (which had a mean egg mortality of 52.4%), and egg and larval mortality significantly increased as PCB concentrations increased in eggs. At the lower dose, egg mortality was not significantly different from controls.

The fecundity LOAEL associated with the lower dose is uncertain because fecundity as measured after the first two spawning seasons was not dose responsive. Fecundity comparisons are complicated by the fact that the higher-dosed fish did not spawn during the first season and whole-body tissue concentrations were not measured 1 year later when the high-dose fish finally spawned. After the second spawning, average fecundity was similar between the high and low doses, but variance in fecundity was greater at the higher dose. In addition, the number of fish exposed at each treatment level and evaluated for effects is unclear. Because of these and additional uncertainties discussed in the uncertainty analysis (Section A.6.2.2.2), the range of effects concentrations reported in this paper for the fecundity and the spawning and egg hatchability endpoints was considered to represent the range of exposures over which the lowest adverse effects may occur in fish. Thus, the selected LOAEL for PCBs was set at a range of 520 to 2,640 µg/kg ww. Additional effects data are discussed below for comparison.

---

[56] The reported whole-body fish tissue PCB concentrations were a weighted average of dry weight PCB concentrations in various tissues, specifically, blood, brain, muscle, skin, liver, gonads, adipose tissue, kidney, digestive tract, and skeleton (Leroy 2007 [pers. comm.]).

In the study reporting the next higher LOAEL, Hansen et al. (1974a) exposed 20 female and 10 male adult sheepshead minnows for four weeks to four concentrations of PCBs ranging from 0.1 to 3.2 µg/L. Eggs from five female fish from each exposure level were fertilized using a male from the same exposure level, and 25 successfully fertilized eggs from each exposure group were raised to the fry stage in PCB-free water and evaluated for survival. Reduced survival was reported for eggs from fish with maternal adult tissue concentrations of 9,300 µg/kg ww and greater; but to enhance egg production, fish were injected with human chorionic gonadotrophic hormone, which may have affected reproduction in the fish. No effects were observed at the next lower exposure level corresponding with a tissue concentration of 1,900 µg/kg ww. Uncertainties in this study are discussed in the uncertainty analysis (Section A.6.2.2.2).

Among the studies reviewed for this ERA, effects concentrations reported in eggs and embryos ranged from 857 to 77,900 µg/kg ww. The lowest value was for reduced growth of Atlantic salmon fry held in PCB-free water for 176 days following exposure of eggs to aqueous PCB concentrations of 625 to 62,500 µg/L for 48 hours (Fisher et al. 1994). The highest value was for brook trout embryos exposed to 200 µg/L of PCBs in water for 21 days (Freeman and Idler 1975). NOAELs were not identified.

Although these egg and embryo effects concentrations were generally lower than effects concentrations reported in the literature for more mature fish, egg/embryo and adult tissue-residue data are not directly comparable. Uncertainties associated with comparison of exposure concentrations to egg and embryo studies are discussed in the uncertainty analysis (Section A.6.2.2).

Whole-body NOAELs ranged from 980 µg/kg ww, at which no effect on growth or survival was reported in juvenile chinook salmon (Powell et al. 2003), to 120,000 µg/kg ww for no effect on survival of juvenile rainbow trout (Mayer et al. 1985). Because there were no NOAELs identified that were lower than the identified LOAEL range from Hugla and Thome (1999), a NOAEL range of 104 to 528 µg/kg ww was estimated by applying an uncertainty factor of 5 to the range of chronic reproductive effects concentrations reported in Hugla and Thome (1999).

A.4.2.1.2 Tributyltin

Three toxicological studies that evaluated the toxicity of TBT to fish were reviewed (Shimasaki et al. 2003; Nirmala et al. 1999; Nakayama et al. 2005). Table A.4-7 summarizes the results of these studies. Critical tissue residues of TBT were reported in Japanese flounder larvae and Japanese medaka associated with adverse effects on growth and reproductive success, respectively, following exposure to TBT in water.

Table A.4-7. TBT critical whole-body tissue-residue toxicity studies for fish

Chemical	Test Species	NOAEL (µg TBT/kg ww)	LOAEL (µg TBT/kg ww)	Exposure Route and Duration	Effect	Source	Notes
Tributyltin oxide	Japanese flounder larvae	**18**	**159**	0.1 and 1 mg/kg dw in food for approximately 65 days	reduced body weight	Shimasaki et al. (2003)	1
Tributyltin oxide	Japanese medaka	na	1,054	maternal exposure to 5 mg/kg dw in food for 3 weeks	reduced swim-up success and hatchability	Nakayama et al. (2005)	2
Tributyltin oxide	Japanese medaka	na	2,390	maternal exposure to 1 mg/kg dw in food for 3 weeks	reduced hatching, swim-up, and embryonic success	Nirmala et al. (1999)	3

Notes:

1. No replication in study. Survival was not significantly affected at any dose; however, some reduced survival was observed in all groups, including the control group.
2. LOAEL is the TBT concentration in adult female fish estimated using an adult:egg conversion factor of 8.57, based on Nirmala et al. (1999).
3. LOAEL is the TBT concentration in adult female fish.

LOAEL – lowest-observed-adverse-effect level
na – not available
NOAEL – no-observed-adverse-effect level
TBT – tributyltin
ww – wet weight
Bold identifies the NOAEL and LOAEL selected as TRVs.

Whole-body tissue-residue[57] LOAELs ranged from 159 µg TBT/kg ww for reduced body weight in Japanese flounder larvae following 65 days of dietary exposure (Shimasaki et al. 2003) to 2,390 µg TBT/kg ww for reduced hatchability and early-life stage mortality of Japanese medaka offspring spawned from exposure of both parents to dietary TBT for three weeks during reproduction (Nirmala et al. 1999). All three toxicological studies were conducted over a chronic exposure duration or critical life stage (reproduction). The lowest LOAEL reported (159 µg TBT/kg ww) was selected as the LOAEL TRV. There is significant uncertainty associated with this LOAEL because there was no replication of test groups and high mortality was observed consistently across the control and low- and high-dose groups. However, because few toxicological data were available (only three studies were identified) and a clear effect on growth was observed in fish exposed to TBT in the study by Shimasaki et al. (2003), the LOAEL TRV was derived from this study to represent a conservatively based TRV. The other two studies by Nirmala et al. (1999) and Nakayama et al. (2005) present adverse effects to Japanese medaka reproduction following exposure to higher TBT concentrations.

Only one NOAEL was reported in the three studies reviewed (Shimasaki et al. 2003). At 18 µg/kg ww, no effect on Japanese flounder growth was observed following 65 days of dietary exposure to tributyltin oxide. This NOAEL was selected as the NOAEL TRV.

[57] Whole-body tissue concentrations were reported in the reviewed study or were modeled from an egg concentration using an egg:adult conversion factor reported in Nirmala et al. (1999).

A.4.2.2 COPCs evaluated using a dietary approach

The following COPCs were evaluated using a dietary approach: arsenic, cadmium, copper, and vanadium. The toxicological studies identified for these COPCs are summarized in the following subsections, and the selected NOAEL and LOAEL TRVs are presented.

A.4.2.2.1 Arsenic

Six toxicity studies that evaluated the effects of dietary arsenic on fish were identified (Blazer et al. 1997; Cockell and Hilton 1988; Cockell and Bettger 1993; Cockell et al. 1991, 1992; Oladimeji et al. 1984). Table A.4-8 summarizes all dietary fish NOAELs and LOAELs for arsenic reported in the reviewed literature.

Table A.4-8. Arsenic dietary toxicity studies for fish

Chemical	Test Species	NOAEL (mg/kg dw)[a]	LOAEL (mg/kg dw)[a]	Exposure Duration	Effect	Source	Notes
Sodium arsenite	juvenile rainbow trout	**20**	**30**	6 weeks	reduced body weight	Oladimeji et al. (1984)	1, 2
Disodium arsenate heptahydrate	juvenile rainbow trout	8	44	16 weeks	reduced body weight	Cockell et al. (1991)	3
Disodium arsenate heptahydrate	juvenile rainbow trout	na	49	24 weeks	reduced body weight	Cockell et al. (1991)	3
Disodium arsenate heptahydrate	juvenile rainbow trout	na	55	8 days	reduced body weight	Cockell et al. (1992)	3
Disodium arsenate	juvenile rainbow trout	na	58	12 days	reduced body weight	Cockell and Bettger (1993)	3
Disodium arsenate heptahydrate	juvenile rainbow trout	32	60	12 days	reduced body weight	Cockell et al. (1992)	3
Disodium arsenate heptahydrate	juvenile rainbow trout	33	65	24 weeks	reduced body weight	Cockell et al. (1991)	4
Disodium arsenate	juvenile rainbow trout	na	137	8 days	reduced body weight	Cockell and Hilton (1988)	3
Arsenic trioxide	juvenile rainbow trout	na	180	8 days	reduced body weight	Cockell and Hilton (1988)	
Disodium arsenate heptahydrate	juvenile striped bass	52.3	188.8	6 days	reduced body weight	Blazer et al. (1997)	3

[a] Concentrations are for elemental arsenic.

Notes:

1. Only the nominal concentration was reported.
2. Concentrations in figure and text in study do not agree: 20 mg/kg dw is mentioned both as an effect level and a no-effect level in the text; however, it is shown in the figure to be not significant. The NOAEL was assumed to be 20 mg/kg dw.
3. Feed refusal accompanied growth effects.
4. Body weight gain reduced at 12 weeks in fish fed 33 mg/kg arsenic in diet but not at 24 weeks (body weight was recovered).

dw – dry weight

LOAEL – lowest-observed-adverse-effect level

na – not available

NOAEL – no-observed-adverse-effect level

Bold identifies the NOAEL and LOAEL selected as TRVs.

No dietary toxicity data were available for survival or reproductive endpoints. All studies reviewed reported reductions in growth among juvenile rainbow trout and striped bass following dietary exposure to arsenic. Dietary LOAELs based on growth ranged from 30 mg/kg dw for juvenile rainbow trout (Oladimeji et al. 1984) to 188.8 mg/kg dw for juvenile striped bass (Blazer et al. 1997). Oladimeji et al. (1984) reported that juvenile rainbow trout exposed for 2, 4, and 6 weeks (a chronic exposure period) to 30 mg/kg of dietary arsenic (as sodium arsenite) had significantly less weight gain than control fish.

Based on the available data, the LOAEL reported in Oladimeji et al. (1984) (30 mg/kg diet) was selected as the LOAEL TRV, representing the lowest reported effect level associated with chronic exposure. Dietary NOAELs ranged from 8 to 52.3 mg/kg dw for growth of juvenile rainbow trout and juvenile striped bass, respectively (Blazer et al. 1997; Cockell et al. 1991). The highest NOAEL (20 mg/kg) that was below the selected LOAEL was selected as the NOAEL TRV. No significant effect on rainbow trout growth was observed in fish fed this dietary concentration.

The results of Oladimeji et al. (1984) indicated slightly greater toxicity of disodium arsenate to juvenile rainbow trout than did studies by Cockell et al. (1991, 1992). Cockell et al. (1991) presented the results of three studies conducted to differentiate effects on growth attributable to arsenic toxicity from those attributable to reduced palatability of arsenic-contaminated food. LOAELs based on Cockell et al. (1991, 1992) ranged from 44 to 65 gm/kg dw. NOAELs based on the same studies range from 8 to 33 mg/kg dw.

A.4.2.2.2 Cadmium

Nine studies that evaluated the effects of dietary cadmium on fish were identified (Baldisserotto et al. 2005; Franklin et al. 2005; Handy 1993; Hatakeyama and Yasuno 1982, 1987; Kang et al. 2005; Kim et al. 2004; Lundebye et al. 1999; Mount et al. 1994; Szebedinsky et al. 2001).[58] Table A.4-9 summarizes the dietary NOAELs and LOAELs for cadmium reported in the reviewed literature.

Table A.4-9. Cadmium dietary toxicity studies for fish

Chemical	Test Species	NOAEL (mg/kg dw)[a]	LOAEL (mg/kg dw)[a]	Exposure Duration	Effect	Source	Notes
Cadmium nitrate	juvenile rockfish	na	**0.5**	60 days	reduced growth rate and condition factor	Kim et al. (2004); Kang et al. (2005)	1, 7
Cadmium chloride	rainbow trout fry	55	na	60 days	no effect on body weight, length, or survival	Mount et al. (1994)	2, 3
Cadmium nitrate	juvenile rockfish	125	na	60 days	no effect on survival	Kim et al. (2004) Kang et al. (2005)	1, 7
Cadmium	guppy	171	na	10 – 30 days	no effect on growth	Hatakeyama and	4, 5

[58] Note that Kim et al. (2004) and Kang et al. (2005) are the same study reported in two separate publications.

Chemical	Test Species	NOAEL (mg/kg dw)[a]	LOAEL (mg/kg dw)[a]	Exposure Duration	Effect	Source	Notes
chloride						Yasuno (1982)	
Cadmium chloride	adult guppy	210	na	2 months	no effect on fry survival or premature embryos	Hatakeyama and Yasuno (1987)	6
Cadmium	Atlantic salmon	250	na	4 weeks	no effect on growth rate (body weight)	Lundebye et al. (1999)	1, 7
Cadmium chloride	guppy (2 months old)	274	na	30 days	no effect on body weight	Hatakeyama and Yasuno (1987)	6
Cadmium chloride	juvenile rainbow trout	294	na	15 – 30 days	no effect on growth rate or survival	Baldisserotto et al. (2005)	7
Cadmium chloride	juvenile rainbow trout	471	na	28 days	no effect on growth rate or survival	Franklin et al. (2005)	
Cadmium chloride	guppy (1 month old)	500	800	7 months	reduced number of fry produced	Hatakeyama and Yasuno (1987)	6, 8
Cadmium chloride	guppy (1 month old)	na	1,250	7 months	reduced female growth and survival	Hatakeyama and Yasuno (1987)	6, 9
Cadmium nitrate	juvenile rainbow trout	786	1,395	30 days	57% survival	Szebedinsky et al. (2001)	7, 10
Cadmium nitrate	juvenile rainbow trout	1,395	2,265	30 days	reduced specific growth rate (weight)	Szebedinsky et al. (2001)	7
Cadmium sulfate	rainbow trout (130 g)	na	10,000	28 days	39% mortality	Handy (1993)	1, 7, 11

[a] Concentrations are for elemental cadmium.

Notes:

1. Only the nominal concentration was reported.
2. Fish were exposed to copper, cadmium, lead, and zinc in water at 23.0, 0.97, 3.32, and 46.3 µg/L, respectively, at the same time as the dietary exposure to cadmium chloride. No effect on growth or survival was observed following exposure; therefore, a dietary NOAEL TRV was obtained from this study.
3. Fish fed live *Artemia* exposed to cadmium chloride in water. Dietary dose corrected for a theoretical 20% loss related to cadmium depuration from the *Artemia* food source.
4. Fish fed live *Moina macrocopa* exposed to cadmium chloride in water.
5. Significant reduction in growth effect was noted on day 10 and recovered at day 20.
6. Fish fed live *Chironomus yoshimatsui* exposed to cadmium chloride in water.
7. Dietary dose was not reported as ww or dw and was assumed to be a dw concentration.
8. Cumulative number of fry produced decreased to about 60% of the control following exposure to 80 and 160 µg/L cadmium chloride. Study did not present statistical significance of data and no replication of treatment was conducted. The LOAEL and NOAEL were estimated using a figure presented in the study.
9. Female body weight decreased to 68% of control on the 48th day of exposure and six of the seven females died in the group exposed to 1,250 mg/kg dw dietary cadmium. No effect on male growth in fish exposed to the same dose. Study did not present statistical significance of data and no replication of treatment was conducted. The LOAEL was estimated using a figure presented in the study.
10. In a separate experiment reported in this study, 92% survival was reported for juvenile rainbow trout exposed to dietary cadmium concentrations of 1,419 mg/kg dw over a 39-day exposure period. Survival data were not statistically analyzed in either experiment. LOAEL – lowest-observed-adverse-effect level.
11. Fish expelled food so the ingested dose is unknown.

na – not available

NOAEL – no-observed-adverse-effect level

Bold identifies the NOAEL and LOAEL selected as TRVs.

LOAELs ranged from 0.5 mg/kg dw for growth of juvenile rockfish fed dietary cadmium for 60 days (Kang et al. 2005; Kim et al. 2004) to 10,000 mg/kg dw for mortality of adult rainbow trout following dietary exposure for 28 days (Handy 1993).

The lowest LOAEL was derived from Kang et al. (2005) and Kim et al. (2004). In this study (reported in two separate publications), juvenile rockfish were treated with 0.5, 5, 25, or 125 mg/kg dw of cadmium as cadmium nitrate for 60 days. Significant effects on growth (identified as condition factor, body weight growth rate, and body length growth rate) were reported for fish exposed to all four dietary concentrations. The lowest LOAEL, 0.5 mg/kg dw, was selected as the LOAEL TRV because it was the most conservative LOAEL reported in the reviewed studies. Reported effects on growth are somewhat uncertain because in one of the two papers in which the results of this study were reported (Kim et al. 2004), the observed growth effect was partially attributed to reduced food intake, which may be the result of food avoidance rather than toxicological effects.

NOAELs ranged from 55 mg/kg dw for rainbow trout fry, as reported in Mount et al. (1994), to 1,395 mg/kg dw for growth of rainbow trout (Szebedinsky et al. 2001). No NOAEL lower than the selected LOAEL was reported, so a NOAEL was estimated by applying an uncertainty factor of 5 to the chronic LOAEL. The resulting NOAEL of 0.1 mg/kg dw was selected.

There was high variability in the toxicological data reviewed for dietary exposure of fish to cadmium. The lowest LOAEL of 0.5 mg/kg dw was two to three orders of magnitude lower than the NOAELs reported in the eight other studies (which ranged from 55 to 1,395 mg/kg dw) and was three to four orders of magnitude lower than the LOAELs reported in the three other studies that reported LOAELs (which ranged from 800 to 10,000 mg/kg dw). The lack of agreement in the toxicological studies results in increased uncertainty associated with the selected TRV.

Of the nine studies available for cadmium, six studies were conducted with salmonid species (Baldisserotto et al. 2005; Franklin et al. 2005; Handy 1993; Lundebye et al. 1999; Mount et al. 1994; Szebedinsky et al. 2001). These studies provide species-specific information for juvenile chinook salmon, which serves as an ROC representing all out-migrating juvenile salmonids. LOAELs were reported in two of these studies, ranging from 1,395 mg/kg dw for mortality of rainbow trout (Szebedinsky et al. 2001) to 10,000 mg/kg dw, also for mortality of rainbow trout (Handy 1993). Salmonid-specific NOAELs were reported in five studies (Baldisserotto et al. 2005; Franklin et al. 2005; Lundebye et al. 1999; Mount et al. 1994; Szebedinsky et al. 2001), ranging from 55 mg/kg dw for growth of rainbow trout (Mount et al. 1994) to 1,395 mg/kg dw, also for growth of rainbow trout (Szebedinsky et al. 2001). These NOAEL TRVs, in particular, suggest that the TRVs selected for fish ROCs in general may result in overestimates of risk for juvenile chinook salmon.[59]

[59] TRVs for salmonids are discussed further in the uncertainty analysis (Section A.6.2.1.2).

A.4.2.2.3 Copper

Fifteen toxicity studies that exposed fish to dietary copper were evaluated for TRV selection (Baker et al. 1998; Berntssen et al. 1999a; 1999b; Gatlin and Wilson 1986; Handy 1992, 1993; Kamunde et al. 2001; Kang et al. 2005; Lanno et al. 1985a; 1985b; Lorentzen et al. 1998; Lundebye et al. 1999; Miller et al. 1993; Mount et al. 1994; Murai et al. 1981). Table A.4-10 summarizes the dietary NOAELs and LOAELs for copper reported in the reviewed literature.

Table A.4-10. Copper dietary toxicity studies for fish

Chemical	Test Species	NOAEL (mg/kg dw)[a]	LOAEL (mg/kg dw)[a]	Exposure Duration	Effect	Source	Notes
Copper sulfate	channel catfish fingerling	8	16	16 weeks	reduced growth	Murai et al. (1981)	1, 2
Copper sulfate pentahydrate	channel catfish fingerling	40	na	13 weeks	no effect on growth	Gatlin and Wilson (1986)	
Copper sulfate	juvenile rockfish	**50**	**100**	60 days	reduced growth rate	Kang et al. (2005)	2
Copper sulfate pentahydrate	Atlantic salmon parr	98	na	12 weeks	no effect on survival or growth	Lorentzen et al. (1998)	
Copper sulfate	rainbow trout (138 g)	200	na	32 days	no effect on survival	Handy (1992)	2
Copper sulfate	juvenile rainbow trout	684	na	42 days	no effect on growth	Miller et al. (1993)	
Copper sulfate pentahydrate	Atlantic salmon parr	691.3	na	4 weeks	no effect on body length, weight, or condition factor	Berntssen et al. (1999b)	
Copper sulfate pentahydrate	Atlantic salmon fry	500	700	3 months	reduced growth	Lundebye et al. (1999)	2
Copper sulfate pentahydrate	juvenile rainbow trout	287	730	8 weeks	reduced growth	Lanno et al. (1985b)	3
Copper sulfate pentahydrate	juvenile rainbow trout	730	na	8 weeks	no effect on survival	Lanno et al. (1985b)	4
Copper sulfate pentahydrate	juvenile rainbow trout	na	796	16 weeks	reduced growth	Lanno et al. (1985a)	
Copper chloride	rainbow trout fry	352	na	60 days	reduced survival	Mount et al. (1994)	5
Copper sulfate	Atlantic salmon fry	638	868	3 months	reduced growth	Berntssen et al. (1999a)	
Copper chloride	rainbow trout fry	800	na	60 days	no effect on body weight or length	Mount et al. (1994)	5
Copper sulfate pentahydrate	juvenile rainbow trout	1,042	na	28 days	no effect on survival or growth	Kamunde et al. (2001)	
Copper sulfate pentahydrate	juvenile grey mullet	na	2,397	67 days	reduced growth	Baker et al. (1998)	6
Copper sulfate	rainbow trout (130 g)	10,000	na	28 days	no effect on survival	Handy (1993)	2

[a] Concentrations are for elemental copper.

Notes:

1. Significant effects on growth were reported following a 4-week dietary exposure to 16 and 32 mg/kg dw. At the NOAEL (8 mg/kg dw), no effect on weight gain was reported.

2. Only the nominal concentration was reported.
3. Significant reduction in growth was reported in fish fed 664 mg/kg dw following 16 weeks of exposure; however, growth reduction was recovered at 24 weeks.
4. Mortality (5.2%) in fish fed 730 mg/kg dw was similar to mortality (3%) reported in fish fed control diet.
5. Fish were fed live *Artemia* previously exposed to copper in water. Dietary concentrations corrected for a theoretical 20% loss resulting from depuration of copper from *Artemia.* Fish were exposed to copper, cadmium, lead, and zinc in water at 23.0, 0.97, 3.32, and 46.3 µg/L, respectively, at the same time as the dietary exposure to copper chloride. No effect on growth or survival was observed following exposure; therefore, a dietary NOAEL TRV was obtained from this study. Increased mortality observed at a dietary concentration of approximately 700 mg/kg dw was attributed to elevated copper concentrations in water; therefore, no dietary LOAEL was identified from this study.
6. Growth effects were associated with a reduction in feeding.

dw – dry weight

LOAEL – lowest-observed-adverse-effect level

na – not available

NOAEL – no-observed-adverse-effect level

Bold identifies the NOAEL and LOAEL selected as TRVs.

Adverse effects on growth or survival were reported in channel catfish, Atlantic salmon, rainbow trout, and grey mullet following exposure to dietary copper; no dietary studies evaluating reproductive effects were found. LOAELs ranged from 16 mg/kg dw for growth of channel catfish (Murai et al. 1981) to 2,397 mg/kg dw for growth of grey mullet (Baker et al. 1998). NOAELs ranged from 8 mg/kg dw for growth of channel catfish (Murai et al. 1981) to 10,000 mg/kg dw for mortality of rainbow trout (Handy 1993).

The lowest LOAEL reported in the reviewed literature was based on Murai et al. (1981). In this study, a significant decrease in body weight was reported for channel catfish fingerlings exposed to 16 mg/kg dw of copper as copper sulfate in a prepared diet for 16 weeks compared to the control group, but a significant reduction in body weight was not observed in fish fed 8 mg/kg dw relative to controls. However, the sensitivity of channel catfish fingerlings documented by Murai et al. (1981) has not been confirmed in subsequent studies using similar exposures and fish of similar age (Erickson et al. 2003; Gatlin and Wilson 1986). Gatlin and Wilson (1986) attempted to reproduce the exposure conditions used by Murai et al. (1981). However, they used larger fingerling catfish (5.5 g/fish versus 1 g/fish in Murai et al.). Gatlin and Wilson (1986) did not report any difference in weight gain in their highest dietary exposure of 40 mg/kg dw. Likewise, Erickson et al. (2003) did not report differences in weight gain following exposure for 30 days to copper-contaminated prey at dietary concentrations of 157 and 246 mg/kg dw using much smaller (0.2 g/fish) fingerling channel catfish. However, that study has not been published in manuscript form and thus was not included in the studies evaluated for TRV derivation. Although the results of Erickson et al. (2003) are not published, they do help bracket the size of fingerlings tested and confirm that the Murai et al. (1981) study results are anomalous. The sensitivity of catfish to copper reported by Murai et al. (1981) has also been characterized as atypical by other studies of copper in fish (Lorentzen et al. 1998).

The next lowest LOAEL was presented in Kang et al. (2005). In that study, juvenile rockfish were exposed to 50, 100, 250, or 500 mg/kg dw of copper as copper sulfate for

60 days. Significant effects on growth (identified as body weight growth rate) were reported for fish exposed to dietary concentrations of 100 mg/kg dw or greater. No adverse effects were observed in fish exposed to 50 mg/kg dw. The NOAEL and LOAEL of 50 and 100 mg/kg dw, respectively, were thus selected as the NOAEL and LOAEL TRVs.

A.4.2.2.4 Vanadium

One study that evaluated the toxicological effects of dietary vanadium to fish (Hilton and Bettger 1988) was identified (Table A.4-11).

Table A.4-11. Vanadium dietary toxicity studies for fish

Chemical	Test Species	NOAEL (mg/kg dw)[a]	LOAEL (mg/kg dw)[a]	Exposure Duration	Effect	Source	Notes
Sodium orthovanadate	juvenile rainbow trout	na	**10.2**	12 weeks	reduced body weight	Hilton and Bettger (1988)	1

[a] Concentrations are for elemental vanadium.

Note:

1. Reduced food intake by fish fed 10.2 mg/kg dw compared to fish fed control diet.

dw – dry weight

LOAEL – lowest-observed-adverse-effect level

na – not available

NOAEL – no-observed-adverse-effect level

Bold identifies the LOAEL selected as the TRV.

Body weights of juvenile rainbow trout fed vanadium were significantly lower than those of the control group following 12 weeks of exposure to 10.2, 80, and 493 mg/kg dw vanadium in a prepared diet (Hilton and Bettger 1988). The LOAEL, 10.2 mg/kg dw, was selected as the LOAEL TRV. Food intake was reduced in fish fed 10.2 mg/kg dw; however, the ratio of food ingestion:body weight was significantly greater for fish in this group than for fish in the control group. Therefore, the reduction in body weight observed at this LOAEL was not attributed to reduced food intake. No NOAEL was reported by Hilton and Bettger (1988); therefore, a NOAEL was extrapolated from the chronic LOAEL using an uncertainty factor of 5. The resulting NOAEL of 2 mg/kg dw was selected as the NOAEL TRV. Because of the paucity of toxicological data on dietary effects of vanadium to fish (only one study was reviewed), there is uncertainty associated with the selected TRVs. Orthovanadate is in a +5 oxidation state, which is more toxic than the +3 redox state. The +5 form is commonly used in toxicity testing; the prevalence of this form in the LDW is unknown.

A.4.3 SUMMARY OF FISH ASSESSMENT

A.4.3.1 Exposure assessment

In Section A.4.1, exposure of fish to COPCs was assessed using one of two approaches depending on the COPC:

- A critical tissue-residue approach was used for COPCs that persist in fish tissue. ROC/COPC pairs evaluated included TBT and total PCBs in Pacific staghorn sculpin tissue and total PCBs in English sole tissue.
- Concentrations in the diet were used for chemicals that are highly regulated or metabolized by fish. COPCs evaluated using this approach include arsenic, cadmium, copper, and vanadium for all fish ROCs.

Whole-body concentrations of TBT and total PCBs were summarized in Table A.4-2 for English sole and Pacific staghorn sculpin.

Concentrations in the diet were estimated based on assumptions regarding ROC-specific prey items and incidental sediment ingestion. Resulting concentrations in the diet were summarized in Tables A.4-4 and A.4-5.

A.4.3.2 Effects assessment

The effects assessment was divided into two sections. Section A.4.2.1 presented the critical tissue-residue approach, and Section A.4.2.2 presented the dietary approach. Each section presented an overview of the available literature from laboratory studies with controlled exposures to COPCs. Ranges of NOAELs and LOAELs from these studies were reviewed, and TRVs were selected. A summary of selected TRVs for COPCs evaluated using the critical tissue-residue approach is presented in Table A.4-12. A summary of selected dietary TRVs is presented in Table A.4-13.

Table A.4-12. TRVs selected for COPCs evaluated using the critical tissue-residue approach

COPC	TEST SPECIES	NOAEL (µg/kg ww)	LOAEL (µg/kg ww)	EFFECT	SOURCE
Total PCBs	common barbel	104 – 528	520 – 2,640	fecundity, egg hatchability, spawning	Hugla and Thome (1999)
TBT	Japanese flounder	18	159	reduced body weight	Shimasaki et al. (2003)

COPC – chemical of potential concern
LOAEL – lowest-observed-adverse-effect level
na – not available
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
TBT – tributyltin
ww – wet weight

Table A.4-13. TRVs selected for COPCs calculated using the dietary approach

COPC	Test Species	NOAEL (mg/kg dw)	LOAEL (mg/kg dw)	Effect	Source
Arsenic	rainbow trout	20	30	reduced body weight	Oladimeji et al. (1984)
Cadmium	rockfish	0.1[a]	0.5	reduced growth rate and condition factor	Kim et al. (2004); Kang et al. (2005)
Copper	juvenile rockfish	50	100	reduced growth rate	Kang et al. (2005)
Vanadium	rainbow trout	2	10.2	reduced body weight	Hilton and Bettger (1988)

[a] NOAEL estimated using an uncertainty factor of 5 (chronic LOAEL to NOAEL).

COPC – chemical of potential concern
dw – dry weight
ROC – receptor of concern
LOAEL – lowest-observed-adverse-effect level
na – not available
NOAEL – no-observed-adverse-effect level

A.5.0 Exposure and Effects Assessment: Wildlife

This section presents the exposure and effects assessment for birds and mammals. In the problem formulation, three bird and two mammal species were selected as ROCs in the LDW: spotted sandpiper, great blue heron, osprey, river otter, and harbor seal. COPCs for these ROCs were selected based on the COPC screen described in Section A.2.5.3, as summarized in Table A.5-1.

Table A.5-1. ROC/COPC pairs evaluated for wildlife receptors

COPC[a]	Receptors of Concern			
	Spotted Sandpiper	Great Blue Heron and Osprey	River Otter	Harbor Seal
Arsenic	X		X	
Cadmium	X			
Chromium	X	X		
Cobalt	X		X	
Copper	X			
Lead	X	X		
Mercury	X	X	X	X
Nickel	X			
Selenium	X		X	
Zinc	X			
Vanadium	X			
PCBs[b]	X	X	X	X

[a] Total DDTs also screened in as a COPC for spotted sandpiper, great blue heron, and osprey, but will be evaluated in the uncertainty analysis because of uncertainties associated with the exposure data for organochlorine pesticides (see Section A.2.4.2.2).

[b] PCBs were evaluated in two ways: as total PCBs and as a toxic equivalent (TEQ) for dioxin-like PCB congeners, as discussed in Section A.5.1.1.

COPC – chemical of potential concern

PCB – polychlorinated biphenyl

ROC – receptor of potential concern

This section presents methods and assumptions for estimating doses of COPCs (Section A.5.1) for each ROC, and provides a review of toxicity data identified in the literature (Section A.5.2). Data presented in these sections are synthesized in the risk characterization, and uncertainties are discussed in Section A.6.3.

A.5.1 EXPOSURE ASSESSMENT

This exposure assessment presents the methods used and results of calculated daily exposure doses of COPCs to wildlife receptors in the LDW.

A.5.1.1 Approach

In this assessment, estimates of daily doses of each chemical for each receptor are calculated for three ingested media: food, water, and sediment. Other pathways considered in the conceptual site model in the problem formulation were determined to be insignificant relative to these primary exposure pathways.[60] The daily doses were estimated using the following equation:

$$\text{Daily Dose} = \frac{[(\text{FIR} \times C_{food}) + (\text{WIR} \times C_{water}) + (\text{SIR} \times C_{sed})] \times \text{SUF}}{\text{BW}} \qquad \textbf{Equation 5-1}$$

Where:

Daily Dose	=	COPCs ingested per day via food, water, and sediment (mg COPC/kg body weight/day)
FIR	=	food ingestion rate (kg food dw/day)
C_{food}	=	concentration in prey items (mg COPC/kg food dw)
WIR	=	water ingestion rate (L water/day)
C_{water}	=	concentration in water (mg COPC/L water)
SIR	=	sediment ingestion rate (kg sediment dw/day)
C_{sed}	=	concentration in sediment (mg COPC/kg dw)
SUF	=	site use factor (unitless); fraction of time that a receptor spends foraging in the LDW relative to the entire home range
BW	=	ROC species body weight (kg ww)

Site use factors, body weights, and food, water, and sediment ingestion rates vary among ROCs. The body weights and food, water, and sediment ingestion rates were obtained from the literature for each receptor, as described in Section A.5.1.2. Incidental sediment ingestion rates are calculated as a specified percentage of the food ingestion rate for each ROC. As shown in Equation 5-1, food alone represents 100% of the food ingestion rate, with sediment comprising an additional component of the diet. All COPCs were conservatively assumed to have the same bioavailability in the field as in the laboratory toxicity study that provides the basis for the TRV in all media. The chemical concentration in each receptor's diet was calculated from concentrations in each component of the ROC diet and estimates of each component's

[60] Direct (or dermal) contact with sediment was considered a complete exposure pathway. However, risks from sediment contact are considered to be insignificant relative to those from ingestion (EPA 2000b). Direct contact with water was also considered a complete exposure pathway, but also was assumed to be insignificant because feathers on birds and fur on mammals limit direct contact of skin with contaminated media.

fraction of the total diet. For example, the concentration in food for an ROC that might ingest both fish and benthic invertebrates would be estimated as follows:

$$C_{food} = (C_f \times F_f) + (C_a \times F_a)$$ **Equation 5-2**

Where:

C_{food} = concentration in prey items (mg COPC/kg food dw)
C_f = concentration in fish tissue (mg COPC/kg tissue dw)
C_a = concentration in benthic invertebrate tissue (mg COPC/kg tissue dw)
F_f = fraction of the ROC diet consisting of fish (kg fish/kg food)
F_a = fraction of the ROC diet consisting of benthic invertebrates (kg benthic invertebrates/kg food)

The dietary fraction of each component in each ROC's diet was based on information from the literature. The dietary fractions assumed for each ROC and the assumptions used to derive them are described in detail in Section A.5.1.2.

For PCBs, dietary risks to birds and mammals were evaluated in two ways: as exposure to total PCBs and as exposure to dioxin-like PCB congeners. Dioxin-like PCB congeners show structural and toxic similarities to dioxins and furans. The potency of each individual dioxin-like PCB congener relative to that of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (2,3,7,8-TCDD) is quantified using TEFs. These factors and the concentrations of each dioxin-like PCB congener are multiplied by the congener-specific TEF, and the products are then summed to calculate a toxic equivalent (TEQ) for the dioxin-like PCB congeners in each sample, as follows:

$$TEQ = \sum_{i=1}^{n} C_i TEF_i$$ **Equation 5-3**

Where:

TEF_i = the TEF for an individual PCB congener
C_i = concentration of an individual PCB congener

Using the assumption that the combined effect of these PCB congeners is additive, the total TCDD-like toxicity of the PCB congeners is estimated by summing the TEF-concentration products for individual PCB congeners.

The TEFs used in this ERA were developed by the World Health Organization (WHO) in 1998, with updated values for mammals developed in 2005, from a database containing all available mammalian, bird, and fish studies involving the relative toxicity of dioxin-like compounds (Van den Berg et al. 1998; 2006).

The TEQ approach accounts for the toxicity of a subset of PCB congeners that have dioxin-like modes of toxic action. All PCB congeners may elicit other toxic responses with different toxic mechanisms. Therefore, the TEQ approach is not used as a surrogate for the total PCB approach, which captures all PCB toxicity mechanisms or modes of action because it contains multiple classes of congeners including coplanar PCB congeners. It should also be noted that the TEQs calculated in this ERA were used

only for assessing the toxicity of the dioxin-like PCB congeners and do not account for TEQ contributions from dioxin and furan congeners; dioxin and furan concentrations in tissue were not available for the LDW. Risk estimates for wildlife based on TEQs calculated on the basis of using only dioxin-like PCB congeners are lower than the actual risk resulting from the cumulative exposure of wildlife to dioxins, furans, and dioxin-like PCB congeners in the LDW; the degree of underestimation is uncertain. Uncertainties in the TEQ approach are discussed in Section A.6.3.1.2.

A.5.1.2 Exposure assumptions

This section presents values used in Equations 5-1 and 5-2 to calculate the daily exposure dose for each ROC, including dietary fractions of prey items, ingestion rates of food, water, and incidental sediment, site use factors, and body weights. Tables A.5-2 and A.5-3 summarize these values and the following sections provide details of exposure factor assumptions and sources of information for each ROC.

Table A.5-2. Dietary fractions of prey items used in wildlife exposure calculations

ROC	COPC	Fraction of Prey Item in Diet (by weight)							
		Shiner Surfperch	English Sole	Juvenile Chinook Salmon	Pacific Staghorn Sculpin	Crab	Benthic Invertebrates	Clams	Mussels
Spotted sandpiper	all COPCs	0	0	0	0	0	1.0	0	0
Great blue heron	total PCBs and mercury	0.24	0.24	0.24	0.24	0.05	0	0	0
	other COPCs	0.32	0.32	na[a]	0.32	0.05	0	0	0
Osprey	total PCBs and mercury	0.34	0.21	0.45	0	0	0	0	0
	other COPCs	0.61	0.39	na[a]	0	0	0	0	0
River otter	total PCBs and mercury	0.22	0.22	0.22	0.22	0.10	0	0.01	0.01
	other COPCs	0.29	0.29	na[a]	0.29	0.10	0	0.01	0.01[b]
Harbor seal	total PCBs and mercury	0.25	0.25	0.25	0.25	0	0	0	0
	other COPCs	0.33	0.33	na[a]	0.33	0	0	0	0

[a] COPCs other than total PCBs and mercury were not analyzed in juvenile chinook salmon; therefore, the dietary fractions of other prey were adjusted to compensate.

[b] Selenium and PCB congeners were not analyzed in mussels, so the clam fraction of the diet was assumed to be 0.02 for selenium and PCB TEQ exposure calculations.

COPC – chemical of potential concern

na – not available

PCB – polychlorinated biphenyl

ROC – receptor of concern

Table A.5-3. Exposure factor values for each ROC

ROC	Gender[a]	Body Weight (kg ww)	Food Ingestion Rate (kg dw/day)	Water Ingestion Rate (kg/day)	Incidental Sediment Ingestion Rate (kg dw/day)	Site Use Factor (unitless)[b]
Spotted sandpiper	M	0.038	0.0060	0.0066	0.0011	1
	F	0.047	0.0074	0.0076	0.0013	
	average	0.043	0.0067	0.0071	0.0012	
Great blue heron	M	2.6	0.11	0.11	0.0022	0.5
	F	2.2	0.093	0.10	0.0019	
	average	2.4	0.10	0.11	0.020	
Osprey	M	1.5	0.076	0.077	0.00076	0.75
	F	1.8	0.091	0.087	0.00091	
	average	1.7	0.083	0.083	0.00083	
River otter	M	9.2	0.30	0.73	0.0060	1
	F	7.9	0.26	0.64	0.0052	
	average	8.6	0.28	0.68	0.0056	
Harbor seal	M	85	0.62	5.4	0.012	0.33
	F	77	0.58	4.9	0.012	
	average	81	0.60	5.1	0.0012	

[a] Female values were used for COPCs with a TRV based on a reproductive endpoint and average values were used for COPCs with a TRV based on a growth or survival endpoint. Average values were used for the COPC screen (Equation 2-2 in Section A.2.5.3).

[b] Site use factor is the fraction of a receptor's total foraging time that is spent in the LDW.

dw – dry weight

M – male

F – female

ROC – receptor of concern

ww – wet weight

A.5.1.2.1 Spotted sandpiper

Body Weight

Representative body weights for adult male and female spotted sandpiper (0.0379 and 0.0471 kg, respectively) were obtained from a study by Maxson and Oring (1980), as cited in EPA (1993).

Food Ingestion Rate

The FIR for spotted sandpiper was estimated as a function of the metabolic rate and the caloric content of the prey using the following equation:

$$\text{FIR} = \frac{\text{FMR}}{\text{ME}} \times \frac{0.001\,\text{kg food}}{\text{g food}} \qquad \textbf{Equation 5-4}$$

Where:

FIR = food ingestion rate (kg food dw/day)
FMR = free-living metabolic rate (kcal/day)
ME = average metabolizable energy of the total diet (kcal/g food dw)

The body-weight-normalized free-living metabolic rate (FMR) for the common sandpiper (0.676 kcal/g bw/day) from Nagy et al. (1999) was multiplied by male and female spotted sandpiper body weights to derive male and female FMRs used in Equation 5-4 for spotted sandpiper (25.7 and 31.8 kcal/day, respectively). An average metabolizable energy (ME) value of 4.3 kcal/g dw was used in Equation 5-4. This value was derived from Nagy (1987) as the average ME of insects that are ingested by birds. Using these FMR and ME values, the calculated male and female food consumption rates are 0.0060 and 0.0074 kg dw/day, respectively (Table A.5-3).

Water Ingestion Rate

The water ingestion rate (WIR) was estimated as a function of the spotted sandpiper's body weight, using an allometric equation developed by Calder and Braun (1983), as recommended in EPA (1993), as follows:

$$WIR = 0.059 \times BW^{0.67}$$ **Equation 5-5**

Where:

WIR = water ingestion rate (L water/day)
BW = body weight (kg)

The calculated male and female WIR values are 0.0066 and 0.0076 L/day, respectively.

Incidental Sediment Ingestion Rate

Site-specific data on incidental sediment ingestion by the spotted sandpiper were not available. However, data were available from Beyer et al. (1994) regarding incidental sediment ingestion by four other sandpiper species that feed on mud-dwelling invertebrates. On a dry-weight basis, the incidental sediment ingestion ranged from 7.3 to 30% of the diet, with an average of 18%, which was used in exposure calculations. The uncertainty in estimated sediment ingestion for spotted sandpiper is addressed in the uncertainty analysis (Section A.6.3.1.2). It was assumed that spotted sandpiper would ingest sediment only from intertidal areas.

Composition of Diet

Spotted sandpipers feed along the sandy or muddy edges of inlets, creeks, and ponds. Their diet is composed primarily of terrestrial and marine invertebrates (Bent 1929), but they also may feed on crustaceans, leeches, mollusks, small fish, and carrion (Oring et al. 1983). In the LDW, it was assumed that spotted sandpipers feed solely on benthic invertebrates, such as amphipods and polychaetes, in the intertidal mudflats along the LDW.

Exposure of spotted sandpipers via ingestion of prey was calculated using benthic invertebrate and amphipod tissue data. The possibility that spotted sandpipers could also ingest fish, mussels, or crab was addressed in the uncertainty assessment in the Phase 1 ERA (Windward 2003b).

Site Use

Spotted sandpiper are a common bird in western Washington, and are known to nest along the LDW. They have been observed in the LDW from late June through September (Cordell et al. 1996), but have been known to overwinter locally (Paulson 1993). Spotted sandpipers breed in open habitats along the margins of water bodies (Oring and Lank 1986). Canning et al. (1979) recorded seven spotted sandpiper nests located on Kellogg Island and at least three additional nest sites were thought to be present on the island based on the behavior of adult or juvenile birds in those areas.

A survey was conducted in June 2004 to identify the presence and quality of spotted sandpiper habitat along the LDW (Windward 2004h). Although no nests were observed during this survey, high-quality nesting habitat was observed at the following areas:

- Fringe of salt marsh on the north side of Kellogg Island (RM 0.7 to RM 0.9)
- T-105 restored salmon rearing channel (western shoreline, RM 0.4)
- Herring's House restored off-channel emergent marsh (western shoreline, RM 0.5)
- Channel beneath First Avenue South Bridge (western shoreline, RM 2.1)
- Hamm Creek restoration site (western shoreline, RM 4.3)
- Western shoreline of the Upper Turning Basin (RM 4.6 to RM 4.9).

The sandpiper survey also found that approximately 65% of the LDW shoreline contained foraging habitat; with about 40% of the shoreline considered to be high-quality habitat and about 25% considered to be poor-quality habitat. The intertidal areas associated with these shoreline segments are identified on Map A.5-1 as high- or poor-quality foraging habitat.

Based on the survey observations of potential nesting and foraging habitat, and an estimated foraging range of about 1 mile from the nest (Norman 2002b), six exposure scenarios for spotted sandpiper exposure were developed. Assuming a nest in the center of each area at RM 0.8, RM 2.1, and RM 4.3, and a foraging range of 1 mile upstream and downstream of the nest, three hypothetical exposure areas were created (Map A.5-1). Within each of the three exposure areas, two foraging scenarios were assessed; one in which spotted sandpipers forage only in high-quality habitat and another in which they forage in both high- and poor-quality habitats. These two foraging scenarios in each of the three areas resulted in a total of six exposure scenarios.

A.5.1.2.2 Great blue heron

Body Weight

Representative body weights of adult male and female great blue heron (2.6 and 2.2 kg, respectively) were obtained from Hartman (1961), as cited in EPA (1993).

Food Ingestion Rate

The FIR values for males and females were calculated using an allometric equation for wading birds (Kushlan 1978):

$$FIR = 10^{0.966\log(BW)-0.64} \times \frac{0.001\ kg}{g}$$ **Equation 5-6**

Where:

FIR = food ingestion rate (kg ww/day)
BW = ROC body weight (g)

The FIR values were converted from wet weight to dry weight assuming a moisture content in prey of 76%, which was the average moisture content in fish collected from the LDW during Phase 2. The body weights and calculated FIR values are presented in Table A.5-3.

Water Ingestion Rate

The WIR was estimated as a function of the great blue heron's body weight, using the allometric equation recommended in *Wildlife Exposure Factors Handbook* (EPA 1993). This equation is the same one presented in Section A.5.1.2.1 (Equation 5-5). WIR values of 0.11 and 0.10 L/day were calculated for male and female great blue heron, respectively.

Incidental Sediment Ingestion Rate

Information on rates of incidental sediment ingestion by herons was not available. Great blue herons may ingest a small amount of sediment while foraging for food. Sediment ingestion was estimated to be 2% of the heron's diet. It was assumed that only intertidal sediment would be ingested.

Composition of Diet

Great blue herons feed in shallow water primarily on fish, but they also ingest crustaceans, insects, amphibians, reptiles, and occasionally small mammals (Kushlan 1978; Butler 1993). Great blue herons hunt by sight and stalk or ambush their prey. They will also feed by probing, quickly moving their bills in and out of the water and substrate. Great blue herons feed on small fish that range in length from 8 to 33 cm (Kirkpatrick 1940; Alexander 1977; Hoffman 1978). Butler (1993) reported that shiner surfperch is a major food source for female and hatchling great blue herons and may be important for juvenile survival. Butler (1997) also reported that sculpin and starry flounder are important prey species for herons along the British Columbia coast, and that shrimp and crabs may also be ingested. The predominant prey items identified at

the West Seattle colony were tails from sculpin (Krausmann 2002a). However, sculpin may not be representative of actual prey captured because the large lateral and dorsal spines make them more likely to be rejected by the heron chicks (Krausmann 2002a). In addition to perch and sculpin, herons may feed on other small fish in the LDW, including juvenile salmonids.

For exposure calculations, it was estimated that crabs account for a small portion of the heron's diet (5%), with shiner surfperch, Pacific staghorn sculpin, juvenile chinook salmon, and English sole/starry flounder accounting for equal amounts of the remaining 95 % of the diet.

Site Use

A great blue heron colony of at least 37 nests (presumably more than 74 birds) occupied a site in West Seattle in 1998 (Norman 2002). In 1999, there were only 21 active nests. No successful nesting occurred in 2000 or 2001, and since then, the colony has remained empty (Norman 2002, 2006). It is suspected that the colony was abandoned because of eagle predation or human disturbance from construction at the Herring's House restoration area (Norman 2006). Nevertheless, because of suitable habitat present in the West Seattle area, another colony may be established (Norman 2006). Other colonies are located 12 km south in Renton (Black River colony) and 12 km northwest between the Ship Canal and Discovery Park (Kiwanis Ravine colony). The Black River colony contains about 120 to 135 active nests (Heron Working Group 2006b). The Kiwanis Ravine colony contained 44 nests in 2005, with an estimated 100 young herons successfully fledged (Heron Habitat Helpers 2005).

Information presented in *Wildlife Exposure Factors Handbook* (EPA 1993) indicates that foraging grounds are generally close to breeding colonies, and that 15 to 20 km is the farthest great blue herons might regularly travel from the colony to the foraging area. Herons from the Black River colony are known to forage in the Black River Riparian Forest, along the Cedar River, the Green-Duwamish River, and the Lake Washington shoreline (Heron Working Group 2006b), but the relative time spent foraging in each of these areas is unknown. Studies have not been conducted to determine where herons from the Kiwanis Ravine colony forage, but they have been observed flying in the direction of both saltwater and freshwater during the most intense feeding periods (Heron Habitat Helpers 2005) and great blue herons are regularly observed along the LDW. Based on observations of individual birds from the West Seattle colony when it was active, it was estimated that at least half of the birds from this colony used the LDW to forage, primarily in the Kellogg Island area (Krausmann 2002b). A site use factor of 0.5 was assumed for herons based on observations of the abandoned West Seattle colony; limited foraging information was available for Black River and Kiwanis Ravine herons. The uncertainty of this site use factor is discussed in the uncertainty analysis.

A.5.1.2.3 Osprey

Body Weight

Representative body weights for adult male and female osprey (1.5 and 1.8 kg, respectively) were obtained from Poole (1989), as cited in Poole et al. (2002).

Food Ingestion Rate

The FIR in wet weight was estimated as 21% of the body weight (Poole 1983; as cited in EPA 1993). FIR values were converted from wet weight to dry weight assuming a moisture content in prey of 76%, which was the average moisture content in fish collected from the LDW during Phase 2. FIR values of 0.076 and 0.091 kg dw/day were calculated for male and female osprey, respectively.

Water Ingestion Rate

The WIR was estimated as a function of the osprey's body weight, using the allometric equation recommended in *Wildlife Exposure Factors Handbook* (EPA 1993). This equation is presented in Section A.5.1.2.1 (Equation 5-5). WIR values of 0.077 and 0.087 L/day were calculated for male and female osprey, respectively.

Incidental Sediment Ingestion Rate

Data on incidental sediment ingestion by osprey were not available. Osprey may ingest a small amount of sediment while foraging for fish in shallow water. Sediment ingestion was estimated to be 1% of the osprey's diet. It was assumed that only intertidal sediment would be ingested.

Composition of Diet

Osprey feed almost exclusively on live fish; at least 99% of their prey items were live fish in most published accounts (Poole et al. 2002). Ospreys can penetrate about 1 m below the water surface. Therefore, they generally catch pelagic fish or those that frequent shallow flats and shorelines. Ospreys may infrequently ingest other types of vertebrate prey, such as birds, reptiles, and small mammals. A west-central Idaho osprey study reported 89% of fish ingested by osprey were 11 to 30 cm long, suggesting a preference for medium-sized fish (Van Daele and Van Daele 1982). During a USGS study, osprey were observed while returning to their nests along the LDW with prey. Seventy-three percent of the prey observed were marine fish species (Henny 2005). Salmon, perch, and sole/flounder accounted for approximately 33, 25, and 15% of total prey, respectively. Twenty-five percent of the total prey was peamouth, a freshwater fish, and 2% were not reported. Preliminary data indicate that the size of some of the salmonids taken by osprey from the LDW during the pre-egglaying period ranged from 18 to 30 cm in length (Henny 2006). This dietary information was collected during only a portion of the nesting season. Additional dietary data have been collected and are currently being analyzed but are not yet available (Kaiser 2006). The types of fish ingested by osprey for this assessment were assumed to be the same as those observed in the USGS study, with the proportions re-

allocated to represent 100% of the diet. Juvenile chinook salmon, shiner surfperch, and English sole/starry flounder were assumed to account for 45, 34, and 21% of the diet, respectively. The juvenile chinook salmon collected in the LDW had an average length of 8 cm, so tissue data for these fish may not represent salmon captured by osprey in the LDW. However, they were used as surrogates for larger juvenile salmon. Uncertainties associated with the composition of the osprey's diet are addressed in the uncertainty analysis (Section 6.3.3.2).

Site Use

There are five known osprey nests along the LDW between RM 0.0 and 5.0, one nest on Harbor Island, and three nests along the Duwamish and Green Rivers within about 4 miles of the Upper Turning Basin. The distance osprey travel from their nests to forage depends on the availability of fish near the nest (Van Daele and Van Daele 1982). Osprey were given a site use factor of 0.75, because preliminary USGS data on prey for LDW osprey indicated that approximately 25% of fish in the diet were freshwater; the remaining 75% marine fish were all assumed to be caught from the LDW.

A.5.1.2.4 River otter

Body Weight

Adult body weights of 9.2 and 7.9 kg were assumed for male and female river otter, respectively (Table A.5-3) based on a study by Melquist and Hornocker (1983), as cited in EPA (1993).

Food Ingestion Rate

The FIR was estimated as a function of the metabolic rate and the caloric content of the prey using the following equation:

$$\text{FIR} = \frac{\text{FMR}}{\text{ME}} \times \frac{0.001\,\text{kg food}}{\text{g food}} \qquad \textbf{Equation 5-7}$$

Where:

FIR	=	food ingestion rate (kg food dw/day)
FMR	=	free-living metabolic rate (kilocalories [kcal]/day)
ME	=	average metabolizable energy of the total diet (kcal/g food dw)

The FMRs for males and females were calculated to be 1,340 and 1,180 and kcal/day, using an equation developed by Nagy (1987), as cited in EPA (1993) for placental mammals:

$$\text{FMR (kcal/day)} = 0.800 \times \text{BW}^{0.813} \qquad \textbf{Equation 5-8}$$

where body weight is expressed in grams. The ME value used for mammals on a diet of fish was that calculated by Nagy (1987), as cited in EPA (1993) (4.47 kcal/g dw). The

calculated FIRs for males and females were 0.30 and 0.26 kg dw/day, respectively (Table A.5-3).

Water Ingestion Rate

The WIR was estimated as a function of the river otter's body weight, using an allometric equation recommended in EPA (1993). This equation was developed by Calder and Braun (1983), as cited in EPA (1993):

$$WIR = 0.099 \times BW^{0.90}$$ **Equation 5-9**

Where:

WIR = water ingestion rate (L water/day)
BW = body weight (kg)

WIR values for males and females (0.73 and 0.64 L/day, respectively) were calculated.

Incidental Sediment Ingestion Rate

Data were not available to estimate the amount of sediment ingested incidentally by river otters. A small amount of sediment could be ingested when river otters forage on crabs and benthic fish species; therefore, the incidental sediment ingestion rate was estimated to be 2% of the diet. It was assumed that river otters incidentally ingest sediment from both intertidal and subtidal areas of the LDW.

Composition of Diet

River otters are opportunistic carnivores that take advantage of food that is most abundant and easiest to catch. Fish are their primary prey (Wise et al. 1981; Kurta 1995; Larsen 1984; Stenson et al. 1984). River otters catch fish by diving and ambushing or chasing, and obtain invertebrates by digging in the substrate (Coulter et al. 1984). Slower-moving fish, such as suckers, carp, chubs, and bullheads, are generally eaten most frequently (Wise et al. 1981; Kurta 1995). Studies in coastal southeast Alaska and British Columbia found that river otters feed primarily on sculpin, surfperch, and flatfish, with greenling, salmon, and rockfish making up lesser portions of the diet (Larsen 1984; Stenson et al. 1984). Other components of the river otter diet include aquatic invertebrates (including crayfish, mussels, clams, and aquatic insects), frogs, snakes, and occasionally mammals and birds (Coulter et al. 1984). River otters generally ingest fish ranging from 7.6 to 41 cm in length (Gilbert and Nancekivell 1982; Greer 1955; both as cited in EPA 1993), although Toweill (1974) found that many of the salmon preyed upon by river otters in western Oregon were up to an estimated 80 cm in length. These salmon were taken in coastal streams where fish enter the rivers to spawn. Local river otters feed largely on fish, but will also feed on crabs and sometimes mussels and clams (Strand 1999).

The proportion of prey types ingested by river otter for this assessment was based on Larsen's (1984) study of river otters in southeastern Alaska. This study was used because it was the only study from the Pacific Northwest that reported remains in scat

on a volume basis rather than as a frequency of occurrence. Larsen (1984) reported the following proportions of prey ingested by river otters: 86% fish, 10% crabs, 2% invertebrates other than crabs, 1% birds, and 1% mammals and plant material. Thus, for this assessment, it is assumed that river otters ingest 88% fish, 10% crabs, and 2% mussels. Based on feeding habits of river otters documented in coastal southeast Alaska and British Columbia (Larsen 1984; Stenson et al. 1984), any of the four types of fish tissue for which chemistry data were available in the LDW might be ingested. Because no site-specific information was available on fish preference of river otters, it was assumed that shiner surfperch, English sole/starry flounder, juvenile chinook salmon, and Pacific staghorn sculpin are ingested in equal proportions of the 88% of their diet that is fish. The juvenile chinook salmon collected in the LDW had an average length of 8 cm, so tissue data for these fish may not represent salmon captured by river otters in the LDW. However, they were used as surrogates for larger salmon; uncertainties associated with this assumption are addressed in the uncertainty analysis (Section 6.3.4.2).

Site Use

Anecdotal information indicates that a river otter family lives year-round on Kellogg Island in the LDW, although otters were not observed during wildlife surveys by Cordell (2001). River otters are almost exclusively aquatic and prefer food-rich habitats such as the lower portions of streams and rivers, estuaries, and lakes and tributaries that feed rivers (Tabor and Wight 1977; Mowbray et al. 1979). In streams, the river otter's home range can average 30 km (19 mi) (Melquist and Hornocker 1983). At any given time, river otters generally occupy only a few kilometers of stream, but often move from one area to another (Nebraska Game and Parks Commission 2000). A radio-tracking study of relocated river otters was conducted as part of the New York State Department of Environmental Conservation river otter reintroduction program. This study showed that river otter ranges were from 1.5 to 22.4 km long, with an average length of 10 km (6 mi) for individuals monitored in western New York State (Spinola et al. 1999; as cited in EPA 2000d).

No studies were found that document usage of the LDW by river otters. Because of the average 10 km linear length documented in the literature, and because the extent of the LDW study area is approximately 10 km, it was assumed that river otters could potentially ingest prey solely from within the LDW. Therefore, a site use factor of 1.0 was assumed.

A.5.1.2.5 Harbor seal

Body Weight

Body weights for adult male and female harbor seals (84.6 and 76.5 kg, respectively) were based on a study by Pitcher and Calkins (1979), as cited in the *Wildlife Exposure Factors Handbook* (EPA 1993).

Food Ingestion Rate

The FIR for harbor seals was calculated using an allometric equation developed by Boulva and McLaren (1979) for harbor seals from eastern Canada, as cited in the *Wildlife Exposure Factors Handbook* (EPA 1993):

$$FIR = 0.089 \times BW^{0.76} \qquad \textbf{Equation 5-10}$$

Where:

FIR = food ingestion rate (kg food ww/day)
BW = ROC body weight (kg)

The calculated wet weight FIR values were converted to dry weight using the average moisture content (76%) in whole-body fish from the LDW. FIRs calculated for males and females were 0.62 and 0.58 kg dw/day, respectively (Table A.5-3).

Water Ingestion Rate

The WIR was estimated as a function of the harbor seal's body weight, using the allometric equation recommended in EPA (1993). This equation is presented in Section A.5.1.2.4 (Equation 5-9). Using the male and female body weights of the harbor seal, the calculated WIR values were 4.9 and 5.4 L/day, respectively.

Incidental Sediment Ingestion Rate

Data on incidental sediment ingestion by harbor seals were not available, but it is possible that a small amount of sediment could be incidentally ingested while foraging on bottom fish. Therefore, the sediment ingestion rate was assumed to be 2% of the FIR. It was assumed that harbor seals ingest sediment from both intertidal and subtidal areas of the LDW.

Composition of Diet

Harbor seals are opportunistic feeders, selecting prey based on availability and ease of capture (Pitcher and Calkins 1979; Pitcher 1980; Schaffer 1989). Their diet can vary seasonally with local abundance and includes bottom-dwelling fishes, invertebrates such as octopus and squid, and species that congregate for spawning (Pitcher and Calkins 1979; Everitt et al. 1981; Lowry and Frost 1981; Roffe and Mate 1984). In Washington, the most important prey include Pacific whiting, tomcod, walleye pollock, flatfish, Pacific herring, shiner surfperch, plainfin midshipman, and sculpin (NMFS 1997). Fish ingested were generally between 4 and 28 cm in length (Brown and Mate 1983). Harbor seals may also prey on salmon during upriver spawning migrations of adults or downriver outmigrations of juveniles, although site-specific data were not available on the dietary importance of migrating salmon to local seal populations. Because site-specific information was not available on the amount of each type of prey ingested, it is assumed that juvenile chinook salmon, English sole/starry flounder, shiner surfperch, and Pacific staghorn sculpin are ingested in equal proportions. The juvenile chinook salmon collected in the LDW had an average length

of 8 cm, so they may or may not be representative of salmon ingested by harbor seals in the LDW. However, they are used as surrogates for larger salmon; the absence of specific information on types of prey ingested by harbor seals is discussed in the uncertainty analysis (Section A.6.3.5.2).

Site Use Factor

Harbor seals are commonly seen in Elliott Bay and occasionally enter the LDW (Kenney 1982). Harbor seals have been shown to forage over large distances, ranging from 5 km (3 mi) (Stewart et al. 1989) to 55 km (34 mi) (Beach et al. 1985). In Puget Sound, harbor seals generally forage within 8 to 13 km (5 to 8 mi) of their haulout areas established as pupping sites (Jeffries 2001). The closest known pupping site to the LDW is located at Blakely Rocks off the southeast end of Bainbridge Island, approximately 12 km (7 mi) from the LDW. Site-specific information on harbor seal usage of the LDW is limited. The WDFW observed harbor seals infrequently in the LDW during an intensive survey conducted from December 1998 to June 1999 (WDFW 1999). This survey monitored the waterway for the presence of sea lions and seals up to the 16th Avenue South Bridge for a total of 307 hours on 52 days. Harbor seals were observed on 17 occasions, and were most frequently seen north of the 1st Avenue South Bridge. While harbor seals have been observed in Elliott Bay and may use log booms to haul out, they are not known to aggregate in large numbers (Jeffries 2001). The LDW may be a preferential feeding area during salmonid outmigration from March through August. For example, in the Columbia River, salmonids appear to be targeted as prey by seals in the spring and fall when they are abundant and available in the river (NMFS 1997).

Data from the WDFW survey discussed above were used to establish a site use factor for risk calculations. The following conservative assumptions were used: 1) a harbor seal was observed in the LDW on 17 of 52 days of observation, and it was assumed that all observations were of a single seal; 2) the seal was assumed to obtain all of its food for those days in the LDW; and 3) site usage from December through June accurately represents usage during other times of the year. Based on these assumptions, the site use factor was equal to 17/52 or 0.33. This approach does not include exposure of harbor seals in Elliott Bay to fish that may have been exposed to chemicals in the LDW or through a food web connected to the LDW.

A.5.1.3 Prey tissue, sediment, and water data

This section presents the COPC concentrations in prey tissue, sediment, and water that were used in Equations 5-1 and 5-2 to calculate exposure doses. Attachment 11 presents summary statistics (i.e., minimum, maximum, and mean COPC concentrations) for the prey tissue, sediment, and water data.

A.5.1.3.1 Prey tissue

Tissue data were available for eight tissue types that are potential prey of wildlife ROCs in the LDW: shiner surfperch, English sole, Pacific staghorn sculpin, juvenile

chinook salmon, benthic invertebrates,[61] clams, crabs, and mussels. These data are described in Section A.2.4.1.2.

For all wildlife ROCs except spotted sandpiper, it was assumed that the foraging area includes the entire LDW based on home range information presented in Section A.5.1.2. Thus, for each prey type ingested by those species (i.e., all prey types except benthic invertebrates), the exposure concentration was calculated using all tissue data from the LDW as one exposure data set. The exposure concentration for each tissue type was estimated as the UCL,[62] which was calculated as described in Attachment 11. For example, for great blue heron, all Pacific staghorn sculpin composite tissue samples collected in the LDW were used to estimate the UCL on the mean for this prey type. The calculated concentrations for COPCs in each tissue type are shown in Table A.5-4.

Table A.5-4. COPC concentrations in tissues of prey species ingested by great blue heron, osprey, river otter, and harbor seal

COPC	Tissue Concentration in Prey Species (mg/kg dw)						
	Shiner Surfperch	English Sole	Juvenile Chinook Salmon	Pacific Staghorn Sculpin	Clams	Crabs	Mussels
Arsenic	4.2	13	na	4.0	25	25	5.9
Chromium	0.82	4.4	na	0.40	5.2	0.30	1.2
Cobalt	0.18	0.41	na	0.13	2.9	0.39	0.41
Lead	0.53	1.7	na	0.30	19	0.53	3.3
Mercury	0.15	0.070	0.14	0.15	0.12	0.29	0.094
Selenium	0.77	0.82	na	0.87	2.0	1.2	na
Total PCBs	14	10	3.4	5.0	4.0	6.0	0.27
PCB TEQs (birds)[a]	7.2×10^{-4}	4.2×10^{-4}	na	1.2×10^{-4}	1.5×10^{-4}	3.6×10^{-4}	na
PCB TEQs (mammals)[b]	1.6×10^{-4}	8.2×10^{-5}	na	3.8×10^{-5}	2.7×10^{-5}	5.0×10^{-5}	na

[a] PCB TEQs were calculated using TEFs for birds presented in Van den Berg et al. (1998).These TEFs are listed in Attachment 3, and uncertainties associated with application of these TEFs are discussed in Section A.6.3.1.

[b] PCB TEQs were calculated using TEFs for mammals presented in Van den Berg et al. (2006). These TEFs are listed in Attachment 3.

COPC – chemical of potential concern

dw – dry weight

na – not available

PCB – polychlorinated biphenyl

TEQ – toxic equivalent

[61] Includes amphipod samples collected for the King County Water Quality Assessment (King County 1999d) and benthic invertebrate samples collected for the Phase 2 RI (Windward 2005b).

[62] The UCL was used even if it was higher than the maximum detected concentration.

For spotted sandpipers, which forage over an area smaller than the entire LDW (see Section A.5.1.2.1), concentrations of COPCs in the fraction of the diet consisting of benthic invertebrate tissues were calculated separately for each of six exposure scenarios described in Section A.5.1.2.1. For COPCs with a linear relationship between concentrations in sediment and tissue (i.e., arsenic and total PCBs), the COPC concentrations in benthic invertebrate tissue were estimated using a linear regression, as described in Section 4.0 of Attachment 11. For arsenic, the arithmetic average surface sediment concentration within each intertidal exposure area was used to estimate the benthic invertebrate tissue concentration for each exposure area using the linear regression. For total PCBs, spatially weighted average concentrations (SWACs) for surface sediments have been developed for each exposure area, so these SWACs were used to estimate the total PCB benthic invertebrate tissue concentrations from the linear regression. The SWACs were used for total PCBs because the arithmetic mean of the sediment concentrations in a specified area may be biased high if more highly contaminated areas were more intensively sampled than less-contaminated areas. In the LDW, this bias is most pronounced for total PCBs, which are the focus of several ongoing early action investigations in the LDW. The 95th UCLs on the total PCB and arsenic tissue concentrations estimated from the linear regression were used as the benthic invertebrate concentrations in the exposure assessment (Table A.5-5). The statistical methods used to calculate the benthic tissue concentrations are described in detail in Attachment 11. The locations of intertidal sediment samples included in each exposure scenario are shown in Map A.5-1.

For the remaining COPCs, a significant statistical correlation was not observed between benthic invertebrate tissue and sediment. Therefore, concentrations of these COPCs in benthic invertebrate tissue were calculated as the UCL of the available tissue data collected from each sandpiper intertidal exposure area to estimate spotted sandpiper exposure. Locations of benthic invertebrate tissue samples included in these calculations are shown on Map A.5-1. COPC concentrations in benthic invertebrate tissues that were used in the spotted sandpiper exposure calculations are summarized in Table A.5-5.

Table A.5-5. COPC concentrations in benthic invertebrate tissues used to estimate spotted sandpiper exposure

COPC	Tissue Concentration in Benthic Invertebrates (mg/kg dw)					
	Exposure Area 1 [a]		Exposure Area 2 [a]		Exposure Area 3 [a]	
	High-Quality Foraging Habitat	High- and Poor-Quality Foraging Habitat	High-Quality Foraging Habitat	High- and Poor-Quality Foraging Habitat	High-Quality Foraging Habitat	High- and Poor-Quality Foraging Habitat
Arsenic	23	21	25	21	18	26
Cadmium	0.56	0.56	0.40	0.49	0.64	0.64
Chromium	3.0	3.0	18	51	5.0	5.0
Cobalt	1.8	1.8	2.0	2.2	2.2	2.2
Copper	120	120	140	140	70	70
Lead	31	31	15	660	5.6	5.6
Mercury	0.10	0.10	0.066	0.070	0.50	0.50
Nickel	3.5	3.5	5.8	5.3	7.4	7.4
Selenium	1.9	1.9	1.6	1.8	1.4	1.4
Zinc	190	190	300	380	270	270
Vanadium	6.4	6.4	10	9.8	9.6	9.6
BEHP	2.9	2.9	14	14	61	61
Total PCBs	1.5	1.5	5.9	3.8	2.7	3.4
PCB TEQs (bird)[b]	1.6×10^{-4}	1.6×10^{-4}	5.6×10^{-4}	5.6×10^{-4}	3.9×10^{-4}	3.9×10^{-4}

[a] Six exposure scenarios were evaluated; in each of three exposure areas, foraging only in high-quality habitats and foraging in both high- and poor-quality habitats were evaluated. These exposure scenarios are described in detail in Section A.5.1.2.1.

[b] PCB TEQ concentrations were calculated using TEFs for birds based on TEFs presented in Van den Berg et al. (1998). These TEFs are listed in Attachment 3, and uncertainties associated with application of these TEFs are discussed in Section A.6.3.1.

BEHP – bis(2-ethylhexyl) phthalate

COPC – chemical of potential concern

dw – dry weight

PCB – polychlorinated biphenyl

TEQ – toxic equivalent

A.5.1.3.2 Sediment

Surface sediment data were used to estimate COPC exposure resulting from incidental sediment ingestion. For river otter and harbor seal, it was assumed that the foraging area includes the entire LDW; thus, COPC concentrations in sediment were calculated as the UCL of all surface sediment data.[63] For osprey and great blue heron, it was assumed that the foraging area includes only intertidal areas (areas exposed during

[63] The UCL was used even if it was higher than the maximum detected concentration.

low tide) of the LDW; thus, sediment COPC concentrations were calculated as the UCL of only intertidal surface sediment data. For spotted sandpiper, surface sediment COPC concentrations were calculated as the UCL for each of the six intertidal exposure scenarios discussed in Section A.5.1.2.1 (Map A.5-1).

For PCBs, the sediment concentration was calculated as the UCL of the SWAC concentration for surface sediments discussed above. Statistical methods for calculating this UCL are presented in Attachment 11. For other COPCs, the concentration in sediment was calculated as the UCL of the point data. The calculated concentrations in sediment are presented in Tables A.5-6 (for all ROCs, except spotted sandpiper) and Table A.5-7 (for spotted sandpiper).

Table A.5-6. COPC concentrations in LDW sediment used to estimate exposure of wildlife ROCs, except spotted sandpiper

COPC	Sediment Concentration (mg/kg dw)	
	Entire LDW[a]	Intertidal Areas[b]
Arsenic	30	na
Chromium	50	70
Cobalt	10	na
Lead	300	600
Mercury	0.30	0.40
Selenium	5.0	na
Total PCBs	0.72	0.98
PCB TEQs (bird)[c]	5.4×10^{-4}	9.2×10^{-4}
PCB TEQs (mammal)[d]	7.2×10^{-5}	na

[a] Used for exposure calculations for river otter and harbor seal.

[b] Used for exposure calculations for osprey and great blue heron.

[c] PCB TEQ concentrations were calculated using TEFs for birds as presented in Van den Berg et al. (1998). These TEFs are listed in Attachment 3, and uncertainties associated with application of these TEFs are discussed in Section A.6.3.1.

[d] PCB TEQ concentrations were calculated using TEFs for mammals presented in Van den Berg et al.(2006).

COPC – chemical of potential concern

dw – dry weight

LDW – Lower Duwamish Waterway

na – not applicable; not a COPC for ROCs using intertidal areas (i.e., great blue heron and osprey) based on the COPC screen presented in Section A.2.5.3.

Table A.5-7. COPC concentrations in sediment used to estimate spotted sandpiper exposure

COPC	Sediment Concentration (mg/kg dw)					
	Exposure Area 1[a]		Exposure Area 2[a]		Exposure Area 3[a]	
	High-Quality Foraging Habitat	High- and Poor-Quality Foraging Habitat	High-Quality Foraging Habitat	High- and Poor-Quality Foraging Habitat	High-Quality Foraging Habitat	High- and Poor-Quality Foraging Habitat
Arsenic	36	18	49	18	20	60
Cadmium	0.45	0.65	1.0	0.93	0.62	7.8
Chromium	32	30	43	32	31	120
Cobalt	9.0	8.0	9.5	8.0	8.0	10
Copper	94	80	120	79	63	730
Lead	90	90	160	96	90	1,000
Mercury	0.17	0.16	0.43	0.29	0.60	0.40
Nickel	21	30	22	20	22	90
Selenium	0.59	0.56	7.2	3.0	7.0	6.6
Zinc	180	170	240	180	150	710
Vanadium	53	52	58	55	57	59
BEHP	0.12	2.5	2.7	2.1	0.41	0.41
Total PCBs	0.34	0.33	2.5	1.5	0.72	1.1
PCB TEQs (bird)[b]	3.7×10^{-5}	3.4×10^{-5}	4.3×10^{-3}	1.8×10^{-3}	9.6×10^{-5}	3.0×10^{-3}

[a] Six exposure scenarios were evaluated; in each of three exposure areas, foraging only in high quality habitat and foraging in both high and poor quality habitat were evaluated. These exposure scenarios are described in detail in Section A.5.1.2.1.

[b] PCB TEQs were calculated using TEFs for birds as presented in Van den Berg et al. (1998). These TEFs are listed in Attachment 3, and uncertainties associated with application of these TEFs are discussed in Section A.6.3.1.

BEHP – bis(2-ethylhexyl) phthalate

COPC – chemical of potential concern

dw – dry weight

PCB – polychlorinated biphenyl

TEQ – toxic equivalent

A.5.1.3.3 Water

Water data used to estimate exposure of wildlife to COPCs in the LDW were collected at three locations in 1996 and 1997 as part of the King County WQA (King County 1999d) and at two locations during an additional sampling event by King County in 2005, in which water data were collected for PCB congeners only (Mickelson and Williston 2006), as described in Section A.2.4.1.3. Surface water ingestion represents a very small proportion of the exposure of wildlife to COPCs in the LDW (e.g., less than 0.01% of the overall dose of PCBs is from water).[64]

Metals and BEHP

Metals and BEHP[65] were the only chemicals detected in water samples collected along transects from the three LDW locations sampled as part of the WQA: RM 1.1 in the vicinity of the Brandon CSO, RM 1.9 in the vicinity of the Southwest Michigan CSO, and RM 4.9 in the vicinity of the Norfolk CSO. The Brandon CSO and Southwest Michigan CSO transects contained three sampling locations and the Norfolk CSO transect contained two sampling locations. Each location was sampled at 1 m below the surface and 1 m above the bottom. For estimates of exposure to ROCs expected to forage throughout the LDW (great blue heron, osprey, river otter, and harbor seal), data from all locations and depths were combined into one dataset and each data point was treated equally. The exposure concentration for each metal COPC was estimated as the UCL of the combined data, as presented in Table A.5-8.

Exposure concentrations of total DDTs in water were not calculated because of the uncertainties in predicting organochlorine pesticide concentrations using data from semi-permeable membrane devices. The contribution of COPCs from water to the total exposure dose is very small and is not expected to affect risk calculations (e.g., the contribution of total PCBs via water exposure was less than 0.01 percent of the contribution from food and sediment exposure).

---

[64] There is some uncertainty associated with exposure to PCBs through water. Exposure to PCBs in water was based on a summation of PCB congener concentrations in grab samples collected from the channel at two locations in the LDW; these samples were collected by King County in 2005. It is very unlikely that risk estimates would be affected by the uncertainty associated with this limited sampling because ingestion of PCBs through water is less than 0.01% of the amount ingested from food and sediment.

[65] BEHP was detected in laboratory blank water samples; therefore, BEHP concentrations in water are likely overestimated.

Table A.5-8. Concentrations of metals and BEHP in surface water used to estimate exposure of great blue heron, osprey, river otter, and harbor seal

COPC	LDW-Wide Concentration in Water (mg/L)
Arsenic	8.8×10^{-4}
Cadmium	4.7×10^{-5}
Chromium	7.1×10^{-4}
Cobalt	2.7×10^{-4}
Copper	1.6×10^{-3}
Lead	5.0×10^{-4}
Mercury	2.7×10^{-6}
Nickel	8.2×10^{-4}
Selenium[c]	2.7×10^{-4}
Vanadium	1.4×10^{-3}
Zinc	3.2×10^{-3}
BEHP	2.1×10^{-3}

BEHP – bis(2-ethylhexyl) phthalate

COPC – chemical of potential concern

For spotted sandpiper, water data collected at the Brandon, Southwest Michigan, and Norfolk locations were used for Exposure Areas 1, 2, and 3, respectively. The UCL was based upon the available data from each location (Table A.5-9).

Table A.5-9. Concentrations of metals and BEHP in surface water used to estimate exposure of spotted sandpiper

COPC	Concentration in Water (mg/L)		
	Exposure Area 1	Exposure Area 2	Exposure Area 3
Arsenic	9.7×10^{-4}	9.3×10^{-4}	5.3×10^{-4}
Cadmium	5.4×10^{-5}	5.5×10^{-5}	1.6×10^{-5}
Chromium	6.1×10^{-4}	7.3×10^{-4}	1.1×10^{-3}
Cobalt	2.2×10^{-4}	2.5×10^{-4}	4.6×10^{-4}
Copper	1.5×10^{-3}	1.6×10^{-3}	2.3×10^{-3}
Lead	3.8×10^{-4}	4.9×10^{-4}	8.9×10^{-4}
Mercury	2.0×10^{-6}	2.0×10^{-4}	4.3×10^{-6}
Nickel	7.1×10^{-4}	8.2×10^{-4}	1.3×10^{-3}
Selenium	2.7×10^{-4}	1.6×10^{-4}	1.6×10^{-4}
Zinc	3.0×10^{-3}	3.0×10^{-3}	4.6×10^{-3}
Vanadium	1.4×10^{-3}	1.5×10^{-3}	1.9×10^{-3}
BEHP	3.5×10^{-3}	2.4×10^{-2}	1.8×10^{-4}

BEHP – bis(2-ethylhexyl) phthalate

COPC – chemical of potential concern

PCBs

PCBs were analyzed in water samples collected from two locations: RM 0.0 (LTKE03) and RM 3.3 (LUTM03), as described in Section A.2.4.1.3. For the ROCs expected to forage throughout the LDW (great blue heron, osprey, river otter, and harbor seal), all data from both locations were combined into one dataset. The PCB water exposure concentration for these four ROCs was estimated as the UCL of these combined data, as presented in Table A.5-10.

Table A.5-10. Total PCB concentrations in surface water used to estimate exposure of wildlife ROCs

COPC	Concentration in Water (mg/L)		
	Osprey, Great Blue Heron, River Otter, and Harbor Seal LDW-Wide	Spotted Sandpiper	
		Exposure Area 1	Exposure Areas 2 and 3
Total PCBs	1.6×10^{-6}	1.7×10^{-6}	1.9×10^{-6}

COPC – chemical of potential concern

LDW – Lower Duwamish Waterway

PCB – polychlorinated biphenyl

ROC – receptor of concern

For spotted sandpiper, PCB data from the LTKE03 location were used for Area 1, and PCB data from the LUTM03 location were used for Areas 2 and 3. The UCLs were based on all data from each location, as presented in Table A.5-10.

PCB TEQs in water were not calculated because PCB exposure via water is a negligible component of overall PCB exposure, based on total PCB exposure calculations. The contribution of total PCBs via water exposure was less than 0.01% of the contribution from food and sediment exposure.[66]

A.5.1.4 Estimated dietary doses

Exposures as dietary doses were estimated for each wildlife ROC based on the information presented in preceding sections. Estimated dietary exposures for spotted sandpiper are presented in Table A.5-11, and those for great blue heron, osprey, river otter, and harbor seal are presented in Table A.5-12. Tables containing all data used in these calculations for each COPC/ROC pair are presented in Attachment 12.

[66] There is some uncertainty in not including undetected PCB congeners in the calculation of total PCBs, but the contribution of total PCBs via water exposure would remain less than 0.01% if undetected congeners were included at one-half the reporting limit; therefore, uncertainty associated with undetected PCB congeners is very low.

Table A.5-11. Estimated dietary doses of COPCs for spotted sandpiper

COPC	Dietary Dose (mg/kg bw/day)					
	Exposure Area 1[a]		Exposure Area 2[a]		Exposure Area 3[a]	
	High-Quality Foraging Habitat	High- and Poor-Quality Foraging Habitat	High-Quality Foraging Habitat	High- and Poor-Quality Foraging Habitat	High-Quality Foraging Habitat	High- and Poor-Quality Foraging Habitat
Arsenic	4.6	3.8	5.3	3.8	3.4	5.8
Cadmium	0.10	0.11	0.090	0.10	0.12	0.32
Chromium	1.4	1.3	4.0	8.8	1.6	4.1
Cobalt	0.53	0.50	0.58	0.57	0.57	0.62
Copper	21	21	25	24	13	31
Lead	7.4	7.4	6.8	110	3.4	29
Mercury	0.020	0.020	0.022	0.019	0.095	0.090
Nickel	1.1	1.4	1.5	1.4	1.8	3.7
Selenium	0.32	0.31	0.45	0.37	0.41	0.40
Zinc	35	34	53	64	46	62
Vanadium	2.5	2.4	3.2	3.1	3.1	3.1
BEHP	0.46	0.53	2.3	2.3	9.6	9.6
Total PCBs	0.25	0.25	1.0	0.64	0.45	0.57
PCB TEQs	2.6×10^{-5}	2.6×10^{-5}	2.1×10^{-4}	1.4×10^{-4}	6.4×10^{-5}	1.4×10^{-4}

[a] Six exposure scenarios were evaluated; in each of three exposure areas, foraging only in high quality habitats and foraging in both high and poor quality habitats were evaluated. These exposure scenarios are described in detail in Section A.5.1.2.1.

BEHP – bis(2-ethylhexyl) phthalate

bw – body weight

COPC – chemical of potential concern

PCB – polychlorinated biphenyl

TEQ – toxic equivalent

Table A.5-12. Dietary exposure doses of COPCs for great blue heron, osprey, river otter, and harbor seal

COPC	Dietary Exposure Dose (mg/kg bw/day)			
	Great Blue Heron	Osprey	River Otter	Harbor Seal
Arsenic	ne	ne	0.32	ne
Chromium	0.067	0.11	ne	ne
Cobalt	ne	ne	0.016	ne
Lead	0.27	0.26	ne	ne
Mercury	0.0031	0.0051	0.0048	3.3×10^{-4}
Selenium	ne	ne	0.032	ne
Total PCBs	0.17	0.32	0.26	0.020
PCB TEQs	9.3×10^{-6}	2.3×10^{-5}	6.4×10^{-6}	5.1×10^{-7}

bw – body weight

COPC – chemical of potential concern

ne – not evaluated; not a COPC for this ROC

PCB – polychlorinated biphenyl

TEQ – toxic equivalent

A.5.2 Effects Assessment

This section presents a summary of the toxicity literature for each of the COPCs and describes the selection of TRVs for birds and mammals. The literature search and guidelines for TRV selection for wildlife ROCs was described in detail in Section A.2.3.3. Toxicological data presented in this section are assessed in combination with exposure data (presented in Section A.5.1) in the risk characterization (Section A.6.3).

In many of the studies reviewed, the toxicity data were presented as a concentration in the diet rather than as an ingested dose. These dietary concentrations were converted to ingested doses using the toxicity test species-specific FIRs and body weights, as follows:

$$\text{ingested dose} = \frac{(C_{food} \times FIR)}{BW} \qquad \textbf{Equation 5-11}$$

Where:

ingested dose	=	COPCs ingested per day via food (mg COPC/kg body weight/day)
FIR	=	food ingestion rate (kg food dw/day)
C_{food}	=	COPC concentration in dietary items (mg COPC/kg food dw)
BW	=	test species body weight (kg ww)

If the FIR and no-effect or effect concentrations were in different units (i.e., wet weight vs. dry weight), the units of the no-effect or effect concentrations were converted to the units of the FIR, using the moisture content in food as presented in table footnotes in this section. The values for FIR, BW, and food moisture content were obtained from the specific toxicity study, if available.

A.5.2.1 TRVs for birds

This section presents results from laboratory toxicity studies for COPCs identified for bird ROCs (Table A.5-1), and selects TRVs for estimating risks. TRVs for PCBs were selected for both total PCBs (generally based on Aroclors) and for 2,3,7,8-TCDD (as TEQs for dioxin-like PCB congeners). Toxicity data are not sufficient to develop unique TRVs for each of the bird species used as ROCs in this risk assessment. Therefore, a single NOAEL and LOAEL TRV was selected for each COPC and used for all bird ROCs.

A.5.2.1.1 Total PCBs

Effects on birds from exposure to dietary PCBs include disruption of normal patterns of growth, reproduction, metabolism, and behavior (Eisler 1986b). The most sensitive effects are related to reproduction, and include egg production, fertility, and hatching success (Eisler 1986b). Of the bird laboratory species used to examine reproductive endpoints, chickens and other galliformes, such as pheasant and quail, have been found to be the most sensitive to PCB toxicity (Kennedy et al. 1996). Because of concerns with poultry laboratory studies, only data from wildlife laboratory studies were considered in choosing a bird PCB TRV. This approach is consistent with an EPA-sponsored peer review panel charged with reviewing an ERA for the Hudson River. This panel evaluated use of PCB TRVs derived from chicken studies to assess risks to wild birds. Reviewers considered PCB TRVs developed from chicken studies to be "unrealistically low and excessively conservative" and found that "using the chicken as a representative species for wild birds was not defensible" (EPA 2000b). The use of chicken reproductive toxicity data to assess risks to ROCs should be considered protective, but it is not likely to predict risks accurately. Therefore, chicken toxicity data for reproductive endpoints were not used in this ERA.

Thirteen studies were identified that evaluated the dietary toxicity of PCBs to birds (Table A.5-13). Various species were studied, including American kestrels, screech owls, turtle doves, Japanese quail, and mallard ducks. All studies reviewed involved the assessment of reproductive endpoints following dietary exposure to PCBs. These endpoints included fertility, hatchability, eggshell thickness, egg production, eggshell weight, embryo development, clutch size, and embryo mortality and viability.

Table A.5-13. PCB dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)[a]	Effect Conc. (mg/kg ww)[a]	Body Weight (kg)	Food Ingestion Rate[b]	Source	Notes
Aroclor 1248	American kestrel	na	0.35	5.5 months	reduced eggshell thickness	na	3	0.13 (Pattee 1984)	0.0136 kg dw/day, Eurasian kestrel, Nagy (2001)	Lowe and Stendell (1991)	1
Aroclor 1248	screech owl	**0.49**	na	two generations	no effect on eggshell thickness, egg production, or hatching and fledging success	3	na	0.181 (Dunning 1993)	0.0266 kg dw/day, carnivorous birds, Nagy (2001)	McLane and Hughes (1980)	1
Aroclor 1242	Japanese quail	na	0.60	45 days	reduced eggshell thickness	na	10	0.09 (Dunning 1993)	0.0048 kg dw/day, galliformes, Nagy (2001)	Hill et al. (1976)	1
Aroclor 1254	ringed turtle dove	na	**1.4**	two generations	reduced hatching success in second generation	na	10	0.155 (Sample et al. 1996)	0.0202 kg dw/day, all birds, Nagy (2001)	Peakall et al. (1972); Peakall and Peakall (1973)	2
Aroclor 1254	mallard	2.5	na	~ 1 month	no effect on reproductive success	25	na	1.082 (Dunning 1993)	0.1082 kg ww/day (Heinz et al. 1987)	Custer and Heinz (1980)	
Aroclor 1254	mallard	3.9	na	4 months	no effect on egg production or eggshell thickness	39	na	1.082 (Dunning 1993)	0.1082 kg ww/day (Heinz et al. 1987)	Risebrough and Anderson (1975)	
Aroclor mixture	American kestrel	na	7	100 days until eggs hatched	reduced egg laying in second generation (exposed *in ovo*); reduced clutch size and fledgling success	na	na	na	na	Fernie et al. (2000; 2001)	3, 4
Aroclor mixture	American kestrel	na	7	three generations	increased incidence of embryo abnormalities and cracked eggs, reduced F1 offspring survival, reduced offspring growth rate in F2 nestlings	na	na	na	na	Fernie et al. (2003a; 2003b; 2003c)	3, 4

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)[a]	Effect Conc. (mg/kg ww)[a]	Body Weight (kg)	Food Ingestion Rate[b]	Source	Notes
Aroclor 1242	mallard	na	15	12 weeks	reduced hatchability, embryo mortality, and egg viability, and increased incidence of embryo abnormalities	na	150	1.082 (Dunning 1993)	0.1082 kg ww/day (Heinz et al. 1987)	Haseltine and Prouty (1980)	

[a] No-effect and effect concentrations are presented in the units given in the studies reviewed. Table notes indicate how units were converted to wet weight or dry weight to correspond to the FIR units for calculating NOAELs and LOAELs.

[b] Ingestion rates are from equations for bird groups presented in Nagy (2001), from data presented for individual bird species (Nagy 2001), or from other sources as noted.

Notes:

1. Effect and/or no-effect concentration converted into dry weight assuming 10% moisture in prepared diet.
2. Effect concentration converted into dry weight assuming 9% moisture contents of seeds (EPA 1993b).
3. Body weight-normalized dose was estimated in study.
4. Fifty-two percent of first-generation offspring died within 3 days of hatching.

F1 – first generation
F2 – second generation
bw – body weight
dw – dry weight
LOAEL – lowest-observed-adverse-effect level
na – not available or not applicable
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
ww – wet weight
Bold identifies the NOAEL and LOAEL selected as the TRVs.

LOAELs ranged from 0.35 mg/kg bw/day for eggshell thickness in American kestrels (Lowe and Stendell 1991) to 15 mg/kg bw/day for reproduction in mallards (Haseltine and Prouty 1980). The lowest calculated LOAELs (0.35 and 0.60 mg/kg bw/day) were based on eggshell thinning in American kestrels and Japanese quail (Lowe and Stendell 1991; Hill et al. 1976). These LOAELs were not selected as TRVs because the eggshell thinning results were not associated with reduced hatchability. The next lowest TRV (1.4 mg/kg bw/day) was based on an endpoint of reduced hatching success, which was measured in second generation offspring of ringed turtle doves following dietary exposure to Aroclor 1254 (Peakall et al. 1972; Peakall and Peakall 1973). This dose (1.4 mg/kg bw/day) was selected as the LOAEL TRV for total PCBs.

NOAELs ranged from 0.49 mg/kg bw/day, at which no adverse effect on reproduction was reported in screech owls (McLane and Hughes 1980), to 3.9 mg/kg kg/d, at which egg production and eggshell thinning was unaffected in mallards (Risebrough and Anderson 1975). The NOAEL of 0.49 mg/kg bw/day was selected as the NOAEL TRV.

A.5.2.1.2 PCB TEQs

PCB TEQs are expressed as 2,3,7,8-TCDD equivalents; therefore, toxicity studies involving exposing birds to 2,3,7,8-TCDD were reviewed. Effects of dioxins and furans reported in laboratory studies with various species of animals include developmental toxicity, hepatotoxicity, endocrine disruption, immunotoxicity, and death (Kennedy et al. 1996).

No studies in the published literature involving dietary exposure of birds to 2,3,7,8-TCDD were found. Two studies that evaluated the exposure of birds to 2,3,7,8-TCDD were identified. One study exposed birds through IP injection and the other study used oral intubation. Although these studies used less relevant forms of exposure, they were the only studies available, and thus data from these studies are presented in Table A.5-14.

The lowest dose resulting in effects was from a study of reproductive effects by Nosek et al. (1992), which exposed ring-necked pheasants to weekly IP injections of 2,3,7,8-TCDD for 10 weeks. These weekly injections were converted to daily doses. There are significant uncertainties in assuming that effects resulting from this acute high-level exposure would be similar to effects from chronic dietary exposure. The other study (Schwetz et al. 1973) resulted in effects to cockerels exposed through oral intubation at a higher dose (0.001 mg/kg bw/day) than the Nosek et al. (1992) study, but did not involve reproductive endpoints. Therefore, the LOAEL from Nosek et al. (1992) (0.00014 mg/kg bw/day) was selected as the LOAEL TRV. The highest NOAEL below the LOAEL was from the same study with the same endpoints. This dose of 0.000014 mg/kg bw/day was selected as the NOAEL TRV. Uncertainties associated with the absence of relevant toxicological data for chronic exposure of birds to 2,3,7,8-TCDD, as well as uncertainties associated with the PCB TEQ approach, are discussed in Section A.6.3.1.2.

Table A.5-14. 2,3,7,8-TCDD toxicity studies for birds for use in PCB TEQ risk analysis

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)	Effect Conc. (mg/kg ww)	Body Weight (kg)	Food Ingestion Rate	Source	Notes
2,3,7,8-TCDD	ring-necked pheasant	**0.000014**	**0.00014**	once per week for 10 weeks	reduced body weight, egg production, and survival of adults and embryos	na	na	na	na	Nosek et al. (1992)	1
2,3,7,8-TCDD	cockerel	0.0001	0.001	20 to 21 days	decreased survival	na	na	na	na	Schwetz et al. (1973)	2

Notes:

1. Exposure route was through IP injection
2. Exposure route was through oral intubation.

bw – body weight

LOAEL – lowest-observed-adverse-effect level

na –not applicable; exposure dose was presented in the study

NOAEL – no-observed-adverse-effect level

PCB – polychlorinated biphenyl

TCDD –tetrachlorodibenzo-*p*-dioxin

TEQ – toxic equivalent

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

A.5.2.1.3 Arsenic

Chronic dietary exposure to arsenic has been shown to cause growth and reproductive effects in birds based on three studies identified that evaluated the toxicity of dietary arsenic to mallards (Stanley et al. 1994; USFWS 1964) (Table A.5-15). All three of these studies exposed the mallards to sodium arsenate in their diet. Although arsenite is known to be more highly toxic in general than arsenate (Eisler 1988a), there are no available data on the reproductive toxicity of arsenite to birds, resulting in uncertainty in the arsenic TRV.

Adverse effects were reported in two of the studies presented in Table A.5-15. Reproduction was affected in a study that dosed both parents at 40 mg/kg bw/day (Stanley et al. 1994), and the survival of young mallards was affected when they were dosed at a rate of 50 mg/kg bw/day (USFWS 1964). Adverse effects reported in the study with the lowest LOAEL (40 mg/kg bw/day) included delayed onset of egg laying, decreased offspring body weight, decreased egg production, and decreased eggshell thickness. This dietary dose (40 mg/kg bw/day) was selected as the LOAEL TRV.

NOAELs ranged from 6.1 to 28 mg/kg bw/day. Mallard growth, survival, or reproduction were unaffected at these doses (Camardese et al. 1990; USFWS 1964; Stanley et al. 1994). The highest NOAEL below the LOAEL with the same endpoint, 10 mg/kg bw/day, was from the same study as the selected LOAEL TRV (Stanley et al. 1994), and was selected as the NOAEL TRV.

A.5.2.1.4 Cadmium

Chronic dietary exposure to cadmium has been reported to cause growth and reproductive effects in birds as well as histopathological effects on kidneys and testes (White and Finley 1978b). Eight studies that evaluated the toxicity of dietary cadmium to birds with growth, reproduction, or survival endpoints were identified (Table A.5-16). Dietary exposure to cadmium resulted in adverse effects on growth or reproduction in chickens, mallards, and Japanese quail.

LOAELs ranged from 2.9 mg/kg bw/day for eggshell thinning in chickens (Leach et al. 1979) to 47 mg/kg bw/day for reduced growth in mallards (DiGiulio and Scanlon 1984). At the lowest LOAEL, eggshell thickness was significantly reduced in chickens fed 2.9 mg/kg bw/day cadmium sulfate for 48 weeks (Leach et al. 1979), but decreased reproductive success was not reported, so this LOAEL was not selected as the TRV. The next lowest LOAEL (4.0 mg/kg bw/day) was associated with a reduction in body weight in male chicks following a 6-week exposure (Richardson et al. 1974). This LOAEL was selected as the TRV. The NOAEL of 1.5 mg/kg bw/day from Cain et al. (1983) was selected as the NOAEL TRV because it was lower than the selected LOAEL TRV and was based on the same growth endpoint.

Table A.5-15. Arsenic dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)	Effect Conc. (mg/kg ww)	Body Weight (kg)	Food Ingestion Rate (kg ww/day)	Source	Notes
Sodium arsenate	mallard (young)	6.1	na	10 weeks	no effect on female growth or survival	300	na	0.6305	0.0128	Camardese et al. (1990)	1
Sodium arsenate	mallard	**10**	**40**	115 to 128 days	delayed egg laying; depressed egg weight, production, and shell thinning; decreased offspring body weight	100	400	1.082 (Dunning 1993)	0.1082 (Heinz et al. 1987)	Stanley et al. (1994)	2
Sodium arsenite	mallard (young)	25	50	154 days	reduced survival	250	500	1.082 (Dunning 1993)	0.1082 (Heinz et al. 1987)	USFWS (1964)	3

Notes:

1. No-effect concentration was converted into dry weight using 13.1% moisture as reported in study. Body weight and FIR were reported in study.
2. Effect and no-effect concentrations were converted into dry weight assuming 10% moisture in prepared diet.
3. Literature values for body weight and FIR for adult mallard were used because average values for the birds used in the study (1 day old at the start of the 154-day study) were not available. Based on body weight data for control birds presented in Camardese et al. (1990), birds reach the adult weight of 1.082 kg (Dunning 1993) at approximately 50 days of age.
4. Sample et al. (1996) selected 250 mg/kg ww as the effect concentration from the USFWS study. However, mortality at this concentration (12%) was similar to or less than mortality in the controls (8-31%). Therefore, in this assessment, the concentration of 500 mg/kg ww with 60% mortality was selected as the effect concentration.

bw – body weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

Table A.5-16. Cadmium dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)	Effect Conc.	Body Weight (kg)	Food Ingestion Rate[a]	Source	Notes
Cadmium chloride	mallard (young females)	**1.5**	na	12 weeks	no effect on body weight	14.6	na	0.8825	0.883 kg ww/day (Heinz et al. 1987)	Cain et al. (1983)	1
Cadmium sulfate	chicken (hens)	0.73	2.9	48 weeks	reduced shell thickness	12.22	48.22 mg/kg ww	0.1019 (NRC 1984)	1.7 kg ww/day (NRC 1984)	Leach et al. (1979)	
Cadmium chloride	Japanese quail (chicks)	na	**4.0**	6 weeks	reduced male body weight	na	75 mg/kg dw	0.093	0.0050 kg dw/day, galliformes (Nagy 2001)	Richardson et al. (1974)	1
Cadmium chloride	mallard	19	na	90 days	no effect on survival	200	na	1.153	0.110 kg ww/day	White and Finley (1978a)	2
Cadmium chloride	mallard	20	na	30 to 90 days	no effect on body weight, adult mortality	210	na	1.153	0.110 kg ww/day	White and Finley (1978b)	2
Cadmium chloride	chicken (chicks)	na	24	21 days	reduced male body weight	na	75 mg/kg dw	0.138	0.0434 kg dw/day (NRC 1994)	Freeland and Cousins (1973)	1
Cadmium chloride	chicken (chicks)	na	40	20 days	reduced male body weight	na	400 mg/kg dw	0.574 (NRC 1994)	0.0577 kg dw/day (NRC 1994)	Pritzl et al. (1974)	
Cadmium chloride	mallard	16	47	42 days	reduced body weight	150	450 mg/kg ww	1.119 (NOAEL), 1.027 (LOAEL)	0.122 (NOAEL), 0.108 (LOAEL) kg ww/day	DiGiulio and Scanlon (1984)	2

[a] Ingestion rates are from equations for bird groups presented in Nagy (2001), from data presented for individual bird species (Nagy 2001), or from other sources as noted.

Notes:

1. Body weight was reported in study.
2. Body weight and FIR were reported in study.

bw – body weight
dw – dry weight
LOAEL – lowest-observed-adverse-effect level
na – not available or not applicable
NOAEL – no-observed-adverse-effect level
ww – wet weight
Bold identifies the NOAEL and LOAEL selected as the TRVs.

A.5.2.1.5 Chromium

In its hexavalent form, chromium is mutagenic, carcinogenic, and teratogenic to a wide variety of organisms under laboratory conditions (Eisler 1986a). Fewer data are available involving the effects of trivalent chromium on birds. Three studies that evaluated the toxicity of dietary chromium to birds were identified (Table A.5-17). Adverse effects were observed in one of the three studies. In that study, Haseltine et al. (unpublished) reported a decrease in offspring survival of adult black ducks treated with 5.0 mg/kg bw/day trivalent chromium over a chronic period (10 months) and critical lifestage (reproduction). This LOAEL (5.0 mg/kg bw/day) was selected as the LOAEL TRV for chromium. The results of this unpublished study were reported in Sample et al. (1996), and the original study could not be obtained. Therefore, there is some uncertainty regarding the quality of the study. The highest NOAEL below the LOAEL with the same endpoint was from the same study. This NOAEL of 1.0 mg/kg bw/day was selected as the NOAEL TRV.

A.5.2.1.6 Cobalt

One study that evaluated the toxicity of dietary cobalt to birds was identified (Table A.5-18). In this study (Diaz et al. 1994), body weights of chickens were significantly less than the control group following 2 weeks of exposure to 116, 251, or 472 mg/kg cobalt in the diet. The lowest dose of 116 mg/kg was used to calculate the LOAEL of 23.1 mg/kg bw/day. There was no lower dose with which to calculate a no-effect dose, so the subchronic LOAEL was divided by an uncertainty factor of 10 to estimate the NOAEL TRV of 2.31 mg/kg dw/day.

Table A.5-17. Chromium dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)[a]	Effect Conc. (mg/kg ww)[a]	Body Weight (kg)	Food Ingestion Rate (kg ww/day)	Source	Notes
Chromium picolinate	chicken (hens)	0.10	na	28 days	no effect on egg weight or shell thickness	1.6	na	0.1019 (NRC 1984)	1.7 (NRC 1984)	Lien et al. (2004)	
Chromium potassium sulfate	black duck	**1.0**	**5.0**	10 months	reduced duckling survival	10	50	1.25 (Dunning 1993)	0.125 (Heinz et al. 1987)	Haseltine et al. (unpublished) as cited in Sample et al. (1996)	1
Sodium chromate	chicken (chicks)	7.7	na	22 days	no effect on adult male survival or male body weight	32.1	na	0.12 (NRC 1994)	0.0286 (NRC 1994)	Romoser et al. (1961)	

[a] No-effect and effect concentrations are presented in the units given in the studies reviewed.

Notes:

1. Original paper (unpublished study) could not be obtained.

bw – body weight
LOAEL – lowest-observed-adverse-effect level
na – not available or not applicable
NOAEL – no-observed-adverse-effect level
ww – wet weight
Bold identifies the NOAEL and LOAEL selected as the TRVs.

Table A.5-18. Cobalt dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc.	Effect Conc. (mg/kg ww)	Body Weight (kg)	Food Ingestion Rate (kg ww/day)	Source	Notes
Cobalt chloride	chicken (chicks)	na	**23.1**	14 days	reduced body weight	na	125	0.1462	0.027	Diaz et al. (1994)	1

Notes:

1. Body weight and FIR reported in study.

bw – body weight
LOAEL – lowest-observed-adverse-effect level
na – not available or not applicable
NOAEL – no-observed-adverse-effect level
ww – wet weight
Bold identifies the LOAEL selected as the TRV. A NOAEL TRV was estimated by dividing the subchronic LOAEL TRV by an uncertainty factor of 10. The resulting NOAEL TRV was 2.31 mg/kg bw/day.

A.5.2.1.7 Copper

No data are available to assess the potential effects of chronic dietary exposure of copper to wild birds (Eisler 1997). Seven studies that evaluated the toxicity of dietary copper to chickens were available (Table A.5-19). In studies with chicks, impaired growth was reported at doses ranging from 29 to 66 mg/kg bw/day dietary copper in various forms (copper sulfate, copper chloride, and copper oxide) for 2 to 10 weeks. The lowest dose (29 mg/kg bw/day) was selected as the LOAEL TRV, although there is uncertainty in this growth effect because of the subchronic exposure period of 25 days. The next lowest LOAEL (62 mg/kg bw/day) resulted in a reduction in growth and survival in chicks after exposure for 10 weeks (Mehring et al. 1960).

NOAELs ranged from 2.1 mg/kg bw/day, at which no effect on chick growth or survival was reported (Dozier et al. 2003), to 47 mg/kg bw/day, at which chick growth was not affected (Mehring et al. 1960). The NOAEL of 21 mg/kg bw/day (Poupoulis and Jensen 1976) was selected as the NOAEL TRV for the subchronic LOAEL, because it was the highest NOAEL that was lower than the selected LOAEL TRV based on the same growth endpoint.

A.5.2.1.8 Lead

The acute effects of lead poisoning as a result of lead shot ingestion by birds have been extensively studied, although this exposure route is not relevant to the LDW. Numerous effects have been reported, including mortality, damage to the nervous system, muscular paralysis, kidney and liver damage, internal lesions, enlarged gall bladder, anemia, reduced brain weight, and abnormal skeletal development (Eisler 1988b). Fewer studies have been conducted on chronic effects of dietary exposure to lead.

Four studies that evaluated the chronic toxicity of dietary lead to birds were identified (Table A.5-20). Adverse effects were reported in two of the four studies. One of the studies (Pattee 1984) used metallic lead powder, which is a form of lead that is not likely found in the LDW; adverse effects were not reported in this study. Edens et al. (1976) reported a significant decrease in egg hatchability of Japanese quail exposed to 20 mg/kg bw/day lead over a chronic period (12 weeks) and critical lifestage (reproduction). Japanese quail chicks fed 28 mg/kg bw/day lead for 6 weeks experienced a reduction in body weight (Morgan et al. 1975). The lowest LOAEL of 20 mg/kg bw/day was selected as the TRV.

NOAELs ranged from 2.0 mg/kg bw/day, at which egg hatchability was unaffected in Japanese quail (Edens et al. 1976), to 5.82 mg/kg bw/day, at which no effect on survival or reproduction was reported in American kestrels (Pattee 1984). The NOAEL of 5.82 mg/kg bw/day was selected as the TRV because it was the highest NOAEL below the selected LOAEL based on a reproductive endpoint.

Table A.5-19. Copper dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc.[a]	Effect Conc.[a]	Body Weight (kg)	Food Ingestion Rate	Source	Notes
Copper sulfate, copper amino acid complex	chicken (chicks)	2.1	na	17 days	no effect on body weight or survival	18 mg/kg dw	na	0.254 (NRC 1994)	0.0295 kg dw/day (NRC 1994)	Dozier et al. (2003)	
Copper sulfate	chicken (hens)	11.2	na	90 days	no effect on damaged egg ratio, egg weight, or adult survival	200 mg/kg ww	na	0.09585	1.71 kg ww/day (NRC 1984)	Balevi and Coskun (2004)	1
Copper sulfate	chicken (hens)	15	na	28 days	no effect on egg weight and shell thickness	250 mg/kg ww	na	0.1019 (NRC 1984)	1.70 kg ww/day (NRC 1984)	Lien et al. (2004)	
Copper sulfate	chicken (1 day old)	16	29	25 days	reduced growth	200 mg/kg ww	350 mg/kg ww	0.534 (EPA 1993)	0.044 kg ww/day (EPA 1993)	Smith (1969)	
Copper sulfate	chicken (chicks)	**21**	41	4 weeks	reduced growth	250 mg/kg ww	500 mg/kg ww	0.534 (EPA 1993)	0.044 kg ww/day (EPA 1993)	Poupoulis and Jensen (1976)	
Copper oxide	chicken (chicks)	47	**62**	10 weeks	reduced growth and survival	570 mg/kg ww	749 mg/kg ww	0.534 (EPA 1993)	0.044 kg ww/day (EPA 1993)	Mehring et al. (1960)	
Copper chloride	chicken (chicks)	na	66	8 to 22 days	reduced body weight	na	500 mg/kg dw	0.3845	0.0509 kg dw/day	Persia et al. (2004)	2

[a] No-effect and effect concentrations are presented in the units given in the studies reviewed.

Notes:

1. Body weight reported in study.
2. Body weight and FIR reported in study.

bw – body weight

dw – dry weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

Table A.5-20. Lead dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc.[a]	Effect Conc.[a]	Body Weight (kg)	Food Ingestion Rate[b]	Source	Notes
Lead nitrate	mallard (first year)	2.5	na	12 weeks	no effect on survival	25 mg/kg ww	na	1.082 (Dunning 1993)	0.1082 kg ww/day (Heinz et al. 1987)	Finley et al. (1976)	
Metallic lead powder	American kestrel	**5.82**	na	5 to 7 months	no effect on survival, fertility, egg production, or eggshell thickness	50 mg/kg ww	na	0.13	0.0136 kg dw/day, Eurasian kestrel, Nagy (2001)	Pattee (1984)	1, 2
Lead acetate	Japanese quail	2.0	**20**	12 weeks	reduced egg hatchability	10 mg/kg ww	100 mg/kg ww	0.155 (Edens and Garlich 1983)	0.031 kg ww/day (Edens and Garlich 1983)	Edens et al. (1976)	
Lead acetate	Japanese quail (chicks)	5.5	28	6 weeks	reduced body weight	100 mg/kg dw	500 mg/kg dw	0.0715	0.0040 kg dw/day, galliformes, Nagy (2001)	Morgan et al. (1975)	2

[a] No-effect and effect concentrations are presented in the units given in the studies reviewed. Table notes indicate how units were converted to wet weight or dry weight to correspond to the FIR units for calculating NOAELs and LOAELs.

[b] Ingestion rates are from equations for bird groups presented in Nagy (2001), from data presented for individual bird species (Nagy 2001), or from other sources as noted

Notes:

1. No-effect concentration converted into dry weight assuming 10% moisture in prepared diet.
2. Body weight reported in study.

bw – body weight

dw – dry weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

A.5.2.1.9 Mercury

Chronic effects of dietary mercury on birds include adverse effects on growth, development, reproduction, metabolism, and behavior (Eisler 1987). Six studies that evaluated the toxicity of dietary mercury to birds were identified (Table A.5-21). When reviewing the toxicity literature for mercury, only forms of mercury relevant to the LDW were considered. Acceptable forms included inorganic mercury salts, such as mercuric chloride, as well as organic forms, such as methyl mercury chloride and dimethylmercury. Mercury-containing fungicides (e.g., Ceresan, methyl mercury dicyandiamide) were not considered relevant because these forms of mercury are not expected to occur in the LDW. The toxicity of these fungicidal formulations is likely highly influenced by the attached anions that are intended to enhance the toxicity of the fungicide because of the additive effects of these non-mercury components. As a result, laboratory bird studies involving mercury-containing fungicides were not considered for TRV selection.

In the studies reviewed, adverse effects on reproduction, early-life-stage growth, or adult survival were reported for various bird species, including great egrets, Japanese quail, zebra finch, and bobwhite quail from dietary exposure to mercury. LOAELs ranged from 0.091 mg/kg bw/day for reduced growth in young great egrets (Heinz 1980) to 62 mg/kg bw/day for offspring mortality of Japanese quail (Hill and Soares 1987). The lowest LOAEL of 0.091 mg/kg bw/day mercury was selected as the TRV.

NOAELs ranged from 0.43 mg/kg bw/day, at which there was no effect on survival of young bobwhite quail (Spann et al. 1986), to 5.24 mg/kg bw/day, at which there was no effect on eggshell thickness in American kestrels (Peakall and Lincer 1972). None of these NOAELs were lower than the lowest LOAEL. Therefore, the chronic LOAEL was divided by an uncertainty factor of 5 to obtain the NOAEL TRV of 0.018 mg/kg bw/day.

A.5.2.1.10 Nickel

Three studies that evaluated the effects of dietary nickel to birds were identified (Table A.5-22). Adverse effects were reported in two studies, ranging from a dose of 33 mg/kg bw/day that affected growth of broiler chicks (Weber and Reid 1968) to a dose of 107 mg/kg bw/day that affected mallard growth and survival (Cain and Pafford 1981). The lowest LOAEL of 33 mg/kg bw/day was selected as the TRV, although there is uncertainty in this growth effect because of the subchronic exposure period of only 4 weeks. The other LOAEL (107 mg/kg bw/day) resulted in a reduction in growth and survival in mallards after exposure for 90 days (Cain and Pafford 1981).

The NOAEL of 77 mg/kg bw/day from Cain and Pafford (1981) was selected as the NOAEL TRV. It was selected because it is the highest NOAEL that was lower than the selected LOAEL TRV and was from the same study based on the same endpoint.

Table A.5-21. Mercury dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc.[a]	Effect Conc.[a]	Body Weight (kg)	Food Ingestion Rate[b]	Source	Notes
Methyl-mercury chloride	great egret (1 day old)	na	**0.091**	14 weeks	reduced growth	na	0.5 mg/kg ww	1.02 (Fish 2002)	0.185 kg ww/day (Kushlan 1978)	Spalding et al. (2000)	
Methyl-mercury chloride	mallard	0.50	na	> 60 days	no effect on eggshell thickness	5 mg/kg ww	na	1.082 (Dunning 1993)	0.1082 kg ww/day (Heinz et al. 1987)	Heinz (1980)	
Methyl-mercury chloride	Japanese quail (chicks)	na	0.9	5 days	reduced hatchling survival of offspring	na	16 mg/kg ww	0.1 (NRC 1994)	0.0053 kg dw/day, galliformes (Nagy 2001)	Hill and Soares (1987)	1
Methyl-mercury chloride	zebra finch	0.72	1.4	76 days	reduced survival	2.5 mg/kg dw	5 mg/kg dw	0.012 (Dunning 1993)	0.0034 kg dw/day, passerines (Nagy 2001)	Scheuhammer (1988)	2
Methyl-mercury chloride	northern bobwhite quail (12 days old)	0.43	1.6	6 weeks	reduced survival	5.4 mg/kg ww	20 mg/kg ww	0.19 (EPA 1993)	0.0150 kg ww/day (EPA 1993)	Spann et al. (1986)	
Mercuric chloride	Japanese quail (1 day old)	0.80	1.6	10 weeks	reduced eggshell thickness	4 mg/kg ww	8 mg/kg ww	0.155 (Edens and Garlich 1983)	0.031 kg ww/day (Edens and Garlich 1983)	Stoewsand et al. (1971)	
Dimethyl mercury	American kestrel	5.24	na	3 months	no effect on eggshell thickness	10 mg/kg ww	na	0.13 (Pattee 1984)	0.0136 kg dw/day, Eurasian kestrel (Nagy 2001)	Peakall and Lincer (1972)	3
Mercuric chloride	Japanese quail (chicks)	na	62	5 days	reduced offspring hatchling survival	na	1,045 mg/kg ww	0.1 (NRC 1994)	0.0053 kg dw/day, galliformes (Nagy 2001)	Hill and Soares (1987)	1

[a] No-effect and effect concentrations are presented in the units given in the studies reviewed. Table notes indicate how units were converted to wet weight or dry weight to correspond to the FIR units for calculating NOAELs and LOAELs.

[b] Ingestion rates are from equations for bird groups presented in Nagy (2001), from data presented for individual bird species (Nagy 2001), or from other sources as noted

Notes:

1. Effect concentration converted into dry weight assuming 10% moisture in prepared diet.
2. No-effect and effect concentration converted into wet weight assuming 10% moisture in prepared diet.
3. Study did not indicate whether the mercury concentration in the diet, which consisted of dead chicks, was reported in wet weight or dry weight. It was assumed to be reported in wet weight and was converted into dry weight using 80% moisture content.

bw – body weight

dw – dry weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the LOAEL selected as the TRV. A NOAEL TRV was estimated by dividing the chronic LOAEL TRV by an uncertainty factor of 5. The resulting NOAEL TRV was 0.018 mg/kg bw/day.

Table A.5-22. Nickel dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)[a]	Effect Conc. (mg/kg ww)[a]	Body Weight (kg)	Food Ingestion Rate (kg ww/day)	Source	Notes
Nickel sulfate	chicken (chicks)	15	na	4 weeks	no effect on body weight gain	na	111.7	0.484	0.067	Weber and Reid (1968)	1, 2
Nickel sulfate	chicken (chicks)	**17**	**33**	4 weeks	reduced body weight	117	156	0.467 (NOAEL); 0.39 (LOAEL)	0.069 (NOAEL); 0.083 (LOAEL)	Weber and Reid (1968)	1, 2, 3
Nickel acetate	chicken (chicks)	na	38	4 weeks	reduced body weight	na	165	0.376	0.086	Weber and Reid (1968)	1, 2
Nickel sulfate	mallard	77	107	90 days	reduced body weight and survival	774	1,069	0.561 (NOAEL); 0.178 (LOAEL)	0.0178 (NOAEL); 0.0561 (LOAEL) (Heinz et al. 1987)	Cain and Pafford (1981)	1, 3
Nickel sulfate	mallard	132	na	90 days	no effect on adult survival, body weight, or offspring hatchling weight	800	na	1.082 (Dunning 1993)	0.178	Eastin and O'Shea (1981)	2

[a] No-effect and effect concentrations are presented in the units given in the studies reviewed.

Notes:

1. Body weight was reported in study.
2. FIR was reported in study.
3. Body weight and ingestion rates used were specific to the no-effect and effect concentration test groups.

bw – body weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

A.5.2.1.11 Selenium

Selenium is an essential nutrient and deficiency in the diet may cause adverse effects in birds; elevated dietary concentrations have also been observed to cause adverse effects (Eisler 1985). Five studies that evaluated the dietary toxicity of selenium to birds were identified (Table A.5-23). Adverse effects were reported on offspring survival and growth, embryo development, adult growth, reproductive success, and adult survival. LOAELs ranged from 0.82 mg/kg bw/day, which affected growth and reproduction in mallards (Heinz et al. 1989), to 10 mg/kg bw/day, which affected survival in mallards (Heinz et al. 1988, 1989). In the study reporting the lowest LOAEL, offspring survival and growth was significantly affected for mallards fed 0.82 mg/kg bw/day selenium, as selenomethionine, for approximately 100 days (Heinz et al. 1989). This dose was selected as the LOAEL TRV.

NOAELs ranged from 0.025 mg/kg bw/day, which did not affect survival and growth in chicks (Choct et al. 2004), to 4.6 mg/kg bw/day, which did not affect survival in mallards (Heinz et al. 1988). The highest NOAEL that was lower than the selected LOAEL TRV and based on a reproductive endpoint was 0.5 mg/kg bw/day. This dose was selected as the NOAEL TRV.

A.5.2.1.12 Vanadium

Two studies that evaluated the toxicity of dietary vanadium to birds were identified (Table A.5-24). One of these studies reported reduced body weight in chickens exposed to 2.3 mg/kg bw/day vanadium in the form of ammonium metavanadate in the diet for 4 weeks (Ousterhout and Berg 1981). This dose was selected as the LOAEL TRV because this was the only LOAEL reported. The NOAEL from the same study and endpoint was selected as the NOAEL TRV (1.2 mg/kg bw/day). The only other dietary study identified was conducted with mallards for a longer period of 12 weeks using a different form of vanadium (vanadium sulfate); no effects were reported in that study at a dose of 11.4 mg/kg bw/day.

Table A.5-23. Selenium dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc.	Effect Conc. (mg/kg ww)	Body Weight (kg)	Food Ingestion Rate	Source	Notes
Sodium selenite or sel-plex 50	chicken (chicks)	0.025	na	~40 days	no effect on body weight or survival	0.25 mg/kg dw	na	0.993	0.0995 (NOAEL) kg dw/day	Choct et al. (2004)	1, 2
Seleno-methionine	mallard	0.42	**0.82**	~100 days	reduced offspring growth and survival	4.15 mg/kg ww	8.15	1.158 (NOAEL); 1.145 (LOAEL)	0.1158 (NOAEL); 0.1145 (LOAEL) kg ww/day (Heinz et al. 1987)	Heinz et al. (1989)	1, 3
Sodium selenite	mallard	**0.50**	1.0	4 weeks before laying to 3 weeks after hatching	increased embryo abnormalities	5 mg/kg ww	10	1.036 (NOAEL); 1.046 (LOAEL)	0.105 kg ww/day	Heinz et al. (1987)	1, 2, 3
Sodium selenite	mallard	1.0	2.5	4 weeks before laying to 3 weeks after hatching	reduced adult growth	10 mg/kg ww	25	1.046 (NOAEL); 0.938 (LOAEL)	0.105 (NOAEL); 0.094 (LOAEL) kg ww/day	Heinz et al. (1987)	1, 2, 3
Seleno-methionine	mallard	1.6	na	~100 days	no effect on body weight or adult survival	16.15 mg/kg ww	na	1.107	0.1107 kg ww/day (Heinz et al. 1987)	Heinz et al. (1989)	1, 3
Seleno-methionine	screech owl	1.0	3.2	~ 3 months	reduced body weight, clutch size, hatching success, offspring survival, egg size, and mass	3.7 mg/kg ww	12.1	0.189	0.050 kg ww/day	Wiemeyer and Hoffman (1996)	1, 2, 3
Sodium selenite	mallard	4.6	na	42 days	no effect on survival	20.1 mg/kg ww	na	0.326	0.075 kg ww/day	Heinz et al. (1988)	1, 2
Sodium selenite	mallard	2.1	4.6	42 days	reduced body weight	10.1 mg/kg ww	20.1	0.326	0.075 kg ww/day	Heinz et al. (1988)	1, 2, 3
Sodium selenite	mallard	2.5	10	4 weeks before laying to 3 weeks after hatching	reduced adult survival	25 mg/kg ww	100	0.718	0.072 kg ww/day	Heinz et al. (1987)	1, 2
Sodium selenite	mallard	na	10	42 days	reduced adult survival	na	40.1	0.146	0.038 kg ww/day	Heinz et al. (1988)	1, 2

Notes:

1. Body weight was reported in study.
2. FIR was reported in study.
3. Body weight and FIRs used were specific to the no-effect and effect concentration test group.

bw – body weight

dw – dry weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

Table A.5-24. Vanadium dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)	Effect Conc. (mg/kg ww)	Body Weight (kg)	Food Ingestion Rate (kg ww/day)	Source	Notes
Ammonium metavanadate	chicken (hens)	**1.2**	**2.3**	4 weeks	reduced body weight	20	40	1.71 (Dunning 1993)	0.0997 (NRC 1984)	Ousterhout and Berg (1981)	
Vanadium sulfate	mallard	11.4	na	12 weeks	no effect on body weight or survival	110	na	1.17	0.121	White and Dieter (1978)	1

Notes:

1. Body weight and FIR provided in study.

bw – body weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

A.5.2.1.13 Zinc

Most laboratory studies investigating zinc toxicity have been conducted with domestic birds such as chickens; only one wildlife study, using mallards, was found in the literature. Reported effects of zinc exposure on ducks and chickens in laboratory studies included reduced food intake and egg production, cessation of egg laying, weight loss, leg paralysis, pancreatic histopathology, and mortality (Eisler 1993). Adverse effects may also be observed if zinc is deficient in the diet because zinc is a nutrient essential for normal growth, development, and function. Effects from zinc deficiency are generally noted at concentrations below 120 mg/kg ww in food (Eisler 1993).

Six studies that evaluated the toxicity of dietary zinc to birds were identified (Table A.5-25). Adverse effects on growth and survival in young birds (mallards or chickens) were reported in four of these studies. LOAELs ranged from 124 mg/kg bw/day, which resulted in reduced growth of young chickens (Roberson and Schaible 1960), to 659 mg/kg bw/day, which resulted in mortality and reduced growth of young broiler chickens (Oh et al. 1979). The lowest LOAEL of 124 mg/kg bw/day was selected as the TRV. The NOAEL of 82 mg/kg bw/day from the same study was selected as the NOAEL TRV because it was the highest NOAEL below the LOAEL based on a growth endpoint.

Table A.5-25. Zinc dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)	Effect Conc.	Body Weight (kg)	Food Ingestion Rate	Source	Notes
Zinc sulfate or copper amino acid complex	chicken (chicks)	17	na	17 days	no effect on body weight or survival	150	na	0.254 (NRC 1994)	0.0295 kg ww/day (NRC 1994)	Dozier et al. (2003)	1
Zinc oxide, zinc sulfate, or zinc carbonate	chicken (chicks)	**82**	**124**	5 weeks	reduced growth	1,000	1,500 mg/kg ww	0.534 (EPA 1993)	0.044 kg ww/day (EPA 1993)	Roberson and Schaible (1960)	
Zinc sulfate	chicken (hens)	133	na	44 weeks	no effect on egg hatchability	2,028	na	1.9	0.125 kg ww/day	Stahl et al. (1990)	2
Zinc carbonate	mallard (7 weeks old)	na	300	60 days	reduced survival	na	3,000 mg/kg ww	1.082 (Dunning 1993)	0.1082 kg ww/day (Heinz et al. 1987)	Gasaway and Buss (1972)	
Zinc acetate	chicken (chicks)	330	659	5 weeks	reduced growth and survival	4,000	8,000 mg/kg ww	0.534 (EPA 1993)	0.044 kg ww/day (EPA 1993)	Oh et al. (1979)	
Zinc chloride	chicken (chicks)	na	344	8 to 22 days	reduced body weight	na	2,500 mg/kg dw	0.2873	0.0395 kg dw/day	Persia et al. (2004)	2

Notes:

1. NOAEL is based on background concentration in prepared food (30 mg/kg) plus exposure concentration added to food (120 mg/kg).
2. Body weight and FIR provided in study.

bw – body weight

dw – dry weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

A.5.2.2 TRVs for mammals

This section presents results from laboratory toxicity studies for COPCs identified for river otter and harbor seal (Table A.5-1), and selects TRVs for estimating risks. TRVs for PCBs were selected for both total PCBs (generally based on Aroclors) and for 2,3,7,8-TCDD (for comparison to PCB TEQs to assess effects from dioxin-like PCB congeners).

A.5.2.2.1 Total PCBs

PCBs have been reported to elicit a broad range of toxic effects in laboratory mammals under controlled exposure conditions, including lethality, hepatotoxicity, porphyria, body weight loss, dermal toxicity, thymic atrophy, immunosuppressive effects, reproductive and developmental effects, carcinogenesis, and neurotoxicity (Safe 1992, 1991, 1984; Seegal 1996; Safe 1990, 1994; Kimbrough 1985, 1987; Silberhorn et al. 1990; WHO 1993; Bolger 1993; Battershill 1994; Delzell et al. 1994). Review of the toxicology literature indicates that the potency of PCB mixtures depends on the chlorine content of the mixture and, in general, mixtures with higher chlorine content (i.e., Aroclors 1242, 1248, 1254, and 1260) are more toxic than mixtures with lower chlorine content (i.e., Aroclors 1221 and 1232). In general, the gastrointestinal tract of most mammals readily absorbs PCBs, but the absorption rate may be affected by the dose level and lipophilicity of the compound (Eisler 1986b; Van den Berg et al. 1998). There is evidence for placental transfer of PCBs in mammals (Eisler 1986b), and PCBs can also accumulate in the lipid portion of milk, resulting in exposure to suckling young.

Adverse reproductive effects (e.g., fertility, litter size, offspring survival) appear to be among the most sensitive *in vivo* endpoints of PCB toxicity in mammals (Golub et al. 1991; Rice and O'Keefe 1995; Hoffman et al. 1996). Reproductive success can be affected directly by toxic action on the differentiated reproductive tract or indirectly on systems that regulate reproduction (e.g., endocrine and central nervous systems). In laboratory studies, PCBs have been reported to elicit a broad range of direct and indirect effects associated with reproductive functions. Direct effects on the gonads and the female reproductive tract have been reported (Fuller and Hobson 1986). The precise mechanism by which PCBs cause reproductive effects in mammals remains unclear, but reproductive success appears to be a sensitive integrated endpoint of *in vivo* toxicity.

The most comprehensive studies of PCB toxicity in a non-domesticated mammal have been conducted with mink. Mink also appears to be one of the most sensitive mammalian species tested (Fuller and Hobson 1986), and is therefore a good surrogate for the assessment of risk to other mammals. Thus, only mink studies were reviewed for the development of PCB TRVs. Monkeys are also sensitive to PCBs, with reproductive effects reported at approximately 0.1 mg/kg bw/day (Allen et al. 1980; Barsotti et al. 1976; Truelove et al. 1982). However, data from mink studies were used instead because of their greater taxonomic similarity to river otter and harbor seal.

Ten studies that evaluated the toxicity of dietary PCBs to mink were identified (Table A.5-26). In the studies reviewed, adverse effects on maternal growth, kit growth, kit survival, whelping success, and reproductive success were reported for captive-bred mink following dietary exposure to PCBs. Reported reproductive effect levels ranged from 0.089 mg/kg bw/day (Brunström et al. 2001) to 2.6 mg/kg bw/day (Bleavins et al. 1980). At the lowest dose, offspring growth was significantly reduced in mink fed 0.089 mg/kg bw/day of a Clophen A50 PCB mixture for 18 months compared to mink in the control group (Brunström et al. 2001). This dose was selected as the LOAEL TRV.

NOAELs ranged from 0.13 mg/kg bw/day, which had no effect on reproduction in mink (Aulerich and Ringer 1977), to 1.5 mg/kg bw/day, which had no effect on growth in mink (Aulerich et al. 1986). There was no NOAEL that was lower than the selected LOAEL TRV in any of the studies reviewed, so a NOAEL TRV was estimated from the selected chronic LOAEL using an uncertainty factor of 2. This uncertainty factor was selected rather than an uncertainty factor of 5 used for other ROC/COPC pairs because of the large toxicity dataset for mink and PCBs that indicates that an uncertainty factor of 5 is high. As shown in Table A.5-26, the LOAELs are higher than the NOAELs from the same studies by factors ranging from 1.5 to 2 in the two studies that had both NOAELs and LOAELs (Aulerich and Ringer 1977; Aulerich et al. 1986). In addition, dose-response plots of toxicity to mink exposed to PCBs show very steep transitions between PCB exposures causing no adverse effects and those resulting in severe adversity (EPA 2003), indicating that an uncertainty factor of 2 is more appropriate than 5. The resulting NOAEL was 0.045 mg/kg bw/day.

Table A.5-26. PCB dietary toxicity studies for mammals

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)	Effect Conc.	Body Weight (kg)	Food Ingestion Rate (kg ww/day)	Source	Notes
Clophen A50	mink	na	**0.089**	18 months	reduced offspring kit growth	na	0.1 mg/day	1.12	na	Brunström et al. (2001)	1, 2
Aroclor 1254	mink	na	0.13	6 months	reduced offspring kit growth rate	na	1 mg/kg ww	1.34 (Bleavins and Aulerich 1981)	0.18 (Bleavins and Aulerich 1981)	Wren et al. (1987)	
Aroclor 1254	mink	na	0.22	4 and 9 months prior to giving birth	reduced number of offspring per female, decrease in offspring kit body weight	na	2 mg/kg ww	1.34 (Bleavins and Aulerich 1981)	0.15	Ringer (1983)	3
Aroclor 1254	mink	0.13	0.26	4 months	no kits born alive at 4 weeks	1	2 mg/kg ww	1.34 (Bleavins and Aulerich 1981)	0.18 (Bleavins and Aulerich 1981)	Aulerich and Ringer (1977)	
Aroclor 1254	mink	na	0.39	88 to 102 days	no kits whelped or born alive	na	2.5 mg/kg ww	0.87 (Bleavins and Aulerich 1981)	0.13 (Bleavins and Aulerich 1981)	Aulerich et al. (1985)	
PCB mixture (compo-sition not reported)	mink	na	0.51	66 days	reduced number of kits born alive	na	3.3 mg/kg ww	0.87 (Bleavins and Aulerich 1981)	0.13 (Bleavins and Aulerich 1981)	Jensen et al. (1977)	
Aroclor 1242	mink	na	0.65	8 months	reduced reproductive success	na	5 mg/kg ww	1.34 (Bleavins and Aulerich 1981)	0.18 (Bleavins and Aulerich 1981)	Bleavins et al. (1980)	
Aroclor 1254	mink	na	1.31	4 weeks	reduced weight gain in adults	na	10 mg/kg ww	1.34 (Bleavins and Aulerich 1981)	0.18 (Bleavins and Aulerich 1981)	Hornshaw et al. (1986)	
Aroclor 1254	mink	na	1.64	3 months	all whelps stillborn	na	na	na	na	Kihlstrom et al. (1992)	4
Aroclor 1254	mink	1.2	1.8	28 days	reduced female growth	na	na	na	na	Aulerich et al. (1986)	4
Clophen A50	mink	na	2.0	3 months	all whelps stillborn	na	na	na	na	Kihlstrom et al. (1992)	4

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)	Effect Conc.	Body Weight (kg)	Food Ingestion Rate (kg ww/day)	Source	Notes
Aroclor 1254	mink	1.5	2.4	28 days	reduced male and female growth	na	na	na	na	Aulerich et al. (1986)	4
Aroclor 1016	mink	na	2.6	8 months	reduced birth weight and growth rate of offspring kits, and 25 % adult female mortality	na	20 mg/kg ww	1.34 (Bleavins and Aulerich 1981)	0.18 (Bleavins and Aulerich 1981)	Bleavins et al. (1980)	

Notes:

1. Dietary dose determined by dividing daily dose by body weight. Female mink were exposed to 0.24 mg Clophen A50 three times a week, or 0.1 mg/day.
2. Body weight provided in study.
3. FIR provided in study.
4. Dietary dose calculated in study.

bw – body weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

PCB – polychlorinated biphenyl

ww – wet weight

Bold identifies the LOAEL selected as the TRV. A NOAEL TRV was not available from the study in which the chronic LOAEL of 0.089 mg/kg bw/day was reported, so it was estimated using an uncertainty factor of 2. The resulting NOAEL TRV is 0.045 mg/kg bw/day.

A.5.2.2.2 PCB TEQs

PCB TEQs are expressed as 2,3,7,8-TCDD equivalents; therefore, toxicity studies involving 2,3,7,8-TCDD effects on mammals were reviewed. Effects of dioxins and furans reported in laboratory studies with various species of mammals include developmental toxicity, hepatotoxicity, endocrine disruption, immunotoxicity, and death (Kennedy et al. 1996).

Seven studies that evaluated the dietary toxicity of 2,3,7,8-TCDD to mammals were identified (Table A.5-27). In these studies, adverse effects on growth, reproduction, and survival of guinea pigs, rats, or mink were reported following exposure to dietary 2,3,7,8-TCDD. The lowest dose at which effects were reported (4.9×10^{-6} mg/kg bw/day) was a subchronic study that resulted in reduced growth in guinea pigs exposed to 2,3,7,8-TCDD for 90 days (DeCaprio et al. 1986). This LOAEL was selected as the LOAEL TRV, although there is some uncertainty associated with this value because it was a short-term growth study. The highest NOAEL below this LOAEL (6.5×10^{-7} mg/kg bw/day) was from the same study with the same endpoint. This dose was selected as the NOAEL TRV.

Table A.5-27. 2,3,7,8-TCDD toxicity studies for mammals

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)	Effect Conc. (mg/kg ww)	Body Weight (kg)	Food Ingestion Rate (kg ww/day)	Source	Notes
2,3,7,8-TCDD	Hartley guinea pig	**6.5×10^{-7}**	**4.9×10^{-6}**	90 days	reduced body weight	1.0×10^{-5}	7.6×10^{-5}	na	na	DeCaprio et al. (1986)	1
2,3,7,8-TCDD	mink	2.6×10^{-6}	9.1×10^{-6}	131 to 132 days	decreased survival in kits at 3 weeks	1.6×10^{-5}	5.3×10^{-5}	1.089 (NOAEL), 1.054 (LOAEL)	0.18 (Bleavins and Aulerich 1981)	Hochstein et al. (2001)	2
2,3,7,8-TCDD	Sprague-Dawley rat	1.0×10^{-6}	1.0×10^{-5}	3 generations	reduced litter size and F2 postnatal survival	na	na	na	na	Murray et al.(1979)	1
2,3,7,8-TCDD	Hartley guinea pig	4.9×10^{-6}	2.85×10^{-5}	90 days	reduced survival	7.6×10^{-5}	4.3×10^{-4}	na	na	DeCaprio et al. (1986)	1
2,3,7,8-TCDD	mink	4.9×10^{-6}	5.0×10^{-5}	125 days	reduced body weight and adult survival	1.0×10^{-4}	1.0×10^{-3}	0.8776 (NOAEL), 0.8183 (LOAEL)	0.049 (NOAEL) 0.050 (LOAEL)	Hochstein et al. (2001)	3
2,3,7,8-TCDD	Sprague-Dawley rat	1.0×10^{-5}	1.0×10^{-4}	2 years	reduced body weight and adult female survival	na	na	na	na	Kociba et al.(1978)	1
2,3,7,8-TCDD	Sprague-Dawley rat	na	3.2×10^{-4}	13 weeks	reduced body weight	na	na	na	na	Van Birgelen et al. (1994)	1
2,3,7,8-TCDD	Sprague-Dawley rat	1.0×10^{-4}	na	3 generations	reduced body weight	na	na	na	na	Murray et al.(1979)	1

Notes:

1. Dietary dose calculated in study.
2. Body weight and ingestion rates used were specific to the no-effect and effect concentration test groups.
3. Body weight and FIR provided in study.

F2 – second generation

bw – body weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

TCDD – tetrachlorodibenzo-*p*-dioxin

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

A.5.2.2.3 Arsenic

Mammalian effects from chronic exposure to inorganic arsenic may include weakness, paralysis, conjunctivitis, dermatitis, decreased growth, liver damage, and developmental effects in offspring (Eisler 1988a). Early developmental stages are most sensitive to arsenic exposure.

One study that evaluated the toxicity of arsenic to mammals from dietary exposure (i.e., via food rather than drinking water or gavage) was identified (Table A.5-28). In this study, female rats fed 5.4 mg/kg bw/day arsenic had reduced body weights following two years of exposure to sodium arsenite (Byron et al. 1967). The results of this study were not statistically evaluated, although the final body weight range reported in rats fed 5.4 mg/kg bw/day (280.4 g ± standard error of 21.38 g) was lower than the final body weight range of the control group (350.9 g ± standard error of 26.36 g). Growth in rats appeared unaffected in rats exposed to 2.6 mg/kg bw/day (body weight at the end of the 2-year study was 322.4 g, with a standard error range of ± 21.38 g).

The LOAEL and NOAEL from Byron et al. (1967) (5.4 and 2.6 mg/kg bw/day, respectively) were selected as the LOAEL and NOAEL TRVs.

A.5.2.2.4 Cobalt

In laboratory studies, chronic exposure of mice and rats to cobalt via food, gavage or drinking water has resulted in adverse effects on reproduction, development, growth, and survival (ATSDR 2004). In addition, cardiovascular, neurological, renal, and endocrine effects have been reported (ATSDR 2004). Three studies that evaluated the toxicity of cobalt to mammals from dietary exposure (i.e., via food rather than drinking water or gavage) were identified for growth, reproductive, or survival endpoints (Table A.5-29). In these studies, adverse effects on survival or growth of laboratory guinea pigs and rats were reported following subchronic exposure to cobalt in food. No chronic studies were available. The lowest dose at which effects were reported (1.0 mg/kg bw/day) was selected as the LOAEL TRV. This LOAEL resulted in reduced growth in rats exposed to cobalt for 4 weeks (Chetty et al. 1979). No NOAELs that were lower than the selected LOAEL TRV were available. Therefore, a NOAEL TRV was estimated by dividing the subchronic LOAEL TRV by an uncertainty factor of 10, resulting in a NOAEL of 0.1 mg/kg bw/day.

Table A.5-28. Arsenic dietary toxicity studies for mammals

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)	Effect Conc. (mg/kg ww)	Body Weight (kg)	Food Ingestion Rate (kg ww/day)	Source	Notes
Sodium arsenite	rat	**2.6**	**5.4**	2 years	reduced female body weight	31.25	62.5	0.302 (NOAEL), 0.278 (LOAEL)	0.025 (NOAEL), 0.024 (LOAEL)	Byron et al. (1967)	1

Notes:

1. Data were not statistically evaluated. Body weight and FIR were reported in study.

bw – body weight

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

Table A.5-29. Cobalt dietary toxicity studies for mammals

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)	Effect Conc. (mg/kg ww)	Body Weight (kg)	Food Ingestion Rate (kg ww/day)	Source	Notes
Cobalt chloride	rat	na	**1.0**	4 weeks	reduced body weight	na	10	0.187	0.018 (EPA 1993)	Chetty et al. (1979)	1
Cobalt sulfate	guinea pig	na	1.4	5 weeks	reduced survival	na	20	0.50	0.035 (EPA 1993)	Mohiudden et al. (1970)	1, 2
Cobalt sulfate	guinea pig	1.4	na	5 weeks	no effect on body weight	20	na	0.50	0.035 (EPA 1993)	Mohiudden et al. (1970)	1
Cobalt chloride	rat	1.9	10	3 days	reduced body weight	20	100	0.196	0.019 (EPA 1993)	Wellman et al. (1984)	1, 3

Notes:

1. Body weight presented in study.
2. Data were not statistically evaluated; 4 of 20 guinea pigs died at LOAEL, and 1 of 20 guinea pigs died in the control group.
3. Data were not statistically evaluated; reduced food intake at LOAEL.

bw – body weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the LOAEL selected as the TRV. A NOAEL TRV was not available from the study in which the subchronic LOAEL of 1.0 mg/kg bw/day was reported, so it was estimated using an uncertainty factor of 10. The resulting NOAEL TRV is 0.01 mg/kg bw/day.

A.5.2.2.5 Mercury

Exposure of mammals to mercury has been reported to adversely affect reproduction, growth, development, behavior, blood and serum chemistry, motor coordination, vision, hearing, histology, and metabolism (Eisler 1987). Three studies that evaluated the toxicity of dietary mercury to mammals were identified for growth, reproduction, and survival endpoints (Table A.5-30). In these studies, adverse effects following dietary ingestion of mercury included mortality and depressed growth in laboratory rats and mink. At the lowest LOAEL, growth was significantly reduced in rats fed 0.0084 mg/kg bw/day of mercury as methylmercuric chloride for three generations (Verschuuren et al. 1976). Adverse effects in mink were reported at concentrations two orders of magnitude higher than the LOAEL measured for rats. Growth was significantly reduced, and mortality was observed in mink fed diets with 0.25 mg/kg bw/day methylmercuric chloride (Wobeser et al. 1976) and 0.64 gm/kg bw/day methylmercury (Aulerich et al. 1974) for a subchronic duration. The lowest LOAEL, 0.0084 mg/kg bw, was selected as the LOAEL TRV. While toxicology data based on mink studies may be more representative to the mammals utilizing the LDW, the LOAEL based on rats was selected because it was the most conservative effects threshold reported in the three studies reviewed and was based on a multi-generational study. No controlled laboratory studies were found where mink were exposed to dietary mercury over a chronic period or during a critical lifestage.

No NOAELs lower than the selected LOAEL TRV were available. Therefore, the NOAEL TRV was estimated by dividing the selected chronic LOAEL TRV by an uncertainty factor of 5, resulting in a NOAEL TRV of 0.0017 mg/kg bw/day.

A.5.2.2.6 Selenium

Selenium is an essential nutrient and deficiency in the diet may cause adverse effects in mammals; elevated dietary concentrations have also been reported to cause adverse effects (Eisler 1985). Primary effects on laboratory mice and rats from chronic exposure to selenium via food, gavage, or drinking water include reduced growth and survival; effects to the reproductive, cardiovascular, and hematological systems have also been reported (ATSDR 2003). Four studies that evaluated the toxicity of selenium to mammals from dietary exposure (i.e., via food rather than drinking water or gavage) were identified (Table A.5-31). In these studies, adverse effects on growth or survival were reported following subchronic exposure of laboratory rats or hamsters to selenium in their diet. Rats exhibited a higher sensitivity to dietary selenium than did hamsters. No studies were identified where mammals were exposed to dietary selenium for a chronic exposure period or during a critical lifestage.

LOAELs ranged from 0.080 mg/kg bw/day, resulting in reduced growth of rats (Halverson et al. 1966), to 5.8 mg/kg bw/day, resulting in reduced survival of hamsters (Julius et al. 1983). The lowest LOAEL of 0.080 mg/kg bw/day was selected as the TRV. The only NOAEL below this LOAEL (0.055 mg/kg bw/day) was from the same study with the same endpoint. This dose was selected as the NOAEL TRV.

Table A.5-30. Mercury dietary toxicity studies for mammals

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)	Effect Conc. (mg/kg ww)	Body Weight (kg)	Food Ingestion Rate (kg ww/day)	Source	Notes
Methylmercuric chloride	rat	na	**0.0084**	three generations	reduced growth	na	0.0799	0.16	0.016 (EPA 1993)	Verschuuren et al. (1976)	1
Methylmercuric chloride	rat	0.19	na	three generations	no effect on survival or reproduction	1.997	na	0.20	0.019 (EPA 1993)	Verschuuren et al. (1976)	1
Methylmercuric chloride	mink	0.16	0.25	93 days	reduced growth, 40% mortality	1.2	1.9	1.34 (Bleavins and Aulerich 1981)	0.18 (Bleavins and Aulerich 1981)	Wobeser et al. (1976)	2
Methylmercury	mink	na	0.64	2 months	reduced growth, 100% mortality	na	5	1.2	0.15	Aulerich et al. (1974)	1, 3

Notes:

1. Body weight presented in study.
2. Two out of five mink died at the LOAEL.
3. FIR presented in study.

bw – body weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the LOAEL selected as the TRV. A NOAEL TRV was not available from the study in which the chronic LOAEL of 0.0084 mg/kg bw/day was reported, so it was estimated using an uncertainty factor of 5. The resulting NOAEL TRV is 0.0017 mg/kg bw/day.

Table A.5-31. Selenium dietary toxicity studies for mammals

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc. (mg/kg ww)	Effect Conc. (mg/kg ww)	Body Weight (kg)	Food Ingestion Rate (kg ww/day)	Source	Notes
Sodium selenite	rat	**0.055**	**0.080**	6 weeks	reduced body weight	3.2	4.8	0.139 (NOAEL), 0.129 (LOAEL)	0.00238 (NOAEL), 0.00215 (LOAEL)	Halverson et al. (1966)	1, 2, 3
Sodium selenite	rat	0.13	0.14	6 weeks	reduced survival	8.0	9.6	0.129 (NOAEL), 0.1255 (LOAEL)	0.00215 (NOAEL), 0.00186 (LOAEL)	Halverson et al. (1966)	1, 2
L-seleno-methionine	rat	na	0.16	110 days	reduced body weight	na	2	0.34	0.027 (EPA 1993)	Behne et al. (1992)	1
Selenite	rat	0.16	na	110 days	no effect on body weight	2	na	0.34	0.027 (EPA 1993)	Behne et al. (1992)	1
Sodium selenite, nano-Se, or organic selenium	rat	0.17	0.28	13 weeks	reduced body weight	na	na	na	na	Jia et al. (2005)	4
Seleno-methionine	hamster	0.36	0.76	21 days	reduced body weight	5.1	10.1	0.092 (NOAEL), 0.091(LOAEL)	0.00655 (NOAEL), 0.0068 (LOAEL)	Julius et al. (1983)	1, 2, 3
Sodium selenite	hamster	na	3.4	21 days	reduced body weight	na	40.25	0.074	0.0062	Julius et al. (1983)	1, 2
Sodium selenite	hamster	na	5.8	21 days	reduced female survival	na	80.24	0.062	0.0045	Julius et al. (1983)	1, 2

Notes:

1. Body weight presented in study.
2. FIR presented in study. Data presented in study were not statistically evaluated.
3. Body weight and ingestion rates used were specific to the no-effect and effect concentration test groups.
4. Dietary dose calculated in study.

bw – body weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

A.5.3 SUMMARY OF WILDLIFE ASSESSMENT

A.5.3.1 Exposure assessment

The exposure assessment provided an estimate of each wildlife ROC's exposure to COPCs through ingestion of prey, water, and incidental sediment ingestion. Exposure doses were calculated for each ROC/COPC pair, and expressed as mg COPC ingested per kg body weight per day. Estimates of dietary composition and site usage were made using site-specific information, if available, along with species life history information. Exposure doses were estimated using UCL concentrations in prey tissue, sediment, and water. Exposure doses for wildlife were presented in Tables A.5-11 and A.5-12.

A.5.3.2 Effects assessment

The effects assessment selected TRVs to represent dietary thresholds of effects for each ROC/COPC pair. The toxicity literature was searched and relevant data for birds and mammals were compiled and screened against a set of guidelines to select the most appropriate TRVs. TRVs for both no-effects and low-effects data were selected, as summarized in Table A.5-32.

Table A.5-32. TRVs for ROC/COPC pairs

	NOAEL AND LOAEL TRVs (mg/kg bw/day)							
	SPOTTED SANDPIPER		GREAT BLUE HERON AND OSPREY		RIVER OTTER		HARBOR SEAL	
COPC	NOAEL	LOAEL	NOAEL	LOAEL	NOAEL	LOAEL	NOAEL	LOAEL
Arsenic	10	40	ne	ne	2.6	5.4	ne	ne
Cadmium	1.5	4.0	ne	ne	ne	ne	ne	ne
Chromium	1.0	5.0	1.0	5.0	ne	ne	ne	ne
Cobalt	2.31	23.1	ne	ne	0.1	1.0	ne	ne
Copper	21	29	ne	ne	ne	ne	ne	ne
Lead	5.82	20	5.82	20	ne	ne	ne	ne
Mercury	0.018	0.091	0.018	0.091	0.0017	0.0084	0.0017	0.0084
Nickel	17	33	ne	ne	ne	ne	ne	ne
Selenium	0.5	0.82	ne	ne	0.055	0.080	ne	ne
Vanadium	1.2	2.3	ne	ne	ne	ne	ne	ne
Zinc	82	124	ne	ne	ne	ne	ne	ne
Total PCBs	0.49	1.4	0.49	1.4	0.018	0.089	0.018	0.089
PCB TEQs	1.4×10^{-5}	1.4×10^{-4}	1.4×10^{-5}	1.4×10^{-4}	2.6×10^{-6}	9.1×10^{-6}	2.6×10^{-6}	9.1×10^{-6}

bw – body weight
COPC – chemical of potential concern
LOAEL – lowest-observed-adverse-effect level
ne – not evaluated; not a COPC for this ROC
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
TEQ – toxic equivalent
TRV – toxicity reference value

A.6.0 Risk Characterization and Uncertainty Analysis

This section presents the risk characterization for each ROC/COPC pair identified in the problem formulation (Section A.2.0) and discussed in the exposure and effects assessments (Sections A.3.0 through A.5.0) of this baseline ERA. The risk characterization section for each receptor group (i.e., benthic invertebrates, fish, and wildlife) consists of a risk estimate, an uncertainty analysis, and a risk conclusion section. The risk estimate section presents the hazard quotients (HQs)[67] calculated for each ROC/COPC pair. Uncertainties inherent in the risk assessment, including calculation of HQs, the problem formulation, and the exposure and effects assessment approach are discussed in the uncertainty analysis. The results of the HQ calculations and the uncertainty analysis are then integrated into risk conclusions.

In ERAs, HQs greater than 1.0 indicate that the exposures of some receptors are estimated to be greater than toxicological benchmarks. Such a finding is generally regarded as indicating a potential for adverse effects, particularly if the benchmark is an effects concentration (or dose) at which adverse effects were observed (i.e., a LOAEL). HQs may also be calculated based on a NOAEL. The potential for adverse effects associated with a NOAEL HQ greater than 1.0 is uncertain unless the LOAEL is also assessed because the true threshold for effects occurs at a concentration (or dose) somewhere between the NOAEL and LOAEL. An exposure falling between the NOAEL and LOAEL may or may not result in any adverse effect. Therefore, both types of HQs are calculated and presented to better describe the potential for adverse effects and to support risk management decisions.

A.6.1 BENTHIC INVERTEBRATES

This section characterizes risks to benthic invertebrates closely associated with sediment, such as amphipods, bivalves, and polychaetes, as well as more mobile, higher-trophic-level benthic invertebrates, such as crabs, that may travel over relatively greater distances than other invertebrates. Results from a direct measure of effect to one specific invertebrate group, gastropods, are also summarized.

Risk characterization for infaunal and epibenthic invertebrates was based primarily on an assessment of effects through the comparison of available surface sediment chemistry data with available sediment chemical criteria and guidelines and through the use of site-specific sediment toxicity tests (Ecology 1995).

Risks to crabs from two COPCs, zinc and PCBs, were characterized using a critical tissue-residue approach. The critical tissue-residue approach was also used to evaluate risks to infaunal invertebrates from TBT. Risks to meso- and neogastropods were

[67] The HQ is the ratio of the exposure concentration (or dose) to a concentration (or dose) associated with adverse effects.

characterized in two site-specific studies using a direct measure of effect (i.e., imposex).

Risks to benthic invertebrates from VOCs in porewater were characterized by comparing detected porewater concentrations to TRVs selected from the available literature. The risk characterization focused on areas where VOCs have been historically detected in groundwater at upland properties immediately adjacent to the LDW.

In this section, the risk characterization for the benthic invertebrate community is presented in Section A.6.1.1, and the risk characterization for crabs is presented in Section A.6.1.2. Risk characterization for the benthic invertebrate community includes risks from sediment-associated chemicals (Section A.6.1.1.1), porewater-associated VOCs (Section A.6.1.1.2), and TBT (Section A.6.1.1.3).

A.6.1.1 Benthic invertebrate community

The risk characterization for the benthic invertebrate community evaluated the following:

- Site-specific sediment toxicity test results and comparisons of surface sediment chemical concentrations to available sediment chemical criteria, guidelines, and TRVs
- Comparisons of VOC concentrations in porewater to toxicity data for benthic invertebrates
- Risks from TBT exposure based on results of the imposex study of meso- and neogastropods and using a critical tissue-residue approach

A.6.1.1.1 Sediment

Risk Estimates

The potential for adverse effects to benthic invertebrate communities resulting from exposure to sediment-associated COPCs was evaluated through site-specific toxicity testing and through a comparison of COPC concentrations in LDW surface sediment to SMS chemical criteria for 41 COPCs for benthic invertebrates. Three COPCs were evaluated based on a comparison of sediment chemical concentrations to toxicologically based sediment guidelines or TRVs.

The SMS regulations (WAC 173-204) provide both chemical- and biological effects-based criteria. The biological effects-based criteria provide an option for conducting site-specific confirmation of the chemical criteria through the use of sediment toxicity tests. Because AETs, which form the basis for the chemical criteria, are based on sediment samples with a mixture of chemicals from various locations in Puget Sound, and exceedance of those criteria is not always an accurate predictor of adverse effects, the regulations state that site-specific toxicity tests supersede site-specific chemistry data. For example, if the concentration of a chemical was greater than the CSL

chemical criteria at a location, but the sample was not toxic in the biological testing, then the location would not be classified as a CSL exceedance.

Table A.6-1 presents a summary of the surface sediment chemistry data for the COPCs identified in Section A.2.5.1.1 with at least one concentration greater than the SQS chemical criteria. The toxicity test results and associated sediment chemistry data from the 46 locations tested in Phase 2 and 7 locations tested in earlier studies are presented in Table A.6-2. The table also presents the final SMS classification for each location tested with the toxicity test results superseding the chemistry results. SQS and CSL exceedances based on toxicity test results and detected chemical concentrations above the SQS and CSL chemical criteria are shown as point locations on Map A.6-1. Chemicals not detected in sediments but with reporting limits greater than the SMS chemical criteria are discussed in the uncertainty analysis.

Table A.6-1. Detection frequencies and frequencies of detected concentrations greater than SQS and CSL for all SMS COPCs

COPC	Sediment Chemistry								
	Detection Frequency		Frequency of Detected Concentrations > SQS			Frequency of Detected Concentrations > CSL			Maximum Detected Conc./CSL
	No. of Samples[a]	Percent	No. of Samples[b]	Percent	No. of Samples with RL > SQS[c]	No. of Samples[d]	Percent	No. of Samples with RL > CSL[e]	
Metals									
Arsenic	754/816	92	13/816	1.6	0	8/816	0.98	0	12
Cadmium	565/799	71	13/799	1.6	0	11/799	1.4	0	18
Chromium	813/813	100	9/813	1.1	0	8/813	0.98	0	4.1
Copper	816/816	100	12/816	1.5	0	12/816	1.5	0	31
Lead	816/816	100	21/816	2.6	0	19/816	2.3	0	43
Mercury	717/833	86	37/833	4.4	0	23/833	2.8	0	7.8
Silver	481/784	61	10/784	1.3	0	10/784	1.3	0	44
Zinc	813/813	100	42/813	5.2	0	16/813	2.0	0	10
PAHs									
2-Methylnaphthalene	139/782	18	3/782	0.38	9	3/782	0.38	3	2.5
Acenaphthene	301/792	38	19/792	2.4	13	3/792	0.38	4	4.6
Anthracene	553/792	70	2/792	0.25	0	0/792	0	0	ne[f]
Benzo(a)anthracene	719/792	91	12/792	1.5	0	3/792	0.38	0	1.6
Benzo(a)pyrene	718/786	91	8/786	1.0	0	3/786	0.38	0	2.0
Benzo(g,h,i)perylene	649/787	82	16/787	2.0	7	7/787	0.89	3	2.3
Total benzofluoranthenes	727/786	92	9/786	1.1	0	4/786	0.51	0	2.0
Chrysene	741/792	94	24/792	3.0	0	1/792	0.13	0	1.3
Dibenzo(a,h) anthracene	400/792	51	19/792	2.4	19	4/792	0.51	9	2.2
Dibenzofuran	246/791	31	10/791	1.3	13	3/791	0.38	4	3.8
Fluoranthene	762/792	96	39/792	4.9	0	8/792	1.0	0	2.7
Fluorene	373/792	47	14/792	1.8	9	3/792	0.38	1	3.7

COPC	Sediment Chemistry								
	Detection Frequency		Frequency of Detected Concentrations > SQS			Frequency of Detected Concentrations > CSL			Maximum Detected Conc./CSL
	No. of Samples[a]	Percent	No. of Samples[b]	Percent	No. of Samples with RL > SQS[c]	No. of Samples[d]	Percent	No. of Samples with RL > CSL[e]	
Indeno(1,2,3-cd)pyrene	694/787	88	23/787	2.9	3	8/787	1.0	2	2.3
Naphthalene	148/782	19	2/782	0.26	2	2/782	0.26	2	1.7
Phenanthrene	727/792	92	27/792	3.4	0	3/792	0.38	0	3.1
Pyrene	755/792	95	4/792	0.51	0	3/792	0.38	0	1.3
Total HPAH	769/792	97	24/792	3.0	0	3/792	0.38	0	1.4
Total LPAH	731/792	92	6/792	0.76	0	3/792	0.38	0	2.9
Phthalates									
Bis(2-ethylhexyl) phthalate	636/796	80	106/796	13	5	58/796	7.3	2	7.4
Butyl benzyl phthalate	390/786	50	77/786	9.8	79	8/786	1.0	4	8.3
Dimethyl phthalate	136/786	17	2/786	0.25	15	2/786	0.25	7	2.6
Other SVOCs									
1,2,4-Trichlorobenzene	5/780	0.64	1/780	0.13	363	1/780	0.13	131	1.4
1,2-Dichlorobenzene	18/780	2.3	3/780	0.38	113	3/780	0.38	113	10
1,4-Dichlorobenzene	35/780	4.5	3/780	0.38	98	3/780	0.38	21	13
2,4-Dimethylphenol	1/773	0.13	1/773	0.13	224	1/773	0.13	224	10
4-Methylphenol	78/795	9.8	4/795	0.50	12	4/795	0.50	12	6.9
Benzoic acid	69/783	8.8	7/783	0.89	107	7/783	0.89	107	6.9
Benzyl alcohol	14/773	1.8	4/773	0.52	112	2/773	0.26	105	9.2
Hexachlorobenzene	46/783	5.9	6/783	0.77	388	2/783	0.26	108	1.7
n-Nitrosodiphenylamine	23/782	2.9	2/782	0.26	70	2/782	0.26	36	2.8
Pentachlorophenol	12/749	1.6	1/749	0.13	120	0/749	0	32	ne[g]
Phenol	254/795	32	25/795	3.1	5	7/795	0.88	0	2.3
PCBs									
Total PCBs	1205/1290	93	474/1290	37	0	173/1290	13	0	150

[a] Number of detected concentrations/number of surface sediment samples analyzed for the COPC.

[b] Number of detected concentrations > SQS/number of surface sediment samples analyzed for the COPC. For individual samples with TOC > 4% or < 0.5%, that sample was tallied as greater than the SQS if the dry weight concentration was greater than the LAET. The number of detected concentrations > SQS includes the number > CSL (i.e., this is not the number of concentrations between the SQS and the CSL).

[c] Number of samples with RL greater than the SQS. The number of samples with RLs > SQS includes the number > CSL (i.e., this is not the number of samples with RLs between the SQS and the CSL).These chemicals are discussed in the uncertainty analysis.

[d] Number of detected concentrations > CSL/number of surface sediment samples analyzed for the COPC. For individual samples with TOC > 4% or < 0.5%, the sample was tallied as greater than the CSL if the dry weight concentration was greater than the 2LAET.

[e] Number of samples with RLs exceeding the CSL. These chemicals are discussed in the uncertainty analysis.

[f] Maximum concentration of anthracene did not exceed CSL; maximum concentration/SQS equals 1.7.

[g] Maximum concentration of pentachlorophenol did not exceed CSL; maximum concentration/SQS equals 1.1.

COPC – chemical of potential concern
CSL – cleanup screening level
HPAH – high-molecular-weight polycyclic aromatic hydrocarbon
LPAH – low-molecular-weight polycyclic aromatic hydrocarbon
ne – not exceeded
PCB – polychlorinated biphenyl
SMS – Washington State Sediment Management Standards
SQS – sediment quality standard
SVOC – semivolatile organic compound

Table A.6-2. Sediment chemistry and toxicity test results for samples from the LDW

Location ID	Conc. > SMS Chemical Criteria	COPC(s) with Detected Concentrations Greater than the SQS or CSL[a]	Toxicity Test Exceedance[b]	Agreement in Classification	Final SMS Classification[c]
DUD200	CSL	**4-methylphenol**, bis(2-ethylhexyl) phthalate, butyl benzyl phthalate, total PCBs	no exceedance	no	no exceedance
DUD201	SQS	bis(2-ethylhexyl) phthalate, total PCBs	no exceedance	no	no exceedance
DUD202	CSL	**bis(2-ethylhexyl) phthalate**, butyl benzyl phthalate, total PCBs	no exceedance	no	no exceedance
DUD203	no exceedance	none	no exceedance	yes	no exceedance
DUD204	CSL	**4-methylphenol**, **bis(2-ethylhexyl) phthalate**, butyl benzyl phthalate	no exceedance	no	no exceedance
DUD205	CSL	**4-methylphenol**, **bis(2-ethylhexyl) phthalate**, butyl benzyl phthalate, total PCBs	no exceedance	no	no exceedance
DUD206	no exceedance	none	CSL	no	CSL
LDW-SS2	SQS	fluoranthene	CSL	no	CSL
LDW-SSB2b	SQS	total PCBs	CSL	no	CSL
LDW-SS6	CSL	arsenic, **bis(2-ethylhexyl) phthalate**, **lead**, **total PCBs**, zinc	CSL	yes	CSL
LDW-SSB6a	SQS	total PCBs	SQS	yes	SQS
LDW-SS15	CSL	**mercury**	SQS	no	SQS
LDW-SS16	SQS	total PCBs	SQS	yes	SQS
LDW-SS17	SQS	bis(2-ethylhexyl) phthalate	SQS	yes	SQS
LDW-SS24	CSL	**benzo(a)anthracene**, benzo(a)pyrene, **benzo(g,h,i)perylene**, **total benzofluoranthenes**, **benzyl alcohol**, **chrysene**, dibenzo(a,h)anthracene, **fluoranthene, indeno(1,2,3-cd)pyrene**, **mercury**, total PCBs, phenanthrene, **pyrene**, **total HPAH**, zinc	CSL	yes	CSL
LDW-SS26	SQS	butyl benzyl phthalate, total PCBs	no exceedance	no	no exceedance
LDW-SS29	no exceedance	none	no exceedance	yes	no exceedance
LDW-SS31	CSL	**arsenic**, **zinc**	CSL	yes	CSL
LDW-SS32	SQS	zinc	SQS	yes	SQS
LDW-SS37	CSL	**mercury**, **total PCBs**	CSL	yes	CSL
LDW-SS39	CSL	**mercury**	CSL	yes	CSL

Location ID	Conc. > SMS Chemical Criteria	COPC(s) with Detected Concentrations Greater than the SQS or CSL[a]	Toxicity Test Exceedance[b]	Agreement in Classification	Final SMS Classification[c]
LDW-SS40	SQS	total PCBs	CSL	no	CSL
LDW-SS49	CSL	**arsenic**, **copper**, zinc	CSL	yes	CSL
LDW-SS50	SQS	total PCBs	CSL	no	CSL
LDW-SS56	CSL	**arsenic**, **total PCBs**, zinc	SQS	no	SQS
LDW-SS57	SQS	total PCBs	CSL	no	CSL
LDW-SS58	SQS	total PCBs	CSL	no	CSL
LDW-SS60	SQS	total PCBs	no exceedance	no	no exceedance
LDW-SS63	no exceedance	none	no exceedance	yes	no exceedance
LDW-SS68	CSL	**hexachlorobenzene**	no exceedance	no	no exceedance
LDW-SS69b	SQS	total PCBs	CSL	no	CSL
LDW-SS70	SQS	bis(2-ethylhexyl) phthalate	SQS	yes	SQS
LDW-SS71	SQS	total PCBs	no exceedance	no	no exceedance
LDW-SS73	CSL	**benzyl alcohol**	SQS	no	SQS
LDW-SS75	SQS	total PCBs	no exceedance	no	no exceedance
LDW-SS77	SQS	arsenic	CSL	no	CSL
LDW-SS85	SQS	total PCBs	no exceedance	no	no exceedance
LDW-SS88	CSL	**mercury**, total PCBs	CSL	yes	CSL
LDW-SS89	CSL	**total PCBs**	no exceedance	no	no exceedance
LDW-SS92	CSL	**total PCBs**	no exceedance	no	no exceedance
LDW-SS106	SQS	total PCBs	no exceedance	no	no exceedance
LDW-SS112	CSL	**arsenic**, butyl benzyl phthalate, fluoranthene, total PCBs	no exceedance	no	no exceedance
LDW-SS114	CSL	**arsenic**, bis(2-ethylhexyl) phthalate, chrysene, fluoranthene, indeno(1,2,3-cd)pyrene, total PCBs	CSL	yes	CSL
LDW-SS115	SQS	chrysene, dibenzo(a,h)anthracene, fluoranthene, phenanthrene, total HPAH	no exceedance	no	no exceedance
LDW-SS119	SQS	butyl benzyl phthalate, total PCBs	no exceedance	no	no exceedance
LDW-SS120	SQS	butyl benzyl phthalate, total PCBs	SQS	yes	SQS
LDW-SS121	CSL	butyl benzyl phthalate, **lead**, total PCBs	no exceedance	no	no exceedance

Location ID	Conc. > SMS Chemical Criteria	COPC(s) with Detected Concentrations Greater than the SQS or CSL[a]	Toxicity Test Exceedance[b]	Agreement in Classification	Final SMS Classification[c]
LDW-SS122	SQS	total PCBs	no exceedance	no	no exceedance
LDW-SS143	CSL	**total PCBs**	no exceedance	no	no exceedance
LDW-SS144	SQS	total PCBs	SQS	yes	SQS
LDW-SS148	SQS	total PCBs	CSL	no	CSL
LDW-SS157	CSL	**benzoic acid**, butyl benzyl phthalate	SQS	no	SQS
LDW-SS158	SQS	total PCBs	no exceedance	no	no exceedance

[a] **Bold** COPCs had detected concentrations greater than the CSL in that sample. Other COPCs had detected concentrations greater than the SQS.

[b] Overall toxicity test exceedance of the SMS (for further details, see Section A.3.2.2).

[c] Overall SMS designation.

COPC – chemical of potential concern

CSL – cleanup screening level

HPAH – high-molecular-weight polycyclic aromatic hydrocarbon

ID – identification

LDW – Lower Duwamish Waterway

PCB – polychlorinated biphenyl

SMS – Washington State Sediment Management Standards

SQS – sediment quality standard

The frequency of COPC concentrations greater than the SQS chemical criteria ranged from 1.1 to 5.2% for metals and trace elements, from 0.25 to 4.9% for PAHs, from 0.25 to 13% for phthalates, and from 0.13 to 3.1% for other SVOCs; the frequency of PCB concentrations greater than the SQS was 37%. All detected chemicals, except anthracene and pentachlorophenol, were above the CSL chemical criteria in at least one sample. The highest ratios of maximum detected concentrations to CSL chemical criteria were for total PCBs (150), silver (44), lead (43), copper (31), cadmium (18), 1,4-dichlorobenzene (13), and arsenic (12). Ratios for all other COPCs were less than or equal to 10.

Table A.6-3 presents the sediment chemistry data for the three COPCs without SMS chemical criteria (nickel, total DDTs, and total chlordane). The frequency of nickel concentrations greater than the SL and ML was 1.0 and 0.52%, respectively. For total DDTs and chlordane, the frequency of concentrations greater than the NOAEL was 0.51 and 9.6%, and the frequency of concentrations greater than the LOAEL was 0.51 and 7.1%, respectively. The ratios of maximum detected concentrations to the ML or NOAEL were 2.5, 2.7, and 48 for nickel, total DDTs, and total chlordane, respectively. The detected chemical concentrations greater than the DMMP guidelines or TRV values are shown as point locations on Map A.6-2.

Application of SMS criteria to assess adverse effects from sediment-associated chemicals requires an assessment of both the magnitude and spatial extent of the contamination. A spatial analysis of potential effects was performed using Thiessen polygons, a method commonly used to illustrate spatial variability in sampling intensity and to extrapolate results from small areas (sample points) to larger areas. The Thiessen polygon associates each point on a plane with the closest sampling location for which an empirical value is available (Burmaster and Thompson 1997). In effect, this algorithm assumes that the concentration at any point where an empirical value is not available has not been made is the same as the concentration in the sample closest to that point. Assumptions regarding spatial homogeneity within polygons introduce uncertainty in all areal percentages discussed in this section. In areas with lower sample density, the uncertainty in this assumption increases (i.e., all points within a polygon are less likely to have the same characteristics as the point that defined the polygon category). The uncertainty section includes a discussion of the spatial analysis using Thiessen polygons relative to another interpolation approach.

Map A.6-3 presents areas of the LDW that are categorized as ≤ SQS (white), > SQS and ≤ CSL (yellow), and > CSL (red) based on a combination of the sediment chemistry data and toxicity test results. The overall SMS designation for each location is presented in Table A.6-2. The percent area in each of those three categories is 75%, 18%, and 7%, respectively.

Table A.6-3. Detection frequencies and frequencies of detected concentrations above the SL/NOAEL and ML/LOAEL for COPCs without SMS chemical criteria

COPC	Sediment Chemistry								
	Detection Frequency		Frequency of Detected Concentrations > SL or NOAEL			Frequency of Detected Concentrations > ML or LOAEL			Maximum Detected Conc./ML or LOAEL
	No. of Samples[a]	Percent	No. of Samples[b]	Percent	No. of Samples with RL > SL or NOAEL[c]	No. of Samples[d]	Percent	No. of Samples with RL > ML or LOAEL[e]	
Nickel	775/775	100	9/775	1.2	0	4/775	0.52	0	2.5
Total DDTs	78/197	40	1/197	0.51	1	1/197	0.51	0	2.7
Total chlordane[f]	33/197	17	19/197	9.6	79	14/ 197	7.1	61	48

[a] Number of detected concentrations/number of surface sediment samples analyzed for the COPC.

[b] [Number of detected concentrations > DMMP SL or NOAEL]/number of surface sediment samples analyzed for the COPC. The number of detected concentrations > SL or NOAEL includes the number > ML or LOAEL (i.e., this is not the number of concentrations between the SL and the ML or between the NOAEL and the LOAEL).

[c] Number of samples with RLs exceeding the DMMP SL or NOAEL. These chemicals are discussed in the uncertainty analysis.

[d] [Number of detected concentrations > DMMP ML or LOAEL]/number of surface sediment samples analyzed for the COPC.

[e] Number of samples with RLs exceeding the DMMP ML or LOAEL. These chemicals are discussed in the uncertainty analysis.

[f] Total chlordane includes the calculated total chlordane for Phase 2 data and chlordane as reported in a subset of historical data (King County 1999b).

COPC – chemical of potential concern

DMMP – Dredged Material Management Program

LOAEL – lowest-observed-adverse-effect level

ML –maximum level (DMMP)

NOAEL – no-observed-adverse-effect level

SL – screening level (DMMP)

SMS – Washington State Sediment Management Standards

TRV – toxicity reference value

The numbers of detected concentrations greater than the SQS and CSL based solely on sediment chemistry are shown in Table A.6-4. As shown in this table, PCBs and BEHP were the two COPCs with the highest numbers of detected concentrations greater than the CSL chemical criteria. Therefore, spatial analyses using Thiessen polygons were also performed for total PCBs and BEHP individually. The spatial distributions of chemical concentrations for these two chemicals relative to SQS and CSL chemical criteria are presented in Maps A.6-4 and A.6-5 and are discussed below.

Table A.6-4. Number of detected concentrations for each COPC greater than SQS and CSL chemical criteria based on sediment chemistry data

COPC	Number of Detected Concentrations > SQS and ≤ CSL	Number of Detected Concentrations > CSL
Total PCBs	301	173
Bis(2-ethylhexyl) phthalate	48	58
Mercury	14	23
Lead	2	19
Zinc	26	16
Copper	0	12
Cadmium	2	11
Silver	0	10
Fluoranthene	31	8
Butyl benzyl phthalate	69	8
Indeno(1,2,3-cd)pyrene	15	8
Chromium	1	8
Arsenic	5	8
Phenol	18	7
Benzo(g,h,i)perylene	9	7
Benzoic acid	0	7
Dibenzo(a,h)anthracene	15	4
Total benzofluoranthenes	5	4
4-Methylphenol	0	4
Phenanthrene	24	3
Total HPAH	21	3
Acenaphthene	16	3
Fluorene	11	3
Benzo(a)anthracene	9	3
Dibenzofuran	7	3
Benzo(a)pyrene	5	3
Total LPAH	3	3
Pyrene	1	3
1,4-Dichlorobenzene	0	3
1,2-Dichlorobenzene	0	3
2-Methylnaphthalene	0	3
Dimethyl phthalate	0	2
Naphthalene	0	2
N-Nitrosodiphenylamine	0	2
Hexachlorobenzene	4	2
Benzyl alcohol	2	2
Chrysene	23	1

COPC	Number of Detected Concentrations > SQS and ≤ CSL	Number of Detected Concentrations > CSL
1,2,4-Trichlorobenzene	0	1
2,4-Dimethylphenol	0	1
Anthracene	2	0
Pentachlorophenol	1	0

COPC – chemical of potential concern

CSL – cleanup screening level

HPAH – high-molecular-weight polycyclic aromatic hydrocarbon

LPAH – low-molecular-weight polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

SQS – sediment quality standard

Using Thiessen polygons and the chemistry data, total PCB concentrations were greater than the SQS and less than or equal to the CSL chemical criteria in approximately 17% of the total LDW area; total PCBs were the only COPC with concentrations greater than the SQS chemical criteria in approximately 14% of the total LDW area. In addition, total PCB concentrations were greater than the CSL chemical criteria in approximately 3% of the total LDW area (Map A.6-3). Total PCB concentrations were less than the SQS or CSL chemical criteria in approximately 80% of the total LDW. Total PCB concentrations greater than the SQS or the CSL chemical criteria were mostly reported between RM 0.4 and RM 0.6, in Slip 4, and between RM 2.9 and RM 3.7 (Map A.6-4). Thiessen polygons associated with locations where total PCBs were not detected are shown on Map A.6-4 in gray.

BEHP concentrations greater than the SQS or the CSL chemical criteria were mostly reported near RM 0.4 (Map A.6-5). BEHP concentrations were greater than the SQS and less than or equal to the CSL in approximately 4% of the total LDW area and greater than the CSL in approximately 2% of the total LDW area. BEHP concentrations were less than the SQS or CSL chemical criteria in approximately 94% of the total LDW. Thiessen polygons associated with locations where BEHP was not detected are shown on Map A.6-5 in gray, regardless of whether the BEHP RL for that location was above or below the SQS or CSL chemical criteria.

Uncertainties Associated with Sediment Risk Estimates

This section presents uncertainties in the sediment-based risk characterization for the benthic invertebrate community. The uncertainties are discussed separately for the problem formulation, exposure assessment, and effects assessment.

Problem Formulation

The benthic invertebrate community as a whole (except for the larger, more mobile species, such as crabs, which were assessed separately) was selected as an ROC because the community encompasses all benthic invertebrates as a functional group,

not as individual species. Because the benthic invertebrate community is the selected receptor, this approach does not address risks or toxicity to each individual species that is or could be present in the sediment environment. Instead, the receptor selection addresses effects at the community level, reflecting the diversity of species and ecological functions that are achieved with various benthic invertebrate assemblages. This receptor group, assessment endpoints (survival, growth and reproduction) and the sediment regulatory framework (SMS criteria) are aimed at protecting community function, not individual species.

AETs, which form the basis for SMS criteria and some of the DMMP guidelines, exist for about 20% of the chemicals that have been detected in the LDW. Therefore, chemicals without such criteria, guidelines, or other relevant toxicity information or chemicals with guidelines not derived on the basis of toxicity were not identified as COPCs during the problem formulation. However, it is likely that locations with the highest potential for adverse effects were adequately identified because criteria and guidelines are available for chemicals within most of the chemical groups (e.g., metals, PAHs, phthalates, SVOCs, PCBs, and pesticides) that are generally considered in CERCLA investigations. The importance of chemicals lacking criteria or other TRVs to the overall risk to benthic invertebrate communities was evaluated by identifying locations where the maximum detected concentration of each of these chemicals was greater than 10 times the mean concentration calculated using only detected values. A factor of 10 was selected as an arbitrary means to identify those chemicals with highly variable concentrations. Sixteen chemicals were identified through this process, including polychlorinated terphenyls (PCTs); thallium, barium, and manganese; and SVOCs such as carbazole and dibenzothiophene. All but three of the locations where the concentrations of these chemicals were elevated relative to the mean were already among the stations at which the concentrations of one or more chemicals exceeded the CSL or SQS. Chemicals without criteria at the three locations without any SQS exceedances were PCTs and barium. Therefore, the lack of criteria or TRVs for several chemicals in sediment is not likely to substantially alter the overall risk conclusions.

Finally, by using sediment criteria to judge the potential for adverse effects on the benthic invertebrate community, the risk assessment does not address the potential for other types of more subtle effects, such as biochemical changes. While the SMS is assumed to address the likelihood of adverse effects, such as reduced survival and growth, assessment of biochemical endpoints that may or may not relate to adverse effects on the benthic invertebrate community as a whole, is beyond the scope of this risk assessment.

Exposure Assessment

Uncertainties in the exposure assessment for the benthic invertebrate community were associated with the following factors.

Depth of biologically active zone. Some benthic invertebrate species (e.g., clams) may burrow deeper than 15 cm, which was the surface sediment threshold used in this

Phase 2 ERA to define the biologically active zone where the majority of the benthic invertebrate community resides. A risk characterization for these animals could have a different outcome than that presented in Section A.6.1.1 if: 1) concentrations in sediment above 15 cm were markedly different than those in sediment between 15 and 40 cm (depth that clams have been found in the LDW), or 2) the chemical sensitivity of animals living below 15 cm is markedly different than the chemical sensitivity of animals living above 15 cm, on which the existing chemical criteria and guidelines are based. The Phase 2 RI presents an evaluation of the potential exposure of these subsurface sediments to human and ecological receptors and assesses risks associated with these exposures.

Relationships among sediment chemistry, toxicity, and actual *in-situ* effects. The use of chemical criteria and toxicity testing to assess *in-situ* effects is uncertain. For example, adverse effects on the benthic invertebrate community may occur in areas with chemical concentrations less than SQS chemical criteria and may not occur in areas with concentrations greater than SQS or CSL chemical criteria. Factors such as site-specific bioavailability, mixtures of chemicals with or without criteria, and species-specific sensitivities may contribute to this uncertainty. Moreover, the SMS provides chemical-specific criteria to assess the risks from individual chemicals. Although these criteria were developed from field data in which mixtures of chemicals are common, the chemical-specific criteria do not assess the cumulative risks to benthic invertebrates from exposure to multiple chemicals with potentially synergistic or antagonistic effects.

Frequency of analyses of chemicals in surface sediment samples. In addition, not all chemicals evaluated in this ERA were analyzed in all surface sediment samples. While most COPCs have been analyzed in surface sediment at more than 700 locations, some, such as total DDTs and chlordane, have been analyzed at fewer locations (e.g., organochlorine pesticides were analyzed at 197 locations; Table A.3-1). The certainty regarding the risk characterization for these chemicals is lower compared to chemicals analyzed more frequently. However, because all COPCs were analyzed throughout the LDW as part of one or more reconnaissance-level sampling event and more recent sampling events focused on specific potential sources, the potential effect of this uncertainty on overall risk conclusions is likely to be low.

Elimination of COPCs based on the 5% detection frequency screen. Nineteen chemicals were not selected as COPCs because they were detected in fewer than 5% of the baseline surface sediment samples. To assess the uncertainty resulting from this approach, a cluster analysis was conducted for each of the chemicals that were screened out. The cluster analysis helped to determine whether the locations where these rarely detected chemicals were detected were clustered together. If detections were clustered, these chemicals might be of greater concern than if they had been widely dispersed or randomly distributed. The majority of the screened-out chemicals (14) had fewer than eight detected concentrations; six of the chemicals were detected only once. Among the other eight chemicals with fewer than eight detected

concentrations, only endrin aldehyde was detected at more than one location within 0.1 RM of each other, all within an early action area. Four of the five chemicals with more than eight detected concentrations (2-dichlorobenzene, benzyl alcohol, n-nitrosodiphenylamine, and pentachlorophenol) did not cluster but were scattered throughout the LDW. Detected concentrations of the fifth remaining chemical (1,4-dichlorobenzene) exceeded the SQS in three areas of the LDW: along the east bank between RM 0.3 and RM 0.7, between RM 3.3 and RM 3.8, and between RM 4.8 and RM 4.9. All of these locations are in early action areas.

RLs greater than criteria or TRVs. Twenty-seven chemicals had non-detected results with RLs greater than the corresponding SQS/CSL chemical criteria in at least one sediment sample in the baseline surface sediment dataset. These chemicals can be divided into three groups: PAHs (9 chemicals), phthalates (5 chemicals), and other SVOCs (13 chemicals). The detected results and RLs for each of these chemicals were compared to the corresponding SQS and CSL chemical criteria, and the results are presented in Table A.6-5. The results are presented separately for the 2004/2005 data collected as part of the Phase 2 RI, which includes all surface sediment samples collected by LDWG in 2004 and 2005, and the non-LDWG data, which includes the portion of the dataset that was presented in the Phase 1 RI (Windward 2003a) and data collected by parties other than LDWG in the time period since the Phase 1 RI.

The sample-specific RL is based on the lowest point of the calibration curve associated with each analytical batch of samples. The most common reason for elevated RL values is sample extract dilution. For example, elevated RLs for some chemicals in some areas reflect the greater degree of analytical dilution required for quantification of other analytes, such as PCBs. In addition, there are analytes known to be analytically difficult. These compounds tend to have chemical characteristics that differ from those of other analytes being analyzed using the same method. For example, benzoic acid, benzyl alcohol, phenols, and n-nitrosodiphenylamine are all more chemically reactive than the other SVOCs analyzed by EPA (2003a). More reactive compounds can be difficult to extract and often degrade during analysis.

PAHs and phthalates were detected relatively frequently in surface sediments, with detection frequencies ranging from 11 to 93% for PAHs and from 3.0 to 76% for phthalates. The majority of the RL values reported for these compounds were below the SQS and CSL chemical criteria. LDWG data and non-LDWG data can be compared in terms of the frequency with which RL values were greater than the chemical criteria. The LDWG data had much lower frequencies of RL values greater than the chemical criteria. No RLs above the SQS were reported for 2-methyl naphthalene, acenaphthylene, benzo(g,h,i)perylene, indeno(1,2,3-cd)pyrene, and naphthalene in the LDWG dataset because the laboratory, where possible, conducted additional analyses to increase the sensitivity of the analyses. RLs greater than the SQS chemical criteria for PAHs and phthalates primarily resulted from analytical dilution of the sample extracts.

Table A.6-5. Summary of chemical data for COPCs with at least one reporting limit greater than SQS/CSL chemical criteria

Chemical	Dataset[a]	Total No. of Samples Analyzed	Detection Frequency (%)	Detected Results		Non-Detected Results			RL > SQS Frequency (%)
				> SQS and ≤ CSL	> CSL	RL ≤ SQS	RL > SQS and ≤ CSL	RL > CSL	
PAHs									
2-Methylnaphthalene	2004/2005 LDWG	193	24	0	1	147	0	0	0.0
	non-LDWG	589	16	0	2	487	6	3	1.5
Acenaphthene	2004/2005 LDWG	193	34	2	2	126	1	0	0.52
	non-LDWG	599	39	14	1	352	8	4	2.0
Acenaphthylene[b]	2004/2005 LDWG	193	30	0	0	136	0	0	0.0
	non-LDWG	589	11	0	0	522	0	3	0.51
Benzo(g,h,i)perylene	2004/2005 LDWG	193	73	2	2	53	0	0	0.0
	non-LDWG	594	86	7	5	78	4	3	1.2
Dibenzo(a,h)anthracene	2004/2005 LDWG	193	31	3	1	130	3	0	1.6
	non-LDWG	599	57	12	3	243	7	9	2.7
Dibenzofuran	2004/2005 LDWG	193	28	0	2	139	1	0	0.52
	non-LDWG	598	32	7	1	393	8	4	2.0
Fluorene	2004/2005 LDWG	193	39	1	2	118	0	0	0.0
	non-LDWG	599	50	10	1	292	8	1	1.5
Indeno(1,2,3-cd)pyrene	2004/2005 LDWG	193	93	4	2	14	0	0	0.0
	non-LDWG	594	87	11	6	76	1	2	0.51
Naphthalene	2004/2005 LDWG	193	26	0	1	143	0	0	0.0
	non-LDWG	589	17	0	1	489	0	2	0.34
Phthalates									
Bis(2-ethylhexyl) phthalate	2004/2005 LDWG	193	76	5	3	47	0	0	0.0
	non-LDWG	603	81	43	55	108	3	2	0.83
Butyl benzyl phthalate	2004/2005 LDWG	193	35	8	0	115	11	4	7.8
	non-LDWG	593	55	61	8	202	64	0	11

Chemical	Dataset[a]	Total No. of Samples Analyzed	Detection Frequency (%)	Detected Results		Non-Detected Results			RL > SQS Frequency (%)
				> SQS and ≤ CSL	> CSL	RL ≤ SQS	RL > SQS and ≤ CSL	RL > CSL	
Diethyl phthalate	2004/2005 LDWG	193	12	0	0	168	1	0	0.52
	non-LDWG	603	3.0	0	0	580	3	2	0.83
Dimethyl phthalate[b]	2004/2005 LDWG	193	14	0	0	165	0	1	0.52
	non-LDWG	593	18	0	2	470	8	6	2.4
Di-n-octyl phthalate	2004/2005 LDWG	193	3.6	0	0	179	0	0	0.0
	non-LDWG	603	6.8	0	0	499	8	0	1.3
Other SVOC									
1,2,4-Trichlorobenzene	2004/2005 LDWG	193	0.0	0	0	155	16	22	20
	non-LDWG	587	0.17	0	1	257	216	109	55
1,2-Dichlorobenzene[b]	2004/2005 LDWG	193	0.5	0	0	174	0	18	9.3
	non-LDWG	587	2.9	0	3	475	0	95	16
1,4-Dichlorobenzene	2004/2005 LDWG	193	0.5	0	0	175	14	3	8.8
	non-LDWG	587	5.8	0	3	472	63	18	14
2,4-Dimethylphenol[b]	2004/2005 LDWG	193	0.0	0	0	162	0	31	16
	non-LDWG	580	0.2	0	1	386	0	193	33
2-Methylphenol	2004/2005 LDWG	193	1.0	0	0	175	0	16	8.3
	non-LDWG	592	0.2	0	0	490	0	101	17
4-Methylphenol[b]	2004/2005 LDWG	193	8.8	0	0	176	0	0	0.0
	non-LDWG	602	10	0	4	529	0	12	2.0
Benzoic acid[b]	2004/2005 LDWG	193	21	0	1	133	0	20	10
	non-LDWG	590	4.9	0	6	474	0	87	15
Benzyl alcohol	2004/2005 LDWG	193	4.7	1	2	166	0	18	9.3
	non-LDWG	580	0.9	1	0	481	7	87	16
Hexachlorobenzene	2004/2005 LDWG	194	6.2	0	2	157	16	9	13
	non-LDWG	589	5.8	4	0	192	264	99	62
Hexachlorobutadiene	2004/2005 LDWG	193	0.0	0	0	178	8	7	7.8
	non-LDWG	589	0.0	0	0	456	49	84	23

Chemical	Dataset[a]	Total No. of Samples Analyzed	Detection Frequency (%)	Detected Results		Non-Detected Results			RL > SQS Frequency (%)
				> SQS and ≤ CSL	> CSL	RL ≤ SQS	RL > SQS and ≤ CSL	RL > CSL	
n-Nitrosodiphenylamine[b]	2004/2005 LDWG	193	7.8	0	0	177	0	1	0.52
	non-LDWG	589	1.4	0	2	512	34	35	12
Pentachlorophenol	2004/2005 LDWG	193	3.6	1	0	168	7	11	9.3
	non-LDWG	556	0.9	0	0	449	81	21	18
Phenol	2004/2005 LDWG	193	22	2	3	151	0	0	0.0
	non-LDWG	602	35	16	4	385	5	0	0.83

[a] Non-LDWG refers to data collected by parties other than LDWG.

[b] The SQS and CSL chemical criteria are the same for these chemicals. Therefore, all values greater than either the SQS or CSL are presented as being above the CSL, except in cases where the AET was used because of TOC values outside the acceptable range.

CSL – cleanup screening level

LDWG – Lower Duwamish Waterway Group

PAH – polycyclic aromatic hydrocarbon

RL – reporting limit

SQS – sediment quality standard

SVOC – semivolatile organic compound

The group of compounds labeled as "other SVOCs" included the following chemicals: chlorobenzenes, phenol, methyl phenols, pentachlorophenol, benzoic acid, benzyl alcohol, hexachlorobutadiene, hexachlorobenzene, and n-nitrosodiphenylamine. This group includes compounds that are analytically difficult to quantify at the levels required for comparison to SQS chemical criteria and are generally very rarely detected. Only benzoic acid and phenol have been detected at frequencies greater than 10% in either dataset. The 2004/2005 LDWG data had consistently lower frequencies of RL values greater than the SQS chemical criteria than did the non-LDWG data because additional selected ion monitoring analyses were conducted to improve the sensitivity of the analyses of these compounds. Except for hexachlorobenzene and n-nitrodiphenylamine, the frequency of RL values above the SQS chemical criteria in the LDWG data was approximately half the frequency observed for the non-LDWG data. For hexachlorobenzene, the frequency of RL values greater than the SQS chemical criteria in the LDWG data was 20% of the frequency in the non-LDWG data. For n-nitrosodiphenylamine, the frequency in the LDWG data was 5% of that observed in the non-LDWG data. Nevertheless, there were many RL values above the SQS and CSL chemical criteria for these compounds in both the 2004/2005 LDWG and the non-LDWG datasets. The highest frequencies of RL values greater than the chemical criteria were reported for 1,2,4-trichlorobenzene, 2,4-dimethylphenol, and hexachlorobenzene. For the LDWG 2004/2005 data, more sensitive analytical methods were used whenever RL values were greater than the SQS with no detected values above the SQS chemical criteria.

Because chemical criteria are not available for organochlorine pesticides, NOAEL and LOAEL values were developed for the two organochlorine pesticide COPCs (total DDTs and total chlordane) for benthic invertebrates.[68] The number of detected results and RLs that exceeded these values are summarized in Table A.6-6. One RL value for total DDTs was greater than the total DDT NOAEL. Sixty-one samples had total chlordane RLs greater than both the NOAEL and LOAEL values. Elevated RL values for organochlorine pesticides generally reflect the presence of probable analytical interference in the analysis because of the presence of PCB congeners. The LDWG dataset represents 91 of the total of 197 samples analyzed for organochlorine pesticides. All of the organochlorine pesticide results for these samples were reported as not detected with elevated RLs because of analytical interference resulting from the presence of PCB congeners. All of the 61 samples with total chlordane RLs above the LOAEL had detected results of other chemicals greater than the SQS. Of the 18 samples with total chlordane RLs above the NOAEL, 15 did not have detected concentrations of other chemicals greater than the SMS.

[68] Chemical criteria were also unavailable for nickel. Nickel was always detected in surface sediment samples, and thus RLs were not an issue for nickel.

Table A.6-6. Summary of detected results and RLs for total DDTs and total chlordane relative to NOAELs and LOAELs

Chemical	Total No. of Samples Analyzed	Detection Frequency (%)	Detected Results > NOAEL and ≤ LOAEL	Detected Results > LOAEL	RL > NOAEL and ≤ LOAEL	RL > LOAEL
Total DDTs	197	40	0	1	1	0
Total chlordane	197	17	5	14	18	61

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

RL – reporting limit

The spatial distributions of the three chemicals with the highest frequencies of RL values greater than the chemical criteria (i.e., 1,2,4-trichlorobenzene, 2,4-dimethyl phenol, and hexachlorobenzene) are presented in Maps A.6-6, A.6-7, and A.6-8. For all three compounds, samples with RLs greater than the SQS chemical criteria occurred throughout the LDW with no spatial relationship between the detected results that were greater than the SQS and CSL chemical criteria and the RLs that were greater than the SQS chemical criteria, although these chemicals were detected infrequently. 1,2,4-trichlorobenzene and 2,4-dimethyl phenol each had one detected concentration above the CSL chemical criteria in the non-LDWG dataset. Hexachlorobenzene had four detected values above the SQS chemical criteria in the non-LDWG dataset and two detected values above the CSL chemical criteria in the LDWG dataset.

Locations where there were no detected concentrations greater than the chemical criteria but RLs were greater than the SQS/CSL chemical criteria are shown on Map A.6-9. Only two of the 294 locations where this occurs were from the LDWG 2004/2005 dataset. The majority of the locations were associated with RL values for 1,2,4-trichlorobenzene, hexachlorobenzene, and 2,4-dimethyl phenol. These three chemicals were responsible for 94% of the locations where only RL values were greater than the SQS or CSL chemical criteria. Two of these chemicals, 1,2,4-trichlorobenzene and 2,4-dimethylphenol, were never reported as detected in the LDWG 2004/2005 surface sediment dataset, despite increased efforts to improve the sensitivity of the analytical method, which resulted in a reduced frequency of RL values above the SQS chemical criteria. Hexachlorobenzene was detected in approximately 6% of the samples in both the LDWG and non-LDWG datasets. The increased sensitivity of the analysis in the LDWG 2004/2005 dataset reduced the frequency of RL values above the SQS chemical criteria. The RL values above SQS chemical criteria for these compounds appear to reflect analytical difficulties with the analysis, and the paucity of detected concentrations suggests that these chemicals are not present in the LDW at the concentrations represented by the RL.

There were 18 locations with RL values greater than the SQS chemical criteria associated with other chemicals. The majority of those locations were associated with n-nitrosodiphenylamine RLs above the SQS chemical criteria (14 locations).

For PAHs and phthalates, the RL values greater than the SQS chemical criteria occur infrequently and are associated with samples that required analytical dilutions because of the presence of other chemicals with detected concentrations above the SQS chemical criteria. Thus, RLs for these chemicals had little effect on risk conclusions.

Spatial analysis using Thiessen polygons. Thiessen polygons were used in the ERA to estimate the area potentially affected by exceedances of SMS at each point for which chemical concentrations were compared to SMS, or for which bioassays were performed, after the chemical and biological analyses of LDW sediment at all points were combined (Section A.6.1.1.1). There is uncertainty associated with methods for interpolation of point values to area values, including the Thiessen polygon method. To assess this uncertainty, results of this interpolation method were compared to results of the same analysis using a different method, inverse distance weighting (IDW). IDW estimates chemical concentrations for each model cell in a grid surface as a weighted average of the sample concentrations that are defined as the neighbors of that cell. The weights are a function of the inverse distance of the cell from each neighboring sample concentration. Thiessen polygons are the result of an algorithm (Voronoi tessellation) that defines polygon boundaries such that any arbitrary location within a polygon is closer to its associated sample location than to any other sample location. The sample concentration within each Thiessen polygon is assumed to be uniformly distributed across the area of the polygon, so that any of the aforementioned arbitrary locations is estimated to have the same chemical concentration as its associated sample location. Each method is known to have a degree of uncertainty regarding the reliability of its estimates; comparison of the results of the two methods provides an indication of the uncertainty associated with application of Thiessen polygons to estimate areas affected when only point values are known. There is no way with the existing dataset to quantify the deviation of either method from reality.

Map A.6-10 illustrates differences in estimates of PCBs generated by the two methods in one area within the LDW. This area is representative of the types of relative over- and underestimation that occur at numerous locations throughout the LDW based on a comparison of the approaches. The differences described here can be observed to varying degrees throughout the waterway. As illustrated on Map A.6-10, the estimated concentration within an individual Thiessen polygon (excluding the specific location of the sample itself where the concentration is known) may be high or low relative to IDW-estimated concentrations for any fraction of the area within the polygon. Moreover, estimates for multiple Thiessen polygons can be wholly overestimated by IDW, depending on the point value for the polygon and the density of the sampling locations.

Individual polygons are marked in Map A.6-10 and discussed here as examples of the degree of uncertainty resulting from the use of each method. In the areas marked as "A," sampling stations with chemical concentrations < SQS are surrounded by or adjacent to stations with concentrations > SQS. Within these polygons, the area of

chemical exceedance estimated using IDW is greater than the area estimated using Thiessen polygons. Two of these areas are notable because the cells representing the specific sampling locations where concentration < SQS are estimated as locations of chemical exceedances by IDW. The areas marked as "B" are areas estimated to have chemical exceedances using IDW that are smaller than the areas estimated to have exceedances by Thiessen polygons because the concentrations at surrounding points are < SQS. Underestimation of chemical exceedance by IDW relative to Thiessen polygon chemical exceedance estimation is less common throughout the LDW because of the positive skew of chemical concentrations and the fixed nature of exceedance criteria. Extreme values have a significant effect on the moving average of the IDW surface, affecting a relatively large area regardless of nearby samples. For Thiessen polygons, the effect of extreme concentration values is limited to the boundaries of the polygons associated with the extreme values.

Effects Assessment

The uncertainty in the effects assessment for the benthic invertebrate community was associated with the use of SMS chemical criteria, DMMP guidelines, or TRVs to assess the potential for a biological effect. Uncertainties associated with using an alternative approach (a tissue-residue approach) to evaluate PAHs and PCBs were also considered as another line of evidence. These uncertainties are discussed below.

Use of SMS criteria, DMMP guidelines, and TRVs to assess biological effects—The likelihood of adverse effects to benthic organisms from chemicals associated with sediments was assessed using two approaches. In the first approach, surface sediment chemical concentrations were compared to SQS/CSL chemical criteria, DMMP guidelines (nickel), or TRVs (total DDTs and total chlordane). In the second approach, site-specific sediment toxicity test results were compared to SMS biological effects criteria.

The chemical criteria, guidelines, and TRVs used in the first approach were based on test species that represent a small portion of the diverse benthic invertebrate community present in the LDW, although the test species included crustaceans, which are considered to represent one of the taxonomic groups most sensitive to chemical exposure (Hyland et al. 1999). In addition, the benthic invertebrate community AETs, which were the basis for several SQS or CSL chemical criteria, incorporated invertebrates with different feeding strategies and habitat requirements and therefore represent COPC concentrations likely to be protective of the benthic invertebrate community as a whole. There is some uncertainty associated with the benthic invertebrate community AETs because these values are based on the abundance of several major benthic infaunal taxa (mollusks, crustaceans, polychaetes) and do not address the potential for effects on species diversity, relative abundances of different taxa, the success of rare species, and other benthic invertebrate community metrics. Thus, potential effects on some LDW benthic species may not be addressed by these

criteria and guidelines; consequently, there is some uncertainty associated with the risk estimates.

The SMS criteria in WAC 173-204 were developed for application to marine sediments. The LDW is an estuarine system with both spatial and temporal variation in salinity. Salinity in the porewater of the surface sediment samples collected for toxicity testing between RM 0.0 and RM 4.9 ranged from 20 to 30 ppt. Thus, although all sediments within LDW are not fully marine, salinity is relatively high because of salt wedge intrusion, and SMS criteria should be applicable.

A fundamental distinction between the two approaches used to estimate risks to the benthic invertebrate community is that one approach (i.e., use of chemical criteria and guidelines) estimates effects based on a comparison to detected chemical concentrations, while the other approach (i.e., sediment toxicity tests) assesses the potential for effects directly through the exposure of test organisms to surface sediment samples collected from the LDW. SMS chemical criteria and some of the DMMP guidelines were developed using the AET approach described in Section A.3.2.1. An AET is the highest "no effect" chemical-specific sediment concentration above which a significant adverse biological effect always occurred among the several hundred samples used for its derivation.

Note that SMS chemical criteria were developed for specific chemicals based on AETs empirically derived from a dataset of field-collected sediment samples that contained diverse chemical mixtures and that were analyzed for both chemistry and toxicity. Therefore, the AETs do not reflect a cause-and-effect relationship for specific chemicals.

Two published studies have assessed the ability of selected AETs to estimate adverse effects in Puget Sound (Barrick et al. 1988; Gries and Waldow 1996). The study by Barrick et al. (1988) calculated overall reliability values[69] between 50 and 96% for benthic, amphipod, Microtox, and oyster larvae AETs. The study by Gries and Waldow (1996) calculated overall reliability values between 65 and 85% for amphipod and echinoderm AETs.

For the LDW, sediment toxicity test results from the 46 Phase 2 locations in the baseline ERA dataset were compared to estimates of toxicity made using only chemical criteria (Table A.6-2). Seven locations assessed by King County (2000a) were also included in the comparison. Site-specific sediment toxicity tests were conducted with *Eohaustorius estuarius* (amphipod), *Neanthes arenaceodentata* (polychaete), and *Mytilus galloprovincialis* (bivalve) using sediment from 46 Phase 2 surface sediment locations within the baseline dataset. The King County sediment toxicity tests were

[69] Overall reliability was calculated as the percentage of all "hit" (i.e., > SQS biological effects criteria) and "no hit" (i.e., ≤ SQS biological effects criteria) samples that were correctly predicted, and thus did not distinguish between SQS and CSL levels of toxicity.

conducted with *Rhepoxynius abronius* (amphipod), *Neanthes arenaceodentata* (polychaete), and echinoderm (*Dendraster excentricus*). Toxicity test responses were evaluated using the toxicity test rules established by SMS to classify responses as exceeding either the SQS (WAC 173-204-320(3)) or the CSL (WAC 173-204-520(3)) (for further details, see Section A.3.2.2).

If the individual SMS classifications are compared considering the level of toxicity (i.e., SQS or CSL), then the chemical and biological criteria concurred at 18 of the 53 locations tested (34%). If the comparison is done by counting the number of locations where the biological effects and chemical predicted either "hits" (either SQS or CSL exceedances) or "no hits" (no SQS exceedances), then the chemical and biological criteria concurred at 31 of the 53 locations tested (58%).

Tissue residue approach for PAHs and PCBs. The numeric standards of SMS and site-specific sediment toxicity tests were used to assess the risks from PAHs and PCBs to the benthic invertebrate community. EPA and Ecology requested that an alternative approach, the tissue-residue approach, also be evaluated in the uncertainty analysis for both PAHs and PCBs. This approach is presented below.

Three studies with mollusks (Borchert et al. 1997; Eertman et al. 1995; Roper et al. 1997) evaluated tissue concentrations of individual PAHs (fluoranthene and benzo(a)pyrene) or a mixture of PAHs associated with adverse effects. Concentrations of benzo(a)pyrene in all benthic invertebrate tissue (market basket samples) did not exceed the LOAEL of 302 µg/kg ww (Eertman et al. 1995), and concentrations of LPAH and HPAH in LDW benthic invertebrate tissue (market basket samples) did not exceed the NOAELs based on a mixture of PAHs (Roper et al. 1997). Fluoranthene in LDW benthic invertebrate tissue (market basket samples) exceeded the LOAEL of 222 µg/kg ww (Eertman et al. 1995) at five locations (B6a, B8a, B3b, B4b, and B6b) in the LDW. Sediment concentrations of several chemicals, including PCBs, PAHs, and metals, exceeded their SMS criteria (PCBs exceeded the SQS and CSL, PAHs exceeded the SQS, and metals exceeded the SQS and CSL) at these five locations, indicating that all locations with potential effects were identified by the primary method for the assessment of risks to the benthic invertebrate community in this ERA.

Seven studies (Boese et al. 1995; Duke et al. 1970; Hansen et al. 1974b; Lowe et al. 1972; Nimmo et al. 1974; Peterson et al. 1994; Sanders and Chandler 1972) evaluated the adverse effects of PCBs on decapods or mollusks. The lowest LOAEL was 1,100 µg/kg ww, reported for Aroclor 1016. Total PCB concentrations in benthic invertebrate tissue samples at two intertidal locations (1,700 µg/kg ww at B5a-2 and 5,900 µg/kg ww at B8a) exceeded this LOAEL, although it should be noted that Aroclor 1016 was never detected in LDW benthic invertebrate tissue samples. Concentrations of several other chemicals exceeded their SMS criteria at these two locations as well (PCBs exceeded the SQS and CSL, PAHs exceeded the SQS, and metals exceeded the SQS and CSL).

Therefore, given both the limits of available TRVs and the fact that tissue concentrations were generally below the available critical tissue residue-based TRVs

for PAHs and PCBs, use of this alternative method for risk evaluation would not have estimated the potential for effects resulting from PAHs or PCBs at additional locations. The absence of TRVs for other chemicals that might occur in the LDW prevents a full exploration of risks to the benthic invertebrate community with this method. Application of the SMS to identify COPCs and areas of concern for the benthic invertebrate community is likely to have captured any areas or chemicals presenting risk to the benthic invertebrate community.

Risk Conclusions

The potential for adverse effects on benthic invertebrate communities was evaluated based on site-specific toxicity tests and on comparisons of chemical concentrations in surface sediment to SQS/CSL chemical criteria, toxicologically based DMMP guidelines, and TRVs. The potential adverse effects included in the existing criteria are reduced survival, abnormal development, and reduced growth at the individual level and altered ecological function at the community level.

In Phase 2, toxicity was tested using three toxicity tests at 48 point locations within the LDW. Nearly all of these locations were selected for toxicity testing because of exceedances of the SQS or CSL chemical criteria.[70] Based on SMS rules, 18 of the 48 sediment samples (37.5%) did not exceed the SQS biological effects criteria, 11 sediment samples (22.9%) exceeded the SQS biological effects criteria, and 19 samples (39.6%) exceeded the CSL biological effects criteria. Because these tests are direct measures of effect, the uncertainty in their interpretation is low at the specific locations tested.

At other point locations in the LDW with chemical concentrations greater than SQS or CSL chemical criteria and other guidelines, the estimation of effects is more uncertain. According to SMS, locations with all chemical concentrations less than or equal to the SQS chemical criteria are defined as having no acute or chronic adverse effects on biological resources, locations with any chemical concentrations greater than CSL chemical criteria are defined as having adverse effects, and locations with any chemical concentrations between the SQS and CSL chemical criteria have adverse effects between these two definitions.

The potential for adverse effects is more uncertain at locations where no detected chemicals exceeded the SQS/CSL chemical criteria or DMMP guidelines but RLs were greater than criteria and guidelines. However, based on an analysis of these elevated

[70] Two locations with no chemical concentrations greater than SQS chemical criteria were selected for toxicity testing at the request of Ecology because they were located near specific potential sources. In addition, toxicity testing was also conducted at one location (B6A) without chemical concentrations greater than criteria based on the rounding rules approved for this baseline surface sediment dataset. This sample was tested for toxicity because using pre-baseline rounding rules, the concentration of PCBs was greater than the SQS chemical criteria.

RLs, the likelihood of risks from non-detected chemicals with RLs that exceeded their respective SQS chemical criteria is low.

Thiessen polygons were used to estimate the areal extent of potential effects based on combined toxicity test results and surface sediment chemistry data. Using this approach, the results indicated that:

- No adverse effects to benthic invertebrates living in intertidal and subtidal sediments are expected for 75% of the LDW area (i.e., the area in which chemical concentrations were less than or equal to the SQS chemical criteria and where sediments were non-toxic according to SQS biological effects criteria).
- There is a higher likelihood for adverse effects in approximately 7% of the LDW area, which was designated as having chemical concentrations or biological effects in excess of the CSL criteria.
- The remaining 18% of the LDW area had chemical concentrations or biological effects between the SQS and CSL criteria, indicating that risks are less certain in these areas than in areas with chemical concentrations greater than one or more CSL values.

There is some uncertainty associated with these area estimates because areas were estimated by interpolating individual points at which sediments were sampled. The spatial extent of individual samples exceeding chemical and biological effects criteria is relevant to the assessment of overall risks to the benthic invertebrate community, both as an ROC and as a food resource. Uncertainty in the areal extent of effects increases as the size of the polygon increases. In some locations, groups of samples exceeded the SQS/CSL chemical criteria and in other locations only an isolated sample exceeded the chemical criteria. In areas with isolated exceedances, the potential impact to the ecological function of the benthic invertebrate community is likely to be less than in areas with numerous locations with exceedances.

A.6.1.1.2 Porewater

This section presents the risk estimates, uncertainties, and risk conclusions for benthic invertebrates exposed to VOCs in porewater.

Risk Estimates

One COPC was identified based on the exposure of benthic invertebrates to VOCs in porewater (cis-1,2-dichloroethene). All other VOCs detected in the porewater samples were at concentrations below levels of concern. COPCs were identified based on a comparison of maximum VOC concentrations detected in porewater samples at two sites (i.e., GWI and Boeing Plant 2/Jorgensen Forge) in the LDW to no-effects concentrations (Section 2.5.1.2). These two sites were selected to represent worst-case exposure areas (Windward 2005i).

The maximum porewater concentration of cis-1,2-dichloroethene (2,900 µg/L), detected in peeper PE-06 (Windward 2005a), was approximately 21 times higher than

the estimated NOEC TRV but was less than the LOEC TRV (Table A.6-7). Other peepers in the vicinity of PE-06 (i.e., PE-03, PE-04, and PE-05) had lower concentrations of cis-1,2-dichloroethene (i.e., 0.5, 2.4, and 630 µg/L, respectively), with only PE-05 exceeding the NOEC TRV. Thus, the spatial extent of groundwater discharge of this chemical into porewater appears to be highly localized. This finding is consistent with the proposed conceptual site model for discharge at the GWI site (Windward 2005i).

Table A.6-7. HQs for and benthic invertebrates exposed to cis-1,2-dichloroethene in porewater

Peeper ID	cis-1,2-Dichloroethene Concentration in Porewater			NOEC-Based HQ	LOEC-Based HQ
	Detected Concentration (µg/L)	NOEC TRV (µg/L)[a]	LOEC TRV (µg/L)[b]		
PE-01	1.6	136	6,785	0.01	< 0.01
PE-02	46	136	6,785	0.34	< 0.01
PE-03	0.5	136	6,785	< 0.01	< 0.01
PE-04	2.4	136	6,785	0.02	< 0.01
PE-05	630	136	6,785	**4.6**	0.09
PE-06	2,900	136	6,785	**21**	0.43
PE-07	18	136	6,785	0.13	< 0.01
PE-08	20[c]	136	6,785	0.15	< 0.01

[a] NOEC was calculated from LC50-based LOEC by dividing by an uncertainty factor of 50.

[b] Concentration is based on an LC50.

[c] Field replicates for PE-08 had detected concentrations of 41 and 27 µg/L.

HQ – hazard quotient

ID – identification

LC50 – concentration that causes the death of 50% of a group of test animals

LOEC – lowest-observed-effect concentration

NOEC – no-observed-effect concentration

TRV – toxicity reference value

Bold identifies NOEC-based HQs greater than 1.0 and LOEC-based HQs greater than or equal to 1.0.

Uncertainties Associated with Porewater Risk Estimates

This section presents uncertainties in the porewater risk characterization for the benthic invertebrate community. The uncertainties are discussed separately for the exposure and effects assessments.

Exposure Assessment

The assessment of porewater-associated chemicals focused on two areas, GWI (RM 2.3 to RM 2.4) and Boeing Plant 2/Jorgensen Forge (RM 3.5 to RM 3.6). These locations were selected as worst-case exposure scenarios based on an evaluation of groundwater VOC data at high-priority sites identified by EPA and Ecology in Phase 1 (Windward 2005i). The groundwater conceptual site models for these locations showed that fresh

groundwater migrating beneath upland areas is likely to discharge primarily into intertidal areas. Analyses of porewater samples collected with piezometers in these areas were consistent with the models. Therefore, even though there is uncertainty associated with the identification of areas with groundwater discharge and resulting porewater exposures, based on this evaluation, the exposure areas sampled with peepers were likely to be the worst-case exposure areas.

Effects Assessment

The primary uncertainty in the porewater assessment is the limited amount of relevant toxicity data available in the literature (i.e., LOECs were rarely reported, NOECs were dependent on the selected test dilution series, and very few studies reported both effect and no-effect concentrations for a single species and endpoint). In addition, no toxicological data were available for isopropylbenzene, and few data were available for 1,1-dichloroethane and vinyl chloride. The NOECs and LOECs for 1,1-dichloroethane and vinyl chloride used in the COPC screen in Section A.2.5.1.2 were obtained by calculating acute and chronic values using the narcosis model from DiToro et al. (2000). Because the calculation of acute and chronic values in the narcosis model followed the EPA water quality criteria guidelines for the protection of aquatic organisms, it is unlikely that the model underestimated effects to benthic invertebrates from 1,1-dichloroethane and vinyl chloride.

There is also uncertainty in the NOEC TRV calculated for cis-1,2-dichloroethene (the single COPC identified). The NOEC TRV was calculated by dividing the LC50-based LOEC TRV by 50. Thus, there is uncertainty regarding the probability of adverse effects in the immediate vicinity of peepers PE-05 and PE-06.

Risk Conclusions

Risks to benthic invertebrates from exposure to VOCs in porewater are expected to be very low for most VOCs and low for one VOC in a localized area in the intertidal area near GWI. These results are based on a risk assessment at two worst-case scenario locations, GWI and Boeing Plant 2/Jorgensen Forge, where only one chemical, cis-1,2-dichloroethene, was identified as a COPC at GWI. Among the peepers sampled, the highest NOEC-based HQ for this compound was 21; the highest LOEC-based HQ was less than 1.0. The NOEC TRV is uncertain because it was derived from the LOEC using a safety factor of 50.

A.6.1.1.3 TBT

This section presents an evaluation of risks to benthic invertebrates from exposure to TBT in the LDW, including uncertainties and risk conclusions. Risks were assessed in two ways:

- Risks to meso- and neogastropods, invertebrates that have been identified as being particularly sensitive to TBT (Meador et al. 2002), were evaluated through direct measurement of effects (i.e., imposex). TBT exposure was based on the TBT concentration in sediment at locations where gastropods were collected.

- Risks to benthic invertebrates other than gastropods were assessed using a critical tissue-residue approach. TBT exposure was based on TBT concentrations in co-located tissue and sediment samples.

Imposex

The potential adverse effects from TBT exposure were assessed for meso- and neogastropods through the direct measurement of imposex in field-collected gastropods from the LDW in two separate investigations. The relative sensitivity of the specific neo- and mesogastropod species found in the LDW to TBT has not been evaluated in the literature.

Meso- and neogastropod imposex data were presented in the *Technical Memorandum: Gastropod Pilot Survey Results* (Windward 2004g) and the *Technical Memorandum: 2005 Gastropod Imposex Study Results* (Windward 2006c) and were summarized in Section A.3.2.4 of this ERA. Imposex was not observed in two of the three neogastropod species examined or in the two mature female mesogastropods collected in the LDW (12 other mesogastropod specimens were collected, but they were either males or immature) (Windward 2004f, 2006c). Imposex was not observed in the abundant neogastropods *Astyris gausapata* or *Olivella baetica*, including specimens collected from areas within the LDW that historically have had high TBT concentrations in surface sediment.

Nassarius mendicus was the only gastropod species among those collected that showed imposex symptoms. All collected females of this species were classified as Stage 2, specifically Stage 2a (large penis with penis duct), according to the methods of Oehlmann et al. (1991). None of the affected specimens had an evident vas deferens. The calculated RPS indices for females of this species ranged from 0.2 to 3.4%. These RPS indices are all below thresholds associated with sterility in female neogastropods (Spence et al. 1990). In all other gastropods that have been studied in Great Britain and other locations, Stage 2 imposex has not interfered with female reproduction (Spence et al. 1990; Gibbs and Bryan 1996).

There is uncertainty in the assessment because: 1) neo-and mesogastropod species may already have been affected by TBT and therefore are no longer present in the LDW, and 2) low numbers of gastropods were collected in the LDW. One to four specimens of *Nassarius mendicus*, the only neogastropod showing imposex characteristics, were collected at five out of the six sampling locations; and females were collected at three of the six locations. No *N. mendicus* females were collected at the two locations with the highest TBT concentrations or at the location with the lowest TBT concentration.

In addition, low numbers (one to six) of specimens from the order Mesogastropoda were assessed for imposex. Because the majority of these specimens were immature, only two females were evaluated for imposex. Neither of these females showed any signs of imposex.

Based on the results of the site-specific imposex studies, it is probable that the level of imposex observed in *N. mendicus* in the LDW does not have an adverse effect on reproduction. The observed imposex was characterized as Stage 2, a stage that is not expected to affect reproduction. Imposex was not observed in any other neo- or mesogastropod collected. Thus, although there were uncertainties in the field studies, TBT risk to gastropods is considered to be low.

Critical Tissue-Residue Approach

The potential adverse effects of TBT exposure were assessed for benthic invertebrates other than gastropods using a critical tissue-residue approach. TBT concentrations in tissue were calculated from the non-linear regression relationship observed between the TBT concentrations in co-located benthic invertebrate tissue and sediment samples, as described in Section A.3.1.3 and Attachment 11. The non-linear regression equation was also used to estimate the TBT concentration in tissue that may be associated with the maximum TBT concentration in surface sediment. The estimated maximum TBT concentration in tissue was 0.61 mg/kg dw. The maximum TBT concentration detected in tissue was 0.55 mg/kg dw. All of the TBT concentrations detected or estimated in benthic invertebrate tissue were less than the NOAEL TRV of 0.97 mg/kg dw (Table A.6-8).

Table A.6-8. TBT risk estimates for benthic invertebrates using the critical tissue-residue approach

Organism	TBT Concentration in Tissue			NOAEL-Based HQ	LOAEL-Based HQ
	Exposure Concentration (mg/kg dw)	NOAEL TRV (mg/kg dw)	LOAEL TRV (mg/kg dw)		
Benthic invertebrates	0.55[a]	0.97	2.36	0.57	0.23
	0.61[b]	0.97	2.36	0.63	0.26

[a] Maximum concentration in benthic invertebrate tissue samples collected from the LDW.

[b] Maximum concentration in benthic invertebrate tissue estimated from the maximum TBT concentration in sediment (3 mg/kg dw) and the non-linear regression equation.

dw – dry weight
HQ – hazard quotient
LOAEL – lowest observed adverse effect level
NOAEL – no observed adverse effect level
TBT – tributyltin
UCL – upper confidence limit on the mean

The largest source of uncertainty associated with the critical tissue-residue evaluation of risk to benthic invertebrates is the TRV value. The selected TRV was based on the response of a single species, the polychaete *Armandia brevis,* in a spiked sediment bioassay. This polychaete is found in marine intertidal mud flats and would not be expected to be present in substantial numbers in the LDW. This species has been shown to bioaccumulate TBT with relatively little ability to metabolize TBT (Meador 1997; Meador et al. 1997). Meador et al. (1997) reported significant differences in TBT accumulation and depuration between *Armandia brevis* and two amphipods, *Rhepoxynius abronius* and *Eohaustorius washingtonianus*. Of the amphipods tested,

E. washingtonianus was most sensitive to sediment-associated TBT, and *R. abronius* was the least sensitive.

The TRV was compared to exposure concentrations represented by benthic invertebrate tissue samples, which contained a wide range of species, including amphipods, mollusks, and polychaetes. It is unknown how similar the species present in these samples are to *Armandia brevis* in terms of their ability to accumulate TBT or their sensitivity to the effects of TBT.

Locations of the co-located benthic invertebrate tissue and sediment samples analyzed to derive the regression relationship were selected to represent the range of TBT concentrations throughout the LDW. The co-located sediments represent 99% of the range of TBT concentrations found throughout the LDW. The use of *in-situ* sediment and tissue concentrations to derive the regression relationship results in reduced uncertainty with regard to the estimate of exposure when compared to using a sediment/tissue relationship from the literature.

The NOAEL-based HQ for benthic invertebrates was less than 1.0. Thus, although there were uncertainties in this analysis, risks to the benthic invertebrate community from exposure to TBT are considered to be very low.

A.6.1.2 Crabs

This section presents the risk estimates, uncertainties, and risk conclusions for crabs.

A.6.1.2.1 Risk estimates

In this section, risks to crabs from exposures to two COPCs, zinc and PCBs, are assessed using a critical tissue-residue approach. Risks were assessed separately for crabs using a critical tissue-residue approach because crabs are more mobile than infaunal organisms, are not specifically covered by SMS criteria, and have a greater potential for exposure through bioaccumulation because of their higher trophic position. HQs were calculated for zinc and PCBs based on the UCL concentration of these COPCs in whole-body crab tissue relative to tissue-based NOAEL and LOAEL TRVs (Table A.6-9). The NOAEL-based HQ was greater than 1.0 for both COPCs (2.5 for zinc and 10 for total PCBs). LOAEL-based HQs were less than 1.0 for zinc and equal to 1.0 for total PCBs.

Table A.6-9. HQs for crabs using whole-body exposure and effects data

Chemical	Unit	Whole-Body Tissue Concentration			NOAEL-Based HQ	LOAEL-Based HQ
		UCL[a]	NOAEL TRV	LOAEL TRV		
Zinc	mg/kg ww	32	12.7	35.2	**2.5**	0.91
Total PCBs	µg/kg ww	1,100	110[b]	1,100	**10**	**1.0**

[a] Whole-body concentrations were estimated by combining hepatopancreas and edible-meat concentrations, assuming 69% by mass edible meat and 31% by mass hepatopancreas.

[b] Calculated by dividing the LOAEL by 10.

HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
UCL – upper confidence limit on the mean
TRV – toxicity reference value

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

A.6.1.2.2 Uncertainty associated with crab tissue risk estimates

This section presents specific areas of uncertainty in the crab risk estimates related to the problem formulation, exposure assessment, effects assessment, and risk characterization.

Problem Formulation

Crabs were selected as an ROC to represent higher-trophic-level benthic invertebrates not addressed by the SMS. There is uncertainty associated with the assumption that COPC concentrations in crab tissue would represent those of other mobile, higher-trophic-level benthic invertebrates in the LDW, which include sea stars and shrimp. Dungeness crabs are scavengers; their diet includes shrimp, mussels, small crabs, clams, and sea urchins. Thus, crabs are likely to be similarly exposed through their diet as sea stars and shrimp, which have comparable diets. Thus, there is relatively little uncertainty in using crabs as representatives of larger, more mobile benthic invertebrates.

The COI screening process is also uncertain because the available toxicity studies for decapods investigated only survival or growth endpoints. Toxicity studies using reproductive endpoints would potentially be more sensitive and thus could have identified additional COPCs had they been available.

Exposure Assessment

Because tissue was used to estimate crab exposure, all potential exposure pathways were integrated. However, because crabs' home ranges can include areas outside of the LDW, the concentrations of COPCs in crab tissue may not be fully reflective of LDW exposure.

The EPCs, which were estimated by combining the Dungeness and slender crab tissue data, are uncertain because the approach assumes similar whole-body concentrations for the two crab species. Combining crab tissue samples likely has little effect on the assessment of risk from zinc because zinc concentrations are similar between the two species. Whole-body concentrations for Dungeness crabs and slender crabs ranged from 22.7 to 33 mg/kg ww and from 33.6 to 35.9 mg/kg ww, respectively. In contrast, combining crab tissue samples may slightly underestimate the risk from PCBs because Dungeness crabs have slightly higher concentrations. Whole-body PCB concentrations for Dungeness crabs and slender crabs ranged from 420 to 1,900 µg/kg ww and from 250 to 838 µg/kg ww, respectively.

There is also uncertainty associated with the LDW crab whole-body tissue-residue data because whole-body concentrations were estimated based on chemical

concentrations in edible meat and the hepatopancreas. It is unknown if whole-body concentrations estimated from these data result in an overestimate or underestimate of the actual tissue residue of COPCs in crabs.

Effects Assessment

The primary uncertainty in the crab effects assessment is the limited number of tissue-based TRVs available in the literature. Effects data for decapods were found for only 14 of the 54 COIs detected in LDW crab tissue. Furthermore, most of these toxicity studies investigated the survival endpoint rather than the potentially more sensitive sublethal endpoints. In some of the studies, tests were conducted only with adults, although juveniles or early life stages may be more sensitive than adults. Additional uncertainties with these studies were associated with exposure durations, exposure pathways (water exposure vs. dietary exposure), and test organism used (decapods other than crabs). The relative uncertainties in the selected TRVs for crabs and the potential effect on the risk estimates are summarized in Table A.6-10.

Table A.6-10. Level of uncertainty associated with crab TRVs

Chemical	Number of Toxicity Studies	Uncertainty in TRV
Zinc	1	High TRV uncertainty because of small dataset with only acute studies that assessed mortality; whole-body tissue concentrations were estimated.
Total PCBs	4	High TRV uncertainty because of small dataset with only acute studies that assessed mortality; whole-body tissue concentrations were estimated.

PCB – polychlorinated biphenyl

TRV – toxicity reference value

The selected LOAEL for zinc and total PCBs and the selected NOAEL for zinc were based on acute exposure studies that assessed survival, which is a less-sensitive endpoint than growth or reproductive endpoints; the NOAEL TRV for PCBs was derived from the LOAEL by dividing by an uncertainty factor of 10. No uncertainty factor was applied to the acute LOAELs to derive chronic LOAELs; therefore, risks from zinc and PCBs may be underestimated.

Arsenic was identified as a COPC for crabs but was not evaluated in the exposure and effects assessments because a LOAEL was not available. The NOAEL-based HQ was 8.6. At EPA's request, a LOAEL for arsenic was derived by multiplying the NOAEL by an uncertainty factor of 10. Using this highly uncertain approach,[71] the LOAEL-based HQ would be 0.86, indicating low risk of adverse effects to crabs.

[71] In contrast to the application of uncertainty factors to LOAELs to estimate NOAELs, no information is available to estimate a LOAEL from a single NOAEL.

Risk Characterization

Total DDTs and methoxychlor were identified as COPCs because their estimated maximum concentrations in crab tissue exceeded the selected NOAELs (Section A.2.5.1.3). Risks to crabs from total DDTs and methoxychlor were not included in the risk estimates because of high uncertainty in the JN-qualified organochlorine pesticide tissue data, which resulted in suspected false identifications of the presence of some organochlorine pesticides as well as overestimates in their concentrations, as discussed in Section A.2.4.2.2. The exposure and effect assessments as well as the risk estimates for these two organochlorine pesticides are discussed in this section.

Table A.6-11 presents whole-body tissue-residue concentrations for total DDTs and methoxychlor in Dungeness and slender crabs using the equation presented in Section A.3.3. Data for Dungeness and slender crabs were combined because insufficient numbers of Dungeness crabs were collected in some of the sampling areas.

Table A.6-11. Estimated total DDT and methoxychlor concentrations in whole-body crab tissue

COPC	Number of Samples	Minimum Concentration (µg/kg ww)	Maximum Concentration (µg/kg ww)	UCL (µg/kg ww)
Total DDTs	19	47.6	150	48
Methoxychlor	19	5.8	90	54

COPC – chemical of potential concern

UCL – upper confidence limit on the mean

ww – wet weight

Three studies assessed the effects of DDTs on decapods (Johnson et al. 1971; Nimmo et al. 1970; Leffler 1975) (Table A.6-12). A mortality rate of 30% was reported in pink shrimp at a whole-body DDT tissue concentration of 60 µg/kg ww after water exposure to DDT and its metabolites for 56 days (Nimmo et al. 1970). No mortality was reported in crayfish exposed to DDT in water for 3 days, resulting in a whole-body DDT tissue concentration of 46 µg/kg ww (Johnson et al. 1971). These two tissue concentrations were selected as the LOAEL and NOAEL TRVs, respectively.

Table A.6-12. Total DDT and methoxychlor critical tissue-residue toxicity studies for crabs and other decapods

Chemical	Test Species	Exposure Conditions	NOAEL (µg/kg ww)	LOAEL (µg/kg ww)	Effect	Source
Total DDTs	crayfish (*Orconectes nais*)	0.8 µg/L in water for 3 days	**46**	na	reduced survival	Johnson et al. (1971)
	pink shrimp (*Penaeus duorarum*)	0.05 µg/L in water for 56 days	na	**60**	reduced survival	Nimmo et al. (1970)
	juvenile blue crab (*Callinectes sapidus*)	dietary exposure for 5 weeks	26	200	increased metabolic rate	Leffler (1975)
Methoxychlor	juvenile Dungeness crab (*Cancer magister*)	0.04 µg/L in water for 18 days	**15**[a]	**150**	reduced survival	Armstrong et al. (1976)
	larval blue crab (*Callinectes sapidus*)	0.7 and 1.0 µg/L in water for a mean of 63 days	< 100	150	reduced survival	Bookhout et al. (1976)
	larval mud crab (*Rhithropanopeus harrisii*)	2.5 µg/L in water for a mean of 20 days	< 100	230	reduced survival	Bookhout et al. (1976)
	adult Dungeness crab (*Cancer magister*)	7.5 µg/L in water for up to 15 days	na	570	reduced survival	Armstrong et al. (1976)

[a] Calculated from LOAEL by dividing by 10.

na – not available

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the NOAELs and LOAELs selected as the TRVs.

Two studies assessed the effects of methoxychlor on decapods (Armstrong et al. 1976; Bookhout et al. 1976) (Table A.6-12). Armstrong et al. (1976) assessed the effects of methoxychlor on juvenile and adult Dungeness crabs. After 36 days of exposure to 0.04 µg/L of methoxychlor in water, juvenile crabs had an approximate 20% increase in mortality over the controls. A similar increase in mortality over the controls was observed in juvenile crabs after an 80-day exposure to 0.04 µg/L of methoxychlor in water. Tissue concentrations were not analyzed in these experiments. However, after exposing juvenile crabs to 0.04 µg methoxychlor/L in water for 18 days, the whole-body concentration was 150 µg/kg. In a similar experiment, adult crabs exposed to 0.4 and 4.0 µg/L of methoxychlor in water for 85 days had survival rates of 100% and 0%, respectively. Concentrations of methoxychlor in tissue were not detected in adult crabs exposed to these concentrations. However, increased mortality was observed in adult crabs exposed to 7.5 µg/L methoxychlor in water for 15 days, and a concentration of 570 µg/kg ww was detected in crab tissue.

Bookhout et al. (1976) assessed the effects of methoxychlor on the larval development of mud and blue crabs. Significant mortality was reported in mud crabs exposed to methoxychlor through the five larval developmental stages. At the end of the last larval stage, an 18% increase in mortality over the controls was reported in organisms, with a whole-body concentration of 230 µg/kg ww. Similarly, significant mortality was reported for blue crabs exposed to methoxychlor through the nine larval developmental stages. High mortality was reported among the later larval stages in the controls. At the end of the last larval stage, a 27% increase in mortality over the controls was reported in organisms, with a whole-body concentration of 150 µg/kg ww. The lowest LOAEL of 150 µg/kg ww reported in the two studies was selected as the LOAEL TRV. Because none of the studies provided a usable NOAEL, a NOAEL of 15 µg/kg ww was calculated using an uncertainty factor of 10 (because the LOAEL was based on an acute endpoint). The NOAELs and LOAELs for total DDTs and methoxychlor were based on survival, which is a less sensitive endpoint than growth or reproduction.

HQs were calculated for total DDTs and methoxychlor based on the UCL concentrations of these COPCs in whole-body crab tissue relative to tissue-based NOAEL and LOAEL TRVs (Table A.6-13). Because of suspected false identifications of the presence of some organochlorine pesticides as well as overestimates in their concentrations, the tissue concentrations are highly uncertain and biased high (Section A.2.4.2.2).

Table A.6-13. Crab HQs using whole-body exposure and effects data

COPC	Whole-Body Tissue Concentration (µg/kg ww)			NOAEL-Based HQ	LOAEL-Based HQ
	Exposure Concentration	NOAEL TRV	LOAEL TRV		
Total DDTs	48	46	60	**1.0**	0.80
Methoxychlor	54	15[a]	150	**3.6**	0.36

[a] Calculated from LOAEL by dividing by 10.

COPC – chemical of potential concern

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

UCL – upper confidence limit on the mean

ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

The NOAEL-based HQ was greater than 1.0 for methoxychlor and equal to 1.0 for total DDTs. LOAEL-based HQs were less than 1.0 for both methoxychlor and total DDTs. The TRVs were based on survival following acute exposures, and the exposure estimates for these organochlorine pesticides were biased high. Therefore, risks to crabs from these pesticides are uncertain.

A.6.1.2.3 Risk conclusions

Risks to crabs from sediment-associated COPCs in the LDW appear to be low. While NOAEL-based HQs were 2.5 for zinc and 10 for total PCBs, the LOAEL-based HQs were less than or equal to 1.0 for the two COPCs (Table A.6-14). Because the true threshold for effects falls somewhere between the NOAEL and the LOAEL and because of the limited toxicity data, risks to crabs are possible but likely to be low because no LOAELs were exceeded.

Table A.6-14. Summary of risk characterization for crab

Chemical	NOAEL-Based HQ	LOAEL-Based HQ	Effect	Primary Uncertainty
Zinc	**2.5**	0.91	reduced survival	High TRV uncertainty because of small dataset with only acute studies; whole-body tissue concentrations were estimated.
Total PCBs	**10**	**1.0**	reduced survival	High TRV uncertainty because of small dataset with only acute studies; whole-body tissue concentrations were estimated.

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

TRV – toxicity reference value

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

HQs for total DDTs and methoxychlor were greater than or equal to 1.0 based on the NOAEL TRV, but less than 1.0 based on the LOAEL TRV (0.80 and 0.36, respectively). Therefore, risks for these pesticides are uncertain. Arsenic had a NOAEL-based HQ of 8.6, but no LOAEL-based TRV was available. Without a LOAEL, effects from arsenic cannot be assessed.

A.6.1.3 Summary of risk conclusions for benthic invertebrates

In summary, results of the benthic invertebrate risk estimates and evaluation of associated uncertainties are as follows:

Benthic invertebrate community. The results of the sediment risk characterization, which used Thiessen polygons and the combination of chemistry and toxicity test results, indicated that:

- No adverse effects to benthic invertebrates living in intertidal and subtidal sediments are predicted for 75% of the LDW area (i.e., the area in which chemical concentrations were less than or equal to the SQS chemical criteria and where sediments were non-toxic according to SQS biological effects criteria).
- There is a higher likelihood for adverse effects in approximately 7% of the LDW area, which was designated as having chemical concentrations or biological effects in excess of the CSL criteria.
- The remaining 18% of the LDW area had chemical concentrations or biological effects between the SQS and CSL criteria, indicating that risks are less certain in these areas than in areas with chemical concentrations greater than one or more CSL values.

There is some uncertainty associated with these area estimates because areas were estimated by interpolating individual points at which sediments were sampled. The spatial extent of individual samples exceeding criteria is relevant to the assessment of overall risks to the benthic invertebrate community, both as an ROC and as a food resource. Uncertainty in the areal extent of effects increases as the size of the polygon increases. In some locations, groups of samples exceeded criteria and in other locations only an isolated sample exceeded criteria. In areas with isolated exceedances, the potential impact to the ecological function of the benthic invertebrate community is likely to be less than in areas with numerous locations with exceedances.

The potential for adverse effects is more uncertain at locations where no detected chemical concentrations were greater than SQS chemical criteria or DMMP guidelines but RLs were greater than criteria and guidelines. However, based on an analysis of these elevated RLs, the likelihood of risks from non-detected chemicals with RLs that exceeded their respective SQS chemical criteria is low.

Based on the porewater risk estimates and uncertainties, risks to benthic invertebrates from exposure to COPCs in LDW porewater are low and localized. Risks to benthic invertebrates from TBT were very low based on NOAEL-based HQs less than 1.0 and

the absence of imposex in all gastropods, except one species of neogastropod with imposex characterized as Stage 2, a stage that is not expected to impact reproduction.

Crabs. Risks to crabs from zinc and total PCBs in the LDW were low based on the critical tissue-residue approach. Risks from arsenic are uncertain because a LOAEL was not available and exposure concentrations were greater than the NOAEL. Risks from pesticides are likely to be low.

A.6.2 RISK CHARACTERIZATION FOR FISH

This section presents a risk characterization and uncertainty analysis for each of the three fish ROCs. The assessment for each ROC estimates risk based on the exposure and effects assessments discussed in Sections A.4.1 and A.4.2, respectively. Following the risk estimates, a detailed evaluation of uncertainty associated with these calculations is presented. Risk conclusions are then presented for each ROC synthesizing risk estimates and uncertainties.

A.6.2.1 Juvenile chinook salmon

This section presents risk estimates, uncertainties, and risk conclusions for juvenile chinook salmon.

A.6.2.1.1 Risk estimates

This section presents the HQ calculations for juvenile chinook salmon. COPCs evaluated for juvenile chinook salmon included arsenic, cadmium, copper, and vanadium, all of which were evaluated using a dietary approach. Two lines of evidence were assessed for this dietary approach: comparison of concentrations in a single composite sample of stomach contents with the TRV, and comparison of estimated prey concentrations using chemistry data for benthic invertebrates (see Section A.4.1.2) with the TRV.

For cadmium, the LOAEL-based HQ was 1.0 based on consumption of benthic invertebrate prey, whereas the LOAEL-based HQ was less than 1.0 based on stomach contents data (Table A.6-15). LOAEL-based HQs were less than 1.0 for all other COPCs. NOAEL-based HQs were greater than 1.0 for arsenic, copper, and vanadium based on consumption of benthic invertebrate prey data but less than 1.0 based on stomach content data. [72]

[72] Vanadium was not analyzed in the composite sample of stomach contents.

Table A.6-15. HQ calculations for juvenile chinook salmon

Approach	COPC	Exposure Concentration (mg/kg dw)	NOAEL TRV (mg/kg dw)	LOAEL TRV (mg/kg dw)	NOAEL-Based HQ	LOAEL-Based HQ
dietary (stomach contents)	arsenic	3.9	20	30	0.20	0.13
	cadmium	0.46	0.1	0.5	**4.6**	0.92
	copper	42	50	100	0.83	0.42
dietary (benthic invertebrates)	arsenic	22	20	30	**1.1**	0.73
	cadmium	0.50	0.1	0.5	**5.0**	**1.0**
	copper	93	50	100	**1.9**	0.93
	vanadium	8.1	2.04	10.2	**4.0**	0.79

COPC – chemical of potential concern
dw – dry weight
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
TRV – toxicity reference value
ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

A.6.2.1.2 Uncertainty analysis

This section presents a discussion of the uncertainty associated with the problem formulation, the exposure and effects assessments, and the risk characterization for juvenile chinook salmon.

Problem Formulation

The primary uncertainties in the problem formulation for juvenile chinook salmon were associated with ROC selection, assessment endpoints, and the COPC screen.

ROC Selection

Uncertainties associated with the representation of other juvenile salmonids in the LDW by juvenile chinook salmon were addressed in the Phase 1 ERA (Windward 2003b). Juvenile chinook salmon are regarded as the most estuarine-dependent juvenile salmonid, so they are likely to have equal or greater exposure to sediment-associated chemicals than other estuarine-dependent salmonids such as juvenile chum salmon. The Phase 1 ERA (Windward 2003a) also concluded that although yearling juvenile chinook salmon may occur in the LDW in limited numbers and eat at a higher trophic level than subyearling chinook salmon (for which tissue data were collected and analyzed), subyearling juvenile chinook salmon are likely to have greater exposure to sediment-associated chemicals than yearling chinook salmon because of their greater residence time in the LDW (subyearling chinook salmon are more estuarine-dependent than are yearling chinook salmon) (Kerwin and Nelson 2000). Therefore, risk estimates for juvenile chinook salmon should be similar to or greater than those for other juvenile salmonids in the LDW.

Assessment Endpoints

Uncertainties associated with how well the endpoints selected to represent potential adverse effects on fish populations in the LDW were addressed in the Phase 1 ERA (Windward 2003b). In that analysis, it was concluded that biomarkers of effect such as cytochrome P450-dependent mono-oxygenase induction (e.g., CYP1a [van der Weiden et al. 1994]), DNA changes (e.g., DNA-xenobiotic adduct formation [Rice et al. 2000]), and sub-organismal immune dysfunction (e.g., antibody formation [Arkoosh et al. 1991]) may provide an early warning of adverse effects on fish at the individual and population levels (e.g., Schmitt et al. 2000). However, from an individual or population level, the overall significance of these changes is unknown. Thus, assessment endpoints were limited to survival and growth in this ERA.[73]

COPC Screen

Uncertainties associated with the COPC screen included a lack of fish tissue data or TRVs for some chemicals detected in sediment and mercury egg LOAELs that were lower than the selected mercury NOAEL TRV. Each of these uncertainties is addressed below.

Of the chemicals screened out as COIs for all fish ROCs using criteria presented in Section A.2.5.2, several chemicals were detected in at least 5% of baseline surface sediment samples but were not analyzed in tissue samples (Table A.6-16). None of these chemicals are defined as bioaccumulative chemicals (EPA 2000a); therefore, risks to fish from exposure to these chemicals are assumed to be very low.

Table A.6-16. Chemicals detected in ≥ 5% of baseline surface sediment samples that were not analyzed in LDW tissue samples

CHEMICALS	
Acetone	Magnesium
Aluminum	Manganese
Barium	Methyl ethyl ketone
Benzaldehyde	Potassium
Beryllium	Retene
Calcium	Sodium
Caprolactam	Tin
Carbon disulfide	Toluene
p-Cymene	Total petroleum hydrocarbons – gasoline, diesel and oil range
Iron	

[73] Effects on reproduction were not evaluated for juvenile chinook salmon because such effects would only be relevant for adult salmon, which acquire nearly all of their body burden of chemical contaminants outside the LDW.

Some chemicals have been detected in LDW sediment and in LDW tissue for which no toxicological data exist. TRVs were available for 47 of the 85 fish COIs. Twenty of these chemicals without toxicological data were PAHs or alkylated PAHs. As discussed in Section A.2.5.2, maximum dietary exposure concentrations for fish (the sum of all alkylated and non-alkylated PAHs) were lower than the available TRVs for benzo(a)pyrene or a PAH mixture. Insufficient effects data were available to evaluate risks to fish from the remaining 18 chemicals.

The NOAEL TRV for mercury used in the COPC screening process was the highest identified whole-body NOAEL below the lowest identified whole-body LOAEL; however, some studies reported adverse effects in fish associated with lower egg or early life stage tissue residues. The lowest whole-body LOAEL was 470 µg/kg ww for reduced survival of male mummichog exposed to dietary methyl mercury for 42 days (Matta et al. 2001). Birge et al. (1979) reported reduced survival of channel catfish embryos exposed to mercury in water associated with a 4-day post-hatch embryo tissue concentration LOAEL of 14 µg/kg ww. Birge et al. (1979) also reported reduced survival of rainbow trout alevins and embryos exposed to mercury at tissue LOAELs of 36 and 41 µg/kg ww, respectively.

Although these egg and embryo effects concentrations were lower than those reported by Matta et al. (2001) in more mature fish, egg/embryo and adult tissue-residue data are not directly comparable. Species-specific ratios relating mercury concentrations in whole bodies of maternal adults to concentrations in unfertilized eggs for yellow perch, smallmouth bass, white bass, white sucker, and rainbow trout ranged from 9.9 to 26 (Niimi 1983). Adjusting for a two-fold weight increase in concentration in fertilized eggs relative to unfertilized eggs (Niimi 1983) results in adult-to-fertilized egg ratios that ranged from 20 to 52 for these species. Therefore, mercury concentrations in whole bodies of adult female fish were likely to be 20 to 52 times higher than concentrations in fertilized eggs from those females. This range is uncertain because available data represent few species, with little or no replication. Based on this range of adult-to-egg mercury ratios, the estimated range of LOAELs in maternal adults associated with the lowest reported egg/embryo LOAEL (14 µg/kg ww) would have been 280 to 728 µg/kg ww. The maximum mercury whole-body fish tissue concentration in the LDW was 39 µg/kg ww; therefore, mercury would have screened out as a COPC for fish even if egg/embryo effects data had been included in the screen.

Exposure Assessment

Uncertainties in the exposure assessment for juvenile chinook salmon were associated with the following factors:

- Water pathway for metals and PAHs
- Dietary composition

- Foraging preferences
- Benthic invertebrate tissue data

Water Pathway for Metals and PAHs

A dietary approach was used in this Phase 2 ERA to evaluate risks to juvenile chinook salmon resulting from exposures to metals and PAHs. This approach does not incorporate exposure from water. To provide an assessment of the risks to fish from metals and PAHs from multiple exposure pathways, this section provides a brief summary of an analysis of water quality (WQA) conducted by King County (King County 1999d), in which risks to fish from estimated water column exposures were evaluated.[74] A more complete summary of the WQA approach was presented in Attachment A.2 of the Phase 1 ERA (Windward 2003b).

Chemical concentrations in surface water in the WQA were estimated based on results of a detailed three-dimensional fate and transport hydrodynamic model calibrated with field data. The model divided the LDW into 129 grid cells, each with 10 water layers and 1 sediment layer, resulting in 1,290 water column and 129 sediment cells. The model estimated chemical concentrations in water in each grid element every 15 minutes for 1 year.

The WQA identified 23 COPCs requiring risk analysis.[75] Risks related to these COPCs were evaluated using a tiered approach. Tier 1 consisted of a comparison of modeled water concentrations to acute and chronic TRVs.[76] For each COPC, the maximum monthly 1-hour moving average and maximum monthly 4-day moving average total recoverable[77] concentration in each cell in the study area for each month of the year were compared to acute and chronic TRVs, respectively. Eight metal and PAH COPCs had concentrations exceeding acute or chronic TRVs in at least one cell, as summarized in Table A.6-17. These COPCs were further evaluated for juvenile chinook salmon by comparing modeled water concentrations to salmonid-specific TRVs. COPCs were subsequently evaluated for all aquatic species in a second tier of the risk assessment.

---

[74] This section discusses only results for the King County (1999d) WQA baseline scenario (includingCSO and stormwater inputs).

[75] Chemicals with the ability to cause cancer in humans were identified as COPCs. For non-carcinogenic chemicals detected more than 5% of the time, COPCs were identified by King County (1999d) based on a comparison of their 95th percentile concentrations in water and sediment to water quality criteria and sediment standards.

[76] TRVs were based on marine Washington State water quality standards or marine federal water quality criteria (WQC) (10 COPCs), toxicity studies from the literature (7 COPCs), and quantitative structure-activity relationships (QSARs) (6 individual PAHs). An uncertainty factor of 20 was applied to the literature- and QSAR-based TRVs by King County (1999d) for this analysis.

[77] Total recoverable concentrations were used as a conservative estimate of bioavailable metals (Prothro 1993).

Table A.6-17. Metals and PAHs with maximum total concentrations in water in any cell exceeding TRVs following the Tier 1 analysis

COPC	Number of Months in the Year in Which TRV was Exceeded	
	Acute TRV	Chronic TRV
Arsenic[a]	2	0
Copper[a]	12	12
Lead[a]	1	5
Nickel[a]	4	4
Zinc[a]	11	0
Benz(a)anthracene[b]	1	0
Benzo(g,h,i)perylene[b]	0	2
Fluoranthene[b]	1	0

[a] TRVs were based on state and federal marine AWQC for protection of aquatic life (state and federal criteria are the same for each of these chemicals).

[b] TRVs were based on QSARs divided by a safety factor of 20.

COPC – chemical of potential concern

PAH – polycyclic aromatic hydrocarbon

TRV – toxicity reference value

In the second tier of the WQA, with an approach equivalent to the Water Environment Research Foundation (WERF) Tier 3 risk assessment methodology (WERF 1996), risks associated with dissolved COPCs (rather than total recoverable concentrations)[78] were evaluated as the probability of affecting a given percentage of species. A logistic regression model was used to fit genus mean acute and chronic toxicity data (based on dissolved concentrations) for each COPC. This method estimates the percent of taxa affected at any given concentration. For each COPC, the percentage of species affected was estimated for each month based on the modeled maximum 1-hour moving average and 4-day moving average dissolved COPC concentrations in each model cell and layer.

Modeled concentrations of metals did not exceed salmonid-specific TRVs. Results of the second tier of the WQA showed that the percentages of representative species that might be subject to acute and chronic risks in the LDW[79] were less than or equal to 1% for all metals, except copper. Maximum monthly acute and chronic exposures to dissolved copper concentrations were estimated to affect 2 to 4% of aquatic species on the west side of Kellogg Island and less than or equal to 2% in all other locations. Based on an EPA-recommended level of protection of at least 95% of species to ensure overall community function (Stephan et al. 1985, as cited in the King County WQA),

[78] Dissolved metal concentrations were used in the Tier 3 evaluations because they more closely represent the bioavailable fraction (Prothro 1993).

[79] Based on all species represented in toxicity studies included in AWQC plus toxicity studies identified in additional literature searches.

these results indicated low risks to the aquatic community from exposure to surface water concentrations of COPCs; 96% or more of the aquatic community were not expected to be affected by copper.

Risks from water exposure to PAHs were not evaluated in the Tier 3 assessment because insufficient toxicity data were available. Based on the Tier 1 assessment, most of the PAH compounds evaluated had acute and chronic HQs less than 1.0, indicating low risk. However, benzo(a)anthracene and fluoranthene had maximum acute HQs of 1.4 and 1.1, respectively. Acute HQs were greater than 1.0 for benzo(a)anthracene and fluoranthene in one month of the year in 0.08 and 0.4%, respectively, of the cells evaluated (i.e., 1 to 5 of the 1,290 cells evaluated had HQs slightly greater than 1.0). For all other months of the year, acute HQs were below 1.0. Chronic HQs were always below 1.0. For benzo(g,h,i)perylene, the highest chronic HQ was 1.4. In two of the modeled months, chronic HQs for this COPC were greater than 1.0 in 0.2 to 3.6% of the cells evaluated (i.e., 3 to 46 of the 1,290 cells evaluated had HQs slightly greater than 1.0). Therefore, overall aquatic life risks from estimated water column PAH exposures were low.

Methodology to combine water and dietary exposures for fish are not currently available. This topic is under study by EPA but remains an uncertainty at the time of this risk assessment.

Dietary Composition

COPCs in the diet of juvenile chinook salmon are a function of the types of prey consumed and their COPC concentrations. It was assumed that juvenile chinook salmon consume only benthic invertebrates, although they could also consume various zooplankton, larval fish, clam siphons, and terrestrial organisms that drift in the current, such as wasps and ants (Cordell et al. 1996; 1997; 1999). Because benthic invertebrates are more closely associated with sediments than other juvenile chinook salmon prey, dietary exposure concentrations calculated based on benthic invertebrate data are likely to be higher for dietary COPCs (which do not biomagnify). Prey that are less closely associated with sediment (e.g., drift organisms and zooplankton) would likely result in lower overall exposures, and therefore, lower risks from sediment-associated COPCs.

Foraging Preferences

In calculating dietary exposure concentrations, juvenile chinook salmon were assumed to forage in intertidal areas of the LDW. Therefore, intertidal benthic invertebrate and sediment data were used to calculate exposure. Although the literature indicates that juvenile chinook salmon primarily use shallow water areas, no studies have been conducted on the foraging depth of juvenile chinook salmon in the LDW. If juvenile chinook salmon were to forage throughout the LDW, their exposure concentrations would be the same as those calculated for English sole assuming 0% sediment ingestion (Table A.6-18). Different foraging assumptions did not substantially affect

the HQs calculated for juvenile chinook salmon for arsenic, cadmium, or copper (Table A.6-18). For vanadium, the LOAEL-based HQ increased from 0.79 to 1.2.

Table A.6-18. Comparison of juvenile chinook salmon HQs assuming that they forage only in intertidal areas versus throughout the LDW

COPC	Intertidal Only		LDW-Wide	
	NOAEL-Based HQ	LOAEL-Based HQ	NOAEL-Based HQ	LOAEL-Based HQ
Arsenic	**1.1**	0.73	**1.2**	0.80
Cadmium	**5.0**	**1.0**	**6.0**	**1.2**
Copper	**1.9**	0.93	**1.8**	0.92
Vanadium	**4.0**	0.79	**5.9**	**1.2**

COPC – chemical of potential concern

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

The location of fish foraging and prey preference may change as a result of potential ecological improvements associated with future habitat restoration projects. This uncertainty was analyzed in the Phase 1 ERA (Windward 2003b). That analysis concluded that fish exposure to sediment-associated chemicals could either increase or decrease depending on changes in foraging behavior and on prey and sediment chemical concentrations at restored sites.

Benthic Invertebrate Tissue Data

There is uncertainty in the benthic invertebrate tissue data used to estimate dietary exposure of juvenile chinook salmon to some COPCs. Sampling of benthic invertebrate tissue was designed to include stations that represented the full range of concentrations of a subset of chemicals (arsenic, PCBs, and lead) in surface sediments (Windward 2004d). Sediment concentrations of other chemicals were not considered in the sampling design.

To evaluate whether the full range of chemical concentrations was sampled for other COPCs, COPC concentrations in sediment samples co-located with the benthic invertebrate tissue samples were compared to the cumulative distribution of that COPC in the full baseline surface sediment dataset. Results for all dietary COPCs are presented in Table A.6-19. The maximum sediment concentrations reported for the co-located sediment samples were greater than 80% of the site-wide concentrations of these COPCs. Therefore, the co-located sediment and tissue data for these COPCs are likely sufficient to assess the relationship between tissue and sediment concentrations for the site.

Table A.6-19. COPC concentrations in sediment samples co-located with benthic invertebrate tissue samples relative to the site-wide baseline surface sediment dataset for dietary COPCs

COPC	Maximum COPC Concentration in Sediment (mg/kg dw)		Percent of Concentrations in LDW-Wide Dataset Less Than or Equal to the Maximum Concentration in Co-Located Samples
	LDW-Wide Baseline Surface Sediment Dataset	Sediment Samples Co-Located with Benthic Invertebrate Tissue Samples	
Arsenic	1,100	725	99.8
Cadmium	120	1.67	93.7
Copper	12,000 J	495	98.8
Vanadium	150	72.6	83.6

COPC – chemical of potential concern

dw – dry weight

J – estimated concentration

LDW – Lower Duwamish Waterway

As discussed in Attachment 11, benthic invertebrate tissue concentrations were estimated from the linear regression of co-located sediment and benthic invertebrate chemical concentrations for chemicals where a significant sediment-tissue relationship existed. Note that because the regressions were not significant for cadmium, copper, and vanadium, the empirical data, rather than a regression, were used in exposure calculations for these COPCs. A significant sediment-tissue relationship existed for arsenic so benthic invertebrate tissue concentrations were estimated using the linear regression. The arsenic sediment concentrations used in the regression were highly skewed, and the highest values were identified as overly influential points using Cook's distance. However, because the arsenic sediment UCL on the mean concentrations in the fish exposure area were intermediate between the majority of the co-located sediment concentrations and the highly influential points, the influential points were retained in the regression model in order to avoid extrapolation outside the dataset. If the influential points had been excluded from the regression model, the regression would have been linear and would have estimated higher arsenic concentrations. Because none of the co-located sediment samples were similar in concentration to the sediment UCL, the regression is uncertain in this concentration range, and effects on the risk estimates are unknown.

If the exposure estimate had been calculated using empirical intertidal benthic invertebrate tissue data rather than the regression, the UCL would have been 14 mg/kg dw, while the UCL was estimated at 22 mg/kg dw using the regression statistics. Because benthic invertebrate tissue sample locations were selected to be representative of the range of sediment arsenic concentrations in the LDW (Table A.6-19), the regression-estimated UCL likely provides a protective estimate of arsenic concentrations in invertebrates representative of intertidal habitats in the LDW. Therefore, using the regression-estimated UCL to assess exposure likely would not have affected risk conclusions.

There is also uncertainty in the tissue data collected for benthic invertebrates because the tissue samples contained various benthic invertebrate species composited together in order to collect sufficient sample for analysis. Different benthic invertebrate species may bioaccumulate chemicals from sediment to different extents. However, without detailed dietary preference information for the fish ROCs and data for individual invertebrate species, the uncertainty introduced through compositing multiple species is unknown.

Effects Assessment

Uncertainties in the effects assessment for juvenile chinook salmon were associated with the following factors:

- Chemical mixtures
- Exclusion of field studies from TRV selection
- Estimation of NOAELs from LOAELs
- Salmonid-specific TRVs for cadmium
- COPCs without LOAEL TRVs
- Regional field studies

In addition, some of the selected TRVs are considered less certain than others if there were a small number of studies, if endpoints were subchronic, or if data quality was questionable. The relative uncertainties in the selected TRVs for each ROC/COPC pair are summarized below.

NOAEL and LOAEL TRVs for arsenic were selected based on review of six studies, several of which reported similar LOAELs. Because only a moderate number of studies were available, toxicity data may not represent the sensitivities of the various fish species in the LDW. The effect of this uncertainty on risk conclusions is unknown.

NOAEL and LOAEL TRVs for cadmium were selected based on review of nine studies. Although a moderate number of studies were available, TRV uncertainty is high because toxicity data varied greatly among studies, and the selected LOAEL TRV was two orders of magnitude lower than next lowest LOAEL identified. In addition, in the study reporting the lowest LOAEL (Kim et al. 2004; Kang et al. 2005), the growth effect reported was partially attributed to reduced feeding. Effects thresholds from salmonid-specific cadmium toxicity studies were higher. These uncertainties are discussed in greater detail below.

NOAEL and LOAEL TRVs for copper were selected based on a review of 15 studies, a reasonable number of studies. Sufficient data were available to show that the lowest LOAEL reported for growth of channel catfish (8 mg/kg dw) (Murai et al. 1981) was lower than NOAELs for channel catfish growth reported in two other studies (Gatlin and Wilson 1986; Erickson et al. 2003), so this study was rejected as the TRV.

NOAEL and LOAEL TRVs for vanadium were highly uncertain because they were based on only one study. The effect of this uncertainty on risk conclusions is unknown.

Chemical Mixtures

Effects from exposure to multiple chemicals that share the same mode of toxic action and other environmental stressors in the LDW that could result in additive, synergistic, or antagonistic effects were not factored into the effects assessment. This uncertainty was analyzed in the Phase 1 ERA (Windward 2003b). That analysis concluded that because the combined effects of complex chemical mixtures and other stressors occurring in the environment have not been sufficiently studied, effects of this uncertainty on risk predictions are unknown.

Exclusion of Field Studies from TRV Selection

TRVs were derived from laboratory studies in which fish were exposed to a single chemical or, in the case of PCBs, DDTs, and PAHs, to well-defined chemical mixtures under controlled conditions. By using such studies, specific chemical exposures associated with toxicity can be determined. By controlling for natural sources of variability, laboratory studies do not address potential implications associated with mixtures of contaminants or the interaction of chemical toxicity with other stressors that occur in the natural environment. A number of studies in which fish from contaminated sites were raised in the laboratory have been conducted to investigate potential adverse effects associated with sites contaminated with PCBs, DDTs, and PAHs (e.g., Arkoosh et al. 1991; Berlin et al. 1981; Hopkins et al. 1969; Mac et al. 1985). Other studies have exposed fish to field-collected contaminated sediments to investigate potential adverse effects associated with specific mixtures of site chemicals(e.g., Roberts et al. 1989). Such studies incorporate conditions and exposure scenarios that provide insight into risks associated with specific sites and the chemical mixtures present at those sites. However, these studies were not used to derive TRVs in this ERA because adverse effects observed in organisms from studies at other sites may be attributed to the presence of multiple chemicals not present in the LDW and/or to other uncontrolled environmental factors, rather than to a single test chemical.

In the problem formulation, PAHs were screened out as a COPC for fish because maximum dietary exposure concentrations were lower than the lowest dietary LOAELs identified from controlled laboratory studies in which fish were exposed to well-defined PAH mixtures. The analysis presented in the problem formulation did not consider studies with a field component in which fish were exposed to PAHs and other chemicals. Because of the uncertainties inherent in field studies described above, the field component of a dietary PAH exposure study conducted by Rice et al. (2000) and several other PAH sediment exposure studies presented in Table A.6-20 are discussed here to further evaluate uncertainties associated with PAH exposure and effects in fish.

Table A.6-20. Toxicity studies for total PAHs and fish, including studies with a field component

Chemical	Test Species	NOAEL (mg/kg dw)	LOAEL (mg/kg dw)	Exposure Duration	Effect	Source	Notes
Total PAHs (in the presence of a mixture of other chemicals in Eagle Harbor sediments)	English sole	11.3	na	28 days	reduced growth rate	Rice et al. (2000)	1
Benzo(a)pyrene	gizzard shad	1	na	22 days	reduced body weight	Kolok et al. (1996)	2
Total PAHs (in the presence of a mixture of other chemicals in Elizabeth River sediments)	spot	81	322	28 days	reduced survival, weight loss	Roberts et al. (1989)	3
Total PAHs (in the presence of a mixture of other chemicals in Eagle Harbor sediments)	English sole	2	4	175 days	reduced growth rate	Kubin (1997)	

na – not available, no LOAEL was identified.

Notes:

1. Dietary exposure to polychaete worms previously exposed to sediment from the Eagle Harbor, Washington Superfund site mixed with 99.9% reference site sediments. Adverse effects were observed in one of the two experimental trials.
2. Dietary exposure of field-collected shad to benzo(a)pyrene-spiked sediments augmented with trout chow.
3. Direct exposure to sediment from the Elizabeth River mixed with sediment from a reference site. Concentrations are those reported for day 0 of exposure. Concentrations at day 15 were approximately half those reported here.

Rice et al. (2000) exposed juvenile English sole to polychaete worms that had previously been exposed in the laboratory to 0.1% of sediments from the Eagle Harbor, Washington Superfund site mixed with 99.9% sediments from a reference site. PAH concentrations were reported for the sediment and worm tissue; other uncharacterized chemicals may also have been present but were not analyzed.[80] The major chemical contaminants present at Eagle Harbor were associated with creosote, which is used as a wood preservative. The major creosote-related chemicals generally associated with toxicity are PAHs, phenols, and cresols (ATSDR 2002). About 300 chemicals have been identified in coal tar creosote, but as many as 10,000 other chemicals may also be in this mixture (ATSDR 2002). Rice et al. (2000) showed that in one experimental trial, fish exposed to contaminated worms that contained a total PAH concentration of 11.3 µg/g dw had a lower daily growth rate than controls. In a second trial of the experiment, a similar trend was observed, but the effect was not

[80] Concentrations were reported as LPAHs and HPAHs; the specific individual PAHs quantitated were not reported.

statistically significant. This study was not selected to derive a LOAEL TRV because the worms used were exposed to field-collected sediments with uncharacterized chemicals, and the significance of the effects observed at this concentration was statistically ambiguous.

The literature was also searched to identify studies in which PAH risks to fish were evaluated based on contaminated sediment exposure. Three studies examined the effects of fish exposed to PAH-contaminated sediments. In the first study, fish were exposed to sediments spiked with benzo(a)pyrene in the laboratory. Kolok et al. (1996) exposed field-collected gizzard shad to sediments spiked with 1 mg/kg dw benzo(a)pyrene for 22 days. Gizzard shad are detritus feeders that incidentally ingest large amounts of sediment. The sediment was augmented with ground trout chow. No effects were reported on body mass between exposed and control fish. The UCL concentration of benzo(a)pyrene in LDW-wide sediment was 0.41 mg/kg dw and ranged from 0.26 to 0.72 mg/kg dw for the four Pacific staghorn sculpin areas. Thus, the UCL concentration of benzo(a)pyrene in LDW sediment is less than the NOAEL reported in Kolok et al. (1996).

In a study of sediment toxicity in the Elizabeth River, Virginia (Roberts et al. 1989), field-collected juvenile spot were exposed for 28 days to field-collected sediments from a clean site amended with 1, 3.2, 10, or 32% Elizabeth River sediments as well as to undiluted Elizabeth River sediment. Resulting sediment total PAH concentrations ranged from 4 (control) to 21,200 mg/kg dw. Fish exposed to 3.2% Elizabeth River sediment (total PAH concentration of 322 mg/kg dw) were reported to exhibit increased mortality and significant weight loss compared to controls.

In a study of sediment toxicity in Eagle Harbor, Washington, Kubin (1997) exposed field-collected juvenile English sole to sediment from a clean site amended with 0.8 and 1.6% Eagle Harbor sediment for 175 days. Sediment concentrations of total PAHs were approximately < 0.50 (control), 2.0, and 4.0 mg/kg dw. Growth was similar for all treatments for the first 3 months; growth was significantly lower in the high-exposure group (4.0 mg/kg dw) for the second 3 months. The percent change in weight was 0.35% per day for fish exposed to the highest concentration of PAHs compared to 0.43% in control fish. The fish exposed to sediments with 2.0 mg/kg dw total PAHs showed no significant decrease in growth rate relative to control fish. It is unknown whether these results are applicable to PAH exposure within the LDW because total PAHs and other chemicals associated with creosote waste may differ considerably from PAH mixtures in the LDW, both in terms of toxicity and bioavailability.

Estimation of NOAELs from LOAELs

There is also uncertainty if NOAELs were estimated from LOAELs because NOAELs were not available. This uncertainty was analyzed in the Phase 1 ERA (Windward 2003b). That analysis concluded that because NOAELs were generally found to be less

than a factor of 5 different from LOAELs in studies reporting both, larger safety factors may overestimate the difference between LOAELs and NOAELs.

Selection of Salmon-Specific TRVs for Cadmium

As discussed in Section A.4.2.2.2, the selected LOAEL and NOAEL TRVs (0.5 and 1.0 mg/kg dw, respectively) were based on reduced growth of rockfish exposed to dietary cadmium; adverse effects in salmonids have been observed only at much higher dietary cadmium concentrations. Because juvenile chinook salmon as an ROC are representative solely of outmigrant juvenile salmonids in the LDW, the selected TRVs may result in the overprediction of risk for this ROC.

Salmonid-specific LOAELs for cadmium were reported in two studies, ranging from 1,395 mg/kg dw for mortality of rainbow trout fry exposed to dietary cadmium for 30 days (Szebedinsky et al. 2001) to 10,000 mg/kg dw, also for mortality of juvenile rainbow trout exposed to dietary cadmium for 28 days (Handy 1993) (Table A.6-21). Salmonid-specific dietary NOAELs were reported in five studies (Baldisserotto et al. 2005; Franklin et al. 2005; Lundebye et al. 1999; Mount et al. 1994; Szebedinsky et al. 2001), ranging from 55 mg/kg dw for growth of rainbow trout fry exposed to dietary and aqueous cadmium for 60 days (Mount et al. 1994) to 1,395 mg/kg dw, also for growth of rainbow trout fry exposed to dietary cadmium for 30 days (Szebedinsky et al. 2001) (Table A.6-21). Because a moderate number of the studies that were available evaluated dietary toxicity of cadmium to salmonids and these studies consistently showed that no adverse effects on growth or survival were observed below dietary concentrations of 55 mg/kg dw, it is unlikely that adverse effects would be observed in salmonids at lower concentrations. Assuming a NOAEL of 55 mg/kg dw, the NOAEL-based HQs would be less than 0.01 calculated using either exposure approach (diet or stomach content data). Based on this analysis, risk to juvenile chinook salmon from cadmium is likely to be very low.

Table A.6-21. Cadmium dietary toxicity studies for fish

Chemical	Test Species	NOAEL (mg/kg dw)	LOAEL (mg/kg dw)	Exposure Duration	Effect	Source	Notes
Cadmium chloride	rainbow trout fry	55	na	60 days	reduced body weight, length, or survival	Mount et al. (1994)	1, 2
Cadmium	Atlantic salmon	250	na	4 weeks	reduced growth rate (body weight)	Lundebye et al. (1999)	3, 4
Cadmium chloride	juvenile rainbow trout	294	na	15 – 30 days	reduced growth rate or survival	Baldisserotto et al. (2005)	4
Cadmium chloride	juvenile rainbow trout	471	na	28 days	reduced growth rate or survival	Franklin et al. (2005)	
Cadmium nitrate	juvenile rainbow trout	786	1,395	30 days	reduced survival	Szebedinsky et al. (2001)	4, 5
Cadmium nitrate	juvenile rainbow trout	1,395	2,265	30 days	reduced growth rate (weight)	Szebedinsky et al. (2001)	4
Cadmium sulfate	rainbow trout (130 g)	na	10,000	28 days	reduced survival	Handy (1993)	3, 4, 6

Notes:

1. Fish were exposed to copper, cadmium, lead, and zinc in water at 23.0, 0.97, 3.32, and 46.3 µg/L, respectively, at the same time as the dietary exposure to cadmium chloride. No effect on growth or survival was observed following exposure; therefore, a dietary NOAEL TRV was obtained from this study.
2. Fish fed live *Artemia* exposed to cadmium chloride in water. Dietary dose corrected for a theoretical 20% loss related to cadmium depuration from the *Artemia* food source.
3. Only the nominal concentration was reported.
4. Dietary dose was not reported as ww or dw and was assumed to be a dw concentration.
5. In a separate experiment reported in this study, 92% survival was reported for juvenile rainbow trout exposed to dietary cadmium concentrations of 1,419 mg/kg dw over a 39-day exposure period. Survival data were not statistically analyzed in either experiment.
6. Fish expelled food so the ingested dose is unknown.

na – not available

NOAEL – no-observed-adverse-effect level

LOAEL – lowest-observed-adverse-effect-level

COPCs Without LOAEL TRVs

As discussed in Section A.2.5.2, one juvenile chinook salmon COPC, chromium, had a maximum exposure concentration exceeding NOAEL TRVs and no LOAEL TRV was identified. Based on benthic invertebrate and juvenile chinook salmon stomach contents, NOAEL-based HQs for chromium were 0.18 and 2.1, respectively). Because no LOAEL toxicity data were available, the low HQs calculated using unbounded NOAEL TRVs are assumed to indicate low to very low risks to fish in the LDW from chromium.

Regional Field Studies

Three studies have been conducted with field-collected juvenile chinook salmon from the LDW. These studies are relevant because the studies are site-specific (i.e., fish collected from the LDW) (Table A.6-22). These studies evaluated effects on survival, growth, and immunocompetence in juvenile chinook salmon. Chemical-specific NOAELs and LOAELs cannot be determined from these studies because the fish were exposed to chemical mixtures.

Table A.6-22. LDW-specific studies of juvenile chinook salmon

STUDY	EXPOSURE	ENDPOINT
Varanasi et al. (1993)	LDW field/Green River hatchery	survival, growth
Arkoosh et al. (1998)	LDW field/Green River hatchery	survival-disease susceptibility (*L. anguillarum*)
Casillas et al. (1995a; 1995b)	LDW field/Green River hatchery	growth, insulin-like growth factor

While survival and growth are widely recognized endpoints of importance in ERAs, immunocompetence of fish can also be an important factor in fish population demographics (Anderson and May 1979). The importance of immune function in a given fish population is influenced by several factors, including the quality of the environment, differential susceptibility of individual fish (as determined by genetics and physiological health), and the presence and virulence of infectious agents (Snieszko 1973).

Varanasi et al. (1993) collected juvenile chinook salmon from hatcheries and the respective estuaries of the Green/Duwamish and Nisqually rivers in 1989 and 1990 and assessed survival and growth of these fish under laboratory conditions. Casillas et al. (1995a; 1995b) reported that a growth experiment similar to that of Varanasi et al. (1993) was carried out in 1993, but the data have not been published. Arkoosh et al. (1998) collected juvenile chinook salmon from the LDW, the Nisqually River, and their respective upstream hatcheries (i.e., Green River and Kalama Creek, respectively) in the spring of 1993 and 1994, and assessed these fish for immunocompetence following acclimation to laboratory conditions.

Arkoosh et al. (1998) reported that LDW fish frequently had the highest cumulative mortality relative to Green River hatchery fish or Nisqually River estuary fish when exposed in the laboratory to the bacterium *Listonella anguillarum* (previously known as *Vibrio anguillarum*). Based on these results, the authors suggested that immunosuppression in juvenile chinook salmon from the LDW may lead to increased susceptibility to infection by a virulent marine bacterium. Because of the implied association between reduced immunocompetence in these fish and exposure of these fish to chemicals from the LDW, toxicity data were reviewed from laboratory studies investigating the immunocompetence of fish exposed to mixtures of either PCBs or PAHs to assess whether single chemical exposures would result in similar effects.

PAHs

The literature search identified two studies in which juvenile chinook salmon were exposed to dietary PAH mixtures and cumulative mortality was assessed following exposure to a pathogen (Palm et al. 2003; Bravo et al., undated draft).

Palm et al. (2003) investigated chinook salmon survival relative to controls following a dietary exposure of total PAHs up to 280 mg/kg dw and subsequent bacterial challenge under freshwater conditions.[81] No difference in mortality was observed between fish exposed to both dietary PAHs and *L. anguillarum* and fish exposed to *L. anguillarum* alone (controls). The highest dietary NOAEL was 280 mg/kg dw.

Bravo et al. (undated draft) exposed juvenile rainbow trout to either a mixture of non-alkylated PAHs (33 and 390 mg/kg dw), benzo(a)pyrene (BaP) (150 mg/kg dw), or benzo(e)pyrene (BeP) (150 mg/kg dw) in their diet for 50 days under freshwater conditions.[82] Following chemical exposure, fish were exposed to two serial concentrations of the pathogen *Aeromonas salmonicida*. At the lower bacterial concentration ($10^{5.5}$ cfu/mL), mortalities of fish fed the PAH mixture at both the 33 and 390 mg/kg dw concentrations were significantly higher than in controls. At the higher bacterial concentration ($10^{5.7}$ cfu/mL), mortality of fish exposed to 390 mg/kg dw was significantly higher than in *A. salmonicida*-exposed controls, and mortality of

[81] Converted from wet weight to dry weight based on reported 10.2% moisture content in food.
[82] Converted from wet weight to dry weight assuming 10% moisture content in food.

fish exposed to 33 mg/kg dw was not significantly higher than in controls. Mortality of BeP-exposed fish was significantly higher than in controls at the lower bacterial concentration but not at the higher bacterial concentration. Mortality of BaP-exposed fish was similar to controls at both bacterial concentrations. Bravo et al. concluded that the potential for reduced immunocompetence for juvenile chinook salmon exposed to dietary PAHs at both exposure levels was high.

In the LDW, the maximum concentration of total PAHs (including all alkylated and non-alkylated PAHs) in LDW benthic invertebrate tissue samples was 36.1 mg/kg dw and the UCL was 11 mg/kg dw (Table A-2.27). Because the UCL was less than the LOAEL of 33 mg/kg dw from Bravo et al. and well below the highest concentration in which no adverse effects were observed in juvenile chinook salmon (280 mg/kg dw from Palm et al. [2003]), risks of reduced immunocompetence for juvenile chinook salmon exposed to dietary PAHs appear to be low.

PCBs

The literature search identified three studies that evaluated the survival of fish following exposure to PCBs and subsequent immunological challenge (Powell et al. 2003; Mayer et al. 1985; Snarski 1982). Powell et al. (2003) investigated juvenile chinook salmon mortality relative to controls following a dietary exposure of up to 17 mg/kg ww Aroclor 1254 for 4 weeks and subsequent 14-day challenge with *Vibrio anguillarum* under freshwater conditions. Powell et al. (2003) reported no increase in juvenile chinook salmon mortality relative to controls. The tissue concentration of Aroclor 1254 in whole bodies of juvenile chinook salmon was 0.98 mg/kg ww at the highest dose.

Mayer et al. (1985) exposed 18-day-old rainbow trout for 90 days to an Aroclor 1254:1260 mixture in water. Fish were exposed to an LD50 concentration of *Yersinia ruckeri* via flush exposure at several times during the PCB exposure. Independent disease challenge trials were conducted at 45, 60, and 90 days of PCB exposure and again at 30 and 60 days following cessation of PCB exposure. In all exposures, time to 50% mortality was not significantly different between PCB-dosed fish and controls. The PCB concentration in tissue associated with the highest NOAEL was 120 mg/kg ww. The authors reported a trend toward higher resistance to disease in fish exposed to PCBs, which they suggested was a result of physical damage (reducing pathogen uptake) to the gills resulting from PCB exposure. In a separate experiment reported in this same paper, juvenile rainbow trout were exposed for 60 days to an Aroclor 1254:1260 mixture in water and then dosed once with *Yersinia ruckeri* via IP injection or flush exposure. No significant difference in survival between PCB-dosed fish and controls was reported for either IP injection or flush pathogen exposures.

Snarski (1982) reported that rainbow trout exposed to Aroclor 1254 in water at 14.7 µg/L for 30 days were less susceptible to the bacteria *Aeromonas hydrophila* than were controls. At this PCB dose, no mortality was observed in any fish following exposure to the pathogen. PCB concentrations in tissue were not reported.

Based on the available data, the tissue-residue NOAELs for survival following PCB exposure and subsequent immunological challenge ranged from 0.98 to 120 mg/kg ww. No LOAELs were identified. The maximum total PCB whole-body tissue concentration in juvenile chinook salmon from the LDW was 1.2 mg/kg ww. Therefore, total PCB tissue concentrations of juvenile chinook salmon within the LDW appear to be below concentrations at which no adverse immunosuppressive effects were observed.

Conclusion and Uncertainties

Based on the results of these laboratory tests, reduced immunocompetence in juvenile chinook salmon from exposure to PCBs or PAHs at concentrations in the LDW is not expected. However, there are a number of uncertainties associated with this conclusion. Uncertainty is associated with:

- The dietary approach for PAHs because water exposure and variation in exposures resulting from specific prey preferences could not be evaluated
- Application of laboratory data to fish in the field because the environmental conditions to which the juvenile salmon in the LDW are exposed, such as variable salinities, were not reflected by the laboratory studies, and the biological conditions of juvenile salmonids in the LDW, including smoltification, were not captured by the laboratory experiments
- Numerous differences between experimental and wild fish that may affect immunocompetence in juvenile chinook salmon such as differences in the lipid content of the fish in the laboratory relative to those in the field, as well as in lipids in their food items
- Interpretation of results from studies with field-exposed fish because field-exposure studies involve uncontrolled variables and include exposure to a mixture of chemicals and other stressors, complicating interpretation of the cause-and-effect relationship

Risk Characterization

Risks to fish from organochlorine pesticides were not included in the risk estimates because of high uncertainty in the JN-qualified tissue pesticide data, resulting in suspected false identifications of some pesticides as well as overestimates in their concentrations, as discussed in Section A.2.4.2. Exposure estimates, TRVs, and risk estimates are discussed in this section for endrin, which was identified as a COPC for juvenile chinook salmon in Section A.2.5.2.

TRVs were searched using the same techniques described in Section A.2.4.5.2. Five endrin toxicity studies were identified for five different species of fish (Table A.6-23). LOAELs ranged from 11.5 µg/kg ww for mortality of fingerling largemouth bass exposed to endrin in water for 120 days (Fabacher 1976) to 1,660 µg/kg ww for golden shiner exposed to endrin in water for 8 hours (Ludke 1976). The LOAEL of 11.5 µg/kg

ww reported in Fabacher (1976) was selected as the LOAEL TRV. No NOAEL lower than the LOAEL was identified, so a NOAEL of 1.15 µg/kg ww was derived using a safety factor of 10 for an acute study because the study evaluated the survival endpoint. The selected endrin TRVs may underestimate toxicity because they are based on a survival endpoint. Although higher growth or reproduction LOAELS and NOAELs are reported for other fish species, it is not known if growth and reproductive endpoints would have resulted in lower TRVs for largemouth bass.

Table A.6-23. Critical tissue-residue toxicity studies of endrin in fish

Test Species	NOAEL (µg/kg ww)	LOAEL (µg/kg ww)	Exposure Route and Duration	Effect	Source	Notes
Largemouth bass fingerlings	**1.15**[a]	**11.5**	water for 120 days	reduced survival	Fabacher (1976)	static tanks treated every 5 days; residues from dead fish
Fathead minnow	na	240	food and water for 300 days	reduced survival	Jarvinen and Tyo (1978)	exposure during reproductive period
Channel catfish fingerling	307	na	food for 198 days	no effect on growth or survival	Argyle et al. (1973)	NOAEL is average residue from day 20 to day 198
Channel catfish fingerling	410	720	water for 55 days	reduced survival (40%)	Argyle et al. (1973)	LOAEL is concentration at 26 days when mortality began; NOAEL is mean concentration when residue levels were maximal from days 49 to 55
Sheepshead minnow	110	880	water for 4 weeks	reduced second generation juvenile survival	Hansen et al. (1977)	embryos spawned from field-collected adults; tissue residues in juvenile fish- effects based on mortality effects in fry
Sheepshead minnow	260	940	water for two generations	reduced female fertility	Hansen et al. (1977)	embryos spawned from field-collected adults
Golden shiner	na	1,660	water for 8 hours	100% mortality	Ludke et al. (1968)	LOAEL is average residue at time of death

[a] NOAEL estimated using uncertainty factor of 10 (acute LOAEL to chronic NOAEL).

LOAEL – lowest-observed-adverse-effect level

na – not available

NOAEL – no-observed-adverse-effect level

ww –wet weight

Bold identifies the NOAEL and LOAEL selected as TRVs.

The NOAEL- and LOAEL-based HQs for endrin and juvenile chinook salmon were 3.7 and 0.37, respectively (Table A.6-24). The selected endrin TRVs may underestimate the potential for sublethal effects because they are based on a survival endpoint. Although higher growth or reproduction LOAELs and NOAELs are reported for other fish species, it is not known if other endpoints would have resulted in lower TRVs for largemouth bass. The exposure concentration is probably an overestimate of actual exposure as a result of analytical interference from PCBs (Section A.2.4.2). Therefore,

although the risks associated with this organochlorine pesticide are uncertain, they are likely lower than these HQs suggest.

Table A.6-24. HQs for juvenile chinook salmon and endrin based on a critical tissue-residue approach

UCL Concentration (µg/kg ww)	NOAEL TRV (µg/kg ww)	LOAEL TRV (µg/kg ww)	NOAEL-Based HQ	LOAEL-Based HQ
4.3	1.15	11.5	**3.7**	0.37

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

TRV – toxicity reference value

UCL – upper confidence limit on the mean

ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

Summary of Uncertainties

Uncertainties in the problem formulation, the effects and exposure assessments, and risk characterization for juvenile chinook salmon were evaluated, as summarized below:

- Uncertainties in ROC selection, assessment endpoints, dietary composition, water exposure to PAHs and metals, benthic invertebrate tissue data, and future habitat changes resulting from restoration are not expected to have an effect on risk conclusions.
- The field component of a dietary PAH exposure study conducted by Rice et al. (2000) and several other studies with field components were discussed to further evaluate uncertainties associated with PAH exposure and effects in fish. No TRVs were selected from these studies because the mixture of chemicals to which the fish were exposed was not fully characterized.
- Risks to juvenile chinook salmon from exposure to vanadium were slightly higher if juvenile chinook salmon were assumed to forage throughout the LDW rather than only in intertidal areas. However, it is not known if juvenile chinook salmon forage in deeper waters, such as the navigation channel, of the LDW.
- The use of safety factors of greater than 5 to estimate NOAELs from LOAELs may overestimate the difference between LOAELs and NOAELs.
- The effect of uncertainties associated with exposures of fish in the LDW to chemical mixtures is not known; it is possible that risks are over- or underestimated.
- Immunocompetence experiments conducted with juvenile chinook salmon collected from the LDW indicated that reduced immunocompetence may occur

in juvenile salmonids migrating through the LDW. However, PAH and PCB toxicity studies conducted in the laboratory did not result in reduced immunocompetence in juvenile chinook salmon from exposure to PAHs or PCBs at levels comparable to dietary concentrations of these chemicals that occur in the LDW. Uncertainties remain regarding the interpretation of the field studies as well as the application of the laboratory results to the field.

- Selection of non-salmonid TRVs for cadmium resulted in a LOAEL-based HQ equal to 1.0; however, salmonid-specific effects data show that risks are likely to be very low.
- Risks from chromium were low but uncertain because a LOAEL TRV was not identified, and the NOAEL-based HQ was 2.1 based on benthic invertebrate tissue data and 0.18 based on stomach contents data.
- An evaluation of risks to juvenile chinook salmon from exposure to endrin indicated a low risk. The selected TRVs are based on survival and may underestimate the potential for sublethal effects. However, because analytical interference from PCBs is likely to have overestimated actual organochlorine pesticide concentrations in LDW tissue samples, actual risks may be lower than the HQs suggest.

A.6.2.1.3 Risk conclusions

Juvenile chinook salmon were evaluated as an ROC to represent all migratory juvenile salmonids in the LDW and because they have been listed as threatened under the Endangered Species Act (Section A.2.2.1). Results of the risk characterization for juvenile chinook salmon are summarized in Table A.6-25.

Two lines of evidence were assessed for juvenile chinook salmon exposure to dietary COPCs: 1) dietary exposure based on a single composite sample of stomach contents (composited from 72 fish), and 2) dietary exposure based on consumption of benthic invertebrates. Dietary exposure based on benthic invertebrate tissue data resulted in higher HQs for all COPCs than did juvenile chinook salmon stomach content data.

No COPC had a LOAEL-based HQ exceeding 1.0. The cadmium LOAEL-based HQs were 1.0 and 0.92 based on concentrations in benthic invertebrates collected from the LDW and in stomach contents, respectively. There is uncertainty in these HQs because the cadmium TRVs were based on juvenile rockfish, and several toxicity studies using salmonids were available with TRVs orders of magnitude higher. Because the salmonid-specific studies likely better represent the sensitivity of juvenile chinook salmon, the selected NOAEL and LOAEL TRVs applied in this ERA likely overestimate the sensitivity of juvenile chinook salmon to cadmium. NOAEL-based HQs for arsenic, copper, and vanadium were greater than 1.0 (1.1, 1.9, and 4.0, respectively), whereas LOAEL-based HQs for these COPCs were less than 1.0. Therefore, risks from these COPCs were low. Uncertainty in the TRV affects the risk conclusion for vanadium. The vanadium HQ is based on a NOAEL TRV extrapolated

from an unbounded LOAEL TRV. No other dietary toxicity studies were available. Because of the high uncertainty in the effects data, risks from vanadium are uncertain and could be higher or lower than the HQs indicate.

Table A.6-25. Summary of risk characterization for juvenile chinook salmon

COPC	Highest NOAEL-Based HQ[a]	Highest LOAEL-Based HQ[a]	LOAEL Endpoint	Primary Uncertainty
Arsenic	**1.1**	0.73	juvenile growth	few toxicity studies available
Cadmium	**5.0**	**1.0**	juvenile growth	high TRV uncertainty[b]
Copper	**1.9**	0.93	juvenile growth	medium uncertainty[c]
Vanadium	**4.0**	0.79	juvenile growth	only one toxicity study available, foraging range exposure assumption

[a] Dietary exposure based on benthic invertebrate tissue data resulted in higher HQs for all COPCs than did juvenile chinook salmon stomach contents data. Thus, benthic invertebrate tissue HQs are shown in this table.

[b] Although a moderate number of toxicity studies were available, the six available dietary toxicity studies conducted with salmonids showed that salmonids in these studies were likely less sensitive to cadmium than suggested by the selected NOAEL and LOAEL TRVs.

[c] A large number of studies that presented a range of effects thresholds were available. Sufficient data from other studies were available to suggest that the lowest LOAEL was not supported by related research; therefore, this LOAEL was not selected. Furthermore, the available toxicity data suggest that the likely dietary threshold for toxic effects in salmonids is higher than the selected dietary LOAEL of 100 mg/kg dw.

COPC – chemical of potential concern

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

TRV – toxicity reference value

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

A.6.2.2 English sole

This section presents risk estimates, uncertainties, and risk conclusions for English sole.

A.6.2.2.1 Risk estimates

COPCs evaluated for English sole included arsenic, cadmium, copper, and vanadium, which were evaluated using a dietary approach, and total PCBs, which were evaluated using a critical tissue-residue approach. Estimated exposure concentrations for English sole were greater than both LOAEL and NOAEL TRVs for cadmium and vanadium, each with a LOAEL-based HQ of 1.2 (Table A.6-26). Exposure concentrations of total PCBs were compared to the range of effect concentrations selected as TRVs. LOAEL-based HQs calculated with this range of LOAELs ranged from 0.98 to 5.0. NOAEL-based HQs ranged from 4.9 to 25 (Table A.6-26). Exposure concentrations of arsenic and copper exceeded NOAEL TRVs, with NOAEL-based HQs of 1.2 and 1.9, respectively; LOAEL-based HQs for arsenic and copper were less than 1.0.

Table A.6-26. HQ calculations for English sole

Approach	COPC	Exposure Concentration	Unit	NOAEL TRV	LOAEL TRV	NOAEL-Based HQ	LOAEL-Based HQ
Tissue residue	total PCBs	2,600	µg/kg ww	104 – 528	520 – 2,640	**4.9 – 25**[a]	0.98 – **5.0**[a]
Dietary	arsenic	24	mg/kg dw	20	30	**1.2**	0.80
	cadmium	0.61	mg/kg dw	0.1	0.5	**6.1**	**1.2**
	copper	93	mg/kg dw	50	100	**1.9**	0.93
	vanadium	12	mg/kg dw	2.04	10.2	**5.9**	**1.2**

[a] Because of the uncertainty in the LOAEL, LOAEL-based HQs were calculated from a range of effects concentrations reported in Hugla and Thome (1999). The NOAEL TRV range was estimated by dividing the LOAEL TRV range by an uncertainty factor of 5.

COPC – chemical of potential concern
dw – dry weight
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
TRV – toxicity reference value
ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

A.6.2.2.2 Uncertainty analysis

This section presents a discussion of the uncertainty associated with the problem formulation, the exposure and effects assessments, and the risk characterization for English sole.

Problem Formulation

Primary uncertainties in the problem formulation for English sole include ROC selection, assessment endpoints, and the COPC screen. Uncertainties associated with fish assessment endpoints are the same as those discussed in Section A.6.2.1.2 for juvenile chinook salmon. Uncertainties associated with ROC selection and the COPC screen are discussed below.

ROC Selection

English sole are benthic fish that live in close contact with sediments and thus have a high likelihood of exposure to sediment-associated chemicals through direct contact and through their diet. Other fish represented by English sole as an ROC have either similar exposure pathways (e.g., starry flounder), or less direct contact with sediments (e.g., shiner surfperch). As part of the Phase 2 RI, shiner surfperch were collected and analyzed to represent prey for various fish and wildlife ROCs. PCB concentrations in shiner surfperch tissue were also compared to TRVs to assess the suitability of English sole as a representative species. Shiner surfperch HQs are compared to English sole HQs for PCBs (the only critical tissue-residue COPC for English sole) in Table A.6-27.

Table A.6-27. HQ calculations for shiner surfperch and English sole for total PCBs

Species	Minimum PCB Conc. (µg/kg ww)	Maximum PCB Conc. (µg/kg ww)	Mean PCB Conc. (µg/kg ww)	UCL Conc. (µg/kg ww)	NOAEL TRV (µg/kg ww)	LOAEL TRV (µg/kg ww)	NOAEL-Based HQ	LOAEL-Based HQ
English sole	450	4,700	2,200	2,600	104 – 528	520 – 2,640	**4.9 – 25**	0.98 – **5.0**
Shiner surfperch	350	18,400	1,800	3,500	104 – 528	520 – 2,640	**6.6 – 34**	**1.3 – 6.7**

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

PCB – polychlorinated biphenyl

TRV – toxicity reference value

UCL –-upper confidence limit on the mean

ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

The exposure concentration for shiner surfperch resulted in a LOAEL-based HQ range of 1.3 to 6.7, whereas the exposure concentration for English sole resulted in a LOAEL-based HQ range of 0.98 to 5.0. Because shiner surfperch and English sole HQs for PCBs were similar and the total PCBs LOAEL TRV is uncertain, shiner surfperch tissue data do not change risk estimates for fish represented by English sole as an ROC.

COPC Screen

Uncertainties associated with chemicals not analyzed in LDW tissue samples, chemicals for which toxicological data were not identified, and the mercury NOAEL TRV are the same as those discussed in Section A.6.2.1.2 for juvenile chinook salmon.

Risks to fish from exposures to BEHP, dimethyl phthalate, and di-n-butyl phthalate in the LDW are uncertain because although these chemicals were not detected in English sole whole-body tissue samples, reporting limits for these chemicals in tissue were greater than the selected NOAEL TRVs (Table A.2-21 in Section A.2.5.2). The reporting limits for BEHP were elevated because of analytical dilutions of the samples. Forty-nine tissue samples with BEHP RLs of 7,200 µg/kg ww were re-analyzed. BEHP was not detected in any of the re-analyzed tissue samples that had RLs from 66 to 130 µg/kg ww.

HQs calculated using minimum and maximum reporting limits relative to NOAEL TRVs for each chemical are presented in Table A.6-28. LOAEL TRVs were not available for any of these chemicals. HQs are also presented for shiner surfperch because BEHP was detected in 5/27 shiner surfperch tissue samples. Dimethyl phthalate and di-n-butyl phthalate detection frequencies were 0/27 and 1/27, respectively, in shiner surfperch.

Table A.6-28. HQs for BEHP, dimethyl phthalate, and di-n-butyl phthalate

Chemical	ROC	Reporting Limit (µg/kg ww)		TRV (µg/kg ww)		NOAEL-Based HQ			
		Min	Max	NOAEL	LOAEL	Min RL	Max RL		
BEHP	English sole	66	3,600	390	na	0.17	**9.2**		
	shiner surfperch	280[a]	2,100[b]	390	na	0.72	**5.4**		
Dimethyl phthalate	English sole	290	580	498	na	0.58	**1.2**		
	shiner surfperch	9.9	2,900	498	na	<0.1	**5.8**		
Di-n-butyl phthalate	English sole	290	1,200	1,170	na	0.25	1.0		
	shiner surfperch	40	2,300[b]	1,170	na	<0.1	**2.0**		

[a] Minimum detected concentration is presented.

[b] Maximum detected concentration is presented.

BEHP – bis(2-ethylhexyl) phthalate

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

na – no TRV available

nc – not calculated because no LOAEL TRV was available

NOAEL – no-observed-adverse-effect level

RL – reporting limit

ROC – receptor of concern

ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0.

All but one of the NOAEL-based HQs calculated using maximum reporting limits as exposure concentrations were greater than 1.0 (Table A.6-28). All NOAEL-based HQs calculated using minimum reporting limits were less than 1.0. Detected concentrations of BEHP and di-n-butyl phthalate in shiner surfperch resulted in HQs similar to English sole HQs based on reporting limits. Based on this analysis, and assuming that English sole and shiner surfperch have similar exposures to phthalates, phthalate risks to English sole are uncertain but appear to be low.

Exposure Assessment

Uncertainties in the exposure assessment for English sole were associated with the following factors:

- Water pathway for metals and PAHs
- Incidental sediment ingestion
- Foraging range
- Benthic invertebrate tissue data and linear regressions
- Future habitat changes resulting from restoration
- Use of starry flounder tissue in Area T4 as a surrogate for English sole

The uncertainties associated with exposure via the water pathway and future habitat changes are similar to those discussed for juvenile chinook salmon in Section A.6.2.1.2.

Incidental Sediment Ingestion

The exposure assessment for English sole assumed 1% of the diet was incidental sediment ingestion, as described in Section A.4.1.2. There is uncertainty in the percentage of sediment ingestion assumed because it is based on subjective observations by experienced fish biologists and not based on empirical data. Estimates ranged from 1% (Lange 2006) to as high as 10% (Johnson 2006). Therefore, uncertainty in dietary exposure estimates calculated using Equation 4-1, as described in Section A.4.1.2 were evaluated assuming 0, 1, and 10% sediment ingestion to bracket the 1% estimate (Table A.6-29).

Table A.6-29. English sole dietary exposure estimates based on three assumed incidental sediment ingestion scenarios

COPC	Dietary Exposure Concentrations Based on Different Incidental Sediment Ingestion Percentages (mg/kg dw)		
	0%	1%	10%
Arsenic	24	24	25
Cadmium	0.60	0.61	0.74
Copper	92	93	100
Vanadium	12	12	17

COPC – chemical of potential concern

dw – dry weight

An assumption of 10% sediment ingestion resulted in slightly higher exposure estimates for cadmium, copper, and vanadium, whereas an assumption of 0% sediment ingestion resulted in slightly lower exposure estimates for these COPCs. However, HQs did not change substantially for any COPC (Table A.6-30), and HQs did not change from less than 1.0 to greater than 1.0 or vice versa, except for copper, for which the LOAEL changed from 0.93 to 1.0.

Table A.6-30. Dietary HQs for English sole as a function of sediment consumption

COPC	0% Sediment consumption		1% Sediment Consumption		10% Sediment Consumption	
	NOAEL-Based HQ	LOAEL-Based HQ	NOAEL-Based HQ	LOAEL-Based HQ	NOAEL-Based HQ	LOAEL-Based HQ
Arsenic	**1.2**	0.80	**1.2**	0.80	**1.3**	0.83
Cadmium	**6.0**	**1.2**	**6.1**	**1.2**	**7.4**	**1.5**
Copper	**1.8**	0.92	**1.9**	0.93	**2.0**	**1.0**
Vanadium	**5.9**	**1.2**	**5.9**	**1.2**	**8.3**	**1.7**

COPC – chemical of potential concern

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

Foraging Range

English sole were assumed to forage exclusively in the LDW even though it is known that they migrate seasonally to spawn in Puget Sound. Uncertainty analysis in the Phase 1 ERA (Windward 2003b) concluded that the assumption of foraging throughout the LDW may potentially overestimate or underestimate exposure depending on the relative magnitude and extent of contamination in other foraging areas.

Benthic Invertebrate Tissue Data and Linear Regressions

The uncertainties associated with benthic invertebrate tissue data in the exposure assessment for English sole are the same as those discussed in Section A.6.2.1.2 for juvenile chinook salmon, with the following exception. More samples were used in the LDW-wide EPC calculations for English sole (n = 20 or 24 depending on the COPC) than for the intertidal EPC calculations for juvenile chinook salmon (n = 13 or 17, depending on the COPC), which improves confidence that the combined sample is representative of English sole exposures in the LDW.

If the exposure estimate had been calculated using empirical benthic invertebrate tissue data rather than the regression, the UCL would have been 37 mg/kg dw, in comparison to the UCL of 24 mg/kg dw estimated using the regression. Assuming an exposure concentration of 37 mg/kg dw, the LOAEL-based HQ would have increased from 0.8 to 1.2, still suggesting a relatively low risk from arsenic exposure.

Use of starry flounder tissue data from Area T4

There is uncertainty associated with the LDW-wide English sole tissue concentrations because three starry flounder composite tissue samples from Area T4 were included in the English sole dataset. These three samples were included because in the 2004 sampling event, it was difficult to catch a sufficient number of English sole to populate six composite tissue samples of English sole. Thus, in 2004, three starry flounder and three English sole composite tissue samples were analyzed. Total PCB concentrations in starry flounder tissue composite samples were lower than those in English sole composite tissue samples from Area T4 in 2004 (mean concentrations of 570 µg/kg ww vs. 1700 µg/kg ww). Therefore, inclusion of starry flounder data may have decreased the exposure estimate because concentrations of total PCBs in starry flounder were lower than those in English sole composite samples from the same area.

Effects Assessment

Uncertainties in the effects assessment for English sole were associated with the following factors:

- Effects from chemical mixtures and estimation of NOAELs from LOAELs
- Exclusion of field studies from TRV selection
- PCB TRV

- Cadmium TRV
- COPCs without LOAEL TRVs
- TEQ approach for PCBs
- Regional field studies
- Critical tissue residue approach

In addition, some of the selected TRVs are considered less certain than others if there were a small number of studies, if endpoints were subchronic, or if data quality was questionable. The relative uncertainties in the selected TRVs for each English sole COPC are the same as those discussed in Section A.6.2.1.2 for juvenile chinook salmon, except for the TRVs for cadmium and total PCBs, which are discussed below.

Effects of Chemical Mixtures and Estimation of NOAELs from LOAELs

Uncertainties associated with effects of chemical mixtures and estimation of NOAELs from LOAELs are the same as those discussed in Section A.6.2.1.2 for juvenile chinook salmon with the following additional uncertainty for PCBs, which were not evaluated as a COPC for juvenile chinook salmon. The laboratory effects studies included in the total PCBs assessment used unweathered Aroclor mixtures. PCBs present in fish tissue have undergone physico-chemical weathering and differential accumulation in the food web, resulting in PCB congener mixtures that are potentially more or less biologically active than the commercial Aroclor mixtures. Because laboratory toxicity tests evaluated for TRV selection generally were conducted with a single Aroclor mixture, the potency of PCB mixtures in LDW fish relative to those in fish from toxicity studies is uncertain.

Exclusion of Field Studies from TRV Selection

Uncertainties discussed in Section A.6.2.1.2 for juvenile chinook salmon associated with the exclusion of field studies from TRV selection are relevant to English sole. Rice et al. (2000) tested the toxicity of sediments contaminated with PCBs, PAHs, and other chemicals using English sole. This study could not be used to derive a PCB or a PAH TRV for English sole because the test sediments were taken from contaminated waterways (Eagle Harbor and Commencement Bay) that contained mixtures of chemicals that included more than PAHs or PCBs, respectively, confounding interpretation of results. The direct sediment exposure studies discussed in Section A.6.2.1.2 are also relevant to English sole, a benthic species.

PCB TRV

As discussed in Section A.4.2.2.1, there is uncertainty in selecting the TRV for total PCBs and fish because: 1) there is uncertainty in interpretation of several studies reporting the lowest concentrations associated with effects, and 2) some effects concentrations were expressed as egg/embryo-tissue concentrations rather than whole-body concentrations.

As noted in Section A.4.2.1.1, there are a number of uncertainties associated with the Hugla and Thome (1999) study reporting the lowest effects concentrations. Because of these uncertainties, rather than selecting a single LOAEL, the range of effects concentrations reported in this paper for the fecundity endpoint (520 and 2,640 µg/kg ww) and for the spawning and egg hatchability endpoints (2,640 µg/kg ww) were considered to represent the lowest exposure levels over which adverse reproductive effects may occur in these fish. This section presents a detailed evaluation of uncertainties associated with this study and implications for risk conclusions. Uncertainties discussed include those associated with the statistical analysis for the fecundity endpoint and the fact that this endpoint was not dose responsive, uncertainties related to test conditions, and uncertainties in the estimation of the whole-body concentration associated with effects.

The number of fish used to evaluate effects endpoints in Hugla and Thome (1999) is unclear. Hugla and Thome (1999) presented the following information:

- A sample size of six was used in the statistical analysis conducted to assess the significance of effects in exposed fish relative to control fish.
- Ten males and six females were exposed in the 2.5-µg/g PCB treatment.
- Six fish were analyzed for PCB concentrations after 50 days of exposure.
- Six ovaries were analyzed 1 year later.

Thus, from the information presented in the Hugla and Thome (1999), it would appear that whole-body concentrations in the fish analyzed at 50 days must be for males only (if any females were analyzed at 50 days, fewer than six would have been available for analysis 1 year later). However, in recent correspondence, the authors have stated that both male and female fish were included in whole-fish tissue analyses conducted at 50 days (Leroy 2007 [pers. comm.]). They also indicated that the total number of fish exposed may have been incorrectly reported in the paper.

Understanding the number of spawnings and the number of fish tested is critical to statistical analysis and interpretation of results. Hugla and Thome (1999) state that a one-way analysis of variance (ANOVA) was used to analyze the data, and ANOVA assumes independence of observations. In recent correspondence (Leroy 2007 [pers. comm.]), the authors reported that three fish were spawned two times each. Under this scenario, the six resulting data points for each female reproductive endpoint would not be independent. Statistical analyses conducted using a sample size of 3 (exposed fish) rather than a sample size of 6 (spawning events), would result in a concomitant reduction of statistical power and potentially different conclusions about the differences in fecundity between the control and the exposed fish.

Another element of uncertainty in the fecundity LOAEL is that observed effects on fecundity were not dose responsive after two spawning seasons, as noted in Section 4.2.1.1. While the PCB-treated fish produced half as many eggs as the control fish, and the number of eggs produced by the control fish was consistent with other studies for

this species (Philippart et al. 1989), the average fecundity after two spawning seasons was similar between the high and low doses; variance in the fecundity endpoint was greater at the higher dose. Furthermore, no barbel PCB toxicity data are available from other studies to compare with the fecundity effect reported in Hugla and Thome (1999).

In addition, the fish holding and exposure conditions used by Hugla and Thome (1999) may have influenced the reproductive effects observed. In recent correspondence, the authors have provided more detailed information than was discussed in the 1999 publication. Fish were kept in artificially heated water and were spawned when 4 years old (Leroy 2007 [pers. comm.]). A separate study indicated that increased water temperatures of 20 to 24°C at the facility where this study was conducted were used to accelerate the sexual maturation of the experimental fish (Philippart et al. 1989). Philippart et al. showed that by manipulating the temperature and/or photoperiod under which fish are reared, barbel are spawned at an earlier age and smaller size than the typical minimum spawning age of 6 years. Because temperature was also used to affect the barbel reproductive cycle in this study, it is uncertain whether these manipulations may have also affected the sensitivity of barbel reproduction to PCBs. The fecundity of the control fish was similar to that found in another study with this species under similar conditions (Philippart et al. 1989); therefore, the effect of temperature, if any, is not known.

Therefore, because of the uncertainties in the statistical analysis, in the effects of elevated fish holding and exposure temperatures, and in the exposure-response relationship for fecundity, the fecundity LOAEL from the low dose is highly uncertain. There is greater confidence in the higher whole-body LOAEL as an effects level for these fish because this treatment resulted in complete reproductive failure following exposure in the first spawning year, and resulted in 96% mortality in offspring 1 year after exposure (relative to 52.4% in controls). Furthermore, increased egg/larval mortality was reported at higher PCB concentrations in eggs (i.e., the response appeared to be exposure-dependent). Given these uncertainties and the lack of confirming studies using the same species, the range of LOAEL TRVs provided by Hugla and Thome (1999) are considered to provide a conservative assessment of PCB risks to fish.

If the next higher LOAEL of 9,300 µg/kg ww for sheepshead minnow egg and larval survival (Hansen et al. 1974a) has been selected as the LOAEL TRV, the LOAEL-based HQ for English sole would have been 0.28. The highest NOAEL below this LOAEL was 1,900 µg/kg ww from the same study. If this NOAEL had been selected as the NOAEL TRV, the NOAEL-based HQ would have been 1.4. Uncertainties in the Hansen et al. (1974a) study include elevated PCB concentrations of 520 to 640 µg/kg ww in control fish and enhancement of egg production by injecting the fish with human chorionic gonadotrophic hormone. Potential confounding effects of hormonal injections on egg survival are unknown.

Four available studies presented PCB effects concentrations in egg and embryo tissues. Effects concentrations ranged from 857 to 77,900 µg/kg ww in egg and embryo tissues (Fisher et al. 1994; Freeman and Idler 1975; Hendricks et al. 1981; McCarthy et al. 2003). The lowest effects concentration was for reduced growth of Atlantic salmon fry held in PCB-free water for 176 days following egg exposure to aqueous PCB concentrations of 625 to 62,500 µg/L for 48 hours (Fisher et al. 1994). The highest effects concentration was for brook trout embryos exposed to 200 µg/L of PCBs in water for 21 days (Freeman and Idler 1975). No NOAELs were identified.

Although these egg and embryo effects concentrations were generally lower than effects concentrations in more mature fish, egg/embryo and adult tissue-residue data are not directly comparable. Species-specific ratios relating PCB concentrations in maternal adults to unfertilized eggs for yellow perch, smallmouth bass, white bass, white sucker, and rainbow trout ranged from 0.83 to 2.35 (Niimi 1983). Adjustment for a 2-fold weight increase in PCB concentration in fertilized eggs relative to unfertilized eggs (Niimi 1983) resulted in adult-to-fertilized egg ratios ranging from 1.7 to 4.7 for these species. Sheepshead minnow adult-to-fertilized-egg ratios were reported to range from 0.90 to 2.3 (Hansen et al. 1974a). Therefore, based on these studies, the ratio of PCB concentrations in fertilized eggs to those in maternal adults would likely range from 0.90 to 4.7. This range is uncertain because data represent only six fish species, with little to no replication.

Using this range of adult-to-egg PCB ratios, the maternal adult PCB concentrations associated with the reported egg LOAELs resulted in an estimated whole-body LOAEL range of 771 to 366,000 µg/kg ww, which is comparable to the range of measured whole-body LOAELs from studies with adults (520 to 429,000 µg/kg ww). Although there are additional uncertainties associated with these studies, based on the likely range of adult tissue concentrations extrapolated from these studies, use of egg LOAELs would not have resulted in risk conclusions different from those based on whole-body effects data.

Cadmium TRV

The lowest cadmium dietary LOAEL and NOAEL TRVs (0.5 and 0.1 mg/kg dw, respectively) for growth of juvenile rockfish were more than an order of magnitude lower than the other cadmium dietary TRVs and the observed effect was partially attributed to reduced food intake (see Section A.4.2.2.2). The next higher dietary LOAEL was 800 mg/kg dw (for growth of guppy) with a corresponding NOAEL of 500 mg/kg dw from the same study (Hatakeyama and Yasuno 1987). Both studies evaluated growth effects, although the fish species and form of cadmium differed. These results show that although few species have been investigated, there is wide variability in dietary toxicity data reported for cadmium.

HQs calculated using the next higher LOAEL and its associated NOAEL TRVs (Hatakeyama and Yasuno 1987) would have been less than 1.0 for English sole. Thus,

risks from cadmium are uncertain for English sole and other fish represented by English sole (Section A.6.2.2.3).

COPCs Without LOAEL TRVs

As discussed in Section A.2.5.2, four chemicals had maximum exposure concentrations exceeding NOAEL TRVs but LOAEL TRVs were not available. One chemical (chromium) was evaluated using a dietary approach, and three chemicals (benzoic acid, dimethyl phthalate, and di-n-butyl phthalate) were evaluated using a critical tissue-residue approach. Risks to English sole from dimethyl phthalate and di-n-butyl phthalate were evaluated in the exposure assessment uncertainty section above because concentrations of these chemicals were generally below detection limits in fish tissue samples.

Dietary concentrations of chromium were calculated using Equation 4-1, as described in Section A.4.1.2. Benzoic acid critical tissue-residue concentrations were calculated as the UCL of the mean of all English sole tissue composite samples from throughout the LDW. NOAEL-based HQs were 1.1 for chromium and 1.5 for benzoic acid (Table A.6-31). Because no LOAEL toxicity data were available for these chemicals, the low HQs calculated using unbounded NOAEL TRVs are assumed to indicate low risks.

Table A.6-31. English sole NOAEL-based HQs for chemicals without LOAEL TRVs

COPC	Units	Exposure Concentration	NOAEL TRV	NOAEL-Based HQ
Chromium	mg/kg dw	10	9.4[a]	**1.1**
Benzoic acid	µg/kg ww	5,100	3,380[b]	**1.5**

[a] No effects on growth of 2-year-old gray mullet exposed to chromium III through diet and sediment for 8 weeks (Walsh et al. 1994).

[b] No effects on survival of mosquitofish exposed in a mesocosm for 24 hours (Lu and Metcalf 1975).

COPC – chemical of potential concern

dw – dry weight

HQ – hazard quotient

NOAEL – no-observed-adverse-effect level

TRV – toxicity reference value

ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0.

Toxic Equivalent Approach for PCBs

Risks from PCBs were evaluated in this ERA based on total PCB concentrations in fish tissue. Risks from PCBs can also be evaluated on a subset of PCB congeners that have dioxin-like properties, using a TEQ approach. English sole tissues were analyzed for dioxin-like PCB congeners and PCB Aroclors. PCB TEQ exposures calculated for

English sole were compared to the lowest LOAEL and NOAEL TRVs for 2,3,7,8-TCDD (2,3,7,8-TCDD toxicity studies are discussed in Attachment 8).

The lowest LOAEL for 2,3,7,8-TCDD was estimated using a study in which hatchery-reared adult female rainbow trout were exposed to dietary 2,3,7,8-TCDD for up to 320 days (Giesy et al. 2002; Jones et al. 2001). Dietary doses were 0, 1.8, 18, and 90 ng/kg ww. Giesy et al. (2002) reported significant effects on adult survival in fish fed 1.8 ng/kg ww.[83] Fillet tissue concentrations of 0.44 ng/kg ww were reported at day 200 in fish fed 1.8 ng/kg ww (Jones et al. 2001). The authors estimated that the fillet concentration reported in fish fed 1.8 ng/kg ww at day 200 was a 28-fold underestimation of the fillet concentration when mortality became significant (Giesy 2006). Therefore, the fillet concentration associated with adverse effects is assumed to be approximately 12 ng/kg ww. Whole-body data were not reported.

Because this study did not meet the TRV selection criteria presented in Section A.2.5.2,[84] but it was the only long-term 2,3,7,8-TCDD toxicity study available for fish, TEQ risks for fish are uncertain. Because a NOAEL lower than the LOAEL was not available, the NOAEL was calculated by dividing the selected LOAEL by 5, resulting in a NOAEL TRV of 2.4 ng/kg ww (for fillets).

TEQs were calculated for dioxin-like PCB congeners in eight English sole composite tissue samples.[85] WHO TEFs for fish reported in Van den Berg et al. (1998) were used to calculate fish TEQs. NOAEL- and LOAEL-based TEQ HQs for English sole were less than 1.0 (Table A.6-32). HQs calculated using a total PCB approach are presented in Table A.6-32 for comparison. HQs using the total PCB approach were higher than the PCB TEQ HQs, indicating that the total PCB approach should provide health-protective risk estimates for fish for PCBs. However, risk estimates for fish based on TEQs were calculated using only the dioxin-like PCB congeners because dioxin and furan tissue data were not available. Thus, risks associated with exposure to all dioxin-like chemicals were likely underestimated; the degree of underestimation is uncertain.

---

[83] There was 2.3 ng TCDD TEQ/kg ww of background PCB contamination in the diet, which was not accounted for in radiometric analyses of 2,3,7,8 TCDD.

[84] Neither whole-body nor egg tissue concentrations were presented for the time period when adverse effects were observed. Instead, fillet concentrations were estimated by the author from data collected at a different time.

[85] One starry flounder was included in the composite sample for Sampling Area 4, where few English sole were captured. Data are described in Section A.2.4.1.2.

Table A.6-32. HQs for English sole total PCBs and PCB TEQs

Basis	UCL Concentration	NOAEL TRV	LOAEL TRV	NOAEL-Based HQ	LOAEL-Based HQ
PCB TEQ	1.88 ng/kg ww	2.4 ng/kg ww	12 ng/kg ww	0.78	0.16
Total PCBs (sum of Aroclors)	2,600 µg/kg ww	104 – 528 µg/kg ww	520 – 2,640 µg/kg ww	**4.9 – 25**	0.98 – **5.0**

HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
TEQ – toxic equivalent
TRV – toxicity reference value
UCL – upper confidence limit on the mean
ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

Regional Field Studies

Several regionally relevant studies have been conducted with field-collected English sole from the LDW (Table A.6-33). Studies with LDW-exposed fish are discussed in detail in the Phase 1 ERA (Section A.4.3) and summarized in this section because of the relevance of the exposure route (i.e., field exposure in the LDW) and mixture of chemicals. However, chemical-specific NOAELs and LOAELs cannot be determined from these studies because the fish were exposed to chemical mixtures under uncontrolled conditions.

Table A.6-33. Growth, reproductive effects, and survival studies using English sole collected from the LDW

Study	Collection location	Effect
Johnson et al. (1988; 1997a; 1998; 1999)	Duwamish Waterway	inhibited gonadal development, altered plasma 17 beta-estradiol levels, reduced fecundity
Casillas et al. (1991a)	Duwamish Waterway	reduced spawning success
Rhodes et al. (1987)	two locations: Upper Duwamish Waterway and Lower Duwamish Waterway	reduced survival, increased lesion incidence
Johnson and Landahl (1994)	four or five locations in LDW	reduced survival, increased lesion incidence

LDW – Lower Duwamish Waterway

English sole collected from the LDW are more likely to have lesions than the sole collected from less-contaminated locations (Johnson and Landahl 1994; Rhodes et al. 1987). Hepatic neoplasms were found to be confined mainly to fish in urban areas (LDW, Commencement Bay [Hylebos Waterway], and the harbor area of Everett, Washington) (Malins et al. 1984).

Potential reproductive effects on English sole exposed to sediment-associated chemicals in the LDW and other sites throughout Puget Sound have been assessed using various endpoints. Concentrations of the female reproductive hormone 17 beta-estradiol in plasma have been analyzed (Johnson et al. 1997a; 1998; 1999), and ovarian

development (Johnson et al. 1988; 1993), spawning success (Casillas et al. 1991a), and fecundity (Johnson et al. 1997a) have been assessed.

English sole from the LDW have been reported to exhibit inhibited gonadal development (Johnson et al. 1988), depressed plasma estradiol and reduced ovarian production *in vitro* (Johnson et al. 1988; 1993), and reduced spawning success (Casillas et al. 1991a). The reports suggest that these effects are a result of elevated concentrations of aromatic and chlorinated hydrocarbons present in LDW and Eagle Harbor sediments. These elevated concentrations have also been suggested to be significant risk factors for development of these reproductive abnormalities (Johnson et al. 1988; Casillas et al. 1991a).

These field studies are supported by laboratory experiments showing that pretreatment of gravid female English sole with extracts of contaminated sediment or crude oil (containing high levels of PAHs) decreased levels of endogenous estradiol (Johnson et al. 1995; Stein et al. 1991). Related experiments suggest that exposure to BaP- or PAH-contaminated sediment may suppress estradiol-induced vitellogenin production in fish, including English sole (Anulacion et al. 1997; Nicolas 1999).

Chemicals implicated as causal factors include PAHs and PCBs; however, linking the results of field studies to risks from specific chemicals is difficult considering, among other factors, the complex mixtures of chemicals in the field and the uncertainties in English sole home range. In addition, interpreting cause and effect of the adverse effects reported in field studies is complicated because of genetic variation, health, and seasonal variation in the spawning cycle. Therefore, although regional studies indicate an increased risk of adverse effects on English sole reproduction in the LDW, these effects cannot be conclusively associated with exposure to specific chemicals or chemical mixtures in the LDW.

Critical Tissue-Residue Approach

Uncertainties inherent in the use of chemical concentrations in tissue to estimate exposure from sediment-associated COPCs in the LDW include: 1) chemical concentrations detected in tissue may not accurately reflect the internal dose at target organs; 2) variability in chemical structure, species sensitivity, and biotransformation may affect critical tissue residues; and 3) the parent compound may be metabolized to more toxic chemicals. The influence of uptake or depuration kinetics on biological responses can also affect tissue residues. For example, short-term exposure of fish to relatively high chemical concentrations that elicited toxicity have been shown to result in lower tissue chemical concentrations than those reported in longer-term exposures to lower chemical concentrations that did not result in adverse effects (e.g., van Wezel et al. 1995).

Risk Characterization

Uncertainties in the risk characterization for English sole were associated with the organochlorine pesticides data.

Risks to fish from organochlorine pesticides were not included in the risk estimates because of high uncertainty in the tissue pesticide data resulting in suspected false identifications of some pesticides as well as overestimates in their concentrations, as discussed in Section A.2.4.2. Therefore, exposure estimates, TRVs, and risk estimates are presented in this uncertainty section for the three organochlorine pesticides (alpha-endosulfan, beta-endosulfan, and endrin) that were identified as COPCs in Section A.2.5.2.

TRVs for endosulfan and endrin were selected using the same techniques described in Section A.2.5.2. One endosulfan toxicity study reported that spot, pinfish, and mullet were exposed to endosulfan in water for 96 hours (Table A.6-34). The lowest LOAEL in the study (31 µg/kg ww) was reported for spot. No lower NOAEL was identified, so a NOAEL of 3.1 was derived using a safety factor of 10 because this was an acute study. As discussed in Section A.6.2.2.2, six endrin toxicity studies were identified for six different species of fish (Table A.6-2). LOAELs ranged from 11.5 µg/kg ww for mortality of fingerling largemouth bass exposed to endrin in water for 120 days (Fabacher 1976) to 1,660 µg/kg ww for golden shiner exposed to endrin in water for 8 hours (Ludke 1976). The LOAEL of 11.5 µg/kg ww reported in Fabacher (1976) was selected as the LOAEL TRV. No NOAEL lower than the LOAEL was identified, so a NOAEL of 1.2 µg/kg ww was derived for endrin using a safety factor of 10 for an acute study. The selected endrin TRVs may underestimate the potential for sublethal effects because they are based on a survival endpoint. Although higher growth or reproduction LOAELS and NOAELs are reported for other fish species, it is not known if other endpoints would have resulted in lower TRVs for largemouth bass.

Table A.6-34. Critical tissue-residue toxicity studies of endosulfan in fish

Test Species	NOAEL (µg/kg ww)	LOAEL (µg/kg ww)	Exposure Route and Duration	Effect	Source	Notes
Spot	**0.62**[a]	**31**	water for 96 hours	reduced survival (65%)	Schimmel et al. (1977a)	tissue residues of surviving fish
Pinfish	195	272	water for 96 hours	reduced survival (65%)	Schimmel et al. (1977a)	tissue residues of surviving fish
Mullet	na	360	water for 96 hours	reduced survival (60%)	Schimmel et al. (1977a)	tissue residues of surviving fish

[a] NOAEL estimated using uncertainty factor of 50 (acute LC50 LOAEL to chronic NOAEL).

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

na – not available

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as TRVs.

HQs for endosulfan and endrin are presented in Table A.6-35. The LOAEL-based HQ for endrin was greater than 1.0. LOAEL-based HQs for alpha-endosulfan and beta-endosulfan were less than 1.0. NOAEL-based HQs were greater than 1.0 for all three COPCs. Endosulfan effects data are uncertain because only three studies were

available, and all evaluated the survival endpoint; thus, risk of sublethal effects may be underestimated. However, because the exposure concentrations are probably overestimates of actual exposure (Section A.2.4.5), it is likely that risks associated with organochlorine pesticides are lower than HQs suggest.

Table A.6-35. HQs for English sole and endosulfan and endrin

COPC	UCL Concentration (µg/kg ww)	NOAEL TRV (µg/kg ww)	LOAEL TRV (µg/kg ww)	NOAEL-Based HQ	LOAEL-Based HQ
alpha-endosulfan	4.2	0.62	31	**6.8**	0.14
beta-endosulfan	18	0.62	31	**29**	0.58
endrin	14	1.2	11.5	**12**	**1.2**

COPC – chemical of potential concern
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
TRV – toxicity reference value
UCL – upper confidence limit on the mean
ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

Summary of Uncertainties

Uncertainties in the problem formulation, the effects and exposure assessments, and risk characterization for English sole were evaluated, as summarized below:

- Uncertainties in ROC selection, incidental sediment ingestion, dietary composition, foraging range, water exposure to PAHs and metals, benthic invertebrate tissue data, and future habitat changes resulting from restoration are not expected to have an effect on risk conclusions.
- Three starry flounder composite tissue samples from Area T4 were included in the LDW-wide estimate of total PCB concentration in English sole because starry flounder served as a surrogate species in this area where English sole were more difficult to catch. Inclusion of starry flounder data may have decreased the exposure estimate because concentrations of total PCBs in starry flounder were lower than those in English sole composite tissue samples from the same area.
- A field study investigating effects on growth of English sole exposed to worms previously exposed in the laboratory to clean sediment amended with 0.1% of sediments from the Eagle Harbor, Washington, Superfund site was not selected as a PAH TRV because the worms used were exposed to field-collected sediments with uncharacterized chemicals, and the significance of the effects observed was statistically ambiguous.
- Effects data for cadmium are highly variable, with most studies showing lower sensitivity than the selected TRV. There is also uncertainty associated with the selected TRV because effects were partially attributed to reduced feeding. Thus, risks from cadmium are low but uncertain.

- Risks from exposure to chromium and benzoic acid were low but uncertain because LOAEL TRVs were not identified and NOAEL-based HQs were close to 1.0.
- Phthalates were not detected in English sole tissue samples. Based on an analysis of RLs and shiner surfperch tissue data, risks from phthalates are likely to be very low.
- Risks from exposure to chromium and benzoic acid, for which no LOAELs were available, are likely to be low because NOAEL-based HQs were just over 1.0.
- Risks to English sole from dioxin-like PCB congeners that were estimated using a TEQ approach were very low and would not change risk conclusions for PCBs.
- Because of numerous uncertainties in the study reporting the lowest PCB effects concentrations, risks to English sole from PCBs are uncertain. LOAEL-based HQs based on toxicity data reported in Hugla and Thome (1999) ranged from 0.98 to 5.0.
- Safety factors of greater than 5 used to estimate NOAELs from LOAELs may overestimate the difference between LOAELs and NOAELs.
- There is uncertainty in the critical tissue-residue approach because concentrations may not reflect the site of action, chemicals may be metabolized, and species- and chemical-specific factors may be important.
- An evaluation of risks from exposure to organochlorine pesticides indicated low risks from endrin and endosulfan. The selected TRVs may underestimate the potential for sublethal effects. However, because of analytical interference from PCBs, the risks are likely to be lower than the HQs suggest.
- Site-specific studies suggest the potential for adverse effects on English sole reproduction; however, these effects have not been conclusively associated with specific chemicals or chemical mixtures.

A.6.2.2.3 Risk conclusions

The English sole ROC was evaluated to represent all fish in the LDW not specifically covered by juvenile chinook salmon or Pacific staghorn sculpin. English sole are more highly exposed to sediment-associated chemicals based on their close sediment proximity and diet of benthic invertebrates. To provide a conservative estimate of risk from COPCs in the calculation of HQs, toxicological data from the most sensitive fish species tested for these COPCs were compared to exposure concentrations detected or estimated in English sole. Results of the risk characterization for English sole are summarized in Table A.6-36.

Table A.6-36. Summary of risk characterization for English sole

APPROACH	COPC	NOAEL-BASED HQ	LOAEL-BASED HQ	EFFECTS	PRIMARY UNCERTAINTY
Tissue residue	total PCBs	**4.9 – 25**[a]	0.98 – **5.0**[a]	reduced fecundity; lack of spawning in first reproductive season; egg and larval mortality[b]	high TRV uncertainty[c]
Dietary	arsenic	**1.2**	0.80	reduced body weight	moderate number of toxicity studies
	cadmium	**6.1**	**1.2**	reduced growth rate and condition factor	high TRV uncertainty[d]
	copper	**1.9**	0.93	reduced growth rate	medium TRV uncertainty[e]
	vanadium	**5.9**	**1.2**	reduced body weight	only one toxicity study available

[a] Because of uncertainty in the LOAEL, LOAEL-based HQs were calculated from a range of effects concentrations reported in Hugla and Thome (1999). The NOAEL TRV range was estimated by dividing the LOAEL TRV range by an uncertainty factor of 5.

[b] LOAEL-based HQ range was calculated based on two effects concentrations. The high end of the HQ range was based on a reduced fecundity endpoint; the low end of the HQ range was based on reduced fecundity, egg/larval mortality, and lack of spawning in the first reproductive season.

[c] Results from the studies reporting the lowest LOAELs were uncertain. The study reporting the lowest effects concentrations was uncertain because of uncertain statistical significance of the fecundity endpoint for the low dose, a lack of dose-response in the fecundity endpoint, uncertain number of fish used in the experiment, and uncertainties associated with fish handling and maintenance protocols.

[d] Selected lowest LOAEL and respective NOAEL TRV were two orders of magnitude lower than the next lowest TRVs, and effects were partially attributed to reduced feeding.

[e] A large number of studies that presented a range of effects thresholds were available; sufficient data were available to show that the lowest LOAEL reported for growth of channel catfish (8 mg/kg dw) (Murai et al. 1981) was inconsistent with NOAELs for channel catfish growth reported in other studies (Gatlin and Wilson 1986; Erickson et al. 2003), and thus that LOAEL was not selected.

COPC – chemical of potential concern

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

TRV – toxicity reference value

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

Both the NOAEL- and LOAEL-based HQs were greater than 1.0 for cadmium (6.1 and 1.2, respectively) based on growth of juvenile fish. There is uncertainty in the cadmium risk because the toxicity data are highly variable, and the effects associated with the lowest LOAEL were partially attributed to reduced feeding. Use of the next higher LOAEL and NOAEL TRVs would result in HQs less than 1.0.

Both the NOAEL- and LOAEL-based HQs for vanadium were greater than 1.0 (5.9 and 1.2, respectively). The vanadium LOAEL-based HQ is based on an unbounded LOAEL reported in the only dietary toxicity study identified. The NOAEL TRV was estimated from the unbounded LOAEL TRV using a safety factor of 5 (Section A.4.2.2.4). Because of the very limited effects data, risk from vanadium is uncertain and could be higher or lower than HQs indicate.

For PCBs, the LOAEL-based HQs ranged from 0.98 to 5.0 based on toxicity information presented in one study reporting the lowest TRVs (Hugla and Thome 1999). A NOAEL was not available from Hugla and Thome (1999), and no NOAELs were reported for any other study below the selected LOAEL range. There is uncertainty in the risk estimates because of uncertainties with the Hugla and Thome (1999) study (see Section 6.2.2.2). If the study reporting the next higher LOAEL TRV had been selected, exposures would have been below the effects thresholds (LOAEL-based HQ of 0.28). PCB risks were also calculated using a PCB TEQ approach and were found to be low, although dioxins and furans were not included in the TEQ assessment. Therefore, the risk associated with exposure to all dioxin-like chemicals was underestimated by an unknown amount.

Of the chemicals evaluated solely in the uncertainty analysis, only the endrin risk analyses resulted in a LOAEL-based HQ greater than 1.0 (LOAEL-based HQ of 1.2). The selected endrin TRVs may underestimate risk because they are based on a survival endpoint. Although higher growth or reproduction LOAELs and NOAELs are reported for other fish species, it is not known if other endpoints would have resulted in lower TRVs for largemouth bass. However, because endrin exposures were likely overestimated because of interference from PCBs in the analysis of organochlorine pesticides in tissue samples (Section A.2.4.2), risks are likely to be low.

Results from several studies suggest that some reproductive functions in English sole collected in the LDW may be impaired relative to fish from reference sites (Johnson et al. 1988; 1993; 1997a; Casillas et al. 1991a) (Section A.6.2.2.3). Chemicals implicated as potential causal factors include PAHs and PCBs; however, linking the results of field studies to risks from specific chemicals is difficult considering, among other factors, the complex mixtures of chemicals in the field and the uncertainties in English sole home range. In addition, interpreting cause and effect of the adverse effects reported in field studies is complicated because of genetic variation, health, or seasonal variation in the spawning cycle. As summarized above for juvenile chinook salmon, PAH risks appear to be low based on dietary and water exposure pathways. However, no effects data were available to evaluate PAH effects on reproduction, and the selected dietary TRVs were based on growth endpoints. Effects data evaluated for PCBs included reproduction endpoints; however, because of high TRV uncertainty, risk estimates for PCBs are uncertain.

A.6.2.3 Pacific staghorn sculpin

This section presents risk estimates, uncertainties, and risk conclusions for Pacific staghorn sculpin.

A.6.2.3.1 Risk estimates

This section presents the HQ calculations for Pacific staghorn sculpin. COPCs were evaluated for two foraging range scenarios because the foraging range of Pacific staghorn sculpin is unknown and may be smaller than the entire LDW. The two

scenarios were: 1) foraging throughout the LDW, and 2) foraging in smaller segments of the LDW corresponding to the modeling areas (M1, M2, M3, and M4) shown in Map A.2-2. COPCs evaluated for Pacific staghorn sculpin included arsenic, cadmium, copper, and vanadium, which were evaluated using a dietary approach, and total PCBs and TBT, which were evaluated using a critical tissue-residue approach (Section A.4.0). Dietary exposure concentrations of arsenic were less than both the NOAEL and LOAEL TRVs (maximum HQs of 0.80 and 0.53, respectively) (Table A.6-37).

Table A.6-37. HQ calculations for Pacific staghorn sculpin

Approach	COPC	Foraging Assumption	Exposure Concen-tration	Unit	NOAEL TRV	LOAEL TRV	NOAEL-Based HQ	LOAEL-Based HQ
Tissue residue	total PCBs	LDW-wide	1,100	µg/kg ww	104 – 528	520 – 2,640	**2 – 11**	0.42 – **2.1**
		M1	800	µg/kg ww	104 – 528	520 – 2,640	**1.5 – 7.7**	0.3 – **1.5**
		M2	920	µg/kg ww	104 – 528	520 – 2,640	**1.7 – 8.8**	0.35 – **1.8**
		M3	2,000	µg/kg ww	104 – 528	520 – 2,640	**3.8 – 19**	0.76 – **3.8**
		M4	940	µg/kg ww	104 – 528	520 – 2,640	**1.8 – 9.0**	0.36 – **1.8**
	TBT	LDW-wide	36	µg/kg ww	18	159	**2.0**	0.23
		M1	37	µg/kg ww	18	159	**2.1**	0.23
		M2	36	µg/kg ww	18	159	**2.0**	0.23
		M3	28	µg/kg ww	18	159	**1.6**	0.18
		M4	53	µg/kg ww	18	159	**2.9**	0.33
Dietary	arsenic	LDW-wide	15	mg/kg dw	20	30	0.75	0.50
		M1	15	mg/kg dw	20	30	0.75	0.50
		M2	16	mg/kg dw	20	30	0.80	0.53
		M3	15	mg/kg dw	20	30	0.75	0.50
		M4	12	mg/kg dw	20	30	0.60	0.40
	cadmium	LDW-wide	0.38	mg/kg dw	0.1	0.5	**3.8**	0.76
		M1	0.49	mg/kg dw	0.1	0.5	**4.9**	0.98
		M2	0.30	mg/kg dw	0.1	0.5	**3.0**	0.60
		M3	0.52	mg/kg dw	0.1	0.5	**5.2**	**1.0**
		M4	0.32	mg/kg dw	0.1	0.5	**3.2**	0.64
	copper	LDW-wide	56	mg/kg dw	50	100	**1.1**	0.56
		M1	65	mg/kg dw	50	100	**1.3**	0.65
		M2	77	mg/kg dw	50	100	**1.5**	0.77
		M3	45	mg/kg dw	50	100	0.90	0.45
		M4	48	mg/kg dw	50	100	**1.0**	0.48

Approach	COPC	Foraging Assumption	Exposure Concen-tration	Unit	NOAEL TRV	LOAEL TRV	NOAEL-Based HQ	LOAEL-Based HQ
Dietary, cont.	vanadium	LDW-wide	8.1	mg/kg dw	2.04	10.2	**4.0**	0.79
		M1	8.8	mg/kg dw	2.04	10.2	**4.3**	0.86
		M2	12	mg/kg dw	2.04	10.2	**5.9**	**1.2**
		M3	12	mg/kg dw	2.04	10.2	**5.9**	**1.2**
		M4	6.6	mg/kg dw	2.04	10.2	**3.2**	0.65

COPC – chemical of potential concern
dw – dry weight
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
TBT – tributyltin
TRV – toxicity reference value
ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

Exposure concentrations of total PCBs resulted in LOAEL-based HQs ranging from 0.30 to 3.8 and NOAEL-based HQs ranging from 1.5 to 19. Dietary exposure concentrations of cadmium and vanadium equaled or exceeded both the LOAEL and NOAEL TRVs in at least one modeling area (maximum LOAEL-based HQ of 1.0 and 1.2 for cadmium and vanadium, respectively). Exposure concentrations of TBT and copper exceeded their NOAEL TRVs (maximum NOAEL-based HQs of 2.9 and 1.5, respectively), but none exceeded their LOAEL TRVs.

With the exception of PCBs, cadmium, and vanadium, HQs were similar for all COPCs in all modeling areas and at the LDW-wide scale. The cadmium LOAEL-based HQ was 1.0 in modeling area M3 but less than 1.0 in all other modeling areas and LDW-wide. The vanadium LOAEL-based HQ was greater than 1.0 in modeling areas M2 and M3 but not in M1, M4, or LDW-wide. The PCB NOAEL- and LOAEL-based HQs for modeling area M3 were approximately two times higher than those for the other modeling areas.

A.6.2.3.2 Uncertainty analysis

This section presents a discussion of the uncertainty associated with the problem formulation, the exposure and effects assessments, and the risk characterization for Pacific staghorn sculpin.

Problem Formulation

Primary uncertainties in the problem formulation for Pacific staghorn sculpin include ROC selection, assessment endpoints, and the COPC screen. Uncertainties associated with assessment endpoints are the same as those discussed in Section A.6.2.1.2 for juvenile chinook salmon. Uncertainties associated with the COPC screen are the same as for English sole, as discussed in Section A.6.2.2.2, except that risks for chemicals with no detected concentrations in tissue are discussed below. Uncertainties associated with ROC selection are also discussed below.

COPC screen

Risks to Pacific staghorn sculpin from exposures to BEHP, dimethyl phthalate, and di-n-butyl phthalate in the LDW are uncertain because these chemicals were not detected in Pacific staghorn sculpin tissue samples. RLs for these chemicals were greater than the selected NOAEL TRVs (Table A.2-21). As discussed for English sole, re-analysis of tissue samples with high BEHP RLs, resulting from sample dilutions, showed that BEHP was not detected at much lower RLs. HQs calculated using minimum and maximum RLs relative to the NOAEL TRVs for each chemical are presented in Table A.6-38. LOAEL TRVs were not available for any of these chemicals.

Table A.6-38. Pacific staghorn sculpin HQs for BEHP, dimethyl phthalate, and di-n-butyl phthalate using reporting limits as exposure concentrations

Chemical	Reporting Limit (µg/kg ww)		TRV (µg/kg ww)		NOAEL-Based HQ			
	Min	Max	NOAEL	LOAEL	Min RL	Max RL		
BEHP	490	5,000	390	na	**1.3**	**13**		
Dimethyl phthalate	40	400	498	na	< 0.1	0.80		
Di-n-butyl phthalate	200	1,300[a]	1,170	na	0.17	**1.1**		

[a] Maximum detected concentration is presented.

BEHP – bis(2-ethylhexyl) phthalate
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
na – no TRV available
nc – not calculated because no LOAEL TRV was available
NOAEL – no-observed-adverse-effect level
ROC – receptor of concern
ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0.

NOAEL-based HQs were greater than 1.0 for BEHP and di-n-butylphthalate using maximum RLs and were either just above 1.0 or much lower than 1.0 using minimum RLs. NOAEL-based HQs were less than 1.0 for all dimethylphthalate RLs. Phthalate risks to Pacific staghorn sculpin were low with some uncertainty.

ROC Selection

In Section A.6.2.2.2, it was found that risk estimates for English sole as an ROC were health protective of risks to shiner surfperch for all critical tissue-residue COPCs, except TBT. To evaluate whether risk estimates for Pacific staghorn sculpin as an ROC are protective of risks from TBT to shiner surfperch, NOAEL- and LOAEL-based HQs calculated using both Pacific staghorn sculpin and shiner surfperch tissue data were compared in Table A.6-39. Pacific staghorn sculpin and shiner surfperch HQs for TBT were similar; therefore, shiner surfperch tissue data do not change risk estimates for fish represented by Pacific staghorn sculpin as an ROC.

Table A.6-39. Shiner surfperch and Pacific staghorn sculpin UCLs and HQs for TBT

Species	COPC	Exposure Concentration (µg/kg ww)	NOAEL TRV	LOAEL TRV	NOAEL-Based HQ	LOAEL-Based HQ
Pacific staghorn sculpin	TBT	36	18	159	**2.0**	0.23
Shiner surfperch	TBT	69	18	159	**3.8**	0.43

COPC – chemical of potential concern
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
Bold identifies NOAEL-based HQs greater than 1.0.
TBT – tributyltin
TRV – toxicity reference value
UCL – upper confidence limit on the mean
ww – wet weight

Exposure Assessment

Uncertainties in the exposure assessment for Pacific staghorn sculpin were associated with the following factors:

- Water pathway for metals and PAHs
- Dietary composition
- Future habitat changes resulting from restoration
- Benthic invertebrate tissue data and linear regressions

The uncertainties associated with exposure via the water pathway and future habitat changes are the same as those discussed in Section A.6.2.1.2 for juvenile chinook salmon.

Dietary Composition

Concentrations of COPCs in the diet of Pacific staghorn sculpin are a function of the types of prey consumed and their COPC concentrations. Pacific staghorn sculpin were assumed to ingest fish and benthic invertebrates. According to regional studies, crabs and shrimp constitute 25 to 32% of Pacific staghorn sculpin diets in Puget Sound (Fresh et al. 1979; Miller et al. 1977c; Wingert et al. 1979). For the exposure calculations in Section A.4.1.2, it was assumed that dietary exposure from crabs and shrimp in the LDW was represented by benthic invertebrate tissue samples containing small (≤ 20 mm) crabs and shrimp in addition to other invertebrate species (Windward 2005b, f).[86] Pacific staghorn sculpin in the LDW are likely to prey on some crabs and shrimp > 20 mm. Tissue data for larger slender and Dungeness crabs (≥ 90 mm) collected from the LDW are also available.[87] Although these crabs are larger than prey

[86] Crabs or shrimp were present at 15 of 26 locations where taxonomic samples were collected and constituted from 0.04 to 2.5% of organisms collected at each location.
[87] Data are described in Section A.2.4.1.2.

that Pacific staghorn sculpin could possibly consume, these data provide another means to represent crabs and shrimp consumed by Pacific staghorn sculpin. For dietary COPCs, the UCL concentrations of the crab and benthic invertebrate tissues were generally within a factor of 2, except for vanadium (Table A.6-40). Vanadium concentrations in crab tissue were lower than those in benthic invertebrate tissue for all areas.

Table A.6-40. Comparison of crab and benthic invertebrate tissue concentrations for Pacific staghorn sculpin dietary COPCs

COPC	Foraging Assumption	UCL Concentration in Crab Tissue (mg/kg dw)	UCL Concentration in Benthic Invertebrate Tissue (mg/kg dw)
Arsenic	LDW-wide	25	24
	M1	36	23
	M2	17	25
	M3	17	23
	M4	27	18
Cadmium	LDW-wide	1.1	0.60
	M1	2.2	0.81
	M2	0.95	0.48
	M3	0.71	0.77
	M4	1.2	0.54
Copper	LDW-wide	90	92
	T1	110	110
	T2	75	130
	T3	63	65
	T4	52	81
Vanadium	LDW-wide	1.0	12
	M1	1.1	12
	M2	1.3	18
	M3	1.0	20
	M4	0.5[a]	10

[a] Maximum RL for single non-detected concentration.

COPC – chemical of potential concern

dw – dry weight

UCL – upper confidence limit on the mean

Risks for Pacific staghorn sculpin were evaluated using both crab and benthic invertebrate tissue data for arsenic, cadmium, copper, and vanadium for locations where exposure concentrations of these tissue types differed by >10% (Table A.6-41). When crab tissue data were included in arsenic risk calculations, risk predictions were similar to those predicted without crab data. For cadmium, the LOAEL-based HQ changed from < 1.0 to > 1.0 for LDW-wide and modeling areas M1 and M4 when crab data were included. The HQs for vanadium changed from > 1.0 to < 1.0 for M2 and

M3 when crab tissue data were included in exposure calculations. The HQs for copper changed from 1.0 to < 1.0 when crab data were included in exposure calculations in M4, but no change occurred in M2. Therefore, inclusion of crab tissue data would increase the Pacific staghorn sculpin risk estimates for cadmium for the LDW as a whole and for modeling areas M1 and M4, and would decrease the risk estimates for vanadium for modeling areas M2 and M3 and for copper for modeling area M4. The inclusion of crab tissue data would not affect risk conclusions for arsenic.

Table A.6-41. HQs for Pacific staghorn sculpin for arsenic, cadmium, copper, and vanadium with and without the use of larger crab tissue data

COPC	Foraging Assumption	Alternative Diet[a]		Diet Assumed in Exposure Assessment[b]	
		NOAEL-Based HQ	LOAEL-Based HQ	NOAEL-Based HQ	LOAEL-Based HQ
Arsenic	M1	0.75	0.50	0.75	0.50
	M2	0.82	0.55	0.80	0.53
	M3	0.73	0.49	0.75	0.50
	M4	0.59	0.39	0.60	0.40
Cadmium	LDW	**5.4**	**1.1**	**3.8**	0.76
	M1	**9.3**	**1.9**	**4.9**	0.98
	M2	**4.5**	0.90	**3.0**	0.60
	M4	**5.3**	**1.1**	**3.2**	0.64
Copper	M2	**1.2**	0.58	**1.5**	0.77
	M4	0.76	0.38	**1.0**	0.48
Vanadium	LDW	**2.3**	0.45	**4.0**	0.79
	M1	**2.6**	0.52	**4.3**	0.86
	M2	**3.3**	0.67	**5.9**	**1.2**
	M3	**3.1**	0.63	**5.9**	**1.2**
	M4	**1.8**	0.35	**3.2**	0.65

[a] Assuming a diet of 32% crabs, 23% benthic invertebrates, 44% fish, and 1% sediment.

[b] Assuming a diet of 55% benthic invertebrates, 44% fish, and 1% sediment.

COPC – chemical of potential concern

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

Benthic Invertebrate Tissue Data and Linear Regressions

The uncertainties associated with the representativeness of benthic invertebrate tissue data are the same as those discussed in Section A.6.2.1.2 for juvenile chinook salmon, with one difference. For Pacific staghorn sculpin, fewer benthic invertebrate tissue samples were available to calculate UCL concentrations for each modeling area (n = 10, 6, 4, and 4 for modeling areas M1, M2, M3, and M4, respectively) because

benthic invertebrate sampling locations were selected to be representative of arsenic, lead, and PCB concentrations in sediment rather than an even distribution of sampling locations throughout the LDW. The relatively low number of benthic invertebrate tissue samples results in higher uncertainty in the UCL in exposure areas where there is high variability in the tissue concentrations. The ProUCL guidelines recommend using the UCL even if it is greater than the maximum concentration (EPA 2004a). In one instance, ProUCL calculated a UCL that was greater than the maximum concentration (for vanadium in modeling area M4); however, this difference was small, and risk conclusions would not have changed if maximum values had been used instead of the UCL.

If the exposure estimate had been calculated using all empirical benthic invertebrate tissue data rather than the regression, the benthic invertebrate UCL would have been 37 mg/kg dw, in comparison to the UCL of 24 mg/kg dw estimated using the regression. Assuming this value in the exposure calculation for benthic invertebrates and 4.2 mg/kg dw for shiner surfperch results in an exposure concentration of 22 mg/kg dw in comparison to 15 mg/kg dw calculated using the regression-estimated benthic invertebrate tissue concentration. Based on this exposure concentration, the LDW-wide NOAEL-based HQ would have increased from 0.75 to 1.1, and the LDW-wide LOAEL-based HQ would have increased from 0.50 to 0.73. Therefore, this uncertainty would not have substantially affected the risk conclusions for arsenic.

Effects Assessment

Uncertainties in the effects assessment for Pacific staghorn sculpin were associated with the following factors:

- Effects from chemical mixtures and estimation of NOAELs from LOAELs
- Exclusion of field studies from TRV selection
- Estimation of NOAELs from LOAELs
- PCB TRV
- Cadmium TRV
- COPCs without LOAEL TRVs
- TEQ approach for PCBs
- Critical tissue residue approach

All of the above uncertainties associated with the effects assessment for Pacific staghorn sculpin were assessed in Section A.6.2.2.2 for English sole, except that COPCs without LOAEL TRVs are assessed below.

In addition, some of the selected TRVs are considered less certain than others if there were a small number of studies, if TRVs were based on less-protective measures such as LC50s, or if data quality was questionable. The relative uncertainties in the selected TRVs for each COPC are the same as those discussed in Section A.6.2.2.2 for English

sole, with the exception of TRVs for TBT. For TBT, only three studies were available. In the study reporting the lowest LOAEL, which was selected as the TRV, treatments were not replicated, and there was high mortality in all treatments and in the controls. The effect of these uncertainties on risk conclusions is unknown.

PCB TRV

Uncertainties associated with the PCB TRV for Pacific staghorn sculpin are the same as those discussed in Section A.6.2.2 for English sole. If the study with the next lowest LOAEL (9,300 µg/kg ww reported in Hansen et al. (1974a)) had been selected as the LOAEL TRV, the LDW-wide LOAEL-based HQ for Pacific staghorn sculpin would have changed from a range of 0.42 to 2.1 based on the selected LOAEL range to a LOAEL-based HQ of 0.12.

COPCs without LOAEL TRVs

As discussed in Section A.2.5.2, four chemicals had maximum exposure concentrations exceeding NOAEL TRVs but LOAEL TRVs were not available. One chemical (chromium) was evaluated using a dietary approach, and three chemicals (benzoic acid, dimethyl phthalate, and di-n-butyl phthalate) were evaluated using a critical tissue-residue approach. Risks to Pacific staghorn sculpin from dimethyl phthalate and di-n-butyl phthalate were evaluated in the exposure assessment uncertainty section above because concentrations of these chemicals were generally below detection limits in fish tissue samples.

Table A.6-42 presents chromium dietary concentrations for Pacific staghorn sculpin, which were calculated using Equation 4-1, as described in Section A.4.1.1.2. Benzoic acid critical tissue-residue concentrations were calculated as the UCL of the mean of all Pacific staghorn sculpin tissue composite samples from throughout the LDW. NOAEL-based HQs ranged from 0.18 to 2.1 for chromium and from 1.5 to 2.1 for benzoic acid (Table A.6-42). Because no LOAEL toxicity data were available for these chemicals, the low NOAEL-based HQs calculated using unbounded NOAEL TRVs are assumed to indicate low risks.

Table A.6-42. Pacific staghorn sculpin NOAEL-based HQs for chemicals for which no LOAEL TRVs were identified

COPC	Units	Dietary Exposure Concentration	NOAEL TRV	NOAEL-Based HQ
Chromium	mg/kg dw	8.0	9.4[a]	0.85
Benzoic acid	µg/kg ww	7,000	3,380[b]	**2.1**

[a] No effects on growth of 2-year-old gray mullet exposed to chromium III through diet and sediment for 8 weeks (Walsh et al. 1994).

[b] No effects on survival of mosquitofish exposed in a mesocosm for 24 hours (Lu and Metcalf 1975).

dw – dry weight
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
TRV – toxicity reference value
ww – wet weight
Bold identifies NOAEL-based HQs greater than 1.0.

Risk Characterization

Risks to Pacific staghorn sculpin from organochlorine pesticides were not included in the risk estimates because of high uncertainty in the JN-qualified tissue pesticide data resulting in suspected false identifications of presence of some pesticides as well as overestimates in their concentrations, as discussed in Section A.2.4.2. Exposure estimates and risk estimates are discussed in this section for the three organochlorine pesticides (alpha-endosulfan, beta-endosulfan, and endrin) that were identified as COPCs in Section A.2.5.2. TRVs for these pesticides are the same as those discussed in Section A.6.2.2.2 for English sole.

HQs for endosulfan and endrin are presented in Table A.6-43. LOAEL-based HQs were greater than 1.0 for endrin and less than 1.0 for alpha-endosulfan and beta-endosulfan. NOAEL-based HQs were greater than 1.0 for all three of these COPCs. The selected endrin TRVs may underestimate risks because they are based on a survival endpoint. Although higher growth or reproduction LOAELs and NOAELs are reported for other fish species, it is not known if other endpoints would have resulted in lower TRVs for largemouth bass. Endosulfan effects data are uncertain because only three studies were available and all evaluated the survival endpoint; thus, risks from sublethal endpoints may be underestimated. However, because the exposure concentrations are likely overestimates of actual exposure (Section A.2.4.2), it is likely that risks associated with organochlorine pesticides are lower than these HQs suggest.

Table A.6-43. HQs for Pacific staghorn sculpin and endosulfan and endrin

COPC	UCL Concentration (µg/kg ww)	NOAEL TRV (µg/kg ww)	LOAEL TRV (µg/kg ww)	NOAEL-Based HQ	LOAEL-Based HQ
alpha-Endosulfan	1.4	0.62	31	**2.3**	<0.1
beta-Endosulfan	4.1	0.62	31	**6.6**	0.13
Endrin	36	1.2	11.5	**31**	**3.1**

COPC – chemical of potential concern
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
TRV – toxicity reference value
UCL – upper confidence limit on the mean
ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

Summary of Uncertainties

Uncertainties in the problem formulation, the effects and exposure assessments, and risk characterization for Pacific staghorn sculpin were evaluated, as summarized below:

- Uncertainties in the COPC screen, ROC selection, water exposure to PAHs and metals, PAH field studies, PCB analysis approach (total PCBs versus PCB TEQ),

and future habitat changes resulting from restoration are not expected to have an effect on risk conclusions.

- Effects data for cadmium are highly variable with most studies showing lower sensitivity than the selected TRV. Because the relative sensitivities of Pacific staghorn sculpin, fish represented by Pacific staghorn sculpin, and fish used in the toxicity studies is unknown, it is not known if the conservative TRV selected overestimates risk.
- When the small crabs and shrimps in the invertebrate portion of Pacific staghorn sculpin diets were represented by Phase 2 crab tissue data rather than benthic invertebrate tissue data (which include crabs and shrimps), the LDW-wide, modeling area M1, and modeling area M4 LOAEL-based HQs increased from slightly less than 1.0 to slightly greater than 1.0 for cadmium; decreased from slightly greater than 1.0 to slightly less than 1.0 for modeling areas M2 and M3 for vanadium; and the NOAEL-based HQ decreased from 1.0 to slightly less than 1.0 for modeling area M4 for copper. For arsenic, no HQs changed from less than 1.0 to greater than 1.0 or vice versa.
- The use of safety factors of greater than 5 to estimate NOAELs from LOAELs may overestimate the difference between LOAELs and NOAELs.
- Because of numerous uncertainties in the study reporting the lowest PCB effects concentrations, the estimated risk to Pacific staghorn sculpin from PCBs is uncertain. LOAEL-based HQs based on toxicity data reported in Hugla and Thome (1999) and sculpin exposures in specific areas in the LDW ranged from 0.30 to 3.8.
- Risks from exposure to chromium and benzoic acid, for which no LOAELs were available, are likely to be very low for chromium (NOAEL-based HQ < 1.0) and low for benzoic acid.
- There is uncertainty in the critical tissue-residue approach because concentrations may not reflect the site of action, chemicals may be metabolized, and species- and chemical-specific factors may be important.
- An evaluation of risks from exposure to organochlorine pesticides indicated a low risk from endosulfan and the potential for adverse effects from endrin (LOAEL-based HQ of 3.1). The selected TRVs are based on a survival endpoint and may underestimate the potential for sublethal effects. However, because of analytical interference from PCBs, the risks are likely to be lower than the HQs suggest.

A.6.2.3.3 Risk conclusions

Pacific staghorn sculpin, a benthic omnivorous fish, was selected to represent upper-trophic-level fish in the LDW. This distinction was made because upper-trophic-level fish may have higher body burdens of biomagnifying chemicals than fish such as English sole that consume primarily small invertebrates at a lower trophic level. Two exposure scenarios were evaluated for Pacific staghorn sculpin because their foraging range is uncertain. The scenarios were: 1) foraging throughout the LDW, and 2) foraging in smaller segments of the LDW corresponding to the modeling areas (M1, M2, M3, and M4) shown in Map A.2-2. Results of the risk characterization for Pacific staghorn sculpin are summarized in Table A.6-44.

Table A.6-44. Summary of risk characterization for Pacific staghorn sculpin

Approach	COPC	Foraging Assumption	NOAEL-Based HQ	LOAEL-Based HQ	LOAEL Endpoint	Primary Uncertainty
Tissue residue	total PCBs	LDW-wide	**2 – 11**[a]	0.42 – **2.1**[a]	reduced fecundity; lack of spawning in first reproductive season; egg and larval mortality[b]	high TRV uncertainty[c]
		M1	**1.5 – 7.7**[a]	0.30 – **1.5**[a]		
		M2	**1.7- 8.8**[a]	0.35 – **1.8**[a]		
		M3	**3.8 – 19**[a]	0.76 – **3.8**[a]		
		M4	**1.8 – 9.0**[a]	0.36 – **1.8**[a]		
	TBT	LDW-wide	**2.0**	0.23	reduced juvenile body weight	few toxicity studies available
		M1	**2.1**	0.23		
		M2	**2.0**	0.23		
		M3	**1.6**	0.18		
		M4	**2.9**	0.33		
Dietary	arsenic	LDW-wide	0.75	0.50	reduced juvenile body weight	moderate number of toxicity studies
		M1	0.75	0.50		
		M2	0.80	0.53		
		M3	0.75	0.50		
		M4	0.60	0.40		
	cadmium	LDW-wide	**3.8**	0.76	reduced juvenile body weight and length growth rate; and condition factor	high TRV uncertainty,[d] inclusion of crab data in exposure calculations[e]
		M1	**4.9**	0.98		
		M2	**3.0**	0.60		
		M3	**5.2**	**1.0**		
		M4	**3.2**	0.64		
	copper	LDW-wide	**1.1**	0.56	reduced juvenile body weight growth rate	medium TRV uncertainty[f]
		M1	**1.3**	0.65		
		M2	**1.5**	0.77		
		M3	0.90	0.45		
		M4	**1.0**	0.48		

Approach	COPC	Foraging Assumption	NOAEL-Based HQ	LOAEL-Based HQ	LOAEL Endpoint	Primary Uncertainty
Dietary, cont.	vanadium	LDW-wide	**4.0**	0.79	reduced juvenile body weight	only one toxicity study available
		M1	**4.3**	0.86		
		M2	**5.9**	**1.2**		
		M3	**5.9**	**1.2**		
		M4	**3.2**	0.65		

[a] Because of uncertainty in the LOAEL, LOAEL-based HQs were calculated from a range of effects concentrations reported in Hugla and Thome (1999). The NOAEL TRV range was estimated by dividing the LOAEL TRV range by an uncertainty factor of 5.

[b] LOAEL-based HQ range was calculated based on two effects concentrations. The high end of the HQ range was based on a reduced fecundity endpoint; the low end of the HQ range was based on reduced fecundity, egg/larval mortality, and lack of spawning in the first reproductive season.

[c] Results from the studies reporting the lowest LOAELs were uncertain. The study reporting the lowest effects concentrations was uncertain because of uncertain statistical significance of the fecundity endpoint for the low dose, a lack of dose-response in the fecundity endpoint, uncertain number of fish used in the experiment, and uncertainties associated with fish handling and maintenance protocols.

[d] Selected LOAEL and respective NOAEL TRV were two orders of magnitude lower than the next lowest TRVs, and effects were partially attributed to reduced feeding.

[e] When the small crabs and shrimps in the invertebrate portion of Pacific staghorn sculpin diets were represented by Phase 2 crab tissue data rather than benthic invertebrate tissue data (which include crabs and shrimps), the cadmium LOAEL-based HQ increased from slightly less than 1.0 to slightly greater than 1.0 for the LDW-wide exposure area and modeling area M1. Vanadium LOAEL-based HQs decreased from slightly greater than 1.0 to slightly less than 1.0 for modeling areas M2 and M3 when crab data were included.

[f] A large number of studies that presented a range of effects thresholds were available, and sufficient data were available to suggest that the lowest LOAEL was not supported by data from other studies.

COPC – chemical of potential concern
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
TBT – tributyltin
TRV – toxicity reference value

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

For PCBs, the LOAEL-based HQs ranged from 0.30 to 3.8 based on toxicity information presented in one study reporting the lowest TRVs (Hugla and Thome 1999) and exposure data from discrete areas in the LDW as well as LDW-wide. A NOAEL was not available from Hugla and Thome (1999), and no NOAELs below the selected LOAEL range were reported for any other study. There is uncertainty in the risk estimate because the study with the lowest effects concentrations was highly uncertain (see Section A.6.2.2.2). If the study reporting the next higher LOAEL TRV had been selected, exposures would have been below the effects threshold (LOAEL-based HQ of 0.28).

Both the NOAEL- and LOAEL-based HQs were greater than 1.0 for vanadium in modeling areas M2 and M3 (NOAEL- and LOAEL-based HQs of 1.2 and 5.9, respectively, in both areas). In modeling areas M1 and M4 and LDW-wide, NOAEL-based HQs were greater than 1.0, but LOAEL-based HQs were less than or equal to 1.0. As discussed above for English sole, because of the very limited effects data, risk from vanadium is uncertain and could be higher or lower than HQs indicate.

The LOAEL based HQ for cadmium was equal to 1.0 in modeling area M3 but lower than 1.0 in all other modeling areas and LDW-wide.

NOAEL-based HQs were greater than 1.0 but LOAEL-based HQs were less than or equal to 1.0 for TBT and copper in at least one modeling area. Uncertainty in data used to represent crabs in sculpin dietary exposure calculations could result in a cadmium LOAEL-based HQ of up to 1.9. Of the chemicals evaluated in the uncertainty analysis, only the endrin risk analysis resulted in a LOAEL-based HQ greater than 1.0 (LOAEL-based HQ of 3.1). The selected endrin TRVs may underestimate risk because they are based on a survival endpoint. Higher growth or reproduction LOAELs and NOAELs are reported for other fish species; it is not known if other endpoints would have resulted in lower TRVs for largemouth bass. However, endrin exposures are likely overestimated because of interference from PCBs in the analysis of organochlorine pesticides in tissue samples (Section A.2.4.2), so risks are likely to be low.

A.6.2.4 Summary of risk conclusions for fish

In summary, results of the risk estimates and evaluation of associated uncertainties for fish are as follows:

- Exposure concentrations of cadmium in the English sole and Pacific staghorn sculpin diets in modeling area M3 were equal to or greater than those associated with adverse effects, although there is uncertainty in selected TRVs. Low risks were estimated for cadmium and Pacific staghorn sculpin and juvenile chinook salmon LDW-wide and in all other modeling areas.
- Exposure concentrations of vanadium in the English sole and Pacific staghorn sculpin diets in modeling areas M2 and M3 were greater than those associated with adverse effects. Risks were very low (NOAEL-based HQs < 1.0) for juvenile chinook salmon and for Pacific staghorn sculpin in modeling areas M1, M4, and LDW-wide. A paucity of effects studies make these risks somewhat uncertain.
- There is high uncertainty in the PCB TRVs used to evaluate risks to English sole and Pacific staghorn sculpin, and thus a range of LOAEL TRVs was selected from the study reporting the lowest effects concentrations. Exposure concentrations of PCBs in English sole and Pacific staghorn sculpin tissue were in between the concentrations selected as LOAELs from this study. Therefore, there is a potential for adverse effects from PCBs, but risk estimates are uncertain. Risks to juvenile chinook salmon from PCBs were estimated to be low. Risks to all three fish ROCs from dioxin-like PCB congeners using the TEQ approach were also low.
- No quantitative ecological risk estimates were calculated for dioxins and furans and thus the level of ecological risk from dioxins and furans is unknown.

- Exposure concentrations of TBT were less than those associated with adverse effects for Pacific staghorn sculpin. TBT was not identified as a COPC for English sole and juvenile chinook salmon in the problem formulation.
- Exposure concentrations of arsenic and copper were less than those associated with adverse effects for all fish ROCs.
- Risks from chromium, organochlorine pesticides, benzoic acid, PAHs, and some phthalates were evaluated in the uncertainty analysis because of uncertainties in exposure or effects data. Considering the uncertainty, risk estimates for these COPCs appear to be low to very low.
- Various exposure assumptions were evaluated in the uncertainty analysis. None of the alternative assumptions had a large effect on risk conclusions.
- Site-specific studies suggest the potential for adverse effects on English sole reproduction; however, these effects have not been conclusively associated with specific chemicals or chemical mixtures.
- Site-specific studies investigating growth or immunocompetence of juvenile chinook salmon are uncertain and also have not been definitively associated with specific chemicals or chemical mixtures.

A.6.3 RISK CHARACTERIZATION FOR WILDLIFE

This section presents a risk characterization and uncertainty analysis for each of the five wildlife ROCs. The assessment for each ROC estimates risk by calculating HQs using estimated ingested doses of COPCs, as described in Section A.5.1, and TRVs, as presented in Section A.5.2. Uncertainties in the exposure and effects data that may result in overestimates or underestimates of risk for each of the COPCs are discussed. Risk conclusions are presented for each ROC that integrate risk estimates with associated uncertainties.

A.6.3.1 Spotted sandpiper

This section presents risk estimates, uncertainties, and risk conclusions for spotted sandpiper.

A.6.3.1.1 Risk estimates

This section presents the HQ calculations for spotted sandpiper. Ingested doses, NOAEL and LOAEL TRVs, and HQs are presented in Table A.6-45.

Table A.6-45. HQ calculations for spotted sandpiper in the six exposure scenarios evaluated

COPC	Exposure Scenario[a]	Ingested Dose (mg/kg bw/day)	Contribution to Ingested Dose (%)[b, c]		TRVs (mg/kg bw/day)		NOAEL-Based HQ	LOAEL-Based HQ
			Prey	Sediment	NOAEL	LOAEL		
Arsenic	Area 1/high	4.6	78	22	10	40	0.46	0.12
	Area 1/high and poor	3.8	87	13	10	40	0.38	0.10
	Area 2/high	5.3	74	26	10	40	0.53	0.13
	Area 2/high and poor	3.8	87	13	10	40	0.38	0.10
	Area 3/high	3.4	84	16	10	40	0.34	<0.1
	Area 3/high and poor	5.8	71	29	10	40	0.58	0.15
Cadmium	Area 1/high	0.10	87	13	1.5	4.0	< 0.1	< 0.1
	Area 1/high and poor	0.10	83	17	1.5	4.0	< 0.1	< 0.1
	Area 2/high	0.090	69	31	1.5	4.0	< 0.1	< 0.1
	Area 2/high and poor	0.10	75	25	1.5	4.0	< 0.1	< 0.1
	Area 3/high	0.12	85	15	1.5	4.0	< 0.1	< 0.1
	Area 3/high and poor	0.32	31	69	1.5	4.0	0.21	< 0.1
Chromium	Area 1/high	1.4	34	66	1.0	5.0	**1.4**	0.28
	Area 1/high and poor	1.3	36	64	1.0	5.0	**1.3**	0.26
	Area 2/high	4.0	70	30	1.0	5.0	**4.0**	0.80
	Area 2/high and poor	8.8	90	10	1.0	5.0	**8.8**	**1.8**
	Area 3/high	1.6	47	53	1.0	5.0	**1.6**	0.32
	Area 3/high and poor	4.1	19	81	1.0	5.0	**4.1**	0.82
Cobalt	Area 1/high	0.53	53	47	2.3	23.1	0.23	<0.1
	Area 1/high and poor	0.50	56	44	2.3	23.1	0.22	< 0.1
	Area 2/high	0.58	54	46	2.3	23.1	0.25	< 0.1
	Area 2/high and poor	0.57	61	39	2.3	23.1	0.25	< 0.1
	Area 3/high	0.57	61	39	2.3	23.1	0.25	< 0.1
	Area 3/high and poor	0.62	55	45	2.3	23.1	0.27	< 0.1

COPC	Exposure Scenario[a]	Ingested Dose (mg/kg bw/day)	Contribution to Ingested Dose (%)[b, c]		TRVs (mg/kg bw/day)		NOAEL-Based HQ	LOAEL-Based HQ
			Prey	Sediment	NOAEL	LOAEL		
Copper	Area 1/high	21	88	12	21	29	1.0	0.72
	Area 1/high and poor	21	89	11	21	29	1.0	0.72
	Area 2/high	25	87	13	21	29	**1.2**	0.86
	Area 2/high and poor	24	91	9	21	29	**1.1**	0.83
	Area 3/high	13	86	14	21	29	0.62	0.45
	Area 3/high and poor	31	35	65	21	29	**1.5**	**1.1**
Lead	Area 1/high	7.4	66	34	5.82	20	**1.3**	0.37
	Area 1/high and poor	7.4	66	34	5.82	20	**1.3**	0.37
	Area 2/high	6.8	35	65	5.82	20	**1.2**	0.34
	Area 2/high and poor	110	98	2	5.82	20	**19**	**5.5**
	Area 3/high	3.4	26	74	5.82	20	0.58	0.17
	Area 3/high and poor	29	3	97	5.82	20	**5.0**	**1.5**
Mercury	Area 1/high	0.020	77	23	0.018	0.091	**1.1**	0.22
	Area 1/high and poor	0.020	78	22	0.018	0.091	**1.1**	0.22
	Area 2/high	0.022	47	53	0.018	0.091	**1.2**	0.24
	Area 2/high and poor	0.019	58	42	0.018	0.091	**1.1**	0.21
	Area 3/high	0.095	83	17	0.018	0.091	**5.3**	**1.0**
	Area 3/high and poor	0.090	88	12	0.018	0.091	**5.0**	0.99
Nickel	Area 1/high	1.1	48	52	17	33	< 0.1	< 0.1
	Area 1/high and poor	1.4	39	61	17	33	< 0.1	< 0.1
	Area 2/high	1.5	60	40	17	33	< 0.1	< 0.1
	Area 2/high and poor	1.4	60	40	17	33	< 0.1	< 0.1
	Area 3/high	1.8	65	35	17	33	0.11	< 0.1
	Area 3/high and poor	3.7	31	69	17	33	0.22	0.11

COPC	Exposure Scenario[a]	Ingested Dose (mg/kg bw/day)	Contribution to Ingested Dose (%)[b, c]		TRVs (mg/kg bw/day)		NOAEL-Based HQ	LOAEL-Based HQ
			Prey	Sediment	NOAEL	LOAEL		
Selenium	Area 1/high	0.32	95	5	0.50	0.82	0.64	0.39
	Area 1/high and poor	0.31	95	5	0.50	0.82	0.62	0.38
	Area 2/high	0.45	56	44	0.50	0.82	0.90	0.55
	Area 2/high and poor	0.37	77	23	0.50	0.82	0.74	0.45
	Area 3/high	0.41	53	47	0.50	0.82	0.82	0.50
	Area 3/high and poor	0.40	55	45	0.50	0.82	0.80	0.49
Vanadium	Area 1/high	2.5	40	60	1.2	2.3	**2.1**	**1.1**
	Area 1/high and poor	2.4	41	59	1.2	2.3	**2.0**	**1.0**
	Area 2/high	3.2	49	51	1.2	2.3	**2.7**	**1.4**
	Area 2/high and poor	3.1	50	50	1.2	2.3	**2.6**	**1.3**
	Area 3/high	3.1	48	52	1.2	2.3	**2.6**	**1.3**
	Area 3/high and poor	3.1	48	52	1.2	2.3	**2.6**	**1.3**
Zinc	Area 1/high	35	85	15	82	124	0.43	0.28
	Area 1/high and poor	34	86	14	82	124	0.41	0.27
	Area 2/high	53	87	13	82	124	0.65	0.43
	Area 2/high and poor	64	92	8	82	124	0.78	0.52
	Area 3/high	46	91	9	82	124	0.56	0.37
	Area 3/high and poor	62	68	32	82	124	0.76	0.50
Total PCBs	Area 1/high	0.25	96	4	0.49	1.4	0.51	0.18
	Area 1/high and poor	0.25	96	4	0.49	1.4	0.51	0.18
	Area 2/high	1.0	93	7	0.49	1.4	**2.0**	0.71
	Area 2/high and poor	0.64	94	6	0.49	1.4	**1.3**	0.46
	Area 3/high	0.45	96	4	0.49	1.4	0.92	0.32
	Area 3/high and poor	0.57	95	5	0.49	1.4	**1.2**	0.41

COPC	Exposure Scenario[a]	Ingested Dose (mg/kg bw/day)	Contribution to Ingested Dose (%)[b, c]		TRVs (mg/kg bw/day)		NOAEL-Based HQ	LOAEL-Based HQ
			Prey	Sediment	NOAEL	LOAEL		
PCB TEQs	Area 1/high	2.6×10^{-5}	96	4	1.4×10^{-5}	1.43×10^{-4}	**1.9**	0.18
	Area 1/high and poor	2.6×10^{-5}	96	4	1.4×10^{-5}	1.43×10^{-4}	**1.9**	0.18
	Area 2/high	2.1×10^{-4}	43	57	1.4×10^{-5}	1.43×10^{-4}	**15**	**1.5**
	Area 2/high and poor	1.4×10^{-4}	64	36	1.4×10^{-5}	1.43×10^{-4}	**10**	0.98
	Area 3/high	6.4×10^{-5}	96	4	1.4×10^{-5}	1.43×10^{-4}	**4.6**	0.45
	Area 3/high and poor	1.4×10^{-4}	43	57	1.4×10^{-5}	1.43×10^{-4}	**10**	0.98

[a] Six exposure scenarios were evaluated; in each of three exposure areas, foraging in high-quality habitat only and foraging in both high- and poor-quality habitat were evaluated. These exposure scenarios are described in detail in Section A.5.1.2.1.

[b] The percent contribution of ingested dose from water is very low.

[c] The percent contribution of ingested dose from prey tissue and sediment is based on both chemical concentration and intake rate.

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

PCB – polychlorinated biphenyl

TEQ – toxic equivalent

TRV – toxicity reference value

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

Twelve COPCs were evaluated for spotted sandpiper: eleven metals and PCBs. Six metals (i.e., arsenic, cadmium, cobalt, nickel, selenium, and zinc) had both NOAEL- and LOAEL-based HQs that were less than 1.0 in all exposure areas. The NOAEL-based HQs ranged from < 0.1 to 0.90 for these six metals.

PCBs were evaluated both as total PCBs and as PCB TEQs. HQs for PCB TEQs were greater than those for total PCBs. For total PCBs, NOAEL-based HQs were greater than 1.0 in some areas, but none of the LOAEL-based HQs were greater than 1.0. The NOAEL-based HQs for total PCBs that were greater than 1.0 ranged from 1.2 to 2.0 in Areas 2 and 3. For PCB TEQs, all NOAEL-based HQs were greater than 1.0, ranging from 1.9 in Area 1 to 15 in Area 2. Only one LOAEL-based HQ exceeded 1.0; this HQ was 1.5 in Area 2.

The remaining five COPCs (chromium, copper, lead, mercury, and vanadium) had LOAEL-based HQs greater than or equal to 1.0 for spotted sandpiper. LOAEL-based HQs ranged from 1.0 to 5.5 for these COPCs in at least one area in the LDW, indicating risks to spotted sandpiper in those areas. LOAEL-based HQs were greater than or equal to 1.0 in Area 2 for chromium (1.8) and lead (5.5), in Area 3 for copper (1.1) and mercury (1.0), and in all three exposure areas for vanadium (1.0 to 1.4). As shown in Table A.6-45, the highest LOAEL-based HQs (1.8 for chromium and 5.5 for lead) were based primarily on elevated concentrations in prey (benthic invertebrates).

A.6.3.1.2 Uncertainties

This section presents a discussion of uncertainties associated with the problem formulation, the exposure and effects assessments, and the risk characterization for spotted sandpiper.

Problem Formulation

The primary uncertainties in the problem formulation for spotted sandpiper are associated with ROC selection and the COPC screen.

ROC Selection

Uncertainties related to how well spotted sandpiper represents other benthivorous birds in the LDW were addressed in the Phase 1 ERA (Windward 2003b). In that analysis, it was concluded that spotted sandpiper is expected to have an exposure that is similar or higher than those of other benthivorous LDW bird species because its diet consists primarily of benthic invertebrates, it has a high sediment ingestion rate and a high body-weight-normalized FIR, and its home range is within the LDW during the nesting season. Thus, risk estimates for spotted sandpiper in the LDW should be higher than would be calculated for other species in the LDW with different diets, lower sediment and FIRs, and less frequent site use.

COPC Screen

Of the chemicals screened out in the first step of the COPC screen (Section A.2.5.3), 20 chemicals were detected in sediment but were not analyzed in tissue samples (Table A.6-16). None of these chemicals are defined as bioaccumulative chemicals (EPA 2000a), so exposure to these chemicals is assumed to be very low.

Eighty-six chemicals were identified as COIs for birds. Effects data for birds were not available for 40 of the COIs, including 20 individual PAHs. Risks to birds from PAHs were evaluated using TRVs for total PAHs and benzo(a)pyrene. The remaining COIs for which there were no TRVs for birds included antimony, silver, three organotin compounds, eleven SVOCs, and four organochlorine pesticides. Risks to birds from exposure to these COIs could not be evaluated.

Two bioaccumulative chemicals, chromium VI and methylmercury, were detected in more than 5% of the surface sediment samples in which they were analyzed;[88] they were not analyzed in tissue. Therefore, these chemicals were evaluated as components of total chromium and total mercury, respectively. The exposure concentrations of total mercury were compared to TRVs for methylmercury for birds and mammals, so risks from methylmercury exposure may have been overestimated. The only avian or mammalian toxicity studies for chromium VI were based on subchronic or drinking water exposures, so the selected TRVs were for chromium III. Therefore, risk could be underestimated for total chromium if chromium VI is the dominant oxidation state of chromium in LDW benthic invertebrates or fish.

Exposure Assessment

Uncertainties in the exposure assessment for spotted sandpiper were associated with the following factors:

- Direct sediment contact
- Incidental sediment ingestion rate
- COPC bioavailability
- Dietary composition
- Site use
- Benthic invertebrate tissue data and linear regressions
- TEQ approach

These uncertainties are discussed in detail below.

[88] Chromium VI was analyzed in seven sediment samples in the vicinity of Harbor Island, and methylmercury was analyzed in 20 sediment samples in the vicinity of Duwamish/Diagonal and Norfolk.

Direct Sediment Contact

Risks to wildlife from direct contact with sediment are considered insignificant relative to risks from incidental sediment ingestion (EPA 2000b). However, the exclusion of this pathway adds a small amount of uncertainty to the risk estimate for spotted sandpiper.

Incidental Sediment Ingestion Rate

Uncertainty in the incidental SIR for spotted sandpiper was discussed in the Phase 1 ERA (Windward 2003b). Using an incidental SIR that is 30% of the FIR would increase the spotted sandpiper HQs by an average of less than 0.1. Three LOAEL-based HQs that were less than 1.0 assuming a SIR that is 18% of the FIR would slightly exceed 1.0 if the SIR were assumed to be 30% of the FIR. In Area 3 high- and poor-quality habitat, the chromium HQ would increase from 0.83 to 1.3, and the mercury HQ would increase from 0.99 to 1.1; in Area 2 high- and poor-quality habitat, the PCB TEQ HQ would increase from 0.97 to 1.2. This conservative analysis indicates that risks could be slightly underestimated for chromium, mercury, and PCBs if the incidental SIR is as high as 30% of the FIR.

COPC Bioavailability

Metals may be less bioavailable in ingested sediment than in ingested prey. In calculating the ingested doses, it was assumed that metals were 100% bioavailable, which may overestimate risk if the primary source of the dose is sediment. Table A.6-45 shows the relative contributions of sediment and prey to the total ingested doses. Up to 97% of the ingested dose is from sediment exposure in some areas, indicating that risks may be overestimated in those areas. Contributions from water are very low.

Dietary Composition

The possibility that spotted sandpipers could consume fish, crabs, or mussels in their diet was considered. To address this uncertainty, HQs were calculated assuming that 25% of the spotted sandpiper diet consisted of the alternative prey type (i.e., fish, crabs, or mussels) with the highest UCL concentration. This assumption resulted in slight changes in HQs, but did not cause a change in risk conclusions.

Site Use

Phase 1 uncertainties regarding site-specific habitat for spotted sandpipers were addressed by the survey of spotted sandpiper presence and habitat conducted in the LDW in 2004 (Windward 2004h). Results of the survey were used to develop various exposure scenarios, which reduced uncertainty in the Phase 2 risk conclusions.

Benthic Invertebrate Tissue Data and Linear Regressions

There is uncertainty in the benthic invertebrate tissue data used to estimate dietary exposure of spotted sandpipers to some COPCs because of the relatively small numbers of samples in some exposure areas. The uncertainty is relatively low for PCBs and

arsenic because a significant linear relationship exists between sediment and benthic invertebrate tissue concentrations (see Attachment 11), and this relationship was used to estimate tissue UCL concentrations from a relatively larger sediment dataset for each exposure area. However, uncertainty is greater for other COPCs that did not have a significant relationship between sediment and tissue concentrations. For these COPCs, the UCL concentration in tissue is more uncertain because it was based on a small tissue dataset.

The lack of a sediment/tissue relationship for a particular COPC may be because a relatively small range of sediment concentrations was sampled for that COPC or because a correlation does not exist. The sampling of benthic invertebrate tissue was designed to include stations that represented the full range of concentrations of arsenic, PCBs, and lead in surface sediments (Windward 2004d). To evaluate whether the full range of chemical concentrations was sampled for other COPCs, each COPC concentration in co-located sediment was plotted against the cumulative distribution of that COPC in the full baseline surface sediment dataset. For example, the results for copper indicate that the smaller co-located dataset is closely representative of the entire LDW, with the highest concentration in the co-located dataset at the 99th percentile for the entire LDW dataset (Figure A.6-1). In contrast, Figure A.6-2 shows a poorer representation for nickel, with the highest concentration in the co-located dataset at the 64th percentile for the entire LDW dataset.



Figure A.6-1. Copper concentrations in sediment samples co-located with benthic invertebrate tissue samples relative to the LDW-wide baseline surface sediment dataset



Figure A.6-2. Nickel concentrations in surface sediment samples co-located with benthic invertebrate tissue samples relative to the LDW-wide baseline surface sediment dataset

Results for all COPCs without a significant sediment/tissue relationship are presented in Table A.6-46. The uncertainty in the relationship between sediment and tissue concentrations for cadmium, chromium, cobalt, copper, mercury, vanadium, and zinc is low because benthic invertebrate tissue samples were analyzed over a relatively wide range of sediment concentrations. For the remaining two COPCs (i.e., nickel and selenium), the range of sediment concentrations in the co-located dataset did not cover the range found in the entire LDW dataset. It is possible that a significant sediment/ tissue relationship was not found because the range of sediment concentrations was not large enough to capture a relationship, if one existed.

Table A.6-46. COPC concentrations in sediment samples co-located with benthic invertebrate tissue samples relative to the site-wide baseline surface sediment dataset for COPCs without a significant sediment/tissue relationship

COPC	Maximum COPC Concentration in Sediment (mg/kg dw)		Percent of Concentrations in LDW-Wide Dataset Less Than or Equal to the Maximum Concentration in Co-Located Samples
	LDW-Wide Baseline Surface Sediment Dataset	Sediment Samples Co-Located with Benthic Invertebrate Tissue Samples	
Cadmium	120	1.67	94
Chromium	1,100 J	42.5	84
Cobalt	140	31.5	99
Copper	12,000 J	495	99
Mercury	4.6 J	0.528	97
Nickel	910	24.8	64
Selenium	28	1.4	14
Vanadium	150	72.6	84
Zinc	9,700	2,080 J	99

COPC – chemical of potential concern

dw – dry weight

J – estimated concentration

LDW – Lower Duwamish Waterway

The relatively low number of benthic invertebrate tissue samples results in higher uncertainty in the UCL in exposure areas where there is high variability in the tissue concentrations. In four instances, ProUCL calculated a UCL that was greater than the maximum concentration. The ProUCL guidelines recommend using the UCL even if it is greater than the maximum concentration (EPA 2004a). In three of these instances, differences between the maximum concentration and the UCL were small. In one instance (lead in Area 2 high- and poor-quality habitats), the UCL tissue concentration was three times higher than the maximum concentration. Nevertheless, in all instances, risk conclusions would not have changed if maximum values had been used instead of the UCL.

Toxic Equivalent Approach

For the calculation of PCB risks using the TEQ approach, fish, benthic invertebrate, and sediment samples were analyzed for PCB congeners, and TEFs were used to account for toxicity relative to 2,3,7,8-TCDD. The TEFs used to calculate TEQs for dioxin-like PCB congeners were WHO consensus values for birds and mammals from Van den Berg et al. (1998; 2006); these TEFs are presented in Attachment 3. The rationale for the use of TEFs is based on evidence that there is a common mechanism of toxicity for certain dioxins, furans, and PCB congeners, which involves binding to the aryl hydrocarbon (Ah) receptor as an initial step. Data on the relative binding affinity of particular PCB congeners compared to 2,3,7,8-TCDD are available from *in*

vivo and *in vitro* studies. These data have been used to derive TEFs for PCB congeners that show structural similarity to dioxins, bind to the Ah receptor, and elicit dioxin-specific biochemical and toxic responses.

An uncertainty in the TEQ approach is related to the derivation of consensus TEF values. Limitations in the underlying data used to derive TEFs, such as the relevance of the endpoints in the studies and the lack of information on interspecies variability, contribute to the uncertainty. Although these uncertainties have been identified by Van den Berg (1998), it was decided at a 1997 WHO expert meeting that an additive TEQ method is the most appropriate risk assessment method for complex mixtures of dioxin-like PCB congeners (EPA 2003b). According to the EPA, the TEQ method is technically appropriate for evaluating risks to birds and mammals, and uncertainties associated with the method are not greater than other sources of uncertainty in the ERA process (EPA 2003b).

The four most potent Ah receptor agonists in birds among PCB congeners are the non-ortho PCBs 77, 81, 126, and 169. The variability in the TEFs appears high for PCB congeners that have been tested on multiple species (Van den Berg et al. 1998). For PCB 77, five studies have been conducted, resulting in a TEF range of < 0.0003 to 0.15 for the various bird species tested for ethoxyresorufin-O-deethylase (EROD) induction or *in ovo* effects. For PCB 81, two identified studies tested several species for EROD induction, with TEFs highly variable, ranging from 0.001 to 0.5. For PCB 126 and 169, data are available from only one study (*in ovo* with chickens). These TEFs derived by EROD induction or *in ovo* studies are most accurate for the assessment of effects based on concentrations in whole embryos (EPA 2003b). Thus, the relevance of applying the bird TEFs to dietary exposure is uncertain. Egg concentrations were not evaluated in this risk assessment because concentrations of dioxin-like compounds in bird eggs were not available, and reliable models for predicting egg concentrations from concentrations in the diet were not available. The absence of data or reliable predictions of dioxin-like compounds in eggs of birds using the LDW results in additional uncertainty in risk estimates for birds. It is not known if the uncertainties discussed above would overestimate or underestimate risks.

It should also be noted that the TEQs calculated in this ERA were used only for assessing the toxicity of the dioxin-like PCB congeners and do not account for TEQ contributions from dioxin and furan congeners; dioxin and furan concentrations in tissue were not available for the LDW. Risk estimates for wildlife based on TEQs calculated considering only dioxin-like PCB congeners are likely lower than the actual risk resulting from the cumulative exposure of wildlife to dioxins, furans, and dioxin-like PCB congeners in the LDW.

Effects Assessment

Uncertainty associated with available toxicity benchmarks for birds may affect risk estimates. These uncertainties were discussed in detail in the Phase 1 ERA (Windward 2003b). The primary uncertainties include:

- None of the laboratory toxicological studies used to derive TRVs were conducted using ROC species.
- The laboratory studies on which TRVs are based were conducted in controlled settings using single-contaminant exposures. Effects associated with multiple-chemical exposure and other environmental stressors present at the site (e.g., habitat loss) were not factored into these studies. It is unknown if these factors would result in additive, synergistic, antagonistic, or neutral effects on overall risk conclusions.
- NOAELs were not available for some COPCs, so they were estimated from LOAELs.

In addition, TRVs are considered less certain if there were a small number of studies, if endpoints were subchronic, or if data quality was questionable. The relative uncertainties in the selected TRVs for birds and the potential effect on the risk estimates are summarized in Table A.6-47.

Table A.6-47. Level of uncertainty associated with TRVs for birds

COPC	Number of TRV Studies	Uncertainty in TRV[a]
Arsenic	3	high; reproductive studies with arsenite (the most toxic form of arsenic) were not available
Cadmium	8	high; the only effect endpoints were eggshell thinning (with no effect on reproductive success) and growth in chicks after subchronic exposure
Chromium	3	high; only one study reported effects, but this study was unpublished and could not be obtained for review of data quality; selected TRV was based on a survival endpoint, so the resulting HQ may underestimate the potential for sublethal effects
Cobalt	1	high; only one toxicity study, which evaluated growth after a 2-week exposure period, was available
Copper	7	medium; selected TRVs were based on subchronic growth and survival endpoints
Lead	4	medium; selected TRVs were based on a chronic reproductive endpoint
Mercury	7	medium; selected TRVs were based on a chronic growth endpoint
Nickel	3	medium; selected TRVs were based on a subchronic growth endpoint
Selenium	5	medium; selected TRVs were based on chronic reproduction and survival endpoints
Vanadium	2	high; only one toxicity study, which evaluated growth after a 4-week exposure period, was available
Zinc	6	high; selected TRVs were based on a subchronic growth endpoint
Total PCBs	13	medium; selected TRVs were based on a chronic reproduction endpoint
PCB TEQs	2	high; no dietary studies were available; selected TRV was based on a study with acute high-level weekly exposure via IP injection

[a] Level of uncertainty key:

Low = large dataset including chronic studies

Medium = moderately sized dataset including chronic studies

High = small dataset with only subchronic studies, unbounded NOAELs/LOAELs, or data with questionable data quality

COPC – chemical of potential concern
HQ – hazard quotient
IP – intraperitoneal
PCB – polychlorinated biphenyl
TEQ – toxic equivalent
TRV – toxicity reference value

In summary, uncertainty associated with the chromium TRV is high because only one study that reported effects was available; this study is unpublished and could not be reviewed. Uncertainty is also high for the PCB TEQ TRV because the selected TRV was based on exposure from acute high-level weekly IP injections. Risks for cadmium, cobalt, nickel, vanadium, and zinc could be under- or overestimated because the endpoints were based on subchronic growth effects and no reproductive endpoints were available. Risks are uncertain for chromium because the TRV was based on a single unpublished study with a survival endpoint. The effect of uncertainties in toxicity data for copper, lead, mercury, selenium and total PCBs on the risk estimates for these COPCs is unknown.

Risk Characterization

Risks to spotted sandpiper from total DDTs were not included in the risk estimates because of high uncertainty in the tissue pesticide data. Probable analytical interference from PCBs in benthic invertebrate, fish, and crab tissue samples collected and analyzed in 2004 resulted in suspected false identifications of presence of some organochlorine pesticides as well as overestimates in their concentrations (JN-qualified). The uncertainty and high bias of these results are discussed in more detail in Section A.2.4.2.2. Ingested dose estimates, TRVs, and risk estimates are presented in this section for total DDTs, which was identified as a spotted sandpiper COPC in Section A.2.5.3.

Methods used to calculate the ingested dose of total DDTs are the same as those used for other COPCs, as described in Section A.5.1.1. The tissue and sediment exposure concentrations are presented in Table A.6-48, and resulting ingested doses are presented in Table A.6-49. Exposure to DDTs via water was not estimated because the contribution of total DDTs via water is expected to be very low based on the low solubility of DDTs and results from PCB exposure calculations, which showed the contribution from water was less than 0.01 percent of the contribution from food and sediment exposure. Tables containing all exposure assumptions and data used in the ingested dose calculations are presented in Attachment 12.

Table A.6-48. Exposure concentrations of total DDTs in benthic invertebrate tissue and sediment for spotted sandpiper

Exposure Area[a]	Exposure Concentration in Benthic Invertebrate Tissue (mg/kg dw)	Exposure Concentration in Sediment (mg/kg dw)
Area 1 – high	0.29	0.017
Area 1 – high and poor	0.29	0.018
Area 2 – high	1.1	1.0
Area 2 – high and poor	0.96	0.52

Exposure Area[a]	Exposure Concentration in Benthic Invertebrate Tissue (mg/kg dw)	Exposure Concentration in Sediment (mg/kg dw)
Area 3 – high	1.8	0.024
Area 3 – high and poor	1.8	0.020

[a] Six exposure scenarios were evaluated for spotted sandpiper. In each of three exposure areas, foraging only in high-quality habitat and foraging in both high- and poor-quality habitat were evaluated. These exposure scenarios are described in detail in Section A.5.1.2.1.

dw – dry weight

Table A.6-49. Ingested doses of total DDTs for spotted sandpiper

Exposure Area	Ingested Dose (mg/kg bw/day)
Area 1 – high	0.046
Area 1 – high and poor	0.046
Area 2 – high	0.20
Area 2 – high and poor	0.17
Area 3 – high	0.28
Area 3 – high and poor	0.28

bw – body weight

Toxicity studies with any form of DDT, including DDD and DDE, were evaluated to select the TRV for total DDTs. The evaluation identified numerous studies that analyzed the dietary toxicity of DDT, DDD, and DDE to birds. Table A.6-50 presents the results from 10 studies with the lowest reported effect concentrations. All of the reviewed studies evaluated reproductive endpoints; eight of the studies reported increased eggshell thinning. A dose of 0.15 mg/kg bw/day resulted in eggshell thinning in Japanese quail exposed to p,p′-DDT in the diet for 194 days. Although eggshell thinning was significantly different in the treated group than in controls, hatchability of the eggs was not affected, therefore, it was concluded that the degree to which eggshell thinning was observed would not affect reproductive success. The next lowest dose of 0.32 mg/kg bw/day resulted in eggshell thinning, eggshell breakage, and nestling mortality in barn owls exposed to dietary p,p′-DDE for 2 years. This dose was selected as the LOAEL TRV because a clear impairment to reproductive success was observed. The highest NOAEL below the LOAEL was a dose of 0.19 mg/kg bw/day, which did not cause eggshell thinning in mallards exposed to DDT for 11 months (Davison and Sell 1974). Because other reproductive endpoints were not assessed in this study, and it is unknown whether the no effect level for eggshell thinning would be the same as the no effect level for direct measures (e.g., hatchability, viability of offspring) of reproductive success, this NOAEL was not selected. Instead, the NOAEL TRV was estimated from the selected LOAEL TRV using an uncertainty factor of 5, resulting in a NOAEL TRV of 0.065 mg/kg bw/day.

Table A.6-50. DDT, DDD, and DDE dietary toxicity studies for birds

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc.[a]	Effect Conc.[a]	Body Weight (kg)	Food Consumption Rate[b]	Source	Notes
p,p'-DDT	quail	na	0.15	194 days	increased eggshell thinning	na	2.5 mg/kg ww	0.09 (Dunning 1993)	0.0048 kg dw/day galliformes (Nagy 2001)	Stickel and Rhodes (1970)	1
p,p'-DDT	mallard	0.19	na	11 months	no effect on eggshell thinning	2 mg/kg ww	na	1.19	0.115 kg ww/day	Davison and Sell (1974)	2
DDE	barn owl	na	**0.32**	2 years	increased eggshell thinning, eggshell breakage, and nestling mortality	na	2.83 mg/kg ww	0.5235 (Dunning 1993)	0.0539 kg dw/day carnivores (Nagy 2001)	Mendenhall et al. (1983)	1
DDE	America n kestrel	na	0.35	14 days	increased eggshell thinning	na	3 mg/kg ww	0.13 (Pattee 1984)	0.0136 kg dw/day Eurasian kestrel (Nagy 2001)	Peakall et al. (1973)	1
Technical DDD	mallard	na	0.90	2 years	decreased hatchling survival and production	na	10 mg/kg dw	1.082 (Dunning 1993)	0.1082 kg ww/day (Heinz et al. 1987)	Heath et al. (1969)	1
p,p'-DDE	mallard	na	0.90	2 years	increased eggshell thinning and number of cracked eggs; reduced hatchling survival and production	na	10 mg/kg dw	1.082 (Dunning 1993)	0.1082 kg ww/day (Heinz et al. 1987)	Heath et al. (1969)	1
p,p-DDE	black duck	na	1.0	7 months	increased eggshell thinning and duckling mortality; reduced hatchability	na	10 mg/kg ww	1.25 (Dunning 1993)	0.125 kg ww/day (Heinz et al. 1987)	Longcore and Samson (1973)	
DDE	mallard	na	1.0	30 days	increased eggshell thinning	na	10 mg/kg ww	1.082 (Dunning 1993)	0.1082 kg ww/day (Heinz et al. 1987)	Kolaja (1977)	1

Chemical	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Duration	Effect	No-Effect Conc.[a]	Effect Conc.[a]	Body Weight (kg)	Food Consumption Rate[b]	Source	Notes
DDT	mallard	na	1.0	30 days	increased eggshell thinning	na	10 mg/kg ww	1.082 (Dunning 1993)	0.1082 kg ww/day (Heinz et al. 1987)	Kolaja (1977)	1
p,p' DDE	America n kestrel	na	1.0	1 year (two clutches)	increased eggshell thinning	na	2.8 mg/kg ww	0.13 (Pattee 1984)	0.0136 kg dw/day Eurasian kestrel (Nagy 2001)	Wiemeyer and Porter (1970); Porter and Wiemeyer (1970)	1

Notes:

1. Effect concentration converted into dry weight assuming 10% moisture in prepared diet
2. Body weight and food consumption rate reported in study

[a] No-effect and effect concentrations are presented in the units given in the studies reviewed. Table notes indicate how units were converted to wet weight or dry weight to correspond to the food consumption rate units for calculating NOAELs and LOAELs.

[b] Consumption rates are from equations for bird groups presented in Nagy (2001), from data presented for individual bird species (Nagy 2001), or from other sources as noted.

dw – dry weight

LOAEL – lowest-observed-adverse-effect level

na – not available or not applicable

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs. A NOAEL TRV was not available for DDE from the study in which the chronic LOAEL of 0.32 mg/kg bw/day was reported, so it was estimated using an uncertainty factor of 5. The resulting NOAEL TRV is 0.065 mg/kg bw/day.

The estimated ingested dose for spotted sandpiper was compared to the NOAEL and LOAEL TRVs, as presented in Table A.6-51. No LOAEL-based HQs were greater than 1.0. In Area 2, the NOAEL-based HQs were 3.1 and 2.6; and in Area 3, the NOAEL-based HQs were both 4.3. The NOAEL-based HQs were less than 1.0 in Area 1.

Table A.6-51. DDT HQs for spotted sandpiper

Receptor	Ingested Dose (mg/kg bw/day)	TRVs (mg/kg bw/day)		NOAEL-Based HQ	LOAEL-Based HQ
		NOAEL	LOAEL		
Area 1 – high	0.046	0.065	0.32	0.71	0.14
Area 1 – high and poor	0.046	0.065	0.32	0.71	0.14
Area 2 – high	0.20	0.065	0.32	**3.1**	0.63
Area 2 – high and poor	0.17	0.065	0.32	**2.6**	0.53
Area 3 – high	0.28	0.065	0.32	**4.3**	0.88
Area 3 – high and poor	0.28	0.065	0.32	**4.3**	0.88

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

TRV – toxicity reference value

Bold identifies NOAEL-based HQs greater than 1.0.

The high bias for DDTs was confirmed by reanalyzing sediment samples co-located with benthic invertebrate tissue samples and fish and crab tissue samples that had high PCB and DDT concentrations, as described in Section A.2.4.2.2. The presence of DDTs was confirmed in six of the eight tissue samples at concentrations ranging from 5 to 34% of the original results and in sediment samples at concentrations ranging from 4 to 60% of the original results. Because of these uncertainties, it is likely that the NOAEL-based HQs for DDTs would have been lower if there had been no analytical interferences.

Summary of Uncertainties

Uncertainties in the problem formulation, the effects and exposure assessments, and risk characterization for spotted sandpiper were evaluated, with the following conclusions:

- Uncertainties in FIR, direct sediment contact, incidental sediment ingestion, dietary composition, and site use are expected to have minimal or no effect on risk conclusions.
- Uncertainties in COPC selection are not expected to have an effect on risk conclusions.
- For COPCs other than PCBs and arsenic, there is some uncertainty associated with risk estimates because of the relatively small number of benthic invertebrate tissue samples in each sandpiper foraging area.

- For COPCs whose sediment concentrations contributed a large portion of the ingested dose, risks could be overestimated if these COPCs are not 100% bioavailable in sediments.
- The PCB TEQs were calculated using TEFs for individual PCB congeners; the derivation of some of these TEFs is uncertain. It is not known if this uncertainty would overestimate or underestimate risk. Risk estimates for wildlife based on TEQs calculated considering only dioxin-like PCB congeners are likely lower than the actual risk resulting from the cumulative exposure of wildlife to dioxins, furans, and dioxin-like PCB congeners in the LDW.
- Uncertainties in toxicity data may affect risk estimates. Risks could be underestimated for arsenic because of the lack of arsenite toxicity data. Risks are uncertain for cadmium, cobalt, nickel, vanadium, and zinc because TRVs were based on subchronic growth endpoints. Risks are uncertain for chromium because the TRV was based on a single unpublished study with a survival endpoint. Risks are also uncertain for PCB TEQs because the TRVs were based on a study using an acute weekly dose via IP injection and TEFs were applied to dietary exposure estimates. Effects of uncertainties in toxicity data on risk estimates for copper, lead, mercury, selenium, and PCBs (based on a total PCB approach) are unknown.
- The LOAEL-based HQs for spotted sandpiper and DDTs were less than 1.0, although NOAEL-based HQs were greater than 1.0 in Areas 2 and 3. It is likely that NOAEL-based HQs would have been closer to 1.0 if there had been no analytical interferences from PCBs because the interference is known to bias the DDT concentrations high.

A.6.3.1.3 Risk conclusions

Spotted sandpiper, a bird which has been observed nesting along the LDW, was selected to represent benthivorous birds such as dunlin, dowitcher, western sandpiper, and dabbling ducks. The risk characterization for sandpiper should be protective of other benthivorous birds because of the spotted sandpiper's high exposure to COPCs through the ingestion of benthic invertebrates and the incidental ingestion of sediment.

Results of the risk characterization for spotted sandpiper are summarized in Table A.6-52.

Table A.6-52. Summary of risk characterization for spotted sandpiper

COPC	Range of NOAEL-Based HQs[a]	Range of LOAEL-Based HQs[a]	Effect	Primary Uncertainty[b, c]
Arsenic	0.34 – 0.58	< 0.1 – 0.15	delayed egg laying; depressed egg weight, production, and shell thinning; decreased offspring body weight[d]	high uncertainty in TRV because reproductive studies using arsenite were not available; risk could be underestimated
Cadmium	< 0.1 – 0.21	<0.1	reduced body weight (subchronic exposure)	high uncertainty in TRV; risk could be under- or overestimated because of endpoint
Chromium	**1.3** – **8.8**	0.26 – **1.8**	reduced survival	high uncertainty in TRV because study was unpublished; risk could be overestimated because of high sediment contribution to ingested dose and 100% bioavailability assumption; risk may be underestimated because TRV was based on survival rather than a sublethal endpoint
Cobalt	0.22 – 0.27	<0.1	reduced body weight (chronic exposure)	high uncertainty in TRV; risk could be under- or overestimated because of endpoint
Copper	0.62 – **1.5**	0.45 – **1.1**	reduced growth and survival (subchronic exposure)	medium uncertainty in TRV; risk could be under- or overestimated because of endpoint
Lead	0.58 – **19**	0.17 – **5.5**	reduced egg hatchability	medium uncertainty in TRV; risk could be overestimated because of high sediment contribution to ingested dose and 100% bioavailability assumption
Mercury	**1.1** – **5.3**	0.21 – **1.0**	reduced growth (chronic exposure)	medium uncertainty in TRV
Nickel	< 0.1 – 0.22	< 0.1– 0.11	reduced body weight (subchronic exposure)	medium uncertainty in TRV; risk could be over- or underestimated because of endpoint; uncertainty in tissue dataset
Selenium	0.62 – 0.90	0.38 – 0.55	reduced offspring growth and survival	medium uncertainty in TRV; uncertainty in tissue dataset
Vanadium	**2.0** – **2.7**	**1.0** – **1.4**	reduced body weight (subchronic exposure)	high uncertainty in TRV; risk could be under- or overestimated because of endpoint
Zinc	0.41 – 0.78	0.27 – 0.52	reduced growth (subchronic exposure)	high uncertainty in TRV; risk could be under- or overestimated because of endpoint
Total PCBs	0.51 – **2.0**	0.18 – 0.71	reduced hatching success	medium uncertainty in TRV

COPC	Range of NOAEL-Based HQs[a]	Range of LOAEL-Based HQs[a]	Effect	Primary Uncertainty[b, c]
PCB TEQs	**1.9** – **10**	0.18 – **1.5**	reduced body weight, egg production, and survival of adults and embryos	high uncertainty in TRV because of high-level weekly exposures and IP injection; risk could be overestimated; derivation of TEFs is uncertain

[a] Range of HQs calculated for the six spotted sandpiper exposure scenarios.

[b] Level of uncertainty key:

Low = large dataset including chronic studies with species taxonomically similar to the ROC

Medium = moderately sized dataset including chronic studies

High = small dataset with only subchronic studies, unbounded NOAELs/LOAELs, or data with questionable data quality

[c] There may also be some uncertainty associated with risk estimates for COPCs other than arsenic and total PCBs because of the relatively small number of benthic invertebrate tissue samples.

[d] Effects were from exposure to sodium arsenate.

COPC – chemical of potential concern
HQ – hazard quotient
IP – intraperitoneal
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
TEF – toxic equivalency factor
TEQ – toxic equivalent
TRV – toxicity reference value

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

The following COPCs had LOAEL-based HQs greater than or equal to 1.0 for spotted sandpiper: chromium, copper, lead, mercury, and vanadium. LOAEL-based HQs ranged from 1.0 to 5.5 for these COPCs. LOAEL-based HQs were greater than or equal to 1.0 in Exposure Area 2 for chromium (1.8) and lead (5.5), in Exposure Area 3 for copper (1.1) and mercury (1.0), and in all three exposure areas for vanadium (1.0 to 1.4). Risks to spotted sandpiper from chromium are uncertain because the TRV was based on a single unpublished study with a survival endpoint. Risks to spotted sandpiper from copper and vanadium may be under- or overestimated because the selected TRVs were based on subchronic growth endpoints. The ingested dose is primarily from sediment for copper and lead in Area 3 and for vanadium in all areas. Bioavailability of metals in sediment is not likely 100%, so the LOAEL-based HQs (ranging from 1.0 to 1.5) for copper, lead, and vanadium may be overestimated. Overall, these findings indicate risk for spotted sandpiper in some areas of the LDW from exposure to chromium, copper, lead, mercury, and vanadium, but these risks are expected to be low.

For total PCBs, the NOAEL-based HQs were greater than 1.0 in some of the spotted sandpiper exposure areas, but LOAEL-based HQs were less than 1.0, indicating low risk with some uncertainty because the true threshold of effects between the NOAEL and the LOAEL is not known. For PCB TEQs, the LOAEL-based HQs were less than 1.0, except in Area 2 under the high-quality habitat scenario, in which the LOAEL-

based HQ was 1.5. Uncertainty is high for PCB TEQ risk estimates because the TRVs were based on a study using an acute weekly dose via IP injection, and TEFs derived in studies of toxicity to eggs were applied to dietary exposure estimates. Therefore, risks to spotted sandpiper from PCBs are low in most areas of the LDW, with some uncertainty.

The remaining COPCs (arsenic, cadmium, cobalt, nickel, selenium, and zinc) had both NOAEL- and LOAEL-based HQs that were less than 1.0 in all sandpiper exposure areas. Exposure of spotted sandpipers to these COPCs is not expected to result in adverse effects.

For total DDT, the NOAEL-based HQs were greater than 1.0 in some of the spotted sandpiper exposure areas, but LOAEL-based HQs were less than 1.0, indicating low risk. It is likely that HQs would have been lower if there had been no analytical interferences from PCBs because the interference is known to bias the DDT concentrations high. Therefore, risks to spotted sandpiper from DDT are expected to be very low in most areas of the LDW, with some uncertainty.

A.6.3.2 Great blue heron

This section present risk characterization, uncertainties, and risk conclusions for great blue heron.

A.6.3.2.1 Risk estimates

Four COPCs were evaluated for great blue heron: chromium, lead, mercury, and PCBs (using both a total PCB and a PCB TEQ approach). As shown in Table A.6-53, all of the NOAEL- and LOAEL-based HQs were less than 1.0.

Table A.6-53. HQ calculations for great blue heron

ROC	COPC	Ingested Dose (mg/kg bw/day)	TRVs (mg/kg bw/day)		NOAEL-Based HQ	LOAEL-Based HQ
			NOAEL	LOAEL		
Great blue heron	chromium	0.067	1.0	5.0	< 0.1	< 0.1
	lead	0.27	5.82	20	< 0.1	< 0.1
	mercury	0.0031	0.018	0.091	0.17	< 0.1
	total PCBs	0.17	0.49	1.4	0.35	0.12
	PCB TEQs	9.3×10^{-6}	1.4×10^{-5}	1.4×10^{-4}	0.66	< 0.1

[a] No PCB congener data were available for juvenile chinook salmon. When calculating risk estimates, the portion of prey ingestion that had been assigned to juvenile chinook salmon (24%) was divided proportionally among the remaining prey categories.

bw – body weight
COPC – chemical of potential concern
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
ROC – receptor of concern
TEQ – toxic equivalent
TRV – toxicity reference value

A.6.3.2.2 Uncertainties

This section presents a discussion of the uncertainty associated with the problem formulation, the exposure and effects assessments, and the risk characterization for great blue heron.

Problem Formulation

Primary uncertainties in the problem formulation for great blue heron are associated with ROC selection and the COPC screen. Uncertainties in the COPC screen for bird ROCs in general were discussed in Section A.6.3.1.2 for spotted sandpiper.

Uncertainties related to the representation of other bird species by great blue heron were addressed in the Phase 1 ERA (Windward 2003b). In that analysis, it was concluded that the great blue heron may have a lower daily FIR relative to its body weight than some other piscivorous species (i.e., common murre, pigeon guillemot, and Caspian tern), but these other species forage in the LDW infrequently. As a result, the overall ingestion of prey from the LDW is likely to be higher for great blue heron than for other piscivorous species.

Exposure Assessment

Uncertainties in the exposure assessment for great blue heron were associated with the following factors:

- Direct sediment contact
- Incidental sediment ingestion rate
- Dietary composition
- Site use
- TEQ approach

The uncertainties associated with direct sediment contact and the TEQ approach are the same as those discussed in Section A.6.3.1.2 for spotted sandpiper.

To address uncertainties in the amount of sediment incidentally ingested by great blue heron while foraging, ingested doses of COPCs were calculated assuming the SIR was 10% of the FIR versus 2% assumed in Section 5.1.2.2. This conservative assumption would result in an increase of HQs by an average of less than 0.1 and would not change risk conclusions.

Dietary Composition

The assessment for great blue heron is assumed to be protective of omnivorous bird species such as bufflehead, Barrow's goldeneye, and surf scoter, which may consume primarily mussels, clams, or crabs. To address the uncertainty that invertebrate species could be ingested, HQs for each COPC were calculated using the conservative assumption that the heron's diet consisted of the invertebrate species (i.e., mussels, clams, or

crabs) with the maximum concentration of that COPC. This conservative assumption did not result in any NOAEL- or LOAEL-based HQs greater than 1.0 for great blue heron.

There are some uncertainties in the proportions of different fish species in the great blue heron's diet. Different preferences for fish could result in greater exposure of great blue heron. To address this uncertainty, HQs were calculated assuming that 75% of the great blue heron diet was composed of the fish species with the highest concentration for each COPC, with the remainder of the diet composed of equal proportions of the other two species.[89] This assumption resulted in changes in the HQs but did not result in a change of any HQ from less than 1.0 to greater than 1.0.

Site Use

There are uncertainties in the site-use factor for great blue herons because of the lack of site-specific information on the foraging of great blue herons from the Black River and Kiwanis Ravine colonies. A site-use factor of 0.5 is likely to be conservative; but if a higher site-use factor of 0.75 had been used, the HQs for great blue heron would still have remained below 1.0.

Effects Assessment

Uncertainties associated with available toxicity benchmarks for birds are discussed in Section A.6.3.1.2.

Risk Characterization

Risk estimates for organochlorine pesticides were not discussed in Section A.6.3.2.1 because the JN-qualified pesticide tissue data are biased high as a result of analytical interference from PCBs, as discussed in Section A.6.3.1.2 for spotted sandpiper. The same methods and TRVs used for estimating risks to spotted sandpiper from total DDTs were used for great blue heron. The tissue and sediment exposure concentrations are presented in Table A.6-54.

Table A.6-54. Exposure concentrations of total DDTs

Ingested Media	Exposure Concentration (mg/kg dw)
Shiner surfperch	1.0
English sole/starry flounder	0.76
Juvenile chinook salmon	0.21
Pacific staghorn sculpin	0.50
Crab	0.61
Intertidal sediment	0.29

dw – dry weight

[89] In the exposure assessment, it is assumed that heron ingest 34% shiner surfperch, 21% English sole, and 45% juvenile chinook salmon.

The calculated ingested dose of total DDTs for great blue heron, based on the exposure concentrations presented in Table A.6-54, was 0.012 mg/kg bw/day. This concentration is lower than the selected NOAEL TRV of 0.056 mg/kg bw/day, indicating that there is very low risk to great blue heron from exposure to total DDTs.

Summary of Uncertainties

Uncertainties in the problem formulation, the effects and exposure assessments, and risk characterization for great blue heron are summarized as follows:

- Uncertainties in incidental sediment ingestion, dietary composition, and site use are expected to have minimal or no effect on risk conclusions.
- The PCB TEQs were calculated using TEFs for individual PCB congeners; the derivation of some of these TEFs is uncertain, with unknown effects on risk conclusions. Risk estimates for wildlife based on TEQs calculated considering only dioxin-like PCB congeners are likely lower than the actual risk resulting from the cumulative exposure of wildlife to dioxins, furans, and dioxin-like PCB congeners in the LDW.
- Gaps in toxicity data result in uncertainty in risk estimates. Risks are uncertain for chromium because the TRV was based on a single unpublished study with a survival endpoint. Risks are uncertain for PCB TEQs because the TRVs were based on a study using an acute weekly dose via IP injection, and TEFs were applied to dietary exposure estimates. Effects of gaps in toxicity data on risk estimates for lead, mercury, and PCBs (based on a total PCB approach) are unknown.
- The NOAEL-based HQ for great blue heron and total DDTs was less than 1.0, indicating very low risk.

A.6.3.2.3 Risk conclusions

Great blue heron, a commonly observed bird along the LDW, was selected as an ROC to represent primarily piscivorous birds in the LDW, such as loons, western grebe, mergansers, double-crested cormorant, pigeon guillemot, Caspian tern, and common murre. Results for the risk characterization for great blue heron are summarized in Table A.6-55.

Table A.6-55. Summary of risk characterization for great blue heron

COPC	NOAEL-Based HQ	LOAEL-Based HQ	Effect	Primary Uncertainty[a]
Chromium	< 0.1	< 0.1	reduced survival	high uncertainty in TRV because study was unpublished; risk may be underestimated because TRV was based on survival rather than a sublethal endpoint
Lead	< 0.1	< 0.1	reduced egg hatchability	medium uncertainty in TRV
Mercury	0.17	< 0.1	reduced growth (chronic exposure)	medium uncertainty in TRV
Total PCBs	0.35	0.12	reduced hatching success	medium uncertainty in TRV
PCB TEQs	0.66	< 0.1	reduced body weight, egg production, and survival of adults and embryos	high uncertainty in TRV because of high-level weekly exposures and IP injection; risk could be overestimated; derivation of TEFs is uncertain

[a] Level of uncertainty key:

Low = large dataset including chronic studies with species taxonomically similar to the ROC

Medium = moderately sized dataset including chronic studies

High = small dataset with only subchronic studies, unbounded NOAELs/LOAELs, or data with questionable data quality

COPC – chemical of potential concern
HQ – hazard quotient
IP – intraperitoneal
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
TEF – toxic equivalency factor
TEQ – toxic equivalent

The NOAEL- and LOAEL-based HQs for great blue heron are all less than 1.0, indicating that very low risks to great blue herons are expected from exposure to chromium, lead, mercury, and PCBs in the LDW. For chromium and PCB TEQs, there is high uncertainty in the TRVs, but the LOAEL TRVs would need to be lower by a factor of 14 for the LOAEL-based HQs to equal 1.0. Uncertainties in the TEFs used to calculate the PCB TEQ concentrations could also result in overestimates or underestimates of risk.

A.6.3.3 Osprey

This section presents risk estimates, uncertainties, and risk conclusions for osprey.

A.6.3.3.1 Risk estimates

Four COPCs were evaluated for osprey: chromium, lead, mercury, and PCBs (using both a total PCB and a PCB TEQ approach). The only HQ for osprey greater than 1.0 was the NOAEL-based HQ of 1.6 for PCB TEQs (Table A.6-56). All other HQs, both NOAEL- and LOAEL-based HQs, were less than 1.0.

Table A.6-56. HQ calculations for osprey

ROC	COPC	Ingested Dose (mg/kg bw/day)	TRVs (mg/kg bw/day)		NOAEL-Based HQ	LOAEL-Based HQ
			NOAEL	LOAEL		
Osprey	chromium	0.11	1.0	5.0	0.11	< 0.1
	lead	0.26	5.82	20	< 0.1	< 0.1
	mercury	0.0051	0.018	0.091	0.28	< 0.1
	total PCBs	0.32	0.49	1.4	0.65	0.23
	PCB TEQs	2.3×10^{-5}	1.4×10^{-5}	1.4×10^{-4}	**1.6**	0.16

[a] No PCB congener data were available for juvenile chinook salmon. When calculating risk estimates, the portion of prey ingestion that had been assigned to juvenile chinook salmon (45%) was divided proportionally among the remaining prey categories.

bw – body weight
COPC – chemical of potential concern
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
ROC – receptor of concern
TEQ – toxic equivalent
TRV – toxicity reference value

Bold identifies NOAEL-based HQs greater than 1.0.

A.6.3.3.2 Uncertainties

This section presents a discussion of the uncertainty associated with the problem formulation, the exposure and effects assessments, and the risk characterization for osprey.

Problem Formulation

Primary uncertainties in the problem formulation for osprey are associated with ROC selection and the COPC screen. Uncertainties in the COPC screen for birds were discussed in Section A.6.3.1.2.

Osprey were selected to represent piscivorous birds in addition to great blue heron, as well as carnivorous birds in the LDW. Osprey may have lower daily food consumption rates relative to their body weight than some other piscivorous species (i.e., common murre, pigeon guillemot, and Caspian tern), but these other species forage in the LDW infrequently. As a result, the overall consumption of prey from the LDW is likely to be higher for osprey than for other piscivorous species.

Some carnivorous birds, such as bald eagles, consume fish as well as other birds, such as grebes, gulls, and other waterfowl. Birds used as prey by the bald eagle may contain higher concentrations of bioaccumulative COPCs in their tissue than fish if the birds consumed are consumers of fish. Thus, if bald eagles forage primarily in the LDW, they may be more exposed to bioaccumulative COPCs than osprey because of the eagles' consumption of birds. Data on chemical concentrations in tissues of birds using the LDW are not available. The potential effect of the inclusion of birds in the bald eagle diet on the risk to bald eagles estimated from the risks to osprey is not

known because of uncertainties in the proportion of birds in the eagle diet, the amount of fish those birds consume from the LDW, and the extent of biomagnification from fish to birds.

Exposure Assessment

Uncertainties in the exposure assessment for osprey were associated with the following factors:

- Direct sediment contact
- Incidental sediment ingestion rate
- Dietary composition
- TEQ approach

The uncertainties associated with direct sediment contact and the TEQ approach are the same as those discussed in Section A.6.3.1.2 for spotted sandpiper.

To address uncertainties in the amount of sediment incidentally ingested by osprey while foraging, ingested doses of COPCs were calculated assuming the SIR was 10% of the FIR versus 1% assumed in Section 5.1.2.3. This conservative assumption would result in an increase of HQs by an average of less than 0.1 and would not change risk conclusions.

There are some uncertainties in the proportions of different types of fish species in osprey's diet, as discussed in Section A.5.1.2.3, which could have an effect on exposure. To address this uncertainty, HQs for each COPC were calculated assuming that 75% of the osprey's diet was composed of the fish species with the highest concentration for that COPC, with the remainder of the diet composed of equal proportions of the other fish species.[90] This assumption resulted in changes in the HQs but did not result in a change of any HQ from less than 1.0 to greater than 1.0, or vice versa.

Effects Assessment

Uncertainties associated with available toxicity benchmarks for birds are discussed in Section A.6.3.1.2.

Risk Characterization

This section estimates risks using two types of data that are uncertain: 1) DDT data, which were affected by analytical interference from PCBs, and 2) PCB data for osprey eggs, which were estimated from biomagnification factors (BMFs) derived from the literature.

[90] In the exposure assessment, it is assumed that heron ingest 24% shiner surfperch, 24% English sole, 24% juvenile chinook salmon, and 24% Pacific staghorn sculpin.

Risk Estimates for DDTs

The same methods and TRVs used for estimating risks to spotted sandpiper from total DDTs (Section A.6.3.1.2) were used to estimate risks to osprey. The tissue and sediment exposure concentrations of total DDTs are presented in Table A.6-57.

Table A.6-57. Exposure concentrations of total DDTs

INGESTED MEDIA	EXPOSURE CONCENTRATION (mg/kg dw)
Shiner surfperch	1.0
English sole	0.76
Juvenile chinook salmon	0.21
Intertidal sediment	0.29

dw – dry weight

The calculated ingested dose of total DDTs for osprey, based on the exposure concentrations presented in Table A.6-57, is 0.022 mg/kg bw/day. This concentration is lower than the selected NOAEL TRV of 0.056 mg/kg bw/day, indicating that there is very low risk to osprey from exposure to total DDTs.

Risk Estimates Using COPC Concentrations in Osprey Eggs

Risks to birds were estimated in this baseline ERA by calculating the daily ingested dose for each COPC individually and comparing exposure estimates for each bird ROC to TRVs. Alternatively, if chemical concentration data were available for bird eggs, these data could have been used to estimate risks to birds using TRVs derived from toxicity studies based on bird egg concentrations. No egg data were available for use in this ERA.[91] Therefore, PCB concentrations in osprey eggs[92] were estimated using a BMF, and risks were estimated. The PCB concentration in LDW fish was multiplied by this BMF[93] to estimate the PCB concentration in LDW osprey eggs.

Two studies provide BMFs for PCBs, relating PCB concentrations in fish to concentrations in bird eggs. One of the studies calculated BMFs for herring gulls feeding from Lake Ontario (Braune and Norstrom 1989), and the other calculated BMFs for osprey feeding from the Willamette River in Oregon (Henny et al. 2003). The data from the osprey study were used in this evaluation because osprey is one of the ROCs in this risk assessment. To estimate BMFs for osprey, Henny et al. (2003)

---

[91] Quality assurance/quality control information for existing data on PCB concentrations in great blue heron eggs collected from the LDW (Krausmann 2002a) was insufficient, precluding the use of these data in this baseline ERA. In addition, although PCBs have been analyzed in osprey eggs collected from nests along the LDW, these data have not yet been made available from USGS.

[92] This analysis was conducted only for osprey, and not for other bird ROCs, because BMFs for osprey from data collected in the Willamette River were available.

[93] The PCB BMF used in this section is the PCB concentration in bird eggs divided by the PCB concentration in fish assumed to be preyed upon by the birds.

collected 10 osprey eggs (one per nest) from various sites along more than 100 miles of the Willamette River. Twenty-five whole-body composite fish samples of three species were collected at 11 locations along the Willamette River. These species were collected because they comprised 91% of the biomass of the osprey's diet, as observed in prey remains collected beneath the osprey nests. Fish and eggs were analyzed for PCBs. The PCB BMF was calculated by Henny et al. (2003) using the geometric mean of total PCB concentrations in eggs and fish tissue.

The calculated PCB BMF relating concentrations in fish and osprey eggs was 11 on a wet-weight basis (Henny et al. 2003). The estimated concentration of PCBs in the fish in the diet of osprey feeding from the LDW was calculated using Equation A.5-2 and the data presented in Table A.6-58. A BMF of 11 and a PCB concentration of 2.0 mg/kg ww in the osprey's diet resulted in an estimated PCB concentration in osprey eggs of 22 mg/kg ww.

Table A.6-58. Estimated PCB concentration in osprey eggs

PCB Concentration in Prey (mg/kg ww)[a]			Fraction of Prey in Diet[b]			PCB Concentration in Diet (mg/kg ww)	BMF	Estimated PCB Concentration in Osprey Eggs (mg/kg ww)
Perch	Sole	Salmon	Perch	Sole	Salmon			
3.5	2.6	0.70	0.34	0.21	0.45	2.0	11	22

[a] Concentrations are UCLs calculated using methods presented in Attachment 11.

[b] Fraction of prey in osprey diet, as discussed in Section A.5.1.2.3.

BMF – biomagnification factor

PCB – polychlorinated biphenyl

ww – wet weight

Five studies that related PCB concentrations in bird eggs to adverse effects were available (Table A.6-59.[94] These studies exposed adult birds to individual PCB Aroclors or Aroclor mixtures in their diet and evaluated a range of reproductive effects. Three studies reported adverse effects on reproduction at PCB concentrations in eggs ranging from 5.6 mg/kg ww (resulting in reduced eggshell thickness in American kestrel) to 34.1 mg/kg ww (resulting in reduced reproductive success in American kestrel). The lowest LOAEL of 5.6 mg/kg ww was not selected as a TRV because this effect was not at a level at which shell breakage would be expected, as discussed in more detail in Section A.5.2.1.1. The next lowest LOAEL (16 mg/kg ww [Aroclor 1254]), which resulted in reduced hatching success in the second generation of ringed turtle-doves, was selected as the LOAEL TRV (Peakall et al. 1972; Peakall and Peakall 1973). The only NOAEL that was lower than the LOAEL was a concentration of 7.1 mg/kg ww (Aroclor 1248), which did not result in any reported

[94] Reproductive studies with chickens were not considered, as discussed in Section A.5.2.1.1.

reproductive effects in the screech owl (McLane and Hughes 1980). This egg concentration was selected as the NOAEL TRV.

Table A.6-59. Toxicity studies for PCBs in bird eggs

Chemical	Test Species	NOAEL (mg/kg ww egg)	LOAEL (mg/kg ww egg)	Exposure Duration	Effect	Source
Aroclor 1248	screech owl	**7.1**	na	two generations	no effect on eggshell thickness, egg production, hatching success, or fledging success	McLane and Hughes (1980)
Aroclor 1248	American kestrel	na	5.6	5.5 months	reduced eggshell weight and thickness	Lowe and Stendell (1991)
Aroclor 1254	ringed turtle-dove	na	**16**	two generations	reduced hatchability and embryo survival	Peakall et al. (1972); Peakall and Peakall (1973)
Aroclor 1254	mallard	23	na	~ 1 month	no effect on number of laying hens, time to first hatch, clutch size, egg fertility, egg hatchability, or duckling survival to 3 weeks	Custer and Heinz (1980)
Mixture of Aroclors 1248, 1254, and 1260	American kestrel	na	34.1	100 days until eggs hatched	reduced reproductive success of parents exposed *in ovo*	Fernie et al. (2000; 2001)

LOAEL – lowest-observed-adverse-effect level

na – not available

NOAEL – no-observed-adverse-effect level

ww – wet weight

Bold identifies the NOAEL and LOAEL selected as the TRVs.

The NOAEL- and LOAEL-based HQs were calculated using the estimated egg concentration and the NOAEL and LOAEL TRVs, resulting in HQs of 3.1 and 1.4, respectively (Table A.6-60). These HQs are higher than NOAEL- and LOAEL-based HQs of 0.65 and 0.23, respectively, for total PCBs calculated based on ingested doses (Section A.6.3.1.1).

Table A.6-60. HQs for PCBs estimated in osprey eggs

Estimated Egg Concentration (mg/kg ww)	NOAEL TRV (mg/kg ww egg)	LOAEL TRV (mg/kg ww egg)	NOAEL-Based HQ	LOAEL-Based HQ
22	7.1	16	**3.1**	**1.4**

HQ – hazard quotient

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

TRV – toxicity reference value

ww – wet weight

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

There are several uncertainties associated with the BMF approach. The BMF used in this assessment was calculated using a relatively small dataset (i.e., 10 eggs) from a large area in the Willamette River. In addition, as noted by Henny et al. (2003), some osprey may capture different fish species than those used to calculate the BMF because they are opportunistic feeders, and some may forage in nearby ponds or lakes nearby in addition to the Willamette River. There are also uncertainties in applying site-specific data from the Willamette River to the LDW, which contains different fish species and different PCB mixtures. It is not known if these uncertainties would underestimate or overestimate risk to osprey.

Summary of Uncertainties

Uncertainties in the problem formulation, the effects and exposure assessments, and risk characterization for osprey are summarized as follows:

- Uncertainties in incidental sediment ingestion and dietary composition are expected to have minimal or no effect on risk conclusions.
- The PCB TEQs were calculated using TEFs for individual PCB congeners; the derivation of some of these TEFs is uncertain, with unknown effects on risk conclusions. Risk estimates for wildlife based on TEQs calculated considering only dioxin-like PCB congeners are likely lower than the actual risk resulting from the cumulative exposure of wildlife to dioxins, furans, and dioxin-like PCB congeners in the LDW.
- Gaps in toxicity data resulted in uncertainty in risk estimates. Risks to spotted sandpiper from chromium are uncertain because the TRV was based on a single unpublished study with a survival endpoint. Risks are uncertain for PCB TEQs because the TRVs were based on a study using an acute weekly dose via IP injection, and TEFs were applied to dietary exposure estimates. Effects of gaps in toxicity data on risk estimates for lead, mercury, and PCBs (based on a total PCB approach) are unknown.
- The NOAEL-based HQ for osprey and total DDTs was less than 1.0, indicating very low risk.
- Estimates of risk to osprey from total PCBs using egg concentrations resulted in NOAEL- and LOAEL-based HQs greater than 1.0, whereas NOAEL-based HQs for total PCBs were less than 1.0 using an ingested dose approach. Risk estimates from the estimated PCB egg concentrations are considered highly uncertain.

A.6.3.3.3 Risk conclusions

Osprey was chosen as an ROC because of its known site use and diet and to represent other piscivorous birds. The results of the risk characterization for osprey are summarized in Table A.6-61.

Table A.6-61. Summary of risk characterization for osprey

COPC	NOAEL-Based HQ	LOAEL-Based HQ	Effect	Primary Uncertainty[a]
Chromium	0.11	< 0.1	reduced survival	high uncertainty in TRV because study was unpublished; risk may be underestimated because TRV was based on survival rather than a sublethal endpoint
Lead	< 0.1	< 0.1	reduced egg hatchability	medium uncertainty in TRV
Mercury	0.28	< 0.1	reduced growth (chronic exposure)	medium uncertainty
Total PCBs	0.65	0.23	reduced hatching success	medium uncertainty in TRV
PCB TEQs	**1.6**	0.16	reduced body weight, egg production, and survival of adults and embryos	high uncertainty in TRV because of high-level weekly exposures and IP injection; risk could be overestimated; derivation of TEFs is uncertain

[a] Level of uncertainty key:

Low = large dataset including chronic studies with species taxonomically similar to the ROC

Medium = moderately sized dataset including chronic studies

High = small dataset with only subchronic studies, unbounded NOAELs/LOAELs, or data with questionable data quality

COPC – chemical of potential concern

HQ – hazard quotient

IP – intraperitoneal

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

PCB – polychlorinated biphenyl

TEF – toxic equivalency factor

TEQ – toxic equivalent

TRV – toxicity reference value

Bold identifies NOAEL-based HQs greater than 1.0.

The NOAEL- and LOAEL-based HQs for osprey for chromium, lead, and mercury were less than 1.0, indicating that there are very low risks to osprey from exposure to these COPCs in the LDW. There is high uncertainty in the chromium TRV, but the LOAEL TRV would need to be lower by a factor of 50 for the LOAEL-based HQ to equal 1.0.

NOAEL- and LOAEL-based HQs for total PCBs were less than 1.0. The LOAEL-based HQ for PCBs using the PCB TEQ approach was less than 1.0, but the NOAEL-based HQ was 1.6, indicating a low risk with high uncertainty. There were no PCB congener data for juvenile chinook salmon, one of the three species assumed to be ingested by osprey in this assessment, so the portion of prey ingestion that had been assigned to juvenile chinook salmon (45%) was divided proportionally among shiner surfperch and English sole. Shiner surfperch had higher PCB concentrations than those of juvenile chinook salmon, so is it possible that PCB TEQ HQs would have been lower if PCB congener data had been available for salmon. There is also high uncertainty in the PCB TEQ TRV because birds in the study were dosed via IP injection rather than via

the diet and were a high-level weekly dose, which may result in an overestimate of risk.

Comparison of the BMF-estimated PCB concentration in osprey eggs, to the LOAEL TRV resulted in an HQ of 1.4. The risk based on this approach is highly uncertain, as discussed in Section A.6.3.3.2, resulting in overestimates or underestimates of risk. Thus, for PCBs, the ingested dose approach with its associated uncertainties indicated that risks to osprey were very low, while risks estimated using estimated concentrations in eggs indicated that risks were low with a high level of uncertainty.

A.6.3.4 River otter

This section present risk estimates, uncertainties, and risk conclusions for river otter.

A.6.3.4.1 Risk estimates

Five COPCs were evaluated for river otter: arsenic, cobalt, mercury, selenium, and PCBs (using both a total PCB and PCB TEQ approach). The only COPC with a LOAEL-based HQ greater than 1.0 was total PCBs, with a LOAEL-based HQ of 2.9. Mercury and PCB TEQs had NOAEL-based HQs of 2.8 and 4.5, respectively; the LOAEL-based HQs for these COPCs were less than 1.0 (Table A.6-62).

Table A.6-62. HQ calculations for river otter

ROC	COPC	Ingested Dose (mg/kg bw/day)	TRVs (mg/kg bw/day)		NOAEL-Based HQ	LOAEL-Based HQ
			NOAEL	LOAEL		
River otter	arsenic	0.32	2.6	5.4	0.12	< 0.1
	cobalt	0.016	0.10	1.0	0.16	< 0.1
	mercury	0.0048	0.0017	0.0084	**2.8**	0.57
	selenium	0.032	0.055	0.080	0.58	0.40
	Total PCBs	0.26	0.045	0.089	**5.8**	**2.9**
	PCB TEQs[a]	2.9×10^{-6}	6.5×10^{-7}	4.9×10^{-6}	**4.5**	0.59

[a] No PCB congener data were available for juvenile chinook salmon or mussels. When calculating risk estimates, the portion of prey ingestion that had been assigned to juvenile chinook salmon (22%) was divided proportionally among the remaining fish species, and mussel portion was added to the clam portion.

bw – body weight
COPC – chemical of potential concern
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
ROC – receptor of concern
TEQ – toxic equivalent
TRV – toxicity reference value

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

Table A.6-63 presents the relative contribution of each of the prey categories to the ingested PCB dose for river otter. For both the total PCB and PCB TEQ approaches, the largest contributions were from shiner surfperch and English sole.

Table A.6-63. Contribution of each prey category to the ingested PCB dose for river otter

Prey Species	Contribution to Ingested Dose (%)	
	Total PCBs	PCB TEQs
Shiner surfperch	39.6	53.5
English sole	28.3	27.4
Juvenile chinook salmon	9.6	0[a]
Pacific staghorn sculpin	14.2	12.7
Crab	7.7	5.7
Clam	0.5	0.6
Mussel	0.03	0[a]

[a] No PCB congener data were available for juvenile chinook salmon or mussels. The portion of prey ingestion that had been assigned to juvenile chinook salmon was divided proportionally among the other fish species, and the mussels portion was added to the clam portion.

PCB – polychlorinated biphenyl

TEQ – toxic equivalent

A.6.3.4.2 Uncertainties

This section presents a discussion of uncertainty associated with the problem formulation, the exposure and effects assessments, and the risk characterization for river otter.

Problem Formulation

Primary uncertainties in the problem formulation for river otter are associated with ROC selection and the COPC screen, as discussed below.

ROC Selection

Uncertainties related to the representation of other mammals in the LDW by river otter were addressed in the Phase 1 ERA (Windward 2003b). In that analysis, it was concluded that river otters are likely to be more highly exposed to COPCs from the LDW than muskrats or raccoons. Muskrats feed on plants, which are not abundant in the LDW, and raccoons feed on a greater proportion of terrestrial prey. Therefore, risk estimates for river otter should provide conservative risk estimates for other mammal species (i.e., muskrat and raccoon).

COPC Screen

Of the chemicals screened out as COIs for river otter in the first step of the COPC screen (Section A.2.5.3), 29 chemicals were detected in sediments but were not analyzed in tissue samples (Table A.6-16). None of these chemicals are defined as

bioaccumulative chemicals (EPA 2000a), so risks to mammals from exposure to these chemicals are assumed to be very low.

Eighty-six chemicals were identified in the problem formulation as COIs for mammals. Effects data for mammals were not available for 27 of the COIs. Of these COIs with no information on toxicity, 17 were individual PAHs. Risks to mammals from PAHs were evaluated using TRVs for four individual PAH compounds (1-methylnaphthalene, 2-methylnapthalene, benzo(a)pyrene, and naphthalene). The remaining COIs for which there were no TRVs for mammals included silver, two organotin compounds, four SVOCs, and three organochlorine pesticides. Risks to mammals from exposure to these COIs could not be evaluated.

Exposure Assessment

Uncertainties in the exposure assessment for river otter were associated with the following factors:

- Direct sediment contact
- Food and incidental sediment ingestion rates
- Dietary composition
- TEQ approach

Direct Sediment Contact

Risks to wildlife from direct contact with sediment are considered insignificant relative to risks from incidental sediment ingestion (EPA 2000b). However, the exclusion of this pathway adds a small amount of uncertainty to the risk estimate for river otter.

Food and Incidental Sediment Ingestion Rates

As discussed in the Phase 1 ERA (Windward 2003b), there is some uncertainty in the FIR for the river otter because it was calculated from an allometric equation for non-herbivorous placental mammals, rather than based on species-specific information. The FIR rate would need to increase by at least a factor of 2 to increase the NOAEL-based HQs that are less than 1.0 (i.e., arsenic, cobalt, and selenium) to greater than 1.0. This scenario is unlikely, so the uncertainty in the river otter's FIR is not expected to have an effect on risk conclusions.

To address uncertainties in the amount of sediment incidentally ingested by river otters while foraging, ingested doses of COPCs were calculated assuming the SIR was 10% of the FIR versus 2% assumed in Section 5.1.2.4. This conservative assumption would result in a slight increase of HQs; however, it would not change risk conclusions.

Dietary Composition

There are some uncertainties in the proportion of different types of fish species in the river otter's diet, which could affect exposure. To address this uncertainty, HQs for each COPC were calculated assuming that 75% of the river otter diet was composed of the fish species with the highest concentration of that COPC, with the remainder of the diet composed of equal proportions of the other fish species.[95] This assumption resulted in changes in HQs but did not result in a change of any HQ from less than 1.0 to greater than 1.0, or vice versa.

Toxic Equivalent Approach

WHO consensus TEF values from Van den Berg et al. (2006) were used to calculate TEQs for dioxin-like PCB congeners for mammals; these TEFs are presented in Attachment 3.

The TEFs for mammals were derived from a large number of studies, with priority given to *in vivo* toxicity data over *in vitro* data. Despite the numerous biological variables such as species, strain, sex, and age included in these studies, the TEF values for a given congener generally fall within a range of about an order of magnitude for mammals (Sanderson and Van den Berg 1999). It is not known if the uncertainties in these TEFs would overestimate or underestimate risk. It should also be noted that the TEQs calculated in this ERA were used only for assessing the toxicity of the dioxin-like PCB congeners and do not account for TEQ contributions from dioxin and furan congeners; dioxin and furan concentrations in tissue were not available for the LDW. Risk estimates for wildlife based on TEQs calculated considering only dioxin-like PCB congeners are likely lower than the actual risk resulting from the cumulative exposure of wildlife to dioxins, furans, and dioxin-like PCB congeners in the LDW.

Effects Assessment

Uncertainty associated with available toxicity benchmarks for mammals may affect risk estimates. The general uncertainties associated with toxicity studies are the same as those discussed in Section A.6.3.1.2 for spotted sandpiper. Specific uncertainties associated with toxicity studies for the mammalian COPCs are presented in Table A.6-64.

[95] In the exposure assessment, it is assumed that river otter ingest 22% shiner surfperch, 22% English sole, 22% juvenile chinook salmon, and 22% Pacific staghorn sculpin.

Table A.6-64. Level of uncertainty associated with TRVs for mammals

COPC	Number of TRV Studies	Uncertainty in TRV[a]
Arsenic	1	medium; selected TRVs were based on a chronic growth study[b]
Cobalt	3	medium; selected TRVs were based on a subchronic growth endpoint
Mercury	3	medium; selected TRVs were based on a chronic growth endpoint
Selenium	4	medium; selected TRVs were based on a subchronic growth endpoint
Total PCBs	10	medium
PCB TEQs	7	medium; selected TRVs were based on a subchronic growth endpoint

[a] Level of uncertainty key:

Low = large dataset including chronic studies with species (e.g., mink) taxonomically similar to the ROC

Medium = moderately sized dataset including chronic studies

High = small dataset with only subchronic studies, unbounded NOAELs/LOAELs, or data with questionable data quality

[b] Although only one study was available, it was a 2-year study with good data quality; the level of uncertainty is considered medium.

COPC – chemical of potential concern

PCB – polychlorinated biphenyl

TRV – toxicity reference value

There are uncertainties associated with the arsenic, cobalt, selenium, and PCB TEQ TRVs because they were based on subchronic or chronic growth endpoints, which could under- or overestimate risk. Uncertainty in the mercury total PCB TRVs could result in either an overestimate or underestimate of risk.

Summary of Uncertainties

Uncertainties in the problem formulation, the effects and exposure assessments, and risk characterization are summarized as follows:

- Uncertainties in food and incidental sediment ingestion rates and dietary composition are expected to have minimal or no effect on risk conclusions.
- The PCB TEQs were calculated using TEFs for individual PCB congeners, and the derivation of some of these TEFs is uncertain, with unknown effects on risk conclusions. Risk estimates for wildlife based on TEQs calculated considering only dioxin-like PCB congeners are likely lower than the actual risk resulting from the cumulative exposure of wildlife to dioxins, furans, and dioxin-like PCB congeners in the LDW.
- Gaps in toxicity data resulted in uncertainty in risk estimates. Risks may be under- or overestimated for arsenic, cobalt, mercury, and selenium, and overestimated for PCB TEQs; the effect of uncertainties associated with the total PCB TRVs is not known.

A.6.3.4.3 Risk conclusions

Risks were characterized for river otter as the most highly exposed semi-aquatic mammal using the LDW. Results from the risk characterization for river otter are summarized in Table A.6-65.

Table A.6-65. Summary of risk characterization for river otter

COPC	NOAEL-Based HQ	LOAEL-Based HQ	Effect	Primary Uncertainty[a]
Arsenic	0.12	< 0.1	reduced body weight (chronic exposure)	medium uncertainty in TRV[b]
Cobalt	0.16	< 0.1	reduced body weight (subchronic exposure)	medium uncertainty in TRV risk could be under- or overestimated because of endpoint
Mercury	**2.8**	0.57	reduced growth (chronic exposure)	medium uncertainty in TRV
Selenium	0.58	0.40	reduced body weight (subchronic exposure)	medium uncertainty in TRV; risk could be under- or overestimated because of endpoint
Total PCBs	**5.8**	**2.9**	reduced offspring growth	medium uncertainty in TRV
PCB TEQs	**4.5**	0.59	reduced body weight (subchronic exposure)	medium uncertainty in TRV; risk could be under- or overestimated because of endpoint; derivation of TEFs

[a] Level of uncertainty key:

Low = large dataset including chronic studies with species taxonomically similar to the ROC (i.e., mink)

Medium = moderately sized dataset including chronic studies

High = small dataset with only subchronic studies, unbounded NOAELs/LOAELs, or data with questionable data quality

[b] Although only one study was available, it was a 2-year study with good data quality, so the level of uncertainty is considered medium

COPC – chemical of potential concern
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
TEF – toxic equivalency factor
TEQ – toxic equivalent
TRV – toxicity reference value

Bold identifies NOAEL-based HQs greater than 1.0 and LOAEL-based HQs greater than or equal to 1.0.

The LOAEL- and NOAEL-based HQs for PCBs of 2.9 and 5.8, respectively, using the total PCB approach, indicate risk to river otters from PCB exposure. The PCB TRVs are based on an 18-month study with mink in which reduced growth of offspring was reported. The NOAEL-based HQ for PCBs of 4.5, using the PCB TEQ approach, indicates that there could be some risk to river otters from PCB exposure, although the LOAEL-based HQ was < 1.0 using this approach. The PCB TEQ TRVs are based on a 90-day study with guinea pigs in which reduced body weights were reported. There are some uncertainties associated with the PCB TRVs and with the TEF approach, but it is not known whether these uncertainties would underestimate or overestimate risk. The next highest NOAEL and LOAEL values for total PCBs were from a chronic study with mink, which resulted in no effects on offspring kit survival at a dose of 2.6×10^{-6}

mg/kg bw/day but decreased survival in kits at 3 weeks at a dose of 9.1 x 10^{-6} mg/kg bw/day (Hochstein et al. 2001). If the NOAEL and LOAEL from this study had been used to estimate risk, the LOAEL-based HQs would have been less than 1.0, and the NOAEL-based HQ would be 2.5. Based on this analysis, risks to river otter from PCBs are low, with some uncertainty.

For mercury, the LOAEL-based HQ was less than 1.0 (0.57), but the NOAEL-based HQ was greater than 1.0 (2.8). There is some uncertainty in the TRV, which was based on a chronic study of rats that reported effects on growth because the NOAEL was estimated from the LOAEL using an uncertainty factor of 5.0. The next highest NOAEL and LOAEL values were from a 93-day study with mink, which resulted in no effects on growth and survival at a dose of 0.16 mg/kg bw/day and effects on growth and survival at a dose of 0.25 mg/kg bw/day (Wobeser et al. 1976). If the NOAEL and LOAEL from this study had been used to estimate risk, the NOAEL- and LOAEL-based HQs would have been substantially less than 1.0. Based on this analysis, risks to river otter from mercury are low, with some uncertainty.

The NOAEL- and LOAEL-based HQs for river otter for arsenic, cobalt, and selenium were less than 1.0, indicating that there are very low risks to river otters from exposure to these COPCs in the LDW.

A.6.3.5 Harbor seal

This section present risk estimates, uncertainties, and risk conclusions for harbor seal.

A.6.3.5.1 Risk estimates

The COPCs evaluated for harbor seal were mercury and PCBs (using both a total PCB and PCB TEQ approach). The NOAEL- and LOAEL-based HQs for both COPCs were less than 1.0 (Table A.6-66).

Table A.6-66. HQ calculations for harbor seal

ROC	COPC	Ingested Dose (mg/kg bw/day)	TRVs (mg/kg bw/day)		NOAEL-Based HQ	LOAEL-Based HQ
			NOAEL	LOAEL		
Harbor seal	mercury	0.00033	0.0017	0.0084	0.19	< 0.1
	total PCBs	0.020	0.045	0.089	0.44	0.22
	PCB TEQs	2.3 x 10^{-7}	6.5 x 10^{-7}	4.9 x 10^{-6}	0.35	< 0.1

[a] No PCB congener data were available for juvenile chinook salmon. When calculating risk estimates, the portion of prey ingestion that had been assigned to juvenile chinook salmon (25%) was divided proportionally among the remaining prey categories.

bw – body weight
COPC – chemical of potential concern
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
ROC – receptor of concern
TEQ – toxic equivalent
TRV – toxicity reference value

A.6.3.5.2 Uncertainties

This section presents a discussion of uncertainty associated with the problem formulation, the exposure and effects assessments, and the risk characterization for harbor seal.

Problem Formulation

Primary uncertainties in the problem formulation for harbor seal are associated with ROC selection and the COPC screen. Uncertainties in the COPC screen for mammals were discussed in Section A.6.3.4.2 for river otter.

Uncertainties related to the representation of other marine mammals in the LDW (i.e., sea lions or harbor porpoises) by harbor seal were addressed in the Phase 1 ERA (Windward 2003b). In that analysis, it was concluded that all three species are opportunistic feeders, primarily feeding on fish and some invertebrates, so harbor seals should have an exposure similar to that of sea lions and harbor porpoises. Furthermore, harbor seals and sea lions are expected to have similar exposure to LDW prey because LWG site usage for these two species is similar. LDW site use by harbor porpoise is unknown because data are not available.

Exposure Assessment

Uncertainties in the exposure assessment for harbor seal were associated with the following factors:

- Direct sediment contact
- Incidental sediment ingestion rate
- Dietary composition
- TEQ approach

The uncertainties associated with direct sediment contact and the TEQ approach are the same as those discussed in Section A.6.3.4.2 for river otter.

Incidental Sediment Ingestion Rate

To address uncertainties in the amount of sediment incidentally ingested by harbor seal while foraging, ingested doses of COPCs were calculated assuming the SIR was 10% of the FIR versus 2% assumed in Section 5.1.2.5. This conservative assumption would result in a slight increase of HQs (by an average of less than 0.1) but would not change risk conclusions.

Dietary Composition

There are some uncertainties in the proportions of different types of fish species in the harbor seal's diet, which could affect exposure. To address this uncertainty, HQs for each COPC were calculated assuming that 75% of the harbor seal diet was composed of the fish species with the highest concentration of that COPC, with the remainder of

the diet composed of equal proportions of the other fish species.[96] This assumption resulted in changes in HQs but did not result in a change of any HQ from less than 1.0 to greater than 1.0, or vice versa.

Effects Assessment

Uncertainties associated with mammalian TRVs are summarized in Table A.6-64.

Summary of Uncertainties

Uncertainties in the problem formulation, the exposure and effects assessments, and the risk characterization are summarized as follows:

- Uncertainties in incidental sediment ingestion and dietary composition are expected to have minimal or no effect on risk conclusions.
- The PCB TEQs were calculated using TEFs for individual PCB congeners, and the derivation of some of these TEFs is uncertain, with unknown effects on risk estimates. Risk estimates for wildlife based on TEQs calculated considering only dioxin-like PCB congeners are likely lower than the actual risk resulting from the cumulative exposure of wildlife to dioxins, furans, and dioxin-like PCB congeners in the LDW.
- Gaps in toxicity data resulted in uncertainty in risk estimates. The effects of uncertainties in the TRVs are not known.

A.6.3.5.3 Risk conclusions

The harbor seal was chosen to represent marine mammals in the LDW. Results of the harbor seal risk characterization are summarized in Table A.6-67.

Table A.6-67. Summary of risk characterization for harbor seal

COPC	NOAEL-Based HQ	LOAEL-Based HQ	Effect	Primary Uncertainty[a]
Mercury	0.19	< 0.1	reduced growth (chronic exposure)	medium uncertainty in TRV
Total PCBs	0.44	0.22	reduced offspring growth	medium uncertainty in TRV
PCB TEQs	0.35	< 0.1	reduced body weight (subchronic exposure)	medium uncertainty in TRV; risk could be under- or overestimated because of endpoint; derivation of TEFs

[a] Level of uncertainty key:

Low = large dataset including chronic studies with species taxonomically similar to the ROC

Medium = moderately sized dataset including chronic studies

High = small dataset with only subchronic studies, unbounded NOAELs/LOAELs, or data with questionable data quality

[96] In the exposure assessment, it is assumed that harbor seal ingest 25% shiner surfperch, 25% English sole, 25% juvenile chinook salmon, and 25% Pacific staghorn sculpin.

COPC – chemical of potential concern
HQ – hazard quotient
LOAEL – lowest-observed-adverse-effect level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
TEF – toxic equivalency factor
TEQ – toxic equivalent
TRV – toxicity reference value

The NOAEL- and LOAEL-based HQs for harbor seal for mercury and PCBs were less than 1.0, indicating that there is very low risk to harbor seals from exposure to these COPCs in the LDW. The primary uncertainties associated with this conclusion are the general limitations in derivation of TRVs from laboratory toxicity tests.

A.6.3.6 Summary of risk conclusions for wildlife

In summary, results of the wildlife risk estimates and evaluation of associated uncertainties are as follows:

- **Spotted sandpiper.** Exposure estimates were greater than doses associated with adverse effects for chromium and lead in Area 2, copper and mercury in Area 3, and for vanadium in all three exposure areas. Risks from PCBs and DDT were low to very low, with some uncertainty. Risks from arsenic, cadmium, cobalt, nickel, selenium, and zinc were very low.
- **Great blue heron.** Risks to great blue heron from exposure to COPCs in the LDW were very low.
- **Osprey.** PCB risks to osprey were low, with high uncertainty associated with the TEQ approach and BMF calculations. Risks to osprey from other COPCs in the LDW were very low.
- **River otter.** PCBs pose a risk to otters in the LDW with exposure concentrations greater than those associated with adverse effects. Risk to river otter from exposure to mercury was low. There were very low risks to river otter from other COPCs.
- **Harbor seal.** Risks to harbor seals from exposure to COPCs in the LDW were very low.

A.7.0 Selection of Primary Ecological Risk Drivers/Indicator Hazardous Substances

This section presents the rationale for the identification of chemicals as risk drivers (CERCLA terminology) or indicator hazardous substances (MTCA terminology) based on estimated risks to ecological receptors.[97] The risk drivers from both this ERA and the HHRA will be the focus of remedial analyses in the FS.

In this ERA, ecological risks from chemicals were assessed consistent with CERCLA (EPA 1998) and SMS guidance (WAC 173-204). As a result, chemicals detected in sediment and tissue samples collected from the LDW were grouped as follows: 1) chemicals for which there is no cause for concern, 2) COPCs, and 3) COPCs with LOAEL-based HQs ≥ 1.0 or that exceed the SQS of the SMS. This section describes additional considerations and documents decisions about which of the chemicals in the third group are considered risk drivers for ecological receptors (Table A.7-1).

Table A.7-1. COPCs, COPCs that exceed relevant risk thresholds, and risk drivers

Designation	Chemical	
	Benthic Invertebrates	Crab, Fish, or Wildlife Receptors
COPCs	**SMS chemicals** – 41 COPCs including metals, PAHs, PCBs, phthalates, and other SVOCs based on detected exceedance of SQS at one or more locations **non-SMS chemicals** – TBT, nickel, total DDTs, total chlordane, cis-1,2-dichloroethene	**Crabs** – zinc and PCBs **Fish** – arsenic, cadmium, copper, vanadium, PCBs, TBT **Birds** – arsenic, cadmium, chromium, cobalt, copper, lead, mercury, nickel, selenium, vanadium, zinc, PCBs **Mammals** – arsenic, cobalt, mercury, selenium, PCBs
COPCs above risk thresholds[a]	**SMS chemicals** – 41 COPCs **non-SMS** chemicals – nickel, total DDTs, total chlordane	**Crabs** – PCBs **Fish** – cadmium, vanadium, PCBs **Birds** –chromium, copper, lead, mercury, vanadium, PCBs **Mammals** – PCBs
Risk drivers	**SMS chemicals** – 41 COPCs	**Mammals** – PCBs

[a] LOAEL-based HQ ≥ 1.0 or exceedance of the SQS of the SMS.

CERCLA – Comprehensive Environmental Response, Compensation, and Liability Act

COPCs – chemicals of potential concern

MTCA – Model Toxics Control Act

PAHs – polycyclic aromatic hydrocarbons

PCBs – polychlorinated biphenyls

SMS – Washington State Sediment Management Standards

SQS – sediment quality standard

SVOCs – semivolatile organic compounds

TBT – tributyltin

[97] For simplicity, these chemicals will be referred to as risk drivers in the remainder of this section.

A.7.1 CONSIDERATIONS FOR IDENTIFICATION OF RISK DRIVERS

In the problem formulation of this ERA, all analyzed chemicals in the LDW were screened to identify COPCs by comparing maximum exposure concentrations either to the SQS or to NOAEL TRVs (Section A.2.5). The resulting COPCs were further evaluated to derive risk estimates based on more realistic exposure assumptions and a range of toxicity data (i.e., from the highest NOAEL TRV to the lowest LOAEL TRV). Uncertainties in risk estimates were discussed in the risk characterization. In this section, the following additional factors are considered to identify which COPCs that exceeded effects thresholds are risk drivers. Considerations include:

- Uncertainty in risk estimates based on quantity and quality of exposure data
- Uncertainty in risk estimates based on quantity and quality of effects data
- Comparison of concentrations in LDW sediment to regional natural background concentrations in sediment
- The likely magnitude of residual risks following planned sediment remediation within early action areas in the LDW

Risk drivers will be the focus of detailed analyses presented in the FS for all remedial alternatives. COPCs that exceeded risk thresholds, also referred to as chemicals of concern (COCs), that were not selected as risk drivers may also be further evaluated, in consultation with EPA and Ecology. This evaluation may include:

- Assessment of the likely reductions in sediment concentrations or residual risks from these COCs following implementation of the preferred alternative selected in the FS
- Review of any new effects or exposure information as part of the 5-year review
- Inclusion of COCs in any post-remedial monitoring program

A.7.2 IDENTIFICATION OF RISK DRIVER CHEMICALS

Based on the four considerations outlined above, and in consultation with EPA and Ecology, 41 chemicals were selected as risk drivers for benthic invertebrates. PCBs were the only chemical identified as a risk driver for ecological receptors other than benthic invertebrates. The rationale for the selection of these risk drivers is summarized in Table A.7-2 and described briefly below.

Table A.7-2. Rationale for risk driver designation

COPC	ROC	Maximum NOAEL-Based HQ	Maximum LOAEL-Based HQ	Additional Considerations	Risk Driver?
Total PCBs	crabs	**10**	**1.0**	Uncertainty in exposure data: whole-body concentrations were estimated Uncertainty in effects data: LOAEL-based HQ was based on a study with Aroclor 1016 and grass shrimp, and NOAEL was estimated using an uncertainty factor; selection of next higher TRV would result in LOAEL-based HQ < 1.0	no
	river otter	**5.8**	**2.9**	Uncertainty in exposure data: low uncertainty in diet assumptions and home range Uncertainty in effects data: low uncertainty in TRV (growth endpoint in kits)	yes
	English sole	**4.9 – 25**[a]	0.98 – **5.0**[a]	Uncertainty in exposure data: low uncertainty in tissue concentrations Uncertainty in effects data: high uncertainty in lowest LOAEL TRV because of uncertain statistical significance of the fecundity endpoint for the low dose, a lack of dose-response in the fecundity endpoint, uncertain number of fish used in the experiment, and uncertainties associated with fish handling and maintenance protocols	no
	Pacific staghorn sculpin	**3.8 – 19**[a]	0.76 **- 3.8**[a]	Same considerations as listed above for English sole	no
PCB TEQ[b]	spotted sandpiper – Area 2 (high-quality foraging habitat)	**15**	**1.5**	Uncertainty in exposure data: low uncertainty in diet assumptions and home range Uncertainty in effects data: high uncertainty in TRV, which was based on study of reproduction with weekly IP injection; high uncertainty in TEFs; effects data for total PCBs are less uncertain than for PCB TEQs and the LOAEL-based HQ for total PCBs was < 1.0	no
Cadmium	juvenile chinook salmon	**5.0**	**1.0**	Uncertainty in exposure data: LOAEL-based HQ < 1.0 if empirical juvenile chinook salmon stomach contents data from the LDW are used to estimate exposure, instead of estimating exposure based on ingestion of benthic invertebrates Uncertainty in effects data: high uncertainty in the lowest TRV because selection of next higher TRV would result in LOAEL-based HQ < 1.0, all salmonid-specific studies for cadmium with NOAELs result in NOAEL-based HQs less than 0.01	no
	English sole	**6.1**	**1.2**	Uncertainty in exposure data: low uncertainty (LDW-collected benthic invertebrate tissue samples) Uncertainty in effects data: high uncertainty in the lowest TRV because selection of next higher TRV would result in LOAEL-based HQ < 1.0; all other NOAELs and LOAELs were orders of magnitude higher than the selected LOAEL	no

Table A-7.2. Rationale for risk driver designation, cont.

COPC	ROC	Maximum NOAEL-Based HQ	Maximum LOAEL-Based HQ	Additional Considerations	Risk Driver?
	Pacific staghorn sculpin	**5.2**	**1.0**	Uncertainty in exposure data: low uncertainty (LDW-collected shiner surfperch and benthic invertebrate tissue samples) Uncertainty in effects data: high uncertainty in the lowest TRV because selection of next higher TRV would result in LOAEL-based HQ < 1.0; all other NOAELs and LOAELs were orders of magnitude higher than the selected LOAEL	no
Chromium	spotted sandpiper –Area 2 (high- and poor-quality foraging habitat)	**8.8**	**1.8**	Uncertainty in exposure data: high uncertainty because LOAEL-based HQ would be less than 1.0 if the single anomalously high benthic invertebrate tissue sample from RM 3.0 west was excluded; chromium concentrations in sediment were low in this area Uncertainty in effects data: high uncertainty; only one study with reported effects, and study was unpublished and could not be obtained for review	no
Copper	spotted sandpiper –Area 3 (high- and poor-quality foraging habitat)	**1.5**	**1.1**	Uncertainty in exposure data: low uncertainty Comparison to natural background: concentration in sediment (SWAC of 57 mg/kg dw) from Area 3 (high- and poor-quality foraging habitat) similar to PSAMP rural Puget Sound concentrations (50 mg/kg dw [90th percentile]) Residual risk: following planned sediment remediation within early action areas, LOAEL-based HQ would be < 1.0	no
Lead	spotted sandpiper –Area 2 (high- and poor-quality foraging habitat)	**19**	**5.5**	Uncertainty in exposure data: high uncertainty because LOAEL-based HQ would be less than 1.0 if the single anomalously high benthic invertebrate tissue sample from RM 3.0 west was excluded; lead concentrations in sediment were low in this area Uncertainty in effects data: low uncertainty (reproductive endpoint)	no
	spotted sandpiper –Area 3 (high- and poor-quality foraging habitat)	**5.0**	**1.5**	Uncertainty in exposure data: low uncertainty Uncertainty in effects data: low uncertainty (reproductive endpoint) Residual risk: following planned sediment remediation within early action area, LOAEL-based HQ would be < 1.0	
Mercury	spotted sandpiper –Area 3 (high- quality foraging habitat)	**5.3**	**1.0**	Uncertainty in exposure data: low uncertainty Uncertainty in effects data: low uncertainty (TRV was based on a growth endpoint) Residual risk: following planned sediment remediation within early action area, LOAEL-based HQ would be < 1.0	no

Table A-7.2. Rationale for risk driver designation, cont.

COPC	ROC	Maximum NOAEL-Based HQ	Maximum LOAEL-Based HQ	Additional Considerations	Risk Driver?
Vanadium	English sole	**5.9**	**1.2**	Uncertainty in exposure data: low uncertainty Uncertainty in effects data: high uncertainty in TRV because only one study was available Comparison to natural background: exposure concentration in LDW sediment (SWAC of 58 mg/kg dw) was less than PSAMP rural Puget Sound concentration (64 mg/kg dw [90th percentile])	no
	Pacific staghorn sculpin	**3.2 – 5.9**	0.65 **– 1.2**	Same considerations as listed for English sole above	no
	spotted sandpiper – all exposure areas	**2.0 – 2.7**	**1.0 – 1.4**	Uncertainty in exposure data: low uncertainty Uncertainty in effects data: TRV was based on a 4-week growth endpoint, with uncertainty (two available studies: one with reduced body weight in chickens after 4 weeks and the other with no effect on body weight in mallards after 10 weeks) Comparison to natural background: mean exposure concentrations in sandpiper exposure areas ranged from 49 to 57 mg/kg dw, compared to PSAMP rural Puget Sound background concentration of 64 mg/kg dw (90th percentile)	no
41 SMS chemicals[c]	benthic invertebrates	range of values	range of values	Each of these 41 chemicals had at least one detected exceedance of SQS in baseline surface sediment dataset.	yes
Nickel	benthic invertebrates	**6.6**	**2.5**	Uncertainty in exposure data: low uncertainty Uncertainty in effects data: medium uncertainty in the TRV (i.e., the ML) because only no-effects data (amphipod mortality and community abundance AETs) were available; no information was available regarding concentrations associated with adverse effects Residual risk: ML was exceeded at four locations in LDW – all within early action areas with planned sediment remediation	no
Total DDTs	benthic invertebrates	**5.1**	**2.7**	Uncertainty in exposure data: medium uncertainty (i.e., likely interference in pesticide analyses from PCBs) Uncertainty in effects data: medium uncertainty; based on a single study with spiked sediment Residual risk: LOAEL was exceeded at only one location in LDW, location is within early action area with planned sediment remediation	no

Table A-7.2. Rationale for risk driver designation, cont.

COPC	ROC	Maximum NOAEL-Based HQ	Maximum LOAEL-Based HQ	Additional Considerations	Risk Driver?
Total chlordane	benthic invertebrates	**82**	**48**	Uncertainty in exposure data: highly uncertain because all total chlordane concentrations in samples from Phase 2 locations were JN-qualified as a result of probable PCB interference; except one location at RM 2.2, all locations with detected total chlordane concentrations co-occurred with elevated PCB concentrations Uncertainty in effects data: TRV is highly uncertain because it was based on a general Canadian sediment guideline (PEL); this guideline is based mainly on field-collected data with complex mixtures of chemicals Residual risk: LOAEL was exceeded at 14 locations in LDW; all but one of these locations are associated with an early action area with planned sediment remediation	no

Note: HQs for fish are the highest HQs in cases where more than one approach was used.

[a] LOAEL-based HQs were calculated from a range of effects concentrations reported in Hugla and Thome (1999) because of uncertainty in the LOAEL. The NOAEL TRV range was estimated by dividing the LOAEL TRV range by an uncertainty factor of 5. Ranges reported for Pacific staghorn sculpin also included the range in exposure estimates for areas smaller than the entire LDW.

[b] Risk estimates based on TEQs were calculated using only tissue data for dioxin-like PCB congeners because dioxin and furan tissue data were not available. Thus, risks associated with exposure to all dioxin-like chemicals were likely underestimated; the degree of underestimation is uncertain.

[c] Arsenic, cadmium, chromium, copper, lead, mercury, silver, zinc, acenaphthene, anthracene, benz(a)anthracene, benzo(a)pyrene, benzo(g,h,i)perylene, chrysene, dibenzo (a,h)anthracene, fluoranthene, fluorene, indeno (1,2,3,-c,d)pyrene, naphthalene, phenanthrene, pyrene, total benzofluoranthenes, HPAH, LPAH, bis(2-ethylhexyl) phthalate, butyl benzyl phthalate, dimethyl phthalate, 1,2-dichlorobenzene, 1,4-dichlorobenzene, 1,2,4-trichlorobenzene, 2-methylnaphthalene, 4-methylphenol, 2,4-dimethylphenol, benzoic acid, benzyl alcohol, dibenzofuran, hexachlorobenzene, n-nitrosodiphenylamine, pentachlorophenol, phenol, total PCBs.

dw – dry weight
ERA – ecological risk assessment
HQ – hazard quotient
IP – intraperitoneal
LDW – Lower Duwamish Waterway
LOAEL – lowest-observed-adverse-effect level
ML – maximum level
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
PEL – probable effects level
RM – river mile
ROC – receptor of concern
SQS – sediment quality standard
SWAC – spatially weighted average concentration
TEQ – toxic equivalent
TEF – toxic equivalency factor
TRV – toxicity reference value

Bold identifies NOAEL-based HQ greater than 1.0 and LOAEL-based HQ greater than or equal to 1.0.

The 41 chemicals identified as risk drivers for benthic invertebrates were selected because the detected concentration of those chemicals in the LDW baseline surface sediment dataset exceeded the SQS of the SMS at one or more locations, and SMS is a key regulation governing sediment remediation in the State of Washington. Chemicals were selected as risk drivers regardless of site-specific toxicity testing results because of the uncertainty in the cause-and-effect relationship on a chemical-specific basis.

Five chemicals (hexachlorobutadiene, 2-methylphenol, acenaphthylene, di-n-octyl phthalate, and diethyl phthalate) had RLs that were greater than the SQS but did not have detected concentrations greater than the SQS. These five chemicals were not identified as risk drivers for the reasons discussed below.

Hexachlorobutadiene was not selected as a risk driver because although some of the RLs for hexachlorobutadiene were above the SQS, hexachlorobutadiene was never detected in the 782 samples that were analyzed for this compound (Table A.7-3). The improved analytical techniques used to develop the RI/FS dataset, relative to earlier studies, reduced the frequency of RLs above the SQS from 23% (non-LDWG data) to 8% (LDWG data) with no increase in detection frequency. The chemical 2-methylphenol was rarely detected (0.4% detection frequency). The frequency of RLs for 2-methylphenol above the SQS was reduced from 17% (non-LDWG data) to 8.3% (LDWG data), with no detected concentrations above the SQS. The RLs above the SQS for these compounds appear to reflect difficulties routinely encountered with the analysis of these compounds, and thus these chemicals were not selected as risk drivers.

Table A.7-3. Chemicals with RLs greater than SQS that were not identified as risk drivers

Chemical	Number of Samples Analyzed	% Detected	% RLs Above SQS
Hexachlorobutadiene	782	0%	18% (148 samples)
2-Methyl phenol	785	0.4%	15% (117 samples)
Acenaphthylene	782	16%	0.4% (3 samples)
Di-n-octyl phthalate	796	6%	1.0% (8 samples)
Diethyl phthalate	796	0.12%	0.7% (6 samples)

RL – reporting limits

SQS – sediment quality standards (SMS)

Acenaphthylene, di-n-octyl phthalate, and diethyl phthalate were detected in 16%, 6%, and 0.12%, respectively, of the samples analyzed for these compounds (Table A.7-3); none of the detected concentrations were greater than the SQS. The frequencies of RLs above the SQS for acenaphthylene, di-n-octyl phthalate, and diethyl phthalate were all very low (Table A.7-3). These chemicals were not selected as risk drivers because they were never detected above the SQS, and the frequencies of RLs above the SQS in the LDWG data were very low. The RLs above the SQS were generally associated with issues of sample dilution.

Nickel, total DDTs, and total chlordane do not have SQS values but did have LOAEL-based HQs for benthic invertebrates greater than 1.0 based on other guidelines or TRVs. However, these three chemicals were not selected as risk drivers for benthic invertebrates primarily because of uncertainties in their TRVs and because areas with concentrations greater than those TRVs were all within planned sediment remediation areas, except for one location with elevated total chlordane at RM 0.85 in the navigation channel. This sample was JN-qualified, indicating that the chlordane identification was uncertain and the concentration was likely biased high because of analytical interference with PCBs.

PCBs were selected as a risk driver for river otter because the LOAEL-based HQ exceeded 1.0 (HQ of 2.9), and the uncertainties associated with the exposure and effects data were relatively low.

COPCs not selected as risk drivers for ecological receptors other than benthic invertebrates but with LOAEL-based HQs greater than or equal to 1.0, included PCBs (crabs, English sole, Pacific staghorn sculpin, and spotted sandpiper), cadmium (juvenile chinook salmon, English sole, and Pacific staghorn sculpin), chromium (spotted sandpiper), copper (spotted sandpiper), lead (spotted sandpiper), mercury (spotted sandpiper), and vanadium (juvenile chinook salmon, English sole, Pacific staghorn sculpin, and spotted sandpiper).

For crabs, PCBs were not selected as a risk driver primarily because of the uncertainty in the effects data and the low risk estimate (LOAEL-based HQ equal to 1.0). The LOAEL-based HQ was based on a study with Aroclor 1016 and grass shrimp and the NOAEL was estimated using an uncertainty factor. The LOAEL-based HQ would have been less than 1.0 if the next highest TRV (based on Aroclor 1254 and grass shrimp) had been used instead.

For English sole and Pacific staghorn sculpin, PCBs were not selected as a risk driver. The risk estimates for PCBs in these two fish species are uncertain both because the exposure concentrations were in between the concentrations selected as the LOAEL range and because of uncertainty in the study that served as the basis for the LOAEL range itself.

For spotted sandpiper, the LOAEL-based HQ for PCB TEQ was 1.5. However, PCBs were not selected as a risk driver for spotted sandpiper primarily because LOAEL-based HQs for total PCBs were less than 1.0 and risk estimates for total PCBs are more certain than risk estimates for PCB TEQs.

Cadmium was not selected as a risk driver for juvenile chinook salmon, English sole, or for Pacific staghorn sculpin primarily because the selected TRV was highly uncertain (i.e., the selected LOAEL was orders of magnitude lower than NOAELs or LOAELs from all other studies and the observed effects were partially attributed to reduced food intake), and risk estimates were low (LOAEL-based HQs of 1.2 for both juvenile chinook salmon and English sole and up to 1.0 for Pacific staghorn sculpin).

Chromium and lead were not selected as risk drivers for spotted sandpiper because elevated risks within Area 2 were driven by elevated concentrations in a single benthic invertebrate tissue sample collected near RM 3.0 on the west side of the LDW. Dry weight concentrations of chromium and lead were much higher in this sample than in the rest of the benthic invertebrate tissue dataset, whereas concentrations of chromium and lead were not elevated in the sediment sample co-located with this benthic invertebrate composite tissue sample.[98]

Copper was not selected as a risk driver for spotted sandpiper primarily because sediment concentrations of copper in Area 3 (57 mg/kg dw) are similar to the 90th percentile copper concentration (50 mg/kg dw) in PSAMP rural Puget Sound sediment and because LOAEL-based HQs will be less than 1.0 following planned sediment remediation within early action areas. Lead (Area 3) and mercury also were not selected for spotted sandpiper primarily because LOAEL-based HQs will be less than 1.0 following planned sediment remediation within early action areas.

Vanadium was not selected as a risk driver for either fish or spotted sandpiper because of high uncertainty in effects data (few toxicity studies) and because sediment concentrations of vanadium in exposure areas (ranging from 49 to 58 mg/kg dw) are less than the 90th percentile vanadium concentration (64 mg/kg dw) in PSAMP rural Puget Sound sediment. Background information is important because the CERCLA program generally does not require clean up to concentrations below natural or anthropogenic background levels (EPA 2002).

No quantitative ecological risk estimates were calculated for dioxins and furans within the LDW and thus the level of ecological risk from dioxins and furans is not known. Risks were not calculated for several reasons. Primarily, human health risks from dioxins and furans through seafood consumption were assumed to be unacceptable; and therefore, neither tissues from the LDW nor from background areas were analyzed for dioxins and furans. Dioxins and furans were determined to be a risk driver based on human health risks from both seafood consumption and direct contact pathways. Risk management decisions to address dioxin and furan contamination in LDW sediment will be based on MTCA and CERCLA regulations and guidance. Remedial decisions to address dioxin and furan contamination in sediment will be made by EPA and Ecology as part of the FS process and will be documented in the Record of Decision. Additional detail on dioxins and furans is provided in Section B.5.5.2 of the HHRA.

---

[98] Of the eight benthic invertebrate tissue samples analyzed in Area 2, the concentration of chromium in sample B7a was 58 mg/kg dw, compared to an average concentration of 8.7 mg/kg dw in the seven other samples. The concentration of lead in sample B7a was 220 mg/kg dw, compared to an average concentration of 11 mg/kg dw in the seven other samples. The chromium and lead concentrations in the co-located sediment sample were 22.9 mg/kg dw and 21.4 mg/kdw dw, respectively (the 90th percentile chromium and lead concentrations in PSAMP rural Puget Sound sediment were 58.8 mg/kg dw and 19.5 mg/kg dw, respectively).

A.8.0 Conclusions

Baseline risks were calculated in this ERA based on chemical concentrations in sediment, water, and tissue samples collected from the LDW to estimate the chemical exposure of benthic invertebrates, crabs, fish, birds, and mammals that may reside or forage in the LDW for at least a portion of their lives. Site-specific sediment toxicity tests and a site-specific investigation of imposex in gastropods were also conducted. Several conservative assumptions were employed in the risk assessment to compensate for a variety of uncertainties. Based on risk estimates and associated uncertainties, the main conclusions regarding risks to ecological receptors in the LDW from COPCs are as follows:

- **Benthic Invertebrate Community —**Sediment chemistry and site-specific toxicity test results indicate that no adverse effects to benthic invertebrates living in intertidal and subtidal sediments are predicted for 75% of the LDW area (i.e., the area in which chemical concentrations were less than or equal to SQS chemical criteria and where sediments were nontoxic according to SQS biological effects criteria). There is a higher likelihood for adverse effects in approximately 7% of the LDW area, which was designated as having chemical concentrations or biological effects in excess of the CSL. The remaining 18% of the LDW area had chemical concentrations or biological effects between the SQS and CSL, indicating that risks to benthic invertebrate communities are less certain in these areas than in areas with concentrations greater than one or more CSL values. Some uncertainty is associated with these area estimates because areas were estimated by interpolating from individual points at which sediments were sampled. The SQS and CSL were exceeded by 39 chemicals; two additional chemicals exceeded only the SQS.[99] Risks to the benthic invertebrate community from all VOCs detected in sediment porewater were very low, except for cis-1,2-dichloroethene, which had concentrations 21 times the no-effects concentration in a small area at RM 2.4; all concentrations of cis-1,2-dichloroethene were less than the concentration associated with adverse effects. Therefore, there is uncertainty whether exposure to cis-1,2-dichloroethene within the LDW is sufficiently high to result in adverse effects in this small area. Risks to benthic invertebrates from TBT were very low based on NOAEL-based HQs less than 1.0 and the absence of imposex in all gastropods, except one species of neogastropod with imposex characterized as Stage 2, a stage that is not expected to impact reproduction.
- **Crabs—**Exposure concentrations of total PCBs in tissue were equal to concentrations associated with adverse effects in crabs, indicating the potential

[99] Total DDTs, nickel, and total chlordane also exceeded their DMMP guidelines or literature-based TRV at one or more locations.

for adverse effects. Exposure concentrations of zinc in tissue were greater than concentrations associated with no effects but less than those associated with adverse effects, indicating there is uncertainty whether exposure within the LDW is sufficiently high to result in adverse effects.

- **Fish —** Risks were evaluated for three fish ROCs (English sole, Pacific staghorn sculpin, and juvenile chinook salmon), which serve as representative surrogates for all other fish species in the LDW. Exposure concentrations for three of the six COPCs for fish (PCBs, cadmium, and vanadium) were greater than concentrations associated with adverse effects for English sole. LOAEL-based HQs for both metals were 1.2, and LOAEL-based HQs for PCBs ranged from 0.98 to 5.0 based on effects concentrations in the study reporting the lowest TRVs. Therefore, there is a potential for adverse effects to fish from PCBs, but risk estimates are uncertain because the exposure concentrations were in between the concentrations selected as the LOAEL range and because the study that served as the basis for the LOAEL range was uncertain. Estimated exposures of English sole to two additional COPCs (arsenic and copper) were greater than their respective no-effects levels but were lower than the adverse effect levels associated with survival, growth, or reproduction, indicating that there is uncertainty whether exposure within the LDW is sufficiently high to result in adverse effects. Site-specific studies of English sole indicate the potential for reproductive effects that correlate with exposure to chemical mixtures in the field. Exposure concentrations of PCBs, cadmium, and vanadium for Pacific staghorn sculpin were greater than or equal to the concentrations associated with adverse effects in at least one area within the LDW. LOAEL-based HQs of up to 1.0 and 1.2 were estimated for cadmium and vanadium, respectively, indicating a potential for adverse effects. LOAEL-based HQs for PCBs ranged from 0.30 to 3.8 based on effects concentrations in the study reporting the lowest TRVs. Therefore, there is a potential for adverse effects from PCBs, but risk estimates are uncertain because the exposure concentrations were in between the concentrations selected as the LOAEL range. The exposure concentrations of TBT and copper were greater than their respective no-effects concentrations for Pacific staghorn sculpin in at least one area within the LDW but less than those associated with adverse effects. Thus, the potential for adverse effects is uncertain. For juvenile chinook salmon, exposure concentrations of cadmium were greater than concentration associated with adverse effects in any fish species but lower than concentrations associated with adverse effects in salmonids. Exposure concentrations of arsenic, copper, and vanadium in the diet of for juvenile chinook salmon were greater than their respective no-effect concentrations but less than concentrations associated with adverse effects.
- **Birds —** Risks were evaluated for three bird ROCs (spotted sandpiper, great blue heron, and osprey), which serve as representative surrogates for all other

bird species that may be exposed in the LDW. Estimated exposures of spotted sandpiper to six of the 12 COPCs (copper, chromium, lead, mercury, PCB TEQ, and vanadium) for spotted sandpiper were greater than the dietary doses associated with adverse effects on survival, growth, or reproduction in at least one area within the LDW (LOAEL-based HQs of up to 1.1, 1.8, 5.5, 1.0, 1.5, and 1.4, respectively). Therefore, there is a potential for adverse effects from these COPCs. Estimated doses to great blue heron of all four COPCs (chromium, lead, mercury, and total PCBs) were less than no-effects levels, indicating very low risk. For osprey, estimated doses of PCBs were greater than no-effect levels using a TEQ approach but less than those levels using a total PCBs approach; the latter risk estimate is less uncertain. Therefore, the potential for adverse effects from PCBs is uncertain for osprey. Estimated doses of the remaining three COPCs to osprey (chromium, lead, and mercury) were less than the doses associated with no-effects, indicating very low risk.

- **Mammals —** Risks were evaluated for two mammalian ROCs (river otters and harbor seals), which serve as representative surrogates for all other mammal species that may be exposed in the LDW. Estimated dietary doses of total PCBs were greater than doses associated with adverse effects for river otters, with a LOAEL-based HQ of 2.9. Estimated exposure of river otters to mercury was greater than a no-effects level but was less than adverse effects levels associated with survival, growth, or reproduction, indicating that the potential for effects is uncertain. Exposures of otter to the remaining three COPCs (arsenic, cobalt, and selenium) and exposures of harbor seals to both COPCs (mercury and total PCBs) were less than their respective no-effects levels, indicating very low risk.

Table A.8-1 provides a summary of COPCs for crabs, fish, or wildlife for which either the NOAEL-based HQ was > 1.0 or the LOAEL-based HQ was ≥ 1.0. The 44 COPCs for benthic invertebrates that exceeded SMS criteria, DMMP guidelines, or TRVs are listed in Table A.6-4. In summary, risk estimates for PCBs indicated a potential for adverse effects to the benthic invertebrate community, crabs, spotted sandpiper, and river otter. There is also a potential for adverse effects to English sole, Pacific staghorn sculpin, and osprey from PCBs, but risk estimates for these ROCs are more uncertain because exposures were greater than no-effect levels but less than levels associated with adverse effects. Other COPCs with exposures greater than or equal to levels associated with adverse effects for at least one fish or wildlife receptor were cadmium, chromium, copper, lead, mercury, and vanadium. Numerous additional chemicals pose a risk to the benthic invertebrate community as shown in Table A.6-4.

Table A.8-1. COPCs and ROCs with HQs ≥ 1.0

COPC	ROC	NOAEL-Based HQ	LOAEL-Based HQ
COPCs with LOAEL-Based HQs ≥ 1.0[a]			
Total PCBs	crabs	**10**	**1.0**
	river otter	**5.8**	**2.9**
	English sole	**4.9** – **25**	0.98 – **5.0**
	Pacific staghorn sculpin	**1.5** – **19**	0.30 – **3.8**
PCB TEQs	spotted sandpiper	**1.9 - 15**	0.18 – **1.5**
Cadmium	juvenile chinook salmon	**5.0**	**1.0**
	English sole	**6.1**	**1.2**
	Pacific staghorn sculpin	**3.0 – 5.2**	0.60 – **1.0**
Chromium	spotted sandpiper	**1.3 – 8.8**	0.26 – **1.8**
Copper	spotted sandpiper	0.62 – **1.5**	0.45 – **1.1**
Lead	spotted sandpiper	0.58 – **19**	0.17 – **5.5**
Mercury	spotted sandpiper	**1.1 – 5.3**	0.21 – **1.0**
Vanadium	English sole	**5.9**	**1.2**
	Pacific staghorn sculpin	**3.2 – 5.9**	0.65 **– 1.2**
	spotted sandpiper	**2.0 – 2.7**	**1.0 – 1.4**
COPCs with NOAEL-Based HQs ≥ 1.0 and LOAEL-Based HQs < 1.0[b]			
Total PCBs	spotted sandpiper	0.51 – **2.0**	0.18 – 0.71
PCB TEQs	osprey	**1.6**	0.16
	river otter	**4.5**	0.59
Arsenic	juvenile chinook salmon	**1.1**	0.73
	English sole	**1.2**	0.80
	crabs	**3.9**	na
Benzoic acid	English sole	**1.5**	na
	Pacific staghorn sculpin	**2.1**	na
Cadmium	Pacific staghorn sculpin	**3.0 – 4.9**	0.60 – 0.98
Chromium	juvenile chinook salmon	**2.1**	na
	English sole	**1.1**	na
Copper	juvenile chinook salmon	**1.9**	0.93
	English sole	**1.9**	0.93
	Pacific staghorn sculpin	0.9 – **1.5**	0.45 – 0.77
Mercury	river otter	**2.8**	0.57
TBT	Pacific staghorn sculpin	**1.6** – **2.9**	0.18 – 0.33
Vanadium	juvenile chinook salmon	**4.0**	0.79
Zinc	crabs	**2.5**	0.91

Note: HQs for fish are the highest HQs in cases where more than one approach was used.

[a] The LOAEL-based HQs for endrin were 1.2 and 3.1 for English sole and Pacific staghorn sculpin, respectively, based on risk calculations discussed in the uncertainty analysis. These calculations were discussed only in the uncertainty analysis because of analytical interferences from PCB Aroclors in the pesticide analyses, resulting in uncertainties in pesticide identification and a high bias in pesticide concentrations.

[b] The NOAEL-based HQs were > 1.0 for the following COPC/ROC pairs based on risk calculations discussed in the uncertainty analysis: 1) total DDTs and spotted sandpiper (2.6 to 4.3), 2) endrin and juvenile chinook salmon (3.6), 3) alpha-endosulfan and English sole (6.8) and Pacific staghorn sculpin (2.3), 4) beta-endosulfan and English sole (29) and Pacific staghorn sculpin (6.6), 5) endrin and juvenile chinook salmon (3.6), and 6) methoxychlor and crab s(3.6). These calculations were discussed in the uncertainty analysis because of analytical interferences from PCB Aroclors in the pesticide analyses, resulting in uncertainties in pesticide identification and a high bias in pesticide concentrations.

COPC – chemical of potential concern
HQ – hazard quotient
LOAEL – low-observed-adverse-effect level
na – not available
NOAEL – no-observed-adverse-effect level
PCB – polychlorinated biphenyl
ROC – receptor of concern
TBT – tributyltin
TEQ – toxic equivalent

Bold identifies NOAEL-based HQs greater than 1.0 or LOAEL-based HQs greater than or equal to 1.0.

Based on the risk estimates, uncertainties discussed in this ERA, natural background concentrations, and residual risks following planned early actions in the LDW, chemicals were identified as risk drivers for ecological receptors in accordance with EPA (1998) and MTCA (WAC 173-340-703) guidance. The risk drivers from both this ERA and the HHRA will be the focus of remedial analyses in the FS.

In consultation with EPA and Ecology, PCBs were identified as a risk driver for river otter because estimated exposure concentrations for river otter were greater than the LOAEL by a factor of 2.9 and uncertainties in the risk estimate were relatively low. In addition, 41 chemicals were selected as risk drivers for benthic invertebrates because concentrations of these 41 chemicals exceed Washington SMS in one or more locations.

Other chemicals that exceeded risk thresholds (LOAEL-based HQ ≥ 1.0) but were not selected as risk drivers may be addressed through focused evaluation in the FS, as part of the 5-year review, or included in the post-remedial monitoring program, as appropriate.

No quantitative ecological risk estimates were calculated for dioxins and furans within the LDW, and thus the level of ecological risk from dioxins and furans is unknown. Ecological risks associated with exposure to dioxins and furans within the LDW were not assessed for several reasons. Human health risks from these chemicals were assumed to be unacceptable,[100] and tissues were not analyzed for dioxins and furans. Tissues were not analyzed because of difficulties associated with assessing site-specific risks from dioxins and furans, the need for a large background dataset with which to compare site-specific data, and the paucity of background tissue data in the Puget Sound area. Remedial decisions to address dioxins/furan contamination in LDW sediment will be based on MTCA and CERCLA regulations, including those

[100] Dioxins and furans are a primary risk driver for human health.

specifically related to background (Ecology 2001; EPA 2002). Remedial decisions to address dioxin and furan contamination in sediment will be made by EPA and Ecology as part of the FS process.

This ERA is based on the baseline surface sediment dataset, which includes sediment data collected prior to early actions in the LDW. Since these data were collected, early actions in the LDW have been conducted at two locations (Duwamish/Diagonal and the Boeing Developmental Center south storm drain in the Norfolk area). Therefore, the risks discussed in this ERA may represent an overestimate of current risks for areas where remediation has already occurred.

A.9.0 References

Abiola FA. 1992. Ecotoxicity of organochloride insecticides: effects of endosulfan on bird reproduction and evaluation of its induction effect in partridge (*Perdix perdix* L.). Revue Med Vet 143(5):443-450.

Adachi A, Asai K, Koyama Y, Matsumoto Y, Okano T. 2000. Subacute vanadium toxicity in rats. J Health Sci 46(6):503-508.

Alexander G. 1977. Food of vertebrate predators on trout waters in north central lower Michigan. Mich Academ 10:181-195.

Allen JR, Barsotti DA, Carstens LA. 1980. Residual effects of polychlorinated biphenyls on adult nonhuman primates and their offspring. J Toxicol Environ Health 6:55-66.

Allison DT, Kollman BJ, Cope OB, Van Valin C. 1964. Some chronic effects of DDT on cutthroat trout. Research report 64. Bureau of Sport Fisheries and Wildlife, US Fish and Wildlife Service, Washington, DC.

Ambrose AM, Larson PS, Borzelleca JF, Hennigar Jr GR. 1976. Long term toxicologic assessment of nickel in rats and dogs. J Food Sci Technol 13(4):181-187.

Ames J, Phinney DE. 1977. Puget Sound summer-fall chinook methodology: escapement estimates and goals, run size forecasts, and in season run size updates. Technical report no. 29. Washington Department of Fisheries, Olympia, WA.

Anderson B. 2006. Personal communication (telephone conversation with Berit Bergquist, Windward Environmental LLC, regarding falcons in the LDW). Falcon Research Group, Bow, WA. June 9.

Anderson RM, May RM. 1979. Population biology of infectious diseases: part I. Nature (London) 280:361-367.

Andrews AK, Van Valin CC, Stebbings BE. 1966. Some effects of heptachlor on bluegills. Trans Am Fish Soc 95:297-309.

Angell CL, Miller BS, Wellings SR. 1975. Epizootiology of tumors in a population of juvenile English sole (*Parophrys vetulus*) from Puget Sound, Washington. J Fish Res Bd Can 32:1723-1732.

Anthony RG, Miles AK, Estes JA, Isaacs FB. 1999. Productivity, diets, and environmental contaminants in nesting bald eagles from the Aleutian Archipelago. Environ Toxicol Chem 18(9):2054-2062.

Anulacion BF, Lomax DP, Bill BD, Johnson LL, Collier TK. 1997. Assessment of antiestrogenic activity and CYP1A induction in English sole exposed to

environmental contaminants. Poster presented at SETAC 18th Annual Meeting, San Francisco, CA, November 16-20, 1997.

Argyle RL, Williams GC, Dupree HK. 1973. Endrin uptake and release by fingerling channel catfish (*Ictalurus punctatus*). J Fish Res Board Can 30:1743-1744.

Arkoosh MR, Casillas E, Clemons E, McCain B, Varanasi U. 1991. Suppression of immunological memory in juvenile shinook salmon (*Oncorhynchus tshawytscha*) from an urban estuary. Fish Shellfish Immunol 1:261-277.

Arkoosh MR, Casillas E, Huffman P, Clemons E, Evered J, Stein JE, Varanasi U. 1998. Increased susceptibility of juvenile chinook salmon from a contaminated estuary to *Vibrio anguillarum*. Trans Am Fish Soc 127:360-374.

Armstrong DA, Buchanan DV, Mallon MH, Caldwell RS, Millemann RE. 1976. Toxicity of the insecticide methoxychlor to the dungeness crab *Cancer magister*. Mar Biol 38:239-252.

Atar HH, Secer S. 2003. Width/length-weight relationships of the blue crab (*Callinectes sapidus* Rathbun 1896) population living in Beymelek Lagoon Lake. Turk J Vet Anim Sci 27:443-447.

ATSDR. 2002. Public health statement: creosote. Agency for Toxic Substances and Disease Registry, Atlanta, GA.

ATSDR. 2003. Toxicological profile for selenium. Agency for Toxic Substances and Disease Registry, Atlanta, GA.

ATSDR. 2004. Toxicological profile for cobalt. Agency for Toxic Substances and Disease Registry, Atlanta, GA.

Aulerich RJ, Ringer RK. 1977. Current status of PCB toxicity to mink, and effect on their reproduction. Arch Environ Contam Toxicol 6:279-292.

Aulerich RJ, Ringer RK, Iwamoto S. 1974. Effects of dietary mercury on mink. Arch Environ Contam Toxicol 2(1):43-51.

Aulerich RJ, Ringer RK, Bleavins MR, Napolitano A. 1982. Effects of supplemental dietary copper on growth, reproductive performance and kit survival of standard dark mink and the acute toxicity of copper to mink. J Anim Sci 55(2):337-343.

Aulerich RJ, Bursian SJ, Breslin WJ, Olson BA, Ringer RK. 1985. Toxicological manifestations of 2,4,5,2',4',5'-, 2,3,6,2',3',6'-, and 3,4,5,3',4',5'-hexachlorobiphenyl and Aroclor 1254 in mink. J Toxicol Environ Health 5:63-79.

Aulerich RJ, Ringer RK, Safronoff J. 1986. Assessment of primary vs secondary toxicity of Aroclor 1254 to mink. Arch Environ Contam Toxicol 15:393-399.

Azar A, Trochimowicz HJ, Maxfield ME. 1973. Review of lead studies in animals carried out at Haskell Laboratory-two-year feeding study and response to

hemorrhage study. In: Barth D, ed, Environmental Health Aspects of Lead, International Symposium, Amsterdam. Commission of European Communities, pp 199-210.

Baker RTM, Handy RD, Davies SJ, Snook JC. 1998. Chronic dietary exposure to copper affects growth, tissue lipid peroxidation, and metal composition of the grey mullet, *Chelon labrosus*. Mar Environ Res 45(4/5):357-365.

Baldisserotto B, Chowdhury MJ, Wood CM. 2005. Effects of dietary calcium and cadmium on cadmium accumulation, calcium and cadmium uptake from the water, and their interactions in juvenile rainbow trout. Aquat Toxicol 72:99-117.

Balevi T, Coskun B. 2004. Effects of dietary copper on production and egg cholesterol content in laying hens. Brit Poult Sci 45(4):530-534.

Bane G, Robinson M. 1970. Studies on the shiner perch, *Cymatogaster aggregata* Gibbons, in upper Newport Bay, California. Wasmann J Biol 28(2):259-268.

Barrick R, Becker S, Brown L, Beller H, Pastorok R. 1988. Sediment quality values refinement: Volume I. 1988 update and evaluation of Puget Sound AET. Prepared for Puget Sound Estuary Program (PSEP), US Environmental Protection Agency, Region 10. PTI Environmental Services, Inc., Bellevue, WA.

Barrows ME, Petrocelli SR, Macek KJ, Carroll JR. 1980. Bioconcentration and elimination of selected water pollutants by bluegill sunfish (*Lepomis macrochirus*). In: Haque R, ed, Dynamics, exposure and hazard assessment of toxic chemicals. Ann Arbor Science Publishers, Ann Arbor, MI.

Barsotti DA, Marlar RJ, Allen JR. 1976. Reproductive dysfunction in rhesus monkeys exposed to low levels of polychlorinated biphenyls (Aroclor 1248). Food Cosmet Toxicol 14:99-103.

Battelle, Pentec, Striplin, Shapiro, KCDNR. 2001. Reconnaissance assessment of the state of the nearshore ecosystem: Eastern shore of central Puget Sound, including Vashon and Maury Islands (WRIAs 8 and 9). Prepared for King County Department of Natural Resources, Seattle, WA. Battelle Marine Sciences Laboratory, Pentec Environmental, Striplin Environmental Associates, Shapiro Associates, Inc., King County Department of Natural Resources, Seattle, WA.

Battershill JM. 1994. Review of the safety assessment of polychlorinated biphenyls (PCBs) with particular reference to reproductive toxicity. Hum Exp Toxicol 13(9):581–597.

Beach RA, Geiger C, Jeffries SJ, Treacy SD, Troutman BL. 1985. Marine mammals and their interactions with fisheries of the Columbia River and adjacent waters, 1980-1982. Third annual report. Wildlife Management Division, Washington Department of Wildlife, Olympia, WA.

Behne D, Kyriakopoulos A, Gessner H, Walzog B, Meinhold H. 1992. Type I iodothyronine deiodinase activity after high selenium intake, and relations between selenium and iodine metabolism in rats. J Nutr 122(7):1542-1546.

Bent AC. 1929. Life histories of North American shore birds: order Limicolae. US National Museum. Bulletin, US Government Printing Office, Washington, DC. Republished, 1963, Dover Press, New York, NY.

Berlin WH, Hesselberg RJ, Mac MJ. 1981. Growth and mortality of fry of Lake Michigan lake trout during chronic exposure to PCBs and DDE. In: Chlorinated hydrocarbons as a factor in the reproduction and survival of lake trout (*Salvelinus namaycush*) in Lake Michigan. Technical Paper 105. US Fish and Wildlife Service, Washington, DC.

Berntssen MHG, Lundebye AK, Maage A. 1999a. Effects of elevated dietary copper concentrations on growth, feed utilization and nutritional status of Atlantic salmon (*Salmo salar* L.) fry. Aquaculture 174:167-181.

Berntssen MHG, Hylland K, Bonga SEW, Maage A. 1999b. Toxic levels of dietary copper in Atlantic salmon (*Salmo salar* L.) parr. Aquat Toxicol 46:87-99.

Beyer WN, Connor EE, Gerould S. 1994. Estimates of soil ingestion by wildlife. J Wildl Manage 58:375-382.

Bianchini A, Gilles R. 1996. Toxicity and accumulation of mercury in three species of crabs with different osmoregulatory capacities. Bull Environ Contam Toxicol 57:91-98.

Birge WJ, Black JA, Westerman AG, Hudson JE. 1979. The effects of mercury on reproduction of fish and amphibians. In: Nriagu JO, ed, The biogeochemistry of mercury in the environment. Elsevier/North-Holland Biomedical Press, New York, NY, pp 629-655.

Blazer VS, Fournie JW, Weeks-Perkins BA. 1997. Macrophage aggregates: biomarker for immune function in fishes? In: Dwyer FJ, Doane TR, Hinman ML, eds, Environmental toxicology and risk assessment: Modeling and risk assessment. Vol 6. STP 1317. American Society for Testing and Materials, Philadelphia, PA, pp 360-375.

Bleavins MR, Aulerich RJ. 1981. Feed consumption and food passage time in mink (*Mustela vison*) and European ferrets (*Mustela putorius furo*). Lab Anim Sci 31(3):268-269.

Bleavins MR, Aulerich RJ, Ringer RK. 1980. Polychlorinated biphenyls (Aroclors 1016 and 1242): Effects on survival and reproduction in mink and ferrets. Arch Environ Contam Toxicol 9:627-635.

Bleavins MR, Aulerich RJ, Ringer RK. 1984. Effects of chronic dietary hexachlorobenzene exposure on the reproductive performance and

survivability of mink and European ferrets. Arch Environ Contam Toxicol 13:357-365.

Blomberg G, Simenstad C, Hickey P. 1988. Changes in Duwamish River estuary habitat over the past 125 years. Proceedings of the First Annual Meeting on Puget Sound Research, Seattle, WA.

Boese BL, Winsor M, Lee H, Echols S, Pelletier J, Randall R. 1995. PCB congeners and hexachlorobenzene biota sediment accumulation factors for *Macoma nasuta* exposed to sediments with different total organic carbon contents. Environ Toxicol Chem 14(2):303-310.

Bolger M. 1993. Overview of PCB toxicology. Proceedings, US Environmental Protection Agency's National Technical Workshop, PCBs in Fish Tissue, May 10-11, 1993. EPA/823-R-93-003. Office of Water, US Environmental Protection Agency, Washington, DC, pp 1-37 to 1-53.

Bookhout CG, Costlow Jr JD, Monroe R. 1976. Effects of methoxychlor on larval development of mud-crab and blue crab. Wat Air Soil Pollut 5:349-365.

Borchert J, Karbe L, Westendorf J. 1997. Uptake and metabolism of benzo(a)pyrene absorbed to sediment by the freshwater invertebrate species *Chironomus riparius* and *Sphaerium corneum*. Bull Environ Contam Toxicol 58:158-165.

Borgmann U, Chau YK, Wong PTS, Brown M, Yaromich J. 1996. The relationship between tributyltin (TBT) accumulation and toxicity in *Hyalella azteca* for use in identifying TBT toxicity in the field. J Aquat Ecosys Health 5:199-206.

Borgmann U, Janssen C, Blust RJP, Brix KV, Dwyer RL, Erickson RJ, Hare L, Luoma SN, Paquin PR, Roberts CA, Wang W-X. 2005. Incorporation of dietborne metals exposure into regulatory frameworks. Chapter 5. In: Meyer JS, Adams WJ, Brix KV, Luoma SN, Mount DR, Stubblefield WA, Wood CM, eds, Toxicity of dietborne metals to aquatic organisms. Society of Environmental Toxicology and Chemistry (SETAC), Pensacola, FL, pp 153-190.

Boulva J, McLaren IA. 1979. Biology of the harbor seal, *Phoca vitulina*, in eastern Canada. Fish Res Board Can Bulletin 200:1-24.

Bowerman WW, Best DA, Kramer VJ. 1995. A review of factors affecting productivity of bald eagles in the Great Lakes region: implications for recovery. Environ Health Perspectives 103:51-59.

Braune BM, Norstrom RJ. 1989. Dynamics of organochlorine compounds in herring gulls: III. Tissue distribution and bioaccumulation in Lake Ontario gulls. Environ Toxicol Chem 8:957-968.

Bravo CF, Curtis LR, Meador JP, Loge FJ, Collier TK, Arkoosh MR. Undated draft. Increased susceptibility of rainbow trout *Oncorhynchus mykiss* to *Aeromonas*

salmonicida after dietary exposure to PAH. In press (2007). J Aquat Anim Health.

Brown RF, Mate BR. 1983. Abundance, movements, and feeding habits of harbor seals, *Phoca vitulina*, at Nearts and Tillamook Bays, Oregon. USNMFS Fish Bull 81:291-301.

Brunström B, Lund BE, Bergman A, Asplund L, Athanassiadis I, Athanasiadou M, Jensen S, Örberg J. 2001. Reproductive toxicity in mink (*Mustela vison*) chronically exposed to environmentally relevant polychlorinated biphenyl concentrations. Environ Toxicol Chem 20(10):2318-2327.

Buck J. 1999. Changes in productivity and environmental contaminants in bald eagles nesting along the lower Columbia River. Study ID 13420-1130-1F16. US Fish and Wildlife Service, Portland, OR.

Buehler DA. 2000. Bald eagle (*Haliaeetus leucocephalus*). In: Poole A, Gill F, eds, The birds of North America, no. 506. Academy of Natural Sciences, Philadelphia, PA, and American Ornithologists' Union, Washington, DC.

Burmaster DE, Thompson KM. 1997. Estimating exposure point concentrations for surface soils for use in deterministic and probabilistic risk assessments. Hum Ecolog Risk Assess 3(3):363-384.

Bury NR, Walker PA, Glover CN. 2003. Nutritive metal uptake in teleost fish. J Exp Biol 206:11-23.

Butler RB. 1997. The great blue heron. UBC Press, Vancouver, BC.

Butler RW. 1992. Great blue heron (*Ardea herodias*). In: Poole A, Stettenheim P, Gill F, eds, The birds of North America, no. 25. Academy of Natural Sciences, Philadelphia, PA, and American Ornithologists' Union, Washington, DC.

Butler RW. 1993. Time of breeding in relation to food availability of female great blue herons (*Ardea herodias*). Auk 110:693-701.

Byron WR, Bierbower GW, Brouwer JB, Hansen WH. 1967. Pathologic changes in rats and dogs from two-year feeding of sodium arsenite or sodium arsenate. Toxicol Appl Pharmacol 10:132-147.

Cain BW, Pafford EA. 1981. Effects of dietary nickel on survival and growth of mallard ducklings. Arch Environ Contam Toxicol 10:737-745.

Cain BW, Sileo L, Franson JC, Moore J. 1983. Effects of dietary cadmium on mallard ducklings. Environ Res 32:286-297.

Calabrese EJ, Baldwin LA. 1993. Performing ecological risk assessments. Lewis Publishers, Chelsea, MI.

Calder WA, Braun EJ. 1983. Scaling of osmotic regulation in mammals and birds. Am J Physiol 244:R601-R606.

Camardese MB, Hoffman DJ, LeCaptain LJ, Pendleton GW. 1990. Effects of arsenate on growth and physiology in mallard ducklings. Environ Toxicol Chem 9:785-795.

Campbell WW. 1996. Procedures to determine intertidal populations of *Protothaca staminea*, *Tapes phillippinarum*, and *Crassostrea gigas* in Hood Canal and Puget Sound, WA. MRD96-01. Point Whitney Shellfish Laboratory, Washington Department of Fish and Wildlife, Brinnon, WA.

Canli M, Furness RW. 1995. Mercury and cadmium uptake from seawater and from food by the Norway lobster *Nephrops norvegicus*. Environ Toxicol Chem 14(5):819-828.

Canning DJ, Herman SG, Shea GB. 1979. Terminal 107 environmental studies, wildlife study. Prepared for Port of Seattle. Oceanographic Institute of Washington and Northwest Environmental Consultants, Inc., Seattle, WA.

Casillas E, Misitano D, Johnson LL, Rhodes LD, Collier TK, Stein JE, McCain BB, Varanasi U. 1991a. Inducibility of spawning and reproductive success of female English sole (*Paraphrys vetulus*) from urban and nonurban areas of Puget Sound, Washington. Mar Environ Res 31:99-122.

Casillas E, Misitano D, Johnson LL, Rhodes LD, Collier TK, Stein JE, McCain BB, Varansasi U. 1991b. Inducibility of spawning and reproductive success of female English sole (*Parophrys vetulus*) from urban and nonurban areas of Puget Sound, Washington. Mar Environ Res 31:99-122.

Casillas E, Arkoosh MR, Clemons E, Hom T, Misitano D, Collier TK, Stein JE, Varanasi U. 1995a. Chemical contaminant exposure and physiological effects in outmigrant chinook salmon from selected urban estuaries of Puget Sound, Washington. In: Keefe M, ed, Salmon ecosystem restoration: myth and reality. Proceedings of the 1994 Northeast Pacific Chinook and Coho Salmon Workshop. Oregon Chapter, American Fisheries Society, Corvallis, OR, pp 86-102.

Casillas E, Arkoosh MR, Clemons E, Hom T, Misitano D, Collier TK, Stein JE, Varanasi U. 1995b. Chemical contaminant exposure and physiological effects in outmigrant Chinook salmon from urban estuaries of Puget Sound, Washington. Proceedings Puget Sound Research 95. Puget Sound Water Quality Authority, Olympia, WA, pp 657-665.

CCME. 2002. Canadian sediment quality guidelines for the protection of aquatic life: summary tables. Canadian Environmental Quality Guidelines, 1999, updated 2001, updated 2002 [online]. Canadian Council of Ministers of the Environment, Winnipeg, MB. Updated 2002. Available from: http://www.ccme.ca/assets/pdf/e7_002.pdf.

Chadwick B, Davidson B, Hampton T, Groves J, Guerrero J, Stang P. 1999. Offshore porewater and flux chamber sampling of San Diego Bay sediments at site 9,

Naval Air Station, North Island. Technical report 1799. US Navy SPAWAR Systems Center, San Diego, CA.

Chakravarty S, Lahiri P. 1986. Effect of lindane on eggshell characteristics and calcium level in the domestic duck. Toxicology 42:245-258.

Chakravarty S, Mandal A, Lahiri P. 1986. Effect of lindane on clutch size and level of egg yolk protein in domestic duck. Toxicology 39:93-103.

Chapman PM. 1996. A test of sediment effects concentrations: DDT and PCB in the Southern California Bight. Environ Toxicol Chem 15(7):1197-1198.

Chetty KY, Rau S, Drummond L, Desaiah D. 1979. Cobalt induced changes in immune response and adenosine triphosphatase activities in rats. J Environ Sci Health B14(5):525-544.

Choct M, Naylor AJ, Reinke N. 2004. Selenium supplementation affects brioler growth performance, meat yield and feather coverage. Brit Poult Sci 45(5):677-683.

Church PE, Vroblesky DA, Lyford FP, Willey RE. 2002. Guidance on the use of passive-vapor-diffusion samplers to detect volatile organic compounds in ground-water-discharge areas, and example applications in New England. Water-resources investigation report 02-4186. Massachusetts-Rhode Island District, US Geological Survey, Northborough, MA.

Clearwater SJ, Farag AM, Meyer JS. 2002. Bioavailability and toxicity of dietborne copper and zinc to fish. Comp Biochem Physiol Part C 132:269-313.

Cockell KA, Bettger WJ. 1993. Investigations of the gallbladder pathology associated with dietary exposure to disodium arsenate heptahydrate in juvenile rainbow trout. Toxicology 77:233-248.

Cockell KA, Hilton JW. 1988. Preliminary investigation on the comparative chronic toxicity of four dietary arsenicals to juvenile rainbow trout (*Salmo gairdneri* R.). Aquat Toxicol 12:73-82.

Cockell KA, Hilton JW, Bettger WJ. 1991. Chronic toxicity of dietary disodium arsenate heptahydrate to juvenile rainbow trout (*Oncorhynchus mykiss*). Arch Environ Contam Toxicol 21:518-527.

Cockell KA, Hilton JW, Bettger WJ. 1992. Hepatobiliary and hematological effects of dietary disodium arsenate heptahydrate in juvenile rainbow trout (*Oncorhynchus mykiss*). Comp Biochem Physiol 103C(3):453-458.

Coenen TMM, Brouwer A, Enninga IC, Koeman JH. 1992. Subchronic toxicity and reproduction effects if tri-n-butyltin oxide in Japanese quail. Arch Environ Contam Toxicol 23:457-463.

Cordell J. 2001. Personal communication (e-mail to Matt Luxon, Windward Environmental, Seattle, WA, regarding observations of juvenile chinook and

other wildlife in the Lower Duwamish River). Researcher, Department of Fisheries, University of Washington, Seattle, WA. July 9.

Cordell J, Toft J, Cooksey M, Gray A. 2006. Fish assemblages and patterns of chinook salmon abundance, diet, and growth at restored sites in the Duwamish River. Funded by the King Conservation District Salmon Recovery Funding Board, Seattle, WA.

Cordell JR, Tear LM, Simenstad CA, Hood WG. 1996. Duwamish river coastal America restoration and reference sites: Results from 1995 monitoring studies. Fish Research Institute, University of Washington, Seattle, WA.

Cordell JR, Tear LM, Jensen K, Luiting V. 1997. Duwamish river coastal America restoration and reference sites: Results from 1996 monitoring studies. Fisheries Research Institute, University of Washington, Seattle, WA.

Cordell JR, Tear LM, Jensen K, Higgins HH. 1999. Duwamish River coastal America restoration and reference sites: Results from 1997 monitoring studies. FRI-UW-9903. Fisheries Research Institute, University of Washington, Seattle, WA.

Cordell JR, Tear LM, Jensen K. 2001. Biological monitoring at Duwamish River coastal America restoration and reference sites: A seven-year retrospective. SAFS-UW-0108. Wetlands Ecosystem Team, School of Aquatic and Fisheries Sciences, University of Washington, Seattle, WA.

Coulter W, Adams CM, Decker DJ. 1984. River otter biology. In: New York's wildlife resources [online]. Department of Natural Resources, New York State College of Agriculture and Life Sciences, State University at Cornell University, Ithaca, NY. [Cited June 6, 2000]. Available from: http://www.nyotter.org/pages/biology.html.

Cropp T. 1985. Personal communication (as cited in Grette and Salo 1986). Washington Department of Game, Olympia, WA.

Crum JA, Bursian SJ, Aulerich RJ, Polin D, Braselton WE. 1993. The reproductive effects of dietary heptachlor in mink (*Mustela vison*). Arch Environ Contam Toxicol 24:156-164.

Custer TW, Heinz GH. 1980. Reproductive success and nest attentiveness of mallard ducks fed Aroclor 1254. Environ Pollut 21:313-318.

Davison KL, Sell JL. 1974. DDT thins shells of eggs from mallard ducks maintained on *ad libitum* or controlled-feeding regimens. Arch Environ Contam Toxicol 2(3):222-232.

Day DE. 1976. Homing behavior and population stratification in central Puget Sound English sole (*Parophrys vetulus*). J Fish Res Board Can 33:287-282.

DeCaprio AP, McMartin DN, O'Keefe PW, Rej R, Silkworth JB, Kaminsky LS. 1986. Subchronic oral toxicity of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in the guinea pig:

comparisons with a PCB-containing transformer fluid pyrolysate. Fund Appl Toxicol 6:454-463.

Delzell E, Giesy JP, Munro I, Doull J, Mackay D, Williams G. 1994. Interpretive review of the potential adverse effects of chlorinated organic chemicals on human health and the environment. Regulat Toxicol Pharmacol 20(1):S1-S1056.

DeWitt JB. 1956. Chronic toxicity to quail and pheasants of some chlorinated insecticides. Agric Food Chem 4(10):863-866.

Dexter RN, Anderson DE, Quinlan EA, Goldstein LS, Stickland RM, Pavlou SP, Clayton JR, Kocan RM, Landolt M. 1981. A summary of knowledge of Puget Sound related to chemical contaminants. NOAA technical memorandum OMPA-13. Office of Marine Pollution Assessment, National Oceanic and Atmospheric Administration, Boulder, CO.

Diaz GJ, Julian RJ, Squires EJ. 1994. Lesions in broiler chickens following experimental intoxication with cobalt. Avian Dis 38(2):308-316.

DiGiulio RT, Scanlon P. 1984. Sublethal effects of cadmium ingestion on mallard ducks. Arch Environ Contam Toxicol 13:765-771.

Dozier WA, III, Davis AJ, Freeman ME, Ward TL. 2003. Early growth and environmental implications of dietary zinc and copper concentrations and sources of broiler chicks. Brit Poult Sci 44(5):726-731.

Duby RT, Travis HF, Terrill CE. 1971. Uterotropic activity of DDT in rats and mink and its influence on reproduction in the rat. Toxicol Appl Pharmacol 18:348-355.

Duke TW, Lowe JL, Wilson AJ, Jr. 1970. A polychlorinated biphenyl (Aroclor 1254) in the water, sediment, and biota of Escambia Bay, Florida. Bull Environ Contam Toxicol 5:171-180.

Dunning JB, ed. 1993. CRC handbook of avian body masses. CRC Press, Boca Raton, FL.

Eastin WC, Jr, O'Shea TJ. 1981. Effects of dietary nickel on mallards. J Toxicol Environ Health 7:883-892.

Easton MDL, Luszniak K, Von der Geest E. 2002. Preliminary examination of contaminant loadings in farmed salmon, wild salmon and commercial salmon feed. Chemosphere 46:1053-1074.

Ecology. 1995. Sediment management standards. WAC 173-204. Washington State Department of Ecology, Toxics Cleanup Program, Sediment Management Unit, Olympia, WA.

Ecology. 2000. WRIA 9 report: Historical and current salmonid populations, life histories, and habitat conditions. Draft report. King County Department of Natural Resources, Seattle, WA.

Ecology. 2001. Model Toxics Control Act Cleanup Regulation, Chapter 173-340 WAC. Publication No. 94-06. Toxics Cleanup Program, Washington State Department of Ecology, Olympia, WA.

Ecology. 2003a. Norfolk combined sewer overflow (Duwamish River) sediment cap recontamination. Phase I investigation. Publ. no. 03-03-004. Environmental Assessment Program, Washington Department of Ecology, Olympia, WA.

Ecology. 2003b. Sediment sampling and analysis plan appendix: guidance on the development of sediment sampling and analysis plans meeting the requirements of the sediment management standards (chapter 173-204 WAC). Ecology Publication No. 03-09-043. PTI Environmental Services, Bellevue, WA.

Edens FW, Garlich JD. 1983. Lead-induced egg production decrease in leghorn and Japanese quail hens. Poult Sci 62:1757-1763.

Edens FW, Benton E, Bursian SJ, Morgan GW. 1976. Effect of dietary lead on reproductive performance in Japanese quail, *Coturnix coturnix japonica*. Toxicol Appl Pharmacol 38:307-314.

Eertman RHM, Groenink CL, Sandee B, Hummel H, Small AC. 1995. Response of the blue mussel *Mytilus edulis* L. following exposure to PAHs or contaminated sediment. Mar Environ Res 39:169-173.

Eisler R. 1985. Selenium hazards to fish, wildlife, and invertebrates: a synoptic review. US Fish and Wildlife Service Biological Report 85(1.5). Patuxent Wildlife Research Center, US Fish and Wildlife Service, Laurel, MD.

Eisler R. 1986a. Chromium hazards to fish, wildlife, and invertebrates: A synoptic review. US Fish and Wildlife Service Biological Report 85(1.6). Patuxent Wildlife Research Center, US Fish and Wildlife Service, Laurel, MD.

Eisler R. 1986b. Polychlorinated biphenyl hazards to fish, wildlife, and invertebrates: a synoptic review. US Fish and Wildlife Service Biological Report 85(1.7). Patuxent Wildlife Research Center, US Fish and Wildlife Service, Laurel, MD.

Eisler R. 1987. Mercury hazards to fish, wildlife, and invertebrates: A synoptic review. US Fish and Wildlife Service Biological Report 85(1.10). Patuxent Wildlife Research Center, US Fish and Wildlife Service, Laurel, MD.

Eisler R. 1988a. Arsenic hazards to fish, wildlife, and invertebrates: a synoptic review. US Fish and Wildlife Service Biological Report 85(1.12). Patuxent Wildlife Research Center, US Fish and Wildlife Service, Laurel, MD.

Eisler R. 1988b. Lead hazards to fish, wildlife, and invertebrates: a synoptic review. US Fish and Wildlife Service Biological Report 85(1.14). Patuxent Wildlife Research Center, US Fish and Wildlife Service, Laurel, MD.

Eisler R. 1993. Zinc hazards to fish, wildlife, and invertebrates: a synoptic review. US Fish and Wildlife Service Biological Report Number 10. Patuxent Wildlife Research Center, US Fish and Wildlife Service, Laurel, MD.

Eisler R. 1997. Copper hazards to fish, wildlife and invertebrates: a synoptic review. Patuxent Wildlife Research Center, US Fish and Wildlife Service, Laurel, MD.

Ema M, Harazono A, Hirose A, Kamata E. 2003. Protective effects of progesterone on implantation failure induced by dibutyltin dichloride in rats. Toxicol Let 143:233-238.

EPA. 1988. Guidance for conducting remedial investigations and feasibility studies under CERCLA. Interim final. EPA/540/G-89/004, OSWER Directive 9355.3. Office of Emergency and Remedial Response, US Environmental Protection Agency, Washington, DC.

EPA. 1989. Risk management guidance for Superfund, volume 2: Environmental evaluation manual. EPA 540/1-89/001. US Environmental Protection Agency, Washington, DC.

EPA. 1990. Superfund guidance on remedial actions for Superfund sites with PCB contamination. OSWER Directive No. 9355.4-01. EPA/540/G-90/007. US Environmental Protection Agency, Washington, DC.

EPA. 1992. Framework for ecological risk assessment. EPA/630/R-92/001. Risk Assessment Forum, US Environmental Protection Agency, Washington, DC.

EPA. 1993. Wildlife exposure factors handbook. EPA/600/R-93/187a. Office of Research and Development, US Environmental Protection Agency, Washington, DC.

EPA. 1997a. Ecological risk assessment guidance for Superfund: Process for designing and conducting ecological risk assessments. EPA/540/R-97/006. Interim final. Environmental Response Team, US Environmental Protection Agency, Edison, NJ.

EPA. 1997b. EPA Region 10 supplemental ecological risk assessment guidance for Superfund. EPA/910/R-97/005. Region 10 Office of Environmental Assessment Risk Evaluation Unit, US Environmental Protection Agency, Seattle, WA.

EPA. 1998. Guidelines for ecological risk assessment. EPA/630/R-95/002 F. Risk Assessment Forum, US Environmental Protection Agency, Washington, DC.

EPA. 1999a. Development of a tissue trigger level for bioaccumulated tributyltin in marine benthic organisms: West Waterway Harbor Island Superfund Site, Seattle, WA. US Environmental Protection Agency Region 10, Seattle, WA.

EPA. 1999b. USEPA contract laboratory program national functional guidelines for organic data review. EPA-540/R-99/008. Office of Emergency and Remedial Response, US Environmental Protection Agency, Washington, DC.

EPA. 2000a. Bioaccumulation testing and interpretation for the purpose of sediment quality assessment: status and needs. EPA-823-R-00-001. Bioaccumulation Analysis Workgroup, US Environmental Protection Agency, Washington, DC.

EPA. 2000b. Evaluation of dermal contact and inhalation exposure pathways for the purpose of setting Eco-SSLs. Draft ecological soil screening level guidance, Exhibit 1-3. US Environmental Protection Agency, Washington, DC.

EPA. 2000c. Guidance for assessing chemical contaminant data for use in fish advisories. Volume 1: Fish sampling and analysis. Third ed. EPA 823-B-00-007. US Environmental Protection Agency, Washington, DC.

EPA. 2000d. Hudson River PCBs reassessment. Revised baseline ecological risk assessment. US Environmental Protection Agency Region 2, New York, NY.

EPA. 2002. Role of background in the CERCLA cleanup program. OSWER 9285.6-07P. US Environmental Protection Agency, Office of Solid Waste and Emergency Response, Office of Emergency and Remedial Response, Washington, DC.

EPA. 2003a. Analyses of laboratory and field studies of reproductive toxicity in birds exposed to dioxin-like compounds for use in ecological risk assessment. EPA/600/R-03/114F. National Center for Environmental Assessment, US Environmental Protection Agency, Cincinnati, OH.

EPA. 2003b. Framework for application of the toxicity equivalence methodology for polychlorinated dioxins, furans and biphenyls in ecological risk assessment. EPA/630/P-03/002A. External review draft. Risk Assessment Forum, US Environmental Protection Agency, Washington, DC.

EPA. 2004a. ProUCL Version 3.0. Statistical software to compute Upper Confidence Limits of the unknown population mean [online]. Technical Support Center for Monitoring and Site Characterization, US Environmental Protection Agency. Updated 5/6/2004. Available from: http://www.epa.gov/nerlesd1/tsc/download.htm.

EPA. 2004b. USEPA contract laboratory program national functional guidelines for inorganic data review. EPA 540-R-04-004. Office of Emergency and Remedial Response, US Environmental Protection Agency, Washington, DC.

EPA. 2005a. EPA and Ecology comments on memorandum on additional shiner surfperch analyses; Lower Duwamish Waterway; Seattle, Washington. Letter to M. Johns, Windward Environmental, from Allison Hiltner, dated 7/22/05. Superfund Site Manager, US Environmental Protection Agency Region 10, Seattle, WA.

EPA. 2005b. Rhône-Poulenc (Rhodia) sediment & porewater investigation Aug/Sept 2004. Data report: volatile organic compounds, metals, and pesticides/PCBs detected in sediment and porewater. Draft. US Environmental Protection Agency Region 10, Seattle, WA.

EPA. 2006. Integrated Risk Information System (IRIS) database [online]. US Environmental Protection Agency, Washington, DC. [Cited 1/2006]. Available from: http://www.epa.gov/iris/.

Erickson RJ, Highland TL, Hocket JR, Leonard EN, Mattson VR, Mount DR. 2003. Effects of dietary copper, zinc, lead, cadmium, and arsenic on growth and survival of juvenile fish using live food organisms. Platform presentation at SETAC 24th annual meeting, Austin TX, 9-13 November 2003. Manuscript in prep.

ESG. 1999. Waterway sediment operable unit, Harbor Island Superfund site. Assessing human health risks from the consumption of seafood: human health risk assessment report. Prepared for Port of Seattle, Todd Shipyards, and Lockheed-Martin for submittal to US Environmental Protection Agency, Region 10, Seattle, WA. Environmental Solutions Group, Inc., Seattle, WA.

Evans ML. 1980. Copper accumulation in the crayfish (*Orconectes rusticus*). Bull Environ Contam Toxicol 24:916-920.

Everitt RD, Gearin P, Skidmore JS. 1981. Prey items of harbor seals and California sea lions in Puget Sound, Washington. Murrelet 62: 83-8.

Exponent. 1998. Duwamish Waterway phase I site characterization report. Prepared for The Boeing Company. Exponent, Bellevue, WA.

Fabacher DL. 1976. Toxicity of endrin and an endrin-methyl parathion fomulation to largemouth bass fingerlings. Bull Environ Contam Toxicol 16(3):376-378.

Fernie K, Smiths J, Bortolotti G. 2003a. Developmental toxicity of in ovo exposure to polychlorinated biphenyls: I. Immediate and subsequent effects on first-generation nestling American kestrels (*Falco sparverius*). Environ Toxicol Chem 22(3):554-560.

Fernie K, Bortolotti G, Driuillard K, Smiths J, Marchant T. 2003b. Developmental toxicity of in ovo exposure to polychlorinated biphenyls: II. Effects of maternal or paternal exposure on second-generation nestling American kestrels (*Falco sparverius*). Environ Toxicol Chem 22(11):2688-2694.

Fernie K, Bortolotti G, Smits J. 2003c. Reproductive abnormalities, teratogenicity, and developmental problems in American kestrels (*Falco sparverius*) exposed to polychlorinated biphenyls. J Toxicol Env Health Part A 66:2089-2103.

Fernie KJ, Bortolotti GR, Smits JE, Wilson J, Drouillard KG, Bird DM. 2000. Changes in egg composition of American kestrels exposed to dietary polychlorinated biphenyls. J Toxicol Environ Health Part A, 60:291-303.

Fernie KJ, Smits JE, Bortolotti GR, Bird DM. 2001. *In ovo* exposure to polychlorinated biphenyls: reproductive effects on second-generation American kestrels. Arch Environ Contam Toxicol 40:544-550.

Feroz M, Khan MAQ. 1979. Fate of ^{14}C-cis-chlordane in goldfish, *Carassius auratus* (L.). Bull Environ Contam Toxicol 23:64-69.

Finley MT, Dieter MP, Locke LN. 1976. Sublethal effects of chronic lead ingestion in mallard ducks. J Toxicol Environ Health 1:929-937.

Fish AG. 2002. Great egret (*Casmerodius albus*) [online]. Arizona Game & Fish Department, Phoenix, AZ. [Cited 12/10/02]. Available from: http://www.gf.state.az.us/frames/fishwild/ngame_f.htm.

Fisher JP, Spitsbergen JM, Bush B, Jahan-Parwar B. 1994. Effect of embryonic PCB exposure on hatching success, survival, growth and developmental behavior in landlocked Atlantic salmon, *Salmo salar*. In: Gorsuch JW, Dwyer FJ, Ingersoll CG, La Point TW, eds, Environmental toxicology and risk assessment. Vol 2. ASTM STP 1216. American Society for Testing and Materials, Philadelphia, PA, pp 298-314.

Floyd Snider McCarthy. 2004. Boeing Plant 2, Seattle/Tukwila, Washington: Phase 1 transformer PCB investigation report. Floyd Snider McCarthy, Inc., Seattle, WA.

Formigli L, Scelsi R, Poggi P, Gregotti C, Di Nucci A, Sabbioni E, Gottardi L, Manzo L. 1986. Thallium-induced testicular toxicity in the rat. Environ Res 40:531-539.

Forrester CR. 1969. Life history on some ground fish species. Fish Res Bd Can Technical Report 105:1-17.

Franklin NM, Glover CN, Nicol JA, Wood CM. 2005. Calcium/cadmium interactions at uptake surfaces in rainbow trout: waterborne versus dietary routes of exposure. Environ Toxicol Chem 24(11):2954-2964.

Freeland JH, Cousins RJ. 1973. Effect of dietary cadmium on anemia, iron absorption, and cadmium binding protein in the chick. Nutr Rep Internat 8(5):337-347.

Freeman HC, Idler DR. 1975. The effect of PCB on steroidogenesis and reproduction in the brook trout (*Salvelinus fontinalis*). Can J Biochem 53:666-670.

Fresh KL, Rabin D, Simenstad CA, Salo EO, Garrison K, Matheson L. 1979. Fish ecology studies in the Nisqually Reach area of southern Puget Sound, Washington. FRI-UW-7904. Prepared for Weyerhauser Company. Fisheries Research Institute, University of Washington, Seattle, WA.

Fuller GB, Hobson WC. 1986. Effect of PCBs on reproduction in mammals. In: Waid JS, ed, PCBs and the environment. Vol II. CRC Press, Inc, Boca Raton, FL.

Galvez F, Wood CM. 1999. Physiological effects of dietary silver sulfide exposure in rainbow trout. Environ Toxicol Chem 18(1):84-88.

Gasaway WC, Buss IO. 1972. Zinc toxicity in the mallard duck. J Wildl Manage 36:1107-1117.

Gatlin DM, III, Wilson RP. 1986. Dietary copper requirement of fingerling channel catfish. Aquacult 54:277-285.

Gee JM, Craig-Veit CB, Millam JR. 2004. Posthatch methoxychlor exposure adversely affects reproduction of adult zebra finches, *Taeniopygia guttata*. Bull Environ Contam Toxicol 73:607-612.

Gibbs PE, Bryan GW. 1996. Reproductive failure in the gastropod *Nucella lapillus* associated with imposex caused by tributyltin pollution: a review. In: Champ MA, Seligman PF, eds, Organotin: environmental fate and effects. Chapman & Hill, London, pp 259-280.

Gibbs PE, Pascoe PL, Burt GR. 1988. Sex change in the female dog-whelk, *Nucella lapidis*, induced by tributyltin from antifouling paints. J Mar Biol Ass UK 68:715-731.

Giesy J. 2006. Personal communication (e-mail to Matt Luxon, Windward Environmental, regarding 2002 Dioxin LOEL TRV study). University of Saskatchewan, Saskatoon, Saskatchewan, Canada. May 19, 2006.

Giesy JP, Jones PD, Kannan K, Newsted JL, Tillitt DE, Williams LL. 2002. Effects of chronic dietary exposure to environmentally relevant concentrations if 2,3,7,8-tetrachlorodibenzo-p-dioxin on survival, growth, reproduction and biochemical responses of female rainbow trout (*Oncorhynchus mykiss*). Aquat Toxicol 59:35-53.

Gilbert FF, Nancekivell EG. 1982. Food habits of mink (*Mustela vison*) and otter (*Lutra canadensis*) in northeastern Alberta. Can J Zool 60:1282-1288.

Goettl JP, Davies PH, Sinley JR. 1976. Water pollution studies. Colorado Fisheries Research Review 1972-1975. CO Div Wildlife Rev 8:68-75.

Golub MS, Donald JM, Reyes JA. 1991. Reproductive toxicity of commercial PCB mixtures: LOAELs and NOAELs from animal studies. Environ Health Perspect 94:245-253.

Good EE, Ware GW. 1969. Effects of insecticides on reproduction in the laboratory mouse. IV. Endrin and dieldrin. Toxicol Appl Pharmacol 14:201-203.

Gordon CD. 1965. Aspects of the life-history of *Cymatogaster aggregata* Gibbons. MS thesis. University of British Columbia, Victoria, BC.

Greer KR. 1955. Yearly food habits of the river otter in the Thompson Lakes region, Northwestern Montana, as indicated by scat analyses. Am Midl Nat 54:299-313.

Grette GB, Salo EO. 1986. The status of anadromous fishes of the Green/Duwamish River system. Prepared for US Army Corps of Engineers, Seattle District. Evans-Hamilton, Inc., Seattle, WA.

Gries TH, Waldow KH. 1996. Progress re-evaluating Puget Sound apparent effects thresholds (AETs). Volume I: 1994 amphipod and echinoderm larval AETs. Draft report. Prepared for Puget Sound Dredged Disposal Analysis (PSDDA) agencies. Washington Department of Ecology, Olympia, WA.

Hack R, Ebert E, Leist KH. 1995. Chronic toxicity and carcinogenicity studies with the insecticide endosulfan in rats and mice. Food Chem Toxicol 33(11):941-950.

Halverson AW, Palmer IS, Guss PL. 1966. Toxicity of selenium to post-weanling rats. Toxicol Appl Pharmacol 9:477-484.

Handy RD. 1992. The assessment of episodic metal pollution. II. The effects of cadmium and copper enriched diets on tissue contaminant analysis in rainbow trout (*Oncorhynchus mykiss*). Arch Environ Contam Toxicol 22:82-87.

Handy RD. 1993. The effect of acute exposure to dietary Cd and Cu on organ toxicant concentrations in rainbow trout, *Oncorhynchus mykiss*. Aquat Toxicol 27:1-14.

Hansen D, Schimmel SC, Forester J. 1974a. Aroclor 1254 in eggs of sheepshead minnows: effect on fertilization success and survival of embryos and fry. Proceedings of 27th Annual Conference, Southeastern Association of Game and Fish Commissioners, Hot Springs, AR, pp 420-426.

Hansen DJ, Parrish PR, Lowe JL, Wilson AJ, Jr, Wilson PD. 1971. Chronic toxicity, uptake, and retention of Aroclor 1254 in two estuarine fishes. Bull Environ Contam Toxicol 6:113-119.

Hansen DJ, Parrish PR, Forester J. 1974b. Aroclor 1016: Toxicity to and uptake by estuarine animals. Environ Res 7:363-373.

Hansen DJ, Schimmel SC, Forester J. 1975. Effects of Aroclor 1016 on embryos, fry, juveniles, and adults of sheepshead minnows (*Cyprinodon variegates*). Trans Am Fish Soc 104:584-588.

Hansen DJ, Schimmel SC, Forester J. 1977. Endrin: effects on the entire life cycle of a saltwater fish, *Cyprinodon variegatus*. J Toxicol Environ Health 3:721-733.

Harazono A, Ema M. 2003. Suppression of decidual cell resonse induced by dibutyltin dichloride in pseudopregnant rats: as a cause of early embryonic loss. Reprod Toxicol 17:393-399.

Harper-Owes. 1983. Water quality assessment of the Duwamish estuary, Washington. Prepared for Municipality of Metropolitan Seattle. Harper-Owes Company, Seattle, WA.

Hart Crowser. 1993. Results of sampling and analysis, sediment monitoring plan, Duwamish Shipyard, Inc., Seattle, Washington. Hart Crowser, Inc., Seattle, WA.

Hart Crowser. 1998. Dredge material characterization, Hurlen Construction Company and Boyer Alaska Barge Lines Berthing Areas, Duwamish Waterway, Seattle,

Washington. Prepared for Peratrovich, Nottingham & Drage. Hart Crowser, Inc., Seattle, WA.

Hart DR, Heddle JA. 1991. Micronucleus assays in peripheral blood of rainbow trout: Timing of response and chemical mutagen sensitivity. Can Tech Rep Fish Aquat Sci 1774(2):993-1010.

Hartman. 1992. Lone Star Northwest west terminal, Duwamish River PSDDA sampling and analysis results. Hartman Associates, Inc., Seattle, WA.

Hartman. 1995. Lone Star Northwest and James Hardie Gypsum, Kaiser Dock Upgrade, Duwamish Waterway PSDDA sampling and analysis results. Hartman Associates, Inc., Seattle, WA.

Hartman FA. 1961. Locomotor mechanisms in birds. Smithsonian Misc Coll 143, Washington, DC.

Haseltine SD, Prouty RM. 1980. Aroclor 1242 and reproductive success of adult mallards (*Anas platyrhynchos*). Environ Res 23:29-34.

Haseltine SD, Sileo L, Hoffman DJ, Mulhern BD. unpublished. Effects of chromium on reproduction and growth in black ducks. As cited in Sample BE, Opresko DM, Suter GW. 1996. Toxicological benchmarks for wildlife. 1996 revision. ES/ERM-86/R3. Office of Environmental Management, US Department of Energy, Washington, DC.

Hatakeyama S, Yasuno M. 1982. Accumulation and effects of cadmium on guppy (*Poecilia reticulata*) fed cadmium-dosed Cladocera (*Moina macrocopa*). Bull Environ Contam Toxicol 29:159-166.

Hatakeyama S, Yasuno M. 1987. Chronic effects of Cd on the reproduction of the guppy (*Poecilia reticulata*) through Cd-accumulated midge larvae (*Chironomus yoshimatsui*). Ecotoxicol Environ Saf 14:191-207.

Hattula ML, Karlog O. 1972. Toxicity of polychlorinated biphenyls (PCBs) to goldfish. Acta Pharmacol Toxicol 31:238-240.

Healy MC. 1991. Life history of chinook salmon (*Oncorhynchus tshawytscha*). In: Groot C, Margolis L, eds, Pacific salmon life histories. UBC Press, Vancouver, BC, pp 311-394.

Heath RG, Spann JW, Kreitzer JF. 1969. Marked DDE impairment of mallard reproduction in controlled studies. Nature 224:47-48.

Heath RG, Spann JW, Hill EF, Kreitzer JF. 1972. Comparative dietary toxicities of pesticides to birds. Wildlife no. 152. Patuxent Wildlife Research Center, Bureau of Sport Fisheries and Wildlife, Laurel, MD.

Heinz GH. 1980. Eggshell thickness in mallards fed methylmercury. Bull Environ Contam Toxicol 25:498-502.

Heinz GH, Hoffman DJ, Krynitsky AJ, Weller DMG. 1987. Reproduction in mallards fed selenium. Environ Toxicol Chem 6:423-433.

Heinz GH, Hoffman DJ, Gold LG. 1988. Toxicity of organic and inorganic selenium to mallard ducklings. Arch Environ Contam Toxicol 17:561-568.

Heinz GH, Hoffman DJ, Gold LG. 1989. Impaired reproduction of mallards fed an organic form of selenium. J Wildl Manage 53(2):418-428.

Hendricks JD, Scott WT, Putnam TP, Sinnhuber RO. 1981. Enhancement of aflatoxin B1 hepatocarcinogenesis in rainbow trout (*Salmo gairdneri*) embryos by prior exposure of gravid females to dietary Aroclor 1254. In: Branson DR, Dickson KL, eds, Aquat Toxicol and Hazard Assessment. Fourth Conference. ASTM STP 737. American Society for Testing and Materials, Philadelphia, PA, pp 203-214.

Hendricks JD, Meyers TR, Shelton DW, Casteel JL, Bailey GS. 1985. Hepatocarcinogenicity of benzo[a]pyrene to rainbow trout by dietary exposure and intraperitoneal injection. J Nat Cancer Inst 74:839-851.

Henny CJ. 2005. Personal communication (telephone conversation with Erika Hoffman, US Environmental Protection Agency Region 10, regarding fish consumption by osprey). Research Zoologist, USGS Forest and Rangeland Ecosystem Science Center, Corvallis, OR. October 20, 2005.

Henny CJ. 2006. Personal communication (e-mail to Bruce Duncan, EPA Region 10, regarding consumption of salmonids by osprey). Research Zoologist, USGS Forest and Rangeland Ecosystem Science Center, Corvallis, OR. May 8, 2006.

Henny CJ, Kaiser JL, Grove RA, Bentley VR, Elliott JE. 2003. Biomagnification factors (fish to osprey eggs from Willamette River, Oregon, USA) for PCDDs, PCDFs, PCBs and OC pesticides. Environ Monit Assess 84:275-315.

Heron Habitat Helpers. 2005. Newsletter July – August 2005 [online]. Heron Habitat Helpers, Seattle, WA. [Cited June 1, 2006]. Available from: http://heronhelpers.org/newsletter.html.

Heron Working Group. 2006a. About herons: habitat, characteristics, human activity [online]. Heron Working Group. [Cited 12/6/2006]. Available from: http://www.heronworkinggroup.org/aboutherons.html.

Heron Working Group. 2006b. The Heron Network: network sites [online]. Heron Working Group. [Cited June 1, 2006]. Available from: http://www.heronworkinggroup.org/networksites.htm.

Herrera. 1997. Seaboard Lumber Site. Phase 2 site investigation. Draft. Prepared for Seattle Department of Parks and Recreation. Herrera Environmental Consultants, Inc., Seattle, WA.

Hext PM, Pinto PJ, Rimmel BA. 1999. Subchronic feeding study of antimony trioxide in rats. J Appl Toxicol 19:205-209.

Hill EF, Soares JH, Jr. 1987. Oral and intramuscular toxicity of inorganic and organic mercury chloride to growing quail. J Toxicol Env Health 20:105-116.

Hill EF, Heath RG, Spann JW, Williams JD. 1975. Lethal dietary toxicities of environmental pollutants to birds. US Fish and Wildlife Service, Laurel, MD.

Hill EF, Heath RG, Williams JD. 1976. Effect of dieldrin and Aroclor 1242 on Japanese quail eggshell thickness. Bull Environ Contam Toxicol 16(4):445-453.

Hilton JW, Bettger WJ. 1988. Dietary vanadium toxicity in juvenile rainbow trout: a preliminary study. Aquat Toxicol 12:63-71.

Hochstein JR, Render JA, Bursian SJ, Aulerich RJ. 2001. Chronic toxicity of dietary 2,3,7,8-tetrachlorodibenzo-*p*-dioxin to mink. Vet Hum Toxicol 43(3):134-139.

Hoffman DJ, Rice CP, Kubiak TJ. 1996. PCBs and dioxins in birds. In: Beyer WN, Heinz GH, Redmon-Norwood AW, eds, Environmental contaminants in wildlife. Lewis Publishers, Boca Raton, FL, pp 165-207.

Hoffman RD. 1978. The diets of herons and egrets in southwestern Lake Erie. In: Sprunt A, Oge J, Winckler S, eds, Wading birds. Research report No. 7. National Audubon Society, New York, NY, pp 365-369.

Holland GA. 1954. A preliminary study of the populations of English sole in Carr Inlet and other locations in Puget Sound. MS thesis. University of Washington, Seattle, WA.

Hoover A. 1988. Harbor seal, *Phoca vitulina*. In: Lentfor JW, ed, Selected marine mammals of Alaska: Species accounts with research and management recommendations. Marine Mammal Commission, Washington, DC, pp 125-157.

Hopkins CL, Solly SRB, Ritchie AR. 1969. DDT in trout and its possible effect on reproductive potential. NZ J Mar Freshw Res 3:220-229.

Hornshaw TC, Safronoff J, Ringer RK, Aulerich RJ. 1986. LC50 test results in polychlorinated biphenyl-fed mink: age, season, and diet comparisons. Arch Environ Contam Toxicol 15:717-723.

Hotchkiss D. 2006. Personal communication (telephone call to Matt Luxon, Windward Environmental, regarding zoning of Kellogg Island as a wildlife refuge in the City of Seattle city shoreline zoning code). Port of Seattle, Seattle, WA. August 18, 2006.

Hough JL, Baird MB, Sfeir GT, Pacini CS, Darrow D, Wheelock C. 1993. Benzo(a)pyrene enhances atherosclerosis in white carneau and show racer pigeons. Arterioscler Thromb 13:1721-1727.

Hudson RH, Tucker RK, Haegele MA. 1984. Handbook of toxicity of pesticides to wildlife. Resource publication 153. 2nd ed. US Fish and Wildlife Service, Washington, DC.

Hugla JL, Thome JP. 1999. Effects of polychlorinated biphenyls on liver ultrastructure, hepatic monooxygenases, and reproductive success in the barbel. Ecotoxicol Environ Saf 42:265-273.

Hyland JL, Van Dolah RF, Snoots TR. 1999. Predicting stress in benthic communities of southeastern U.S. estuaries in relation to chemical contamination of sediments. Environ Toxicol Chem 18(11):2557-2564.

Ingersoll CG, Wang N, Hayward JMR, Jones JR, Jones SB, Ireland DS. 2005. A field assessment of long-term laboratory sediment toxicity tests with the amphipod *Hyalella azteca*. Environ Toxicol Chem 24(11):2853-2870.

Integral. 2004. Lower Duwamish Waterway slip 4 early action area: cruise and data report. Prepared for City of Seattle and King County. Integral Consulting, Inc., Mercer Island, WA.

Ishida M, Suyama K, Adachi S, Hoshino T. 1982. Distribution of orally administered di-ethylhexyl phthalate in laying hens. Poult Sci 61:262-267.

Ivankovic S, Preussman R. 1975. Absence of toxic and carcinogenic effects after administration of high doses of chromic oxide pigment in subacute and long-term feeding experiments in rats. Food Cosmet Toxicol 13:347-351.

Jarvinen AW, Ankley GT. 1999. Linkage of effects to tissue residues: Development of a comprehensive database for aquatic organisms exposed to inorganic and organic chemicals. SEATAC Press, Pensacola, FL.

Jarvinen AW, Tyo RM. 1978. Toxicity to fathead minnows of endrin in food and water. Arch Environ Contam Toxicol 7:409-421.

Jeanes ED, Hilgert PJ. 2000. Juvenile salmonid use of lateral stream habitats in the Middle Green River, Washington. Report prepared for US Army Corps of Engineers, Seattle District and City of Tacoma Public Utilities. R2 Resource Consultants, Redmond, WA.

Jeffries SJ. 2001. Personal communication (telephone conversation with Berit Bergquist, Windward Environmental LLC, Seattle, WA). Research Scientist, Washington Department of Fish and Wildlife, Olympia, WA. October 29.

Jensen S, Kihlstrom JE, Olsson M, Lundberg C, Orberg J. 1977. Effects of PCB and DDT on mink (*Mustela vison*) during the reproductive season. Ambio 6(4):239.

Jia X, Lia N, Chen J. 2005. A subchronic toxicity study of elemental nano-Se in Sprague-Dawley rats. Life Sci 76:1989-2003.

Johnson BT, Saunders CR, Sanders HO. 1971. Biological magnification and degradation of DDT and aldrin by freshwater invertebrates. J Fish Res Board Can 28:705-709.

Johnson L. 2006. Personal communication (telephone conversation with Matt Luxon, Windward Environmental LLC, regarding incidental sediment ingestion by English sole and Pacific staghorn sculpin). Zoologist, Ecotoxicology and Environmental Fish Health Program, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, WA. May 17, 2006.

Johnson LL, Landahl JT. 1994. Chemical contaminants, liver disease, and mortality rates in English sole (*Pleuronectes vetulus*). Ecol Applic 4:59-68.

Johnson LL, Casillas E, Collier TK, McCain BB, Varanasi U. 1988. Contaminant effects on ovarian development in English sole (*Pleuronectes vetulus*) from Puget Sound, Washington. Can J Fish Aquat Sci 45:2133-2146.

Johnson LL, Casillas E, Sol SY, Collier TK, Stein JE, Varanasi U. 1993. Contaminant effects on reproductive success in selected benthic fish species. Mar Environ Res 35:165-170.

Johnson LL, Stein JE, Hom T, Sol S, Collier TK, Varanasi U. 1995. Effects of exposure to Prudhoe Bay crude oil on reproductive function in gravid female flatfish. Environ Sci 3:67-81.

Johnson LL, Sol SY, Lomax DP, Nelson GM, Sloan CA, Casillas E. 1997a. Fecundity and egg weight in English sole, *Pleuronectes vetulus*, from Puget Sound, Washington: influence of nutritional status and chemical contaminants. Fish Bull 95:231-249.

Johnson LL, Casillas E, Myers MS, Rhodes LD, Olson OP. 1997b. Patterns of oocyte development and related changes in plasma 17-b estradiol, vitellogenin, and plasma chemistry in English sole *Parophrys vetulus* Girard. Mar Environ Res 44(1):41-49.

Johnson LL, Landahl JT, Kubin LA, Horness BH, Myers MS, Collier TK, Stein JE. 1998. Assessing the effects of anthropogenic stressors on Puget Sound flatfish populations. J Sea Res 39:125-137.

Johnson LL, Sol SY, Ylitalo GM, Hom T, French B, Olson OP, Collier TK. 1999. Reproductive injury in English sole (*Pleuronectes vetulus*) from the Hylebos Waterway, Commencement Bay, Washington. J Aquat Ecosys Stress Recov 6:289-310.

Jones PD, Kannan K, Newsted JL, Tillitt DE, Williams LL, Giesy JP. 2001. Accumulation of 2,3,7,8-tetrachlorodibenzo-p-dioxin by rainbow trout (*Oncorhynchus mykiss*) at environmentally relevant dietary concentrations. Environ Toxicol Chem 20(2):344-350.

Julius AD, Davies MH, Birt DF. 1983. Toxic effects of dietary selenium in the Syrian hamster. Ann Nutr Metab 27:296-305.

Kaiser J. 2006. Personal communication (telephone conversation with Erika Hoffman, US Environmental Protection Agency Region 10, regarding osprey diet). Scientist, USGS Forest and Rangeland Ecosystem Science Center, Corvallis, OR. September 27, 2006.

Kaiser KLE, Dixon DG, Hodson PV. 1984. QSAR studies on chlorophenols, chlorobenzenes and para-substituted phenols. In: Kaiser KLE, ed, QSAR in environmental toxicology. D. Reidel Publishing Co., Dordrecht, the Netherlands, pp 189-206.

Kamunde CN, Grosell M, Lott JNA, Wood CM. 2001. Copper metabolism and gut morphology in rainbow trout (*Oncorhynchus mykiss*) during chronic sublethal dietary copper exposure. Can J Fish Aquat Sci 58:293-305.

Kang J-C, Kim S-G, Jang S-W. 2005. Growth and hematological changes of rockfish, *Sebastes schlegeli* (Hilgendorf) exposed to dietary Cu and Cd. J World Aquacult Soc 36(2):188-195.

Kennedy SW, Lorenzen A, Jones SP, Hahn ME, Stegeman JJ. 1996. Cytochrome P4501A induction in avian hepatocyte cultures: a promising approach for predicting the sensitivity of avian species to toxic effects of halogenated aromatic hydrocarbons. Toxicol Appl Pharmacol 141:214-230.

Kenney MJ. 1982. East, West and Duwamish Waterways navigation improvement study: fish and wildlife coordination act report. Division of Ecological Services, US Fish and Wildlife Service, Olympia, WA.

Kerwin J, Nelson TS, eds. 2000. Habitat limiting factors and reconnaissance assessment report, Green/Duwamish and Central Puget Sound watersheds (WRIA 9 and Vashon Island) [online]. Washington Conservation Commission, Lacey, WA, and King County Department of Natural Resources, Seattle, WA. Available from: http://dnr.metrokc.gov/WRIAS/9/Recon.htm.

Khasawinah AM, Grutsch JF. 1989. Chlordane: 24-month tumorigenicity and chronic toxicity test in mice. Regul Toxicol Pharmacol 10:244-254.

Kihlstrom JE, Olsson M, Jensen S, Johansson A, Ahlbom J, Bergman A. 1992. Effects of PCB and different fractions of PCB on the reproduction of the mink (*Mustela vison*). Ambio 2(8):563-569.

Kim S-G, Kim J-W, Kang J-C. 2004. Effect of dietary cadmium on growth and haematological parameters of juvenile rockfish, *Sebastes schlegeli* (Hilgendorf). Aquacult Res 35:80-86.

Kimbrough RD. 1985. Laboratory and human studies on polychlorinated biphenyls (PCBs) and related compounds. Environ Health Perspect 59:99–106.

Kimbrough RD. 1987. Human health effects of polychlorinated biphenyls (PCBs) and polybrominated biphenyls (PBBs). Ann Rev Pharmacol Toxicol 27:87–111.

King County. 1996. Norfolk CSO sediment cleanup study. Prepared for Elliott Bay/Duwamish Restoration Program. King County Department of Natural Resources, Seattle, WA.

King County. 1999a. King County combined sewer overflow water quality assessment for the Duwamish River and Elliott Bay. Vol 1, Appendix A: Problem formulation, analysis plan, and field sampling work plan. King County Department of Natural Resources, Seattle, WA.

King County. 1999b. King County combined sewer overflow water quality assessment for the Duwamish River and Elliott Bay. Vol 1, Appendix B2, B3, & B4: human health, wildlife, and aquatic life risk assessments. King County Department of Natural Resources, Seattle, WA.

King County. 1999c. King County combined sewer overflow water quality assessment for the Duwamish River and Elliott Bay. Vol 1, Appendix C: Issue papers. King County Department of Natural Resources, Seattle, WA.

King County. 1999d. King County combined sewer overflow water quality assessment for the Duwamish River and Elliott Bay. Vol 1: Overview and interpretation, plus appendices. King County Department of Natural Resources, Seattle, WA.

King County. 1999e. Norfolk CSO sediment remediation project five-year monitoring program: April 1999 monitoring report. Prepared for Elliott Bay/Duwamish Restoration Program. King County Department of Natural Resources, Seattle, WA.

King County. 2000a. Duwamish/Diagonal CSO/SD site assessment report. Draft. Prepared for Elliott Bay/Duwamish Restoration Program. King County Department of Natural Resources, Seattle, WA.

King County. 2000b. King County environmental laboratory quality assurance review for estuarine sediment analytical data. Norfolk CSO sediment remediation project five-year monitoring program. February 2000 sampling event. Environmental Laboratory, King County Department of Natural Resources, Seattle, WA.

King County. 2000c. Norfolk CSO sediment remediation project five-year monitoring program. Annual monitoring report-year one, April 2000. Prepared for Elliott Bay/Duwamish Restoration Program. King County Department of Natural Resources, Seattle, WA.

King County. 2000d. Norfolk CSO sediment remediation project five-year monitoring program. Six-month post-construction monitoring report, October 1999. Prepared for Elliott Bay/Duwamish Restoration Program. King County Department of Natural Resources, Seattle, WA.

King County. 2001a. Duwamish/Diagonal CSO/SD cleanup study report. Draft. Prepared for Elliott Bay/Duwamish Restoration Program panel. King County Department of Natural Resources, Seattle, WA.

King County. 2001b. Norfolk CSO sediment remediation project five-year monitoring program. Annual monitoring report-year two, April 2001. Prepared for Elliott Bay/Duwamish Restoration Program. King County Department of Natural Resources, Seattle, WA.

King County. 2002. Norfolk CSO sediment remediation project five-year monitoring program: Annual monitoring report - year 3, April 2002. Prepared for the Elliott Bay/Duwamish Restoration Panel. King County Department of Natural Resources and Parks, Seattle, WA.

Kirkpatrick CM. 1940. Some foods of young great blue herons. Am Midl Nat 24:594-601.

Knight RL, Randolph PJ, Allen GT, Young LS, Wigen RJ. 1990. Diets of nesting bald eagles, *Haliaeetus leucocephalus*, in western Washington. Can Field-Nat 104(4):545-551.

Kociba RJ, Keyes DG, Beyer JE, Carreon RM, Wade CE, Dittenber DA, Kalnins RP, Frauson LE, Park CN, Barnard SD, Hummel RA, Humiston CG. 1978. Results of a two-year chronic toxicity and oncogenicity study of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in rats. Toxicol Appl Pharmacol 46:279-303.

Kolaja GJ. 1977. The effects of DDT, DDE and their sulfonated derivatives on eggshell formation in the mallard duck. Bull Environ Contam Toxicol 17:697-701.

Kolok AS, Huckins JN, Petty JD, Oris JT. 1996. The role of water ventilation and sediment ingestion in the uptake of benzo(a)pyrene in gizzard shad (*Dorosoma cepedianum*). Environ Toxicol Chem 15(10):1752-1759.

Krausmann JD. 2002a. Preliminary exposure assessment of dioxin-like chlorobiphenyls in great blue herons of the lower Duwamish River in Seattle, Washington. US Fish and Wildlife Service, Lacey, WA.

Krausmann JD. 2002b. Personal communication (telephone conversation with Berit Bergquist, Windward Environmental, regarding great blue heron use of the LDW). Fish and Wildlife Biologist, US Fish and Wildlife Service, Lacey, WA. March 26.

Kubin LA. 1997. Growth of juvenile English sole exposed to sediments amended with aromatic compounds. Master's thesis. Western Washington State University, Bellingham, WA. 99 pp.

Kurta A. 1995. Mammals of the Great Lakes region. University of Michigan Press, Ann Arbor, MI.

Kushlan JA. 1978. Feeding ecology of wading birds. In: Sprunt A, Oge J, Winckler S, eds, Wading birds, research report No. 7. National Audubon Society, New York, NY, pp 249-297.

Lamb JC, Chapin RE, Teague J, Lawton AD, Reel JR. 1987. Reproductive effects of four phthalic acid esters in the mouse. Toxicol Appl Pharmacol 88:255-269.

Lange J. 2006. Personal communication (telephone conversation with Matt Luxon, Windward Environmental LLC, regarding incidental sediment ingestion by English sole and Pacific staghorn sculpin). Zoologist, Ecotoxicology and Environmental Fish Health Program, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, WA. May 18, 2006.

Lanno RP, Slinger SJ, Hilton JW. 1985a. Effect of ascorbic acid on dietary copper toxicity in rainbow trout (*Salmo gairdneri* Richardson). Aquaculture 49:269-287.

Lanno RP, Slinger SJ, Hilton JW. 1985b. Maximum tolerable and toxicity levels of dietary copper in rainbow trout (*Salmo gairdneri* Richardson). Aquaculture 49:257-268.

Larsen DN. 1984. Feeding habits of river otters in coastal southeastern Alaska. J Wildl Manage 484:1460-1452.

Leach RM, Jr, Wang KW-L, Baker DE. 1979. Cadmium and the food chain: the effect of dietary cadmium on tissue composition in chicks and laying hens. J Nutr 109:437-443.

Leffler CW. 1975. Effects of ingested mirex and DDT on juvenile *Callinectes sapidus* Rathbun. Environ Pollut 8:283-300.

Leighton FA. 1989. Acute oral toxicity of dibenzothiophene for male CD-1 mice: LD50, lesions, and the effect of preinduction of mixed-function oxidases. Fund Appl Toxicol 12:787-792.

Leon H. 1980. Final Report: Terminal 107 environmental studies. Benthic community impact study for terminal 107 (Kellogg Island) and vicinity. Prepared for Port of Seattle Planning and Research Department. Pacific Rim Planners, Inc., Seattle, WA.

Lepore PD, Miller RF. 1965. Embryonic viability as influenced by excess molybdenum in chicken breeder diets. Proc Soc Exper Biol Med 118:155-157.

Leroy D. 2007. Personal communication (e-mail to E. Hoffman, US Environmental Protection Agency Region 10, regarding J.L. Hugla barbel study [Hugla and Thome 1999]). Research Assistant, University of Liege, Liege, Belgium. February 8, 2007.

Levinton JS. 1982. Marine ecology. Prentice-Hall, Englewood Cliffs, NJ.

Lieb AJ, Bills DD, Sinnhuber RO. 1974. Accumulation of dietary polychlorinated biphenyls (Aroclor 1254) by rainbow trout (*Salmo gairdneri*). J Agr Food Chem 22:638-642.

Lien TF, Chen KL, Wu CP, Lu JJ. 2004. Effects of supplemental copper and chromium on the serum and egg traits of laying hens. Brit Poult Sci 45(4):535-539.

Lindsay DM, Sanders JG. 1990. Arsenic uptake and transfer in a simplified estuarine food chain. Environ Toxicol Chem 9:391-395.

Longcore JR, Samson FB. 1973. Eggshell breakage by incubating black ducks fed DDE. J Wildl Manage 37(3):390-394.

Lorentzen M, Maage A, Julshamn K. 1998. Supplementing copper to a fish meal based diet fed to Atlantic salmon parr affects liver copper and selenium concentrations. Aquacult Nutrit 4:67-72.

Lotufo GR, Farrar JD, Bridges TS. 2000. Effects of exposure source, worm density, and sex on DDT bioaccumulation and toxicity in the marine polychaete *Neanthes arenaceodentata*. Environ Toxicol Chem 19(2):472-484.

Lotufo GR, Farrar JD, Duke BM, Bridges TS. 2001a. DDT toxicity and critical body residue in the amphipod *Leptocheirus plumulosus* in exposures to spiked sediment. Arch Environ Contam Toxicol 41:142-150.

Lotufo GR, Landrum PF, Gedeon ML. 2001b. Toxicity and bioaccumulation of DDT in freshwater amphipods in exposures to spiked sediments. Environ Toxicol Chem 20(4):810-825.

Lowe JI, Parrish PR, Patrick JMJ, Forester J. 1972. Effects of the polychlorinated biphenyl Aroclor 1254 on the American oyster *Crassostrea virginica*. Mar Biol 17:209-214.

Lowe TP, Stendell RC. 1991. Eggshell modifications in captive American kestrels resulting from Aroclor 1248 in the diet. Arch Environ Contam Toxicol 20:519-522.

Lowry LF, Frost KJ. 1981. Feeding and trophic relationships of phocid seals and walruses in the Eastern Bering Sea. In: Hood DW, Calder JA, eds, The Eastern Bering Sea shelf: oceanography and resources. Vol 2. Department of Commerce, Washington, DC, pp 813-824.

Lu P-Y, Metcalf RL. 1975. Environmental fate and biodegradability of benzene derivatives as studied in a model aquatic ecosystem. Environ Health Persp 10:268-284.

Ludke JL. 1976. Organochlorine pesticide residues associated with mortality: additivity of chlordane and endrin. Bull Environ Contam Toxicol 16:253-260.

Ludke JL, Ferguson DE, Burke WD. 1968. Some endrin relationships in resistant and susceptible populations of golden shiners, *Notemigonus crysoleucas*. Trans Am Fish Soc 97:260-263.

Lundebye AK, Berntssen MHG, Bonga WSE, Maage A. 1999. Biochemical and physiological responses in Atlantic salmon (*Salmo salar*) following dietary exposure to copper and cadmium. Mar Pollut Bull 39:137-144.

Lundholm CE. 1997. DDE-induced eggshell thinning in birds: effects of p,p′-DDE on the calcium and prostaglandin metabolism of the eggshell gland. Comp Biol Physiol 118C(2):113-128.

Luxon M. 2004. Personal communication (direct observation of Caspian terns foraging at the north end of Kellogg Island during a fish sampling event in July 2004). Environmental scientist, Windward Environmental LLC, Seattle, WA.

Mabey WR, Smith JH, Podoll RT, Johnson HL, Mill T. 1982. Aquatic fate process data for organic priority pollutants. EPA 440/4-81-014. Office of Water Regulations and Standards, US Environmental Protection Agency, Washington, DC.

Mac MJ, Edsall CC, Seelye JG. 1985. Survival of lake trout eggs and fry reared in water from the upper Great Lakes. J Great Lakes Res 11(4):520-529.

Machemer L, Lorke D. 1981. Embryotoxic effect of cadmium on rats upon oral administration. Toxicol Appl Pharmacol 58:438-443.

MacKenzie KM, Angevine DM. 1981. Infertility in mice exposed *in utero* to benzo(a)pyrene. Biol Reprod 24:183-191.

Malins DC, McCain BB, Brown DW, Sparks AK, Hodgins HO. 1980. Chemical contaminants and biological abnormalities in central and southern Puget Sound. NOAA Technical Memorandum OMPA-2. Environmental Conservation Division, National Marine Fisheries Service, Seattle, WA.

Malins DC, McCain BB, Brown DW, Sparks AK, Hodgins HO, Chan S. 1982. Chemical contaminants and abnormalities in fish and invertebrates from Puget Sound. Environmental Conservation Division, National Marine Fisheries Service, Seattle, WA.

Malins DC, McCain BB, Brown DW, Chan SL, Myers MS, Landahl JT, Prohaska PG, Friedman AJ, Rhodes LD, Burrows DG, Gronlund WD, Hodgins HO. 1984. Chemical pollutants in sediments and diseases of bottom-dwelling fish in Puget Sound, Washington. Environ Sci Tech 18:705-713.

Malins DC, Krahn MM, Brown DW, Rhodes LD, Myers MS, McCain BB, Chan S-L. 1985a. Toxic chemicals in marine sediment and biota from Mukilteo, Washington: Relationships with hepatic neoplasms and other hepatic lesions in English sole (*Parophrys vetulus*). J Nat Cancer Inst 74(2):487-494.

Malins DC, Krahn MM, Myers MS, Rhodes LD, Brown DW, Krone CA, McCain BB, Chan S-L. 1985b. Toxic chemicals in sediments and biota from a creosote-polluted harbor: relationships with hepatic neoplasms and other hepatic lesions in English sole (*Parophrys vetulus*). Carcinogenesis 6:1463-1469.

Marine Mammal Center. 2006. Marine mammal information: California sea lion (*Zalophus californianus*) [online]. Marine Mammal Center, Sausalito, CA. Updated February 2002. [Cited August 10, 2006]. Available from: http://www.marinemannalcenter.org/learning/education/pinnipeds/casealion.asp.

Masutomi N, Shibutani M, Takagi H, Uneyama C, Takahashi N, Hirose M. 2003. Impact of dietary exposure to methoxychlor, genestein, or diisononyl phthalate during the perinatal period on the development of the rat endocrine/reproductive systems in later life. Toxicology 192:149-170.

Matsuda RI, Isaac GW, Dalseg RD. 1968. Fishes of the Green-Duwamish River. Water Quality Series No. 4. Municipality of Metropolitan Seattle, Seattle, WA.

Matta MB, Linse J, Cairncross C, Francendese L, Kocan RM. 2001. Reproductive and transgenerational effects of methylmercury or Aroclor 1268 on *Fundulus heteroclitus*. Environ Toxicol Chem 20(2):327-335.

Mauck WL, Mehrle PM, Mayer FL. 1978. Effects of the polychlorinated biphenyl Aroclor 1254 on growth, survival, and bone development in brook trout (*Salvelinus fontinalis*). J Fish Res Board Can 35:1084-1088.

Maxson SJ, Oring LW. 1980. Breeding season time and energy budgets of the polyandrous spotted sandpiper. Behaviour 74:200-263.

Mayer FL, Mehrle PM, Sanders HO. 1977. Residue dynamics and biological effects of polychlorinated biphenyls in aquatic organisms. Arch Environ Contam Toxicol 5:501-511.

Mayer KS, Mayer FL, Witt A, Jr. 1985. Waste transformer oil and PCB toxicity to rainbow trout. Trans Am Fish Soc 114(6):869-886.

McCain BB, Malins DC, Krahn MM, Brown DW, Gronlund WD, Moore LK, Chan SL. 1990. Uptake of aromatic and chlorinated hydrocarbons by juvenile chinook salmon (*Onchorhynchus tshawytscha*) in an urban estuary. Arch Environ Contam Toxicol 19:10-16.

McCarthy ID, Fuiman LA, Alvarez MC. 2003. Aroclor 1254 affects growth and survival skills of Atlantic croaker *Micropogonias undulatus* larvae. Mar Ecol Prog Ser 252:295-301.

McKim JM, Schmeider PK. 1990. Bioaccumulation: does it reflect toxicity? In: Nagel R, Loskill R, eds, Bioaccumulation in aquatic systems: Contributions to the

assessment. Proceedings, International Workshop, Berlin, 1990. VCG Publishers, New York, NY, pp 161-188.

McLane MAR, Hughes DL. 1980. Reproductive success of screech owls fed Aroclor 1248. Arch Environ Contam Toxicol 9:661-665.

Meador JP. 1997. Comparative toxicokinetics of tributyltin in five marine species and its utility in predicting bioaccumulation and acute toxicity. Aquat Toxicol 37:307-326.

Meador JP. 2000. An analysis in support of tissue and sediment based threshold concentrations of polychlorinated biphenyls (PCBs) to protect juvenile salmonids listed by the Endangered Species Act. Northwest Fisheries Science Center, National Oceanic and Atmospheric Administration, Seattle, WA.

Meador JP, Rice CA. 2001. Impaired growth in the polychaete *Armandia brevis* exposed to tributyltin in sediment. Mar Environ Res 51:113-129.

Meador JP, Krone CA, Dyer DW, Varanasi U. 1997. Toxicity of sediment-associated tributyltin to infaunal invertebrates: species comparison and the role of organic carbon. Mar Environ Res (43):219-241.

Meador JP, Collier TK, Stein JE. 2002. Determination of a tissue and sediment threshold for tributyltin to protect prey species of juvenile salmonids listed under the US Endangered Species Act. Aquat Conserv: Mar Freshw Ecosys 12:539-551.

Meador JP, Sommers FC, Ylitalo GM, Sloan CA. 2006. Altered growth and related physiological responses in juvenile chinook salmon (*Oncorhynchus tshawytscha*) from dietary exposure to polycyclic aromatic hydrocarbons (PAHs). Can J Fish Aquat Sci 63:2364-2376.

Mehring AL, Brumbaugh JH, Sutherland AJ, Titus HW. 1960. The tolerance of growing chickens for dietary copper. Poult Sci 39:713-719.

Mehrle PM, Mayer FL. 1976. Di-2-ethylhexyl phthalate: residue dynamics and biological effects in rainbow trout and fathead minnows. Proceedings of University of Missouri's 10th Annual Conference on Trace Substances in Environmental Health, June 8-10, Columbia, MO, pp 519-524.

Melquist WE, Dronkert AE. 1987. River otter. In: Novak M, Bailer JA, Obbarel ME, eds, Wild furbearer management and conservation in North America. Ontario Trappers' Association and Ontario Ministry of Natural Resources, Ottawa, ON, pp 627-641.

Melquist WE, Hornocker MG. 1983. Ecology of river otters in west central Idaho. In: Kirkpatrick RL, ed, Wildlife monographs. Vol 83. The Wildlife Society, Bethesda, MD, p 60.

Mendenhall VM, Klaas EE, McLane MAR. 1983. Breeding success of barn owls (*Tyto alba*) fed low levels of DDE and dieldrin. Arch Environ Contam Toxicol 12:235-240.

Meyer JH, Pearce TA, Patlan SB. 1981. Distribution and food habits of juvenile salmonids in the Duwamish Estuary. Prepared for Seattle District, US Army Corps of Engineers. US Fish and Wildlife Service, Olympia, WA.

Meyer JS, Adams WJ, Brix KV, Luoma SN, Mount DR, Stubblefield WA, Wood CM, eds. 2005. Toxicity of dietborne metals to aquatic organisms. Chapter 1, introduction, pp. 1-11. Society of Environmental Toxicology and Chemistry (SETAC), Pensacola, FL.

Michelsen TC, Bragdon-Cook K. 1993. Technical information memorandum: Organic carbon normalization of sediment data. Washington Department of Ecology, Olympia, WA.

Mickelson S, Williston D. 2006. Technical memorandum: Duwamish River/Elliott Bay/Green River water column PCB congener survey: transmittal of data and quality assurance documentation. King County Department of Natural Resources, Seattle, WA.

Millam JR, Craig-Veit CB, Batchelder ME, Viant MR, Herbeck TM, Woods LW. 2002. An avian bioassay for environmental estrogens: the growth response of zebra finch (*Taeniopygia guttata*) chick oviduct to oral estrogens. Environ Toxicol Chem 21(12):2663-2668.

Miller BS, Wingert RC, Borton SF. 1975. Ecological survey of demersal fishes in the Duwamish River and at West Point 1974. Prepared for Municipality of Metropolitan Seattle. Report no. FRI-UW-7509. Fisheries Research Institute, University of Washington, Seattle, WA.

Miller BS, McCain BB, Wingert RC, Borton SF, Pierce KV, Griggs DT. 1977a. Ecological and disease studies of demersal fishes in Puget Sound near METRO-operated sewage treatment plants and in the Duwamish River. FRI-UW-7721. Fisheries Research Institute, University of Washington, Seattle, WA.

Miller BS, Wingert RC, Borton SF, Griggs DT. 1977b. Ecological survey of demersal fishes in the Duwamish River and at West Point, 1975. Prepared for Municipality of Metropolitan Seattle. Fisheries Research Institute, University of Washington, Seattle, WA.

Miller BS, Simenstad CA, Moulton LL, Fresh KL, Funk FC, Karp WA, Borton SF. 1977c. Puget Sound baseline program nearshore fish survey. Final report, July 1974-June 1977. Prepared for Washington Department of Ecology. Fisheries Research Institute, University of Washington, Seattle, WA.

Miller PA, Lanno RP, McMaster ME, Dixon DG. 1993. Relative contributions of dietary and waterborne copper to tissue copper burdens and waterborne-copper

tolerance in rainbow trout (*Oncorhynchus mykiss*). Can J Fish Aquat Sci 5(8):1683-1689.

Miramand P, Guary JC, Fowler SW. 1981. Uptake, assimilation, and excretion of vanadium in the shrimp, *Lysmata seticaudata* (Risso), and the crab, *Carcinus maenas* (L.). J Exp Mar Biol Ecol 49:267-287.

Mirenda RJ. 1986a. Acute toxicity and accumulation of zinc in the crayfish, *Orconectes virilis* (Hagen). Bull Environ Contam Toxicol 37:387-394.

Mirenda RJ. 1986b. Toxicity and accumulation of cadmium in the crayfish, *Orconectes virilis* (Hagen). Arch Environ Contam Toxicol 15:401-407.

Mohiuddin SM, Taskar PK, Rheault M, Roy P-E, Chenard J, Morin Y. 1970. Experimental cobalt cardiomyopathy. Amer Heart J 80(4):532-543.

Moore DW, Dillon TM, Suedel BC. 1991. Chronic toxicity of tributyltin to the marine polychaete worm, *Neanthes arenaceodentata*. Aquat Toxicol 21:181-198.

Moore R, Toro E, Stanton M, Khan MAQ. 1977. Absorption and elimination of ^{14}C-*alpha*- and *gamma*-chlordane by a freshwater alga, daphnid, and goldfish. Arch Environ Contam Toxicol 6:411-420.

Morgan GW, Edens FW, Thaxton P, Parkhurst CR. 1975. Toxicity of dietary lead in Japanese quail. Poult Sci 54:1636-1642.

Morrow JE. 1980. The freshwater fishes of Alaska. Animal Resources Ecology Library, University of British Columbia, Vancouver, BC.

Mortimer MR, Miller GJ. 1994. Susceptibility of larval and juvenile instars of the sand crab, *Portunus pelagicus* (L.) to sea water contaminated by chromium, nickel or copper. Aust J Mar Freshwater Res 45:1107-1121.

Mount DR, Barth AK, Garrison TD, Barten KA, Hockett JR. 1994. Dietary and waterborne exposure of rainbow trout (*Oncorhynchus mykiss*) to copper, cadmium, lead and zinc using a live diet. Environ Toxicol Chem 13(12):2031-41.

Mowbray EE, Pursley D, Chapman JA. 1979. The status, population characteristics, and harvest of river otters in Maryland. Maryland Wildlife Administration. Waverly Press, Bethesda, MD.

Mulsow SG, Landrum P. 1995. Bioaccumulation of DDT in a marine polychaete, the conveyor-belt deposit feeder *Heteromastus filiformis* (Claparede). Chemosphere 31(4):3141-3152.

Murai T, Andrews JW, Smith RG, Jr. 1981. Effects of dietary copper on channel catfish. Aquaculture 22:353-357.

Murata Y, Denda A, Maruyama H, Konishi Y. 1993. Chronic toxicity and carcinogenicity studies of 1-methylnaphthalene in $B6C3F_1$ mice. Fund Appl Toxicol 21:44-51.

Murata Y, Denda A, Maruyama H, Nakae D, Tsutsumi M, Tsujiuchi T, Konishi Y. 1997. Chronic toxicity and carcinogenicity studies of 2-methylnaphthalene in $B6C3F_1$ mice. Fund Appl Toxicol 36:90-93.

Murdoch MH, Chapman PM, Norman DM, Quintino VM. 1997. Spiking sediment with organochlorines for toxicity testing. Environ Toxicol Chem 16(7):1504-1509.

Murie DJ. 1995. Comparative feeding ecology of two sympatric rockfish congeners, *Sebastes caurinus* (copper rockfish) and *S. maliger* (quillback rockfish). Mar Biol 124:341-353.

Murray FJ, Smith FA, Nitschke KD, Humiston CG, Kociba RJ, Schwetz BA. 1979. Three-generation reproduction study of rats given 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) in the diet. Toxicol Appl Pharmacol 50:241-252.

Myers MS, Johnson LL, Hom T, Collier TK, Stein JE, Varanasi U. 1998. Toxicopathic hepatic lesions in subadult English sole (*Pleuronectes vetulus*) from Puget Sound, Washington, USA: Relationships with other biomarkers of contaminant exposure. Mar Environ Res 45(1):47-67.

Nagy KA. 1987. Field metabolic rate and food requirement scaling in mammals and birds. Ecol Monogr 57(2):111-128.

Nagy KA. 2001. Food requirements of wild animals: predictive equations for free-living mammals, reptiles, and birds. Nutrition Abstracts and Reviews Series B: Livestock Feeds and Feeding 71(10):21R-31R.

Nagy KA, Girard IA, Brown TK. 1999. Energetics of free-ranging mammals, reptiles, and birds. Annu Rev Nutr. 247-277.

Nakayama K, Oshima Y, Nagafuchi K, Hano T, Shimasaki Y, Honjo T. 2005. Early-life-stage toxicity in offspring from exposed parent medaka, *Oryzias latipes*, to mixtures of tributyltin and polychlorinated biphenyls. Environ Toxicol Chem 24(3):591-596.

Nebeker AV, Puglisi FA, DeFoe DL. 1974. Effect of polychlorinated biphenyl compounds on survival and reproduction of the fathead minnow and flagfish. Trans Am Fish Soc 103:562-568.

Nebeker AV, Schuytema GS, Griffis WL, Barbitta JA, Carey LA. 1989. Effect of sediment organic carbon on survival of *Hyalella azteca* exposed to DDT and endrin. Environ Toxicol Chem 8:705-18.

Nebraska Game and Parks Commission. 2000. The river otter [online]. [Cited June 6, 2000]. Available from: http://ngp.ngpc.state.ne.us/wildlife/otters.html.

Nehlsen W, Williams JE, Lichatowich JA. 1991. Pacific salmon at the crossroads: Stock at risk from California, Oregon, Idaho, and Washington. Fisheries 16(2):4-21.

Nelson B. 2006. Personal communication (e-mail to Suzanne Replinger, Windward Environmental, regarding orca diet and the frequency of orca visits to the Seattle area). Community Conservation Liaison, Seattle Aquarium, Seattle, WA. May 26, 2006.

Nelson TS, Ruggerone G, Kim H, Schaefer R, Boles M. 2004. Juvenile chinook migration, growth and habitat use in the Lower Green River, Duwamish River and nearshore of Elliott Bay, 2001–2003. Draft. King County Department of Natural Resources and Parks Seattle, WA.

Nicolas JM. 1999. Vitellogenesis in fish and the effects of polycyclic aromatic hydrocarbon contaminants. Aquat Toxicol 45:77-90.

Niimi AJ. 1983. Biological and toxicological effects of environmental contaminants in fish and their eggs. Can J Fish Aquat Sci 40:306-312.

Nimmo DR, Wilson Jr AJ, Blackman RR. 1970. Localization of DDT in the body organs of pink and white shrimp. Bull Environ Contam Toxicol 5(4):333-341.

Nimmo DR, Forester J, Heitmuller PT, Cook GH. 1974. Accumulation of Aroclor 1254 in grass shrimp (*Palaemonetes pugio*) in laboratory and field exposures. Bull Environ Contam Toxicol 11(4):303-308.

Nirmala K, Oshima Y, Lee R, Imada N, Honjo T, Kobayashi K. 1999. Transgenerational toxicity of tributyltin and its combined effects with polychlorinated biphenyls on reproductive processes in Japanese medaka (*Oryzias latipes*). Environ Toxicol Chem 18(4):717-721.

NMFS. 1997. Impacts of California sea lions and Pacific harbor seals on salmonids and on the coastal ecosystems of Washington, Oregon, and California. NOAA Technical Memorandum NMFS-NWFSC-28. National Marine Fisheries Service, Seattle, WA.

NMFS. 2002. Unpublished data on PCB concentrations in juvenile chinook salmon captured in the Lower Duwamish Waterway during 1993 and 2000. Environmental Conservation Division, National Marine Fisheries Service, Seattle, WA.

NOAA. 1997. Duwamish River sediment study: sampling and analysis plan, quality assurance plan. Environmental Conservation Division, Northwest Fisheries Science Center, National Oceanic and Atmospheric Administration, Seattle, WA.

NOAA. 1998. Duwamish Waterway sediment characterization study report. Damage Assessment Center, National Oceanic and Atmospheric Administration, Seattle, WA.

NOAA, Ecology. 2000. Sediment quality in Puget Sound. Year 2 - central Puget Sound. Ecology publication no. 00-03-055. National Oceanic and Atmospheric

Administration, Silver Spring, MD and Washington Department of Ecology, Olympia, WA.

Norman D. 1999. Observation. In: King County combined sewer overflow water quality assessment for the Duwamish River and Elliott Bay. Appendix C: Issue Papers. King County Department of Natural Resources, Seattle, WA.

Norman D. 2002. Personal communication (email on 4/5/02 to Berit Bergquist, Windward Environmental LLC, regarding great blue herons in the LDW). Norman Wildlife Consulting, Shoreline, WA. April 5.

Norman D. 2006. Personal communication (telephone conversation with Berit Bergquist, Windward Environmental LLC, regarding use of the LDW by great blue heron). Norman Wildlife Consulting, Shoreline, WA. June 1.

Nosek JA, Craven SR, Sullivan JR, Hurley SS, Peterson RE. 1992. Toxicity and reproductive effects of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in ring-necked pheasant hens. J Toxicol Environ Health 35:187-198.

NRC. 1984. Nutrient requirements of poultry. National Research Council. National Academy Press, Washington, DC.

NRC. 1994. Nutrient requirements of poultry. Ninth revised edition. National Research Council. National Academy Press, Washington, DC.

Nysewander D, Evenson JR, Murphie BL, Cyra TA. 2001. Status and trends for a suite of key diving marine bird species characteristic of greater Puget Sound, as examined by the marine bird component, Puget Sound Ambient Monitoring Program (PSAMP). In: Puget Sound Research 2001: Abstracts & Biographies. Proceedings, meeting of the Puget Sound Water Quality Action Team, Bellevue, WA, February 12-14, 2001. Puget Sound Water Quality Action Team, Office of the Governor, Olympia, WA.

Oehlmann J, Stroben E, Fioroni P. 1991. The morphological expression of imposex in *Nucella lapillus* (Linnaeus) (Gastropoda: Muricidae). J Moll Stud 57:375-390.

Oh SH, Nakaue H, Deagen JT, Whanger PD, Arscott GH. 1979. Accumulation and depletion of zinc in chick tissue metallothioneins. J Nutr 109: 1720-1729.

Oladimeji AA, Leduc G. 1975. Effects of dietary methoxychlor on the food maintenance requirements of brook trout. Prog Wat Tech 7(3/4):587-598.

Oladimeji AA, Qadri SU, DeFreitas ASW. 1984. Long-term effects of arsenic accumulation in rainbow trout, *Salmo gairdneri*. Bull Environ Contam Toxicol 32:732-741.

Omura M, Ogata R, Kubo K, Shimasaki Y, Aou S, Oshima Y, Tanaka A, Hirata M, Makita Y, Inoue N. 2001. Two-generation reproductive toxicity study ot tributyltin chloride in male rats. Toxicol Sci 64:224-232.

Oring LW, Lank DB. 1986. Polyandry in spotted sandpipers: the impact of environment and experience. In: Rubenstein DO, Wrangham RW, eds, Ecological aspects of social evolution - birds and mammals. pp 21-42.

Oring LW, Lank DB, Maxson SJ. 1983. Population studies of polyandrous sandpiper. Auk 100:272-285.

Oring LW, Gray EM, Reed JM. 1997. Spotted sandpiper (*Actitis macularia*). In: Poole A, Gill F, eds, The birds of North America, No. 289. Academy of Natural Sciences, Philadelphia, PA, and American Ornithologists' Union, Washington, DC.

Ousterhout LE, Berg LR. 1981. Effects of diet composition on vanadium toxicity in laying hens. Poult Sci 60:1152-1159.

Pallson W. 2001. Personal communication (conversation with Windward Environmental LLC staff regarding English sole population dynamics in Elliott Bay). Washington Department of Fish and Wildlife, Olympia, WA.

Palm RC, Jr, Powell DG, Skillman A, Godtfredsen K. 2003. Immunocompetence of juvenile chinook salmon against *Listonella anquillarum* following dietary exposure to polycyclic aromatic hydrocarbons. Environ Toxicol Chem 22(12):2986-2994.

Parrish PR, Schimmel SC, Hansen DJ, Patrick Jr JM, Forester J. 1976. Chlordane: effects on several estuarine organisms. J Toxicol Environ Health 1:485-494.

Pattee OH. 1984. Eggshell thickness and reproduction in American kestrels exposed to chronic dietary lead. Arch Environ Contam Toxicol 13:29-34.

Patton JF, Dieter MP. 1980. Effects of petroleum hydrocarbons on hepatic function in the duck. Comp Biochem Physiol 65C:33-36.

Pauley GB, Bortz BM, Shepard MF. 1986. Species profiles: Life histories and environmental requirements of coastal fishes and invertebrates (Pacific Northwest). Steelhead trout. USFW biological report 82(11.62). Coastal Ecology Group, US Army Corps of Engineers, Vicksburg, MS and National Wetlands Research Center, US Fish and Wildlife Service, Slidell, LA.

Pauley GB, Armstrong DA, Heun TW. 1988. Species profiles: Life histories and environmental requirements of coastal fishes and invertebrates (Pacific Northwest). Dungeness crab. USFW biological report 82(11.63). Coastal Ecology Group, US Army Corps of Engineers, Vicksburg, MS and National Wetlands Research Center, US Fish and Wildlife Service, Slidell, LA.

Paulson D. 1993. Shorebirds of the Pacific Northwest. University of Washington Press, Seattle, WA.

Payne PM, Selzer LA. 1989. The distribution, abundance and selected prey of the harbor seal, *Phoca vitulina concolor*, in southern New England. Mar Mammal Sci 5:173-192.

Peakall DB, Lincer JL. 1972. Methyl mercury: its effect on eggshell thickness. Bull Environ Contam Toxicol 3(2):89-90.

Peakall DB, Peakall ML. 1973. Effect of a polychlorinated biphenyl on the reproduction of artificially and naturally incubated dove eggs. J Appl Ecol 10:863-868.

Peakall DB, Lincer JL, Bloom SE. 1972. Embryonic mortality and chromosomal alterations caused by Aroclor 1254 in ring doves. Environ Health Perspect 1:103-104.

Peakall DB, Lincer JL, Risebrough RW, Pritchard JB, Kinter WB. 1973. DDE-induced eggshell thinning; structural and physiological effects in three species. Comp Gen Pharmac 4:305-313.

Persia ME, Baker DH, Parsons CM. 2004. Tolerance for excess basic zinc chloride and basic copper chloride in chicks. Brit Poult Sci 45(5):672-676.

Peterson MJ, Southworth GR, Ham HD. 1994. Effect of sublethal chlorinated discharges on PCB accumulation in transplanted Asiatic clams (*Corbicula fluminea*). Wat Air Soil Pollut 73:169-178.

Philippart JC, Mélard C, Poncin P. 1989. Intensive culture of the common barbel, *Barbus barbus* (L.) for restocking. In: DePauw N, Jaspers E, Ackefors H, Wilkins N, eds, Aquaculture - a biotechnology in progress. European Aquaculture Society, Bredene, Belgium, pp 483-491.

Pitcher KW. 1980. Food of the harbor seal, *Phoca vitulina richardsi*, in the Gulf of Alaska. USNMFS Bull 78:544-549.

Pitcher KW, Calkins OG. 1979. Biology of the harbor seal, Phoca vitulina richardsi, in the Gulf of Alaska. Final. Outer Continental Shelf Environmental Assessment Program Research Unit 229. Bureau of Land Management, US Department of the Interior, Washington, DC.

Plasterer MR, Bradshaw WS, Booth GM, Carter MW. 1985. Developmental toxicity of nine selected compounds following prenatal exposure in the mouse: naphthalene, *p*-nitrophenol, sodium selenite, dimethyl phthalate, ethylenethiourea, and four glycol ether derivatives. J Toxicol Environ Health 15:25-38.

Poole A. 1989. Ospreys: a natural and unnatural history. Cambridge University Press, Cambridge, UK.

Poole AF. 1983. Courtship feeding, clutch size, and egg size in ospreys: a preliminary report. In: Bird DM, Seymour NR, Gerrard JM, eds, Biology and management of bald eagles and ospreys. Harpell Press, St. Anne de Bellvue, Quebec, pp 243-256.

Poole AF, Bierregaard RO, Martell MS. 2002. Osprey (*Pandion haliaetus*). In: Poole AF, Gill F, eds, The Birds of North America, No. 683. The Birds of North America, Inc., Philadelphia, PA.

Porter RD, Wiemeyer SN. 1970. Propagation of captive American kestrels. J Wildl Manage 34(3):594-604.

Poupoulis C, Jensen LS. 1976. Effect of high dietary copper on gizzard integrity of the chick. Poult Sci 55:113-121.

Powell DB, Palm RC, Jr, Skillman A, Godtfredsen K. 2003. Immunocompetence of juvenile chinook salmon against *Listonella anguillarum* following dietary exposure to Aroclor 1254. Environ Toxicol Chem 22(2):285-295.

Pritzl MC, Lie YH, Kienholz EW, Whiteman CE. 1974. The effect of dietary cadmium on development of young chickens. Poult Sci 53:2026-2029.

Prothro MG. 1993. Office of Water policy and technical guidance on interpretation and implementation of aquatic life metals criteria. Office of Water, US Environmental Protection Agency, Washington, DC.

PSDDA. 1988a. PSDDA Reports - Management plan report: Unconfined open-water disposal of dredged material, phase I (Central Puget Sound). Prepared for Puget Sound Dredged Disposal Authority. US Army Corps of Engineers, Seattle District; US Environmental Protection Agency, Region 10, Seattle, WA; Washington Department of Ecology; and Washington Department of Natural Resources.

PSDDA. 1988b. PSDDA Reports: Evaluation procedures technical appendix - phase I (central Puget Sound). Prepared for Puget Sound Dredged Disposal Authority. US Army Corps of Engineers, Seattle District; US Environmental Protection Agency, Region 10, Seattle, WA; Washington Department of Ecology; and Washington Department of Natural Resources.

PTI. 1990. Duwamish channel and settling basin sediment bioassays. Prepared for US Army Corps of Engineers. PTI Environmental Services, Bellevue, WA.

PTI. 1993. Recommendations for assessing adverse benthic effects in Puget Sound. Prepared for Washington Department of Ecology. PTI Environmental Services, Bellevue, WA.

PTI. 1996. Proposed dredging of slip no. 4, Duwamish River, Seattle, Washington: data report. Prepared for Puget Sound Dredged Disposal Analysis Program. PTI Environmental Services, Bellevue, WA.

Reuber MD. 1980. Significance of acute and chronic renal disease in Osborne-Mendel rats ingesting dieldrin or aldrin. Clin Toxicol 17(2):159-170.

Rhodes LD, Myers MS, Gronlund WD, McCain BB. 1987. Epizootic characteristics of hepatic and renal lesions in English sole, *Parophrys vetulus*, from Puget Sound. J Fish Biol 31:395-407.

Rhône-Poulenc. 1995. RCRA facility investigation (RFI) report for the Marginal Way facility. Vol 1: RFI results and conclusions. Prepared for US Environmental Protection Agency, Region 10. Rhône-Poulenc, Tukwila, WA.

Rhône-Poulenc. 1996. RCRA facility investigation (RFI) report for the Marginal Way facility. Round 3 data and sewer sediment technical memorandum. Prepared for US Environmental Protection Agency, Region 10. Rhône-Poulenc, Tukwila, WA.

Rice CA, Myers MS, Willis ML, French BL, Casillas E. 2000. From sediment bioassay to fish biomarker – connecting the dots using simple trophic relationships. Mar Environ Res 50:527-533.

Rice CP, O'Keefe P. 1995. Sources, pathways, and effects of PCBs, dioxins, and dibenzofurans. In: Hoffman DJ, Rattner BA, Burton G, Jr, Cairns J, Jr, eds, Handbook of ecotoxicology. Lewis Publishers, Boca Raton, FL, pp 424-468.

Rice SD, Short JW, Stickle WB. 1989. Uptake and catabolism of tributyltin by blue crabs fed TBT contaminated prey. Mar Environ Res 27:137-145.

Richardson ME, Fox MRS, Bry BE, Jr. 1974. Pathological changes produced in Japanese quail by ingestion of cadmium. J Nutr 104:323-338.

Ringer RK. 1983. Toxicology of PCBs in mink and ferrets. In: D'Itri FM, Kamrin MA, eds, PCBs: human and environmental hazards. Butterworth Publishers, Boston, pp 227-240.

Risebrough RW, Anderson DW. 1975. Some effects of DDE and PCB on mallards and their eggs. J Wildl Manage 39(3):508-513.

Roberson RH, Schaible PJ. 1960. The tolerance of growing chicks for high levels of different forms of zinc. Poult Sci 39:893-895.

Roberts MH, Hargis WJ, Strobel CJ, DeLisle PF. 1989. Acute toxicology of PAH contaminated sediments to the estuarine fish, *Leiostomus xanthurus*. Bull Environ Contam Toxicol 42:142-149.

Robertson B. 2004. Personal communication (telephone conversation with Matt Luxon, Windward Environmental, regarding rockfish observations in the LDW during SCUBA perch collection effort). Diver, The Seattle Aquarium, Seattle, WA. 4/16/2004.

Roffe TJ, Mate BR. 1984. Abundances and feeding habits of pinnipeds in the Rogue River, Oregon. J Wildl Manage 48:1262-1274.

Romoser GL, Dudley WA, Machlin LJ, Loveless L. 1961. Toxicity of vanadium and chromium for the growing chick. Poult Sci 40:1171-1173.

Roper JM, Cherry DS, Simmers JW, Tatem HE. 1997. Bioaccumulation of PAHs in the zebra mussel at Times Beach, Buffalo, New York. Environ Monit Assess 46:267-277.

Ruggerone GT, Volk E. 2004. Residence time and growth of natural and hatchery Chinook salmon in the Duwamish Estuary and Elliott Bay, Washington, based on otolith chemical and structural attributes. Draft. Prepared for US Army Corps of Engineers, Seattle District and Port of Seattle. Natural Resources Consultants, Inc., Seattle, WA.

Ruggerone GT, Nelson TS, Hall J, Jeanes E. 2006. Habitat utilization, migration timing, growth, and diet of juvenile chinook salmon in the Duwamish River and Estuary. King Conservation District and Salmon Recovery Funding Board, Seattle, WA.

Rule JH, Alden RW. 1996. Interactions of Cd and Cu in anaerobic estuarine sediments. II. Bioavailability, body burdens and respiration effects as related to geochemical partitioning. Environ Toxicol Chem 15(4):466-471.

Safe S. 1990. Polychlorinated biphenyls (PCBs), dibenzo-p-dioxins (PCDDs), dibenzofurans (PCDFs), and related compounds: environmental and mechanistic considerations which support the development of toxic equivalency factors (TEFs). Crit Rev Toxicol 21(1):51-88.

Safe S. 1991. Polychlorinated dibenzo-p-dioxins and related compounds: sources, environmental distribution and risk assessment. Environ Carcin Ecotoxicol Rev C9 (2):261–302.

Safe S. 1992. Toxicology, structure—function relationship, and human and environmental health impacts of polychlorinated biphenyls: progress and problems. Environ Health Perspect 100:259–268.

Safe S. 1994. Polychlorinated biphenyls (PCBs): environmental impact, biochemical and toxic responses, and implications for risk management. Crit Rev Toxicol 24(2):87-149.

Safe S. 1984. Polychlorinated biphenyls (PCBs) and polybrominated biphenyls (PBBs): biochemistry, toxicology, and mechanism of action. Crit Rev Toxicol 13.

SAIC. 1992. PSDDA bioassays for Duwamish Channel sediments. Prepared for US Army Corps of Engineers, Seattle District. Science Applications International Corporation, Bothell, WA.

Sample BE, Opresko DM, Suter GW. 1996. Toxicological benchmarks for wildlife. 1996 revision. ES/ERM-86/R3. Office of Environmental Management, US Department of Energy, Washington, DC.

Sanborn HR, Malins DC. 1977. Toxicity and metabolism of naphthalene: a study with marine larval invertebrates. Proc Soc Exper Biol Med 154:151-155.

Sanchez-Fortun S, Sanz, Santa-Maria A, Ros JM, De Vicente ML, Encinas MT, Vinagre E, Barahona MV. 1997. Acute sensitivity of three age classes of *Artemia salina* larvae to seven chlorinated solvents. Bull Environ Contam Toxicol 59:445-451.

Sanders HO, Chandler JH. 1972. Biological magnification of a polychlorinated biphenyl (Aroclor 1254) from water by aquatic invertebrates. Environ Contam Toxicol 7(5):257-263.

Sanderson JT, Van den Berg M. 1999. Toxic equivalency factors (TEFs) and their use in ecological risk assessment: a successful method when used appropriately. Human Ecolog Risk Assess 5:43-52.

Sato M. 1997. The price of taming a river: the decline of Puget Sound's Green/Duwamish Waterway. The Mountaineers, Seattle, WA.

Schaffer KE. 1989. Seasonal and size variations in diets of harbor seals. 8th Biennial Conference on the Biology of Marine Mammals, December 7-11, 1989, Pacific Groves, CA, p 62.

Scheuhammer AM. 1988. Chronic dietary toxicity of methylmercury in the zebra finch, *Poephila guttata*. Bull Environ Contam Toxicol 40:123-130.

Schiewe M, Crecelius EA, Myers MS. 1989. 1988 Reconnaissance survey of environmental conditions in 13 Puget Sound locations. Battelle.

Schimmel SC, Patrick JM, Jr, Forester J. 1976. Heptachlor: Uptake, depuration, retention, and metabolism by spot. J Toxicol Environ Health 2:169-178.

Schimmel SC, Patrick Jr JM, Wilson Jr AJ. 1977a. Acute toxicity to and bioconcentration of endosulfan by estuarine animals. In: Mayer FL, Hamelink JL, eds, Aquatic toxicology and hazard evaluation, ASTM STP 634. American Society for Testing and Materials, Philadelphia, PA, pp 241-252.

Schimmel SC, Patrick Jr JM, Forester J. 1977b. Toxicity and bioconcentration of BHC and lindane in selected estuarine animals. Arch Environ Contam Toxicol 6:355-363.

Schlicker SA, Cox DH. 1968. Maternal dietary zinc, and development and zinc, iron, and copper content of the rat fetus. J Nutr 95:287-294.

Schmitt CJ, Blazer VS, Dethloff GM, Tillitt DE, Gross TS, Bryant WL, DeWeese LR, Smith SB, Goede RW, Bartish TM, Kubiak TJ. 2000. Biomonitoring of environmental status and trends (BEST) program: field procedures for assessing the exposure of fish to environmental contaminants.

Schroeder HA, Balassa JJ. 1967. Arsenic, germanium, tin and vanadium in mice: effects on growth, survival and tissue levels. J Nutr 92:245-252.

Schroeder HA, Mitchener M. 1971. Toxic effects of trace elements on the reproduction of mice and rats. Arch Environ Health 23:102-106.

Schuytema GS, Krawczyk DF, Griffis WL, Nebeker AV, Robideaux ML. 1990. Hexachlorobenzene uptake by fathead minnows and macroinvertebrates in recirculating sediment/water systems. Arch Environ Contam Toxicol 19:1-9.

Schwetz BA, Norris JM, Sparschu GL, Rowe VK, Gehring PJ, Emerson JL, Gerbig CG. 1973. Toxicology of chlorinated dibenzo-p-dioxins. Environ Health Perspect 5:87-99.

Schwetz BA, Norris JM, Kociba RJ, Keeler PA, Cornier RF, Gehring PJ. 1974. Reproduction study in Japanese quail fed hexachlorobutadiene for 90 days. Toxicol Appl Pharmacol 30:255-265.

SEA. 1996. PSDDA chemical characterization of Duwamish Waterway and upper turning basin. FY97 operations and maintenance dredging. Prepared for US Army Corps of Engineers, Seattle District. Striplin Environmental Associates, Inc., Olympia, WA.

Seegal RF. 1996. Epidemiological and laboratory evidence of PCB-induced neurotoxicity. Crit Rev Toxicol 26(6)):709–737.

Shannon J. 2001. Personal communication (telephone conversation with Matt Luxon, Windward Environmental LLC, regarding bull trout captured in LDW). Fisheries Biologist, Taylor and Associates, Seattle, WA. April 11.

Shannon J. 2006. Personal communication (e-mails to Matt Luxon, Windward Environmental, regarding Duwamish-Elliott beach seine data collected by Taylor Associates in 1998, 2000, and 2002-2003. Biologist, Taylor Associates, Seattle, WA. March 30, 2006.

Shimasaki Y, Kitano T, Oshima Y, Inoue S, Imada N, Honjo T. 2003. Tributyltin causes masculinization in fish. Environ Toxicol Chem 22(1):141-144.

Shopp GM, White Jr KL, Holsapple MP, Barnes DW, Duke SD, Anderson AC, Condie Jr LW, Hayes JR, Borzelleca JF. 1984. Naphthalene toxicity in CD-1 mice: general toxicology and immunotoxicology. Fund Appl Toxicol 4:407-419.

Shubat PJ, Curtis LR. 1986. Ration and toxicant preexposure influence dieldrin accumulation by rainbow trout (*Salmo gairdneri*). Environ Toxicol Chem 5:69-77.

Silberhorn EM, Glauert HP, Robertson LW. 1990. Carcinogenicity of polyhalogenated biphenyls: PCBs and PBBs. Crit Rev Toxicol 20(6):397-426.

Smith MS. 1969. Responses of chicks to dietary supplements of copper sulphate. Brit Poult Sci 10:97-108.

Smith RT. 1936. Report on the Puget Sound otter trawl investigations. Wash Dep. Fish. Rep 36. Washington Department of Fisheries, Olympia, WA.

Snarski VM. 1982. The response of rainbow trout *Salmo gairdneri* to *Aeromonas hydrophilia* after sublethal exposures to PCB and copper. Environ Pollut Series A(28):219-232.

Snieszko SF. 1973. Recent advances in scientific knowledge and development pertaining to diseases of fishes. Adv Vet Sci Compar Med 17:291-314.

Spalding MG, Frederick PC, McGill HC, Bouton SN, McDowell LR. 2000. Methylmercury accumulation in tissues and its effects on growth and appetite in captive great egrets. J Wildl Dis 36(3):411-422.

Spann JW, Heinz GH, Camardese B, Hill EF, Moore JF, Murray HC. 1986. Differences in mortality among bobwhite fed methylmercury chloride dissolved in various carriers. Environ Toxicol Chem 5:721-724.

Spearman JW. 1991a. Sediment sampling and analysis, Brown Morton properties, Duwamish Waterway, Seattle, Washington. Jay W. Spearman, Consulting Engineer, Kirkland, WA.

Spearman JW. 1991b. Sediment sampling and analysis, South Park Marina, Duwamish Waterway, Seattle, Washington. Jay W. Spearman, Consulting Engineer, Kirkland, WA.

Spearman JW. 1999. Sediment sampling and analysis, James Hardie Gypsum Inc., Duwamish Waterway, Seattle, Washington. Jay W. Spearman, Consulting Engineer, Kirkland, WA.

Spence SK, Bryan GW, Gibbs PE, Masters D, Morris L, Hawkins SJ. 1990. Effects of TBT contamination on *Nucella* populations. Func Ecol 4:425-432.

Spinola RM, Serfass TL, Brooks RP. 1999. Radiotelemetry study: river otters reintroduction at Letchworth State Park. Progress report no. 1. New York State Department of Environmental Conservation, New York River Otter Project, and New York State Office of Parks, Recreation, and Historic Preservation, Albany, NY.

Srinivasan K, Mahadevappa KL, Radhakrishnamurty R. 1991. Toxicity of beta-and gamma-hexachlorocyclohexane in rats of different ages. Bull Environ Contam Toxicol 47:623-627.

Stahl JL, Greger JL, Cook ME. 1990. Breeding-hen and progeny performance when hens are fed excessive dietary zinc. Poult Sci 69:259-263.

Stanley TR, Jr, Spann JW, Smith GJ, Rosscoe R. 1994. Main and interactive effects of arsenic and selenium on mallard reproduction and duckling growth and survival. Arch Environ Contam Toxicol 26:441-451.

Stein JE, Hom T, Sanborn HR, Varanasi U. 1991. Effects of exposure to a contaminated-sediment extract on the metabolism and disposition of 17B-estrafiol in English sole (*Parophrys vetulus*). Comp Biochem Physiol 99C(1/2):231-240.

Stein JE, Hom T, Collier TK, Brown DW, Varanasi U. 1995. Contaminant exposure and biochemical effects in outmigrant juvenile chinook salmon from urban and nonurban estuaries of Puget Sound, Washington. Environ Toxicol Chem 14(6):1019-1029.

Stenson GB, Badgero GA, Fisher HD. 1984. Food habits of the river otter *Lutra canadensis* in the marine environment of British Columbia. Can J Zool 62:88-91.

Stephan CE, Mount DI, Hansen DJ, Gentile JH, Chapman GA, Brungs WA. 1985. Guidelines for deriving numerical national water quality criteria for the protection of aquatic organisms and their uses. PB85-227049. Office of Research and Development, US Environmental Protection Agency, Washington, DC.

Stern JH, Hennessy DP, Patmont CR. 2003. Improving estimates of contaminant exposure for mobile organisms: an assessment of area-weighted home range exposure estimates applied to the relationship between sediment chemistry and liver lesions in English sole. Puget Sound Research Conference 2003, Vancouver, BC.

Stewart BS, Leatherwood SL, Yochem PK. 1989. Harbor seal tracking and telemetry by satellite. Mar Mam Sci 5:361-375.

Stickel LF, Rhodes LI. 1970. The thin eggshell problem. In: Gillett JW, ed, Proceedings of the symposium, The Biological Impact of Pesticides in the Environment. Oregon State University, Corvallis, OR, pp 31-35.

Stoewsand GS, Anderson JA, Gutenmann WH, Bache CA, Lisk DJ. 1971. Eggshell thinning in Japanese quail fed mercuric chloride. Science 173:1030-1031.

Stoner JD. 1967. Prediction of salt-water intrusion in the Duwamish River Estuary, King County, Washington. Geological Survey professional paper 575-D. Geological Survey, US Department of the Interior, Washington, DC.

Stoner JD. 1972. Determination of mass balance and entrainment in the stratified Duwamish River Estuary, King County, Washington. Geological Survey water supply paper 1873-F. Geological Survey, US Department of the Interior, Washington, DC.

Strand J. 1999. Personal observation. In: King County combined sewer overflow water quality assessment for the Duwamish River and Elliott Bay. Appendix B2, B3, & B4: Human Health, Wildlife, and aquatic life risk assessments. King County Department of Natural Resources, Seattle, WA.

Swartz RC, Schultz DW, Ditsworth GR, DeBen WA, Cole FA. 1985. Sediment toxicity, contamination, and macrobenthic communities near a large sewage outfall. In: Boyle TT, ed, Validation and predictability of laboratory methods for assessing the fate and effects of contaminants in aquatic ecosystems. STP 865. American Society for Testing and Materials, Philadelphia, PA, pp 152-175.

Szebedinsky C, McGeer JC, McDonald DG, Wood CM. 2001. Effects of chronic Cd exposure via the diet or water on internal organ-specific distribution and subsequent gill Cd uptake kinetics in juvenile rainbow trout (*Oncorhynchus mykiss*). Environ Toxicol Chem 20(3):597-607.

Tabor JE, Wight HM. 1977. Population status of river otter in western Oregon. J Wildl Manage 41:692-699.

Tabor R, McCoy C, Camacho S, Celedonia M, Vecht S. 2004. Habitat use and movements of juvenile chinook salmon in south Lake Washington, 2003 investigations. Presentation at 2004 Greater Lake Washington Chinook Workshop, Shoreline Center, Shoreline, Washington, February 2, 2004 [online]. City of Seattle, Seattle, WA. Available from: http://www.cityofseattle.net/salmon/docs/workshop04/abstracts/Tabor2.pdf.

Takeda H, Shimma Y. 1977. Effects of toxic amounts of dietary zinc on the growth and body components of rainbow trout at two levels of calcium. Bull Freshw Fish Res 27:103-109.

Tanner CD. 1991. Potential intertidal habitat restoration sites in the Duwamish River Estuary. EPA 910/9-91-050. US Environmental Protection Agency, Region 10, Seattle, WA.

Toweill DE. 1974. Winter food habits of river otters in western Oregon. J Wildl Manage 38(1):107-111.

Traxler A. 2006. Personal communication (e-mail to Suzanne Replinger, Windward Environmental, regarding orca sightings in Elliott Bay and the Seattle area). Assistant Research Curator, The Whale Museum, Friday Harbor, WA. May 26, 2006.

Treon JF, Cleveland FP. 1955. Toxicity of certain chlorinated hydrocarbon insecticides for laboratory animals, with special reference to aldrin and dieldrin. Agric Food Chem 3:402-408.

Treon JF, Cleveland FP, Cappel J. 1955. Toxicity of endrin for laboratory animals. Agric Food Chem 3:842-848.

Truelove JD, Grant D, Mes J, Tryphonas H, Tryphonas L, Zawidzka Z. 1982. Polychlorinated biphenyl toxicity in the pregnant cynomolgus monkey: A pilot study. Arch Environ Contam Toxicol 11:583-588.

Tyl RW, Price CJ, Marr MC, Kimmel CA. 1988. Developmental toxicity evaluation of dietary di(2-ethylhexyl) phthalate in Fischer 344 rats and CD-1 mice. Fund Appl Toxicol 10:395-412.

Tyl RW, Myers CB, Marr MC, Fail PA, Seely JC, Brine DR, Barter RA, Butala JH. 2004. Reproductive toxicity evaluationof dietary butyl benzyl phthalate (BBP) in rats. Reprod Toxicol 18:241-264.

USACE. 2004. Quality assurance project plan, Lower Duwamish triad sampling event. Prepared for US Environmental Protection Agency. US Army Corps of Engineers, Seattle District, Seattle, WA.

USACE, EPA, WDNR, Ecology. 2000. Dredged material evaluation and disposal procedures. A user's manual for the Puget Sound Dredged Disposal Analysis (PSDDA) Program. US Army Corps of Engineers, Seattle District, Seattle, WA; US Environmental Protection Agency, Region 10, Seattle, WA; Washington Department of Natural Resources; and Washington Department of Ecology.

USEPA. 2000. Bioaccumulation testing and interpretation for the purpose of sediment quality assessment: status and needs. EPA-823-R-00-001. Bioaccumulation Analysis Workgroup, US Environmental Protection Agency, Washington, DC.

USFWS. 1964. Pesticide-wildlife studies, 1963: a review of Fish and Wildlife Service investigations during the calendar year. Circular 199. Fish and Wildlife Service, US Department of the Interior, Washington, DC.

USFWS. 2000a. GIS coverage for Duwamish intertidal habitats. Carol Langston, GIS Analyst, Curtis Tanner, Fish and Wildlife Biologist, US Fish and Wildlife Service, Lacey, WA.

USFWS. 2000b. Intertidal habitat projects monitoring plan. EBDRP, Seattle.

USFWS. 2000c. LDW intertidal metadata (Intertidal-hab.shp), electronic file dated April 18, 2000. Follows FGDC content standards for digital geospatial metadata standard. US Fish and Wildlife Service, Lacey, WA.

Van Birgelen APJM, van der Kolk J, Fase KM, Bol I, Poiger H, van den Berg M, Brouwer A. 1994. Toxic potency of 2,3,3',4,4',5-hexachlorobiphenyl relative to and in combination with 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in a subchronic feeding study in the rat. Toxicol Appl Pharmacol 126:202-213.

Van Daele LJ, Van Daele HA. 1982. Factors affecting the productivity of ospreys nesting in west-central Idaho. Condor 84:292-299.

Van den Berg M, Birnbaum L, Bosveld ATC, Brunström B, Cook P, Feeley M, Giesy JP, Hanberg A, Hasegawa R, Kennedy S, Kubiak T, Larsen JC, van Leeuwen FXR, Djien Liem AK, Nolt C, Peterson RE, Poellinger L, Safe S, Schrenk D, Tillitt D, Tysklind M, Younes M, Waern F, Zacharewski T. 1998. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. Environ Health Perspec 106(12):775-792.

Van den Berg M, Birnbaum LS, Denison M, De Vito M, Farland W, Feeley M, Fiedler H, Hakansson H, Hanberg A, Haws L, Rose M, Safe S, Schrenk D, Tohyama C,

Tritscher A, Tuomisto J, Tysklind M, Walker N, Peterson RE. 2006. The 2005 World Health Organization reevaluation of human and mammalian toxic equivalency factors for dioxins and dioxin-like compounds. Tox Sci 93(2):223-241.

van der Weiden MEJ, Hanegraaf FHM, Eggens ML, Celander M, Seinen W, van den Berg M. 1994. Temporal induction of cytochrome P450 1A in the mirror carp (*Cyprinus carpio*) after administration of several polycyclic aromatic hydrocarbons. Environ Toxicol Chem 13(5):797-802.

Van Velsen FL, Danse LHJC, van Leeuwen FXR, Dormans JAMA, van Logten MJ. 1986. The subchronic oral toxicity of the beta-isomer of hexachlorocyclohexane in rats. Fund Appl Toxicol 6:697-712.

van Wezel AP, Sijm DTHM, Seinen W, Opperhuizen A. 1995. Use of lethal body burdens to indicate species differences in susceptibility to narcotic toxicants. Chemosphere 31(5):3201-3209.

Varanasi U. 1989. Metabolism of polycyclic aromatic hydrocarbons in the aquatic environment. CRC Press, Boca Raton, FL.

Varanasi U, Casillas E, Arkoosh MR, Hom T, Misitano D, Brown DW, Chan S-L, Collier TK, McCain BB, Stein JE. 1993. Contaminant exposure and associated biochemical effects in outmigrant juvenile chinook salmon (*Onchorynchus tschawytscha*) from urban and nonurban estuaries of Puget Sound, Washington. NOAA technical memorandum NMFS-FWFSC-8. Northwest Fisheries Science Center, National Marine Fisheries Service, Seattle, WA.

Vernberg WB, DeCoursey PJ, Kelly M, Johns DM. 1977. Effects of sublethal concentrations of cadmium on adult *Palaemonetes pugio* under static and flow-through conditions. Bull Environ Contam Toxicol 17(1):16-24.

Verschuuren HG, Kroes R, Den Tonkelaar EM, Berkvens JM, Helleman PW, Rauws AG, Schuller PL, Van Esch GJ. 1976. Toxicity of methylmercury chloride in rats. II. Reproduction study. Toxicology 6:97-106.

Vos JG, van der Maas HL, Musch A, Ram E. 1971. Toxicity of hexachlorobenzene in Japanese quail with special reference to porphyria, liver damage, reproduction, and tissue residues. Toxicol Appl Pharmacol 18:944-957.

Walsh AR, O'Halloran J, Gower AM. 1994. Some effects of elevated levels of chromium (III) in sediments to the mullet *Chelon labrosus* (R). Ecotoxicol Environ Saf 27:168-176.

Ware GW, Good EE. 1967. Effects of insecticides on reproduction in the laboratory mouse. II. Mirex, telodrin, and DDT. Toxicol Appl Pharmacol 10:54-61.

Warner EJ, Fritz RL. 1995. The distribution and growth of Green River chinook salmon (*Oncorhynchus tshawytscha*) and chum salmon (*Oncorhynchus keta*) outmigrants

in the Duwamish estuary as a function of water quality and substrate. Water Resources Division, Muckleshoot Indian Tribe, Auburn, WA.

WDF. 1993. 1992 Washington state salmon and steelhead stock inventory. Washington Department of Fisheries, Washington Department of Wildlife, and Western Washington Treaty Indian Tribes, Olympia, WA.

WDFW. 1993. 1992 Washington state salmon and steelhead stock inventory. Washington Department of Fisheries, Washington Department of Wildlife, and Western Washington Treaty Indian Tribes, Olympia, WA.

WDFW. 1999. Bull trout in the Snohomish River system. Washington Department of Fish and Wildlife, Mill Creek, WA.

WDFW. 2000. Bull trout status/distribution table: bulltrout.evt.e00. GIS layer [online]. Washington State Department of Fish and Wildlife, Olympia, WA. [Cited 7/21/00]. Available from: ftp://ftp.streamnet.org/pub/streamnet/gisdata/FishDistribution_March2006/sp1498_BullTrout.htm.

WDFW. 2002. Final peregrine falcon status report [online]. Endangered Species Program, Washington Department of Fish and Wildlife, Olympia, WA. Available from: (http://wdfw.wa.gov/wlm/diversty/soc/status/peregrine/).

WDFW. 2006. State listed species. Revised March 2006. Washington Department of Fish and Wildlife, Olympia, WA.

WDFW. 2007. Species of concern in Washington State (current through May 24, 2007) [online]. Washington Department of Fish and Wildlife, Olympia, WA. [Cited 5/30/07]. Available from: http://wdfw.wa.gov/wlm/diversty/soc/soc.htm.

Webber HM, Haines TA. 2003. Mercury effects on predator avoidance behavior of a forage fish, golden shiner (*Notemigonus crysoleucas*). Environ Toxicol Chem 22(7):1556-1561.

Weber CW, Reid BL. 1968. Nickel toxicity in growing chicks. J Anim Sci 95:612-616.

Weitkamp DE, Campbell RF. 1980. Port of Seattle Terminal 107 fisheries study. Parametrix, Inc., Bellevue, WA.

Weitkamp DE, Ruggerone GT. 2000. Factors affecting chinook populations: background report. Prepared for City of Seattle. Parametrix, Inc., Kirkland, WA.

Weitkamp DE, Schadt TH. 1982. 1980 Juvenile salmonid study. 82-0415-012F. Prepared for Port of Seattle. Parametrix, Inc, Bellevue, WA.

Wellman PJ, Watkins PA, Nation JR, Clark DE. 1984. Conditioned taste aversion in the adult rat induced by dietary ingestion of cadmium or cobalt. Neurotoxicology 5(2):81-90.

WERF. 1996. Methodology for aquatic ecological aquatic ecological risk assessment. Project No. RP91-AER-1. Cited in King County Water Quality Assessment, 1999. Water Environment Research Foundation

West JE. 2001. Personal communication (data transmittal to Tad Deshler, Windward Environmental LLC, Seattle, WA). Marine Resources Division, Washington Department of Fish and Wildlife, Olympia, WA. March 14.

West JE, O'Neill SM, Lippert G, Quinnell S. 2001. Toxic contaminants in marine and anadromous fishes from Puget Sound, Washington. Results of the Puget Sound ambient monitoring program fish component 1989-1999. Washington Department of Fish and Wildlife, Olympia, WA.

Weston. 1993. Harbor Island remedial investigation report (part 2-sediment). Vol 1-report. Prepared for US Environmental Protection Agency, Region 10. Roy F. Weston, Inc., Seattle, WA.

Weston. 1998. Comprehensive RCRA facility investigation report, Boeing Plant 2, Seattle/Tukwila, Washington. Roy F. Weston, Inc., Seattle, WA.

Weston. 1999a. Site inspection report, Lower Duwamish River (RK 2.5-11.5), Seattle, Washington. Vol 1-Report and appendices. Prepared for US Environmental Protection Agency, Region 10. Roy F. Weston, Inc., Seattle, WA.

Weston. 1999b. Site inspection report, Lower Duwamish River (RK 2.5 to 11.5), Seattle, Washington. Volume 2 - map folio. Prepared for US Environmental Protection Agency, Region 10. Roy F. Weston, Inc., Seattle, WA.

Weston. 2000. Outfall and nearshore sediment sampling report, Duwamish facility. Prepared for James Hardie Gypsum, Seattle, Washington. Roy F. Weston, Inc., Seattle, WA.

White DH, Dieter MP. 1978. Effects of dietary vanadium in mallard ducks. J Toxicol Environ Health 4:43-50.

White DH, Finley MT. 1978a. Effects of dietary cadmium in mallard ducks. In: Hemphill DD, ed, Trace substances in environmental health - XII. University of Missouri, Columbia, MO, pp 220-223.

White DH, Finley MT. 1978b. Uptake and retention of dietary cadmium in mallard ducks. Environ Res 17:53-59.

WHO. 1993. Polychlorinated biphenyls and terphenyls. In: Dobson S, van Esch GJ, eds, Environmental health criteria 140. 2nd ed. World Health Organization, Geneva, Switzerland.

Widdows J, Page DS. 1993. Effects of tributyltin and dibutyltin on the physiological energetics of the mussel, *Mytilus edulis*. Mar Environ Res 35:233-249.

Wiemeyer SN, Hoffman DJ. 1996. Reproduction in eastern screech-owls fed selenium. J Wildl Manage 60(2):332-341.

Wiemeyer SN, Porter RD. 1970. DDE thins eggshells of captive American kestrels. Nature 227:737-738.

Wiemeyer SN, Bunck CM, Krynitsky AJ. 1988. Organochlorine pesticides, polychlorinated biphenyls, and mercury in osprey eggs--1970-79--and their relationships to shell thinning and productivity. Arch Environ Contam Toxicol 17:767-787.

Williams MS. 1990. Port of Seattle Terminal 107 (Kellogg Island), biological assessment - 1989. Parametrix, Inc, Bellevue, WA.

Williams RW, Laramie RM, Ames JJ. 1975. Puyallup basin: water resource inventory area 10. In: Catalog of Washington streams and salmon utilization. Vol 1: Puget Sound region. Washington Department of Fisheries, Olympia, WA.

Windward. 2000. Lower Duwamish Waterway remedial investigation. Results of second phase of clam reconnaissance survey. Memorandum to Doug Hotchkiss, Port of Seattle, July 19, 2000. Windward Environmental LLC, Seattle, WA.

Windward. 2002. East Waterway, Harbor Island Superfund site: Technical memorandum: Tissue chemistry results for juvenile chinook salmon collected from Kellogg Island and East Waterway. Prepared for the Port of Seattle. Windward Environmental LLC, Seattle, WA.

Windward. 2003a. Lower Duwamish Waterway remedial investigation. Phase 1 remedial investigation report. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2003b. Lower Duwamish Waterway remedial investigation. Phase 1 remedial investigation report. Appendix A: Ecological risk assessment. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2003c. Lower Duwamish Waterway remedial investigation. Phase 1 remedial investigation report. Appendix B: Human health risk assessment. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2004a. Lower Duwamish Waterway remedial investigation. Crab and shrimp pilot study and quarterly surveys data report. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2004b. Lower Duwamish Waterway remedial investigation. Intertidal clam survey data report. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2004c. Lower Duwamish Waterway remedial investigation. Juvenile chinook salmon data report. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2004d. Lower Duwamish Waterway remedial investigation. Quality assurance project plan: Benthic invertebrate sampling of the Lower Duwamish Waterway. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2004e. Lower Duwamish Waterway remedial investigation. Task 8: Phase 2 RI work plan. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2004f. Lower Duwamish Waterway remedial investigation. Technical memorandum: Gastropod pilot survey of the Lower Duwamish Waterway. Windward Environmental LLC, Seattle, WA.

Windward. 2004g. Lower Duwamish Waterway remedial investigation. Technical memorandum: Gastropod pilot survey results. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2004h. Lower Duwamish Waterway remedial investigation. Technical memorandum: LDW sandpiper presence and habitat survey results. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2004i. Lower Duwamish Waterway remedial investigation. Technical memorandum: Rockfish in the Lower Duwamish Waterway. Windward Environmental LLC, Seattle, WA.

Windward. 2004j. Memorandum: March 31, 2004, LDWG tissue meeting summary memo. Windward Environmental LLC, Seattle, WA.

Windward. 2005a. Lower Duwamish Waterway remedial investigation. Data and analysis report: porewater sampling of Lower Duwamish Waterway. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005b. Lower Duwamish Waterway remedial investigation. Data report: Chemical analyses of benthic invertebrate and clam tissue samples and co-located sediment samples. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005c. Lower Duwamish Waterway remedial investigation. Data report: Fish and crab tissue collection and chemical analyses. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005d. Lower Duwamish Waterway remedial investigation. Data report: Round 1 surface sediment sampling for chemical analyses and toxicity testing.

Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005e. Lower Duwamish Waterway remedial investigation. Data report: Round 2 surface sediment sampling for chemical analyses and toxicity testing. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005f. Lower Duwamish Waterway remedial investigation. Data report: Taxonomic identifications of benthic invertebrate communities. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005g. Lower Duwamish Waterway remedial investigation. Fish and crab data report addendum: PCB congener data, MS/MSD analyses, and DDT confirmation. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005h. Lower Duwamish Waterway remedial investigation. NOAA juvenile chinook salmon tissue data. Memorandum to EPA prepared on behalf of the Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005i. Lower Duwamish Waterway remedial investigation. Quality assurance project plan: Porewater sampling of Lower Duwamish Waterway. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005j. Lower Duwamish Waterway remedial investigation. Technical memorandum: 2005 gastropod imposex study. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2006a. Lower Duwamish Waterway remedial investigation. Data and analysis report: porewater sampling of Lower Duwamish Waterway. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2006b. Lower Duwamish Waterway remedial investigation. Data report: chemical analyses of fish and crab tissue samples collected in 2005. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2006c. Lower Duwamish Waterway remedial investigation. Technical memorandum: 2005 gastropod imposex study results. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2006d. Lower Duwamish Waterway remedial investigation. Technical memorandum: Criteria for defining the baseline surface sediment dataset for

use in the Lower Duwamish Waterway Phase 2 RI/FS. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward, DOF, Onsite. 2004a. Lower Duwamish Waterway Superfund site, Terminal 117 early action area. Sediment field sampling, cruise and data report. Prepared for the Port of Seattle. Windward Environmental LLC, Dalton, Olmsted & Fuglevand, Inc., and Onsite Enterprises, Inc., Seattle, WA.

Windward, DOF, Onsite. 2004b. Lower Duwamish Waterway Superfund site, Terminal 117 early action area. T-117 sediment and soil supplemental data report. Prepared for the Port of Seattle. Windward Environmental LLC, Dalton, Olmsted & Fuglevand, Inc., and Onsite Enterprises, Inc., Seattle, WA.

Wine RN, Li LH, Barnes LH, Gulati DK, Chapin RE. 1997. Reproductive toxicity of di-n-butylphthalate in a continuous breeding protocol in Sprague-Dawley rats. Environ Health Perspect 105:101-107.

Wingert RC, Terry CB, Miller BS. 1979. Food and feeding habits of ecologically important nearshore and demersal fishes in central Puget Sound. FRI-UW-7903. Prepared for Washington Department of Ecology. Fisheries Research Institute, University of Washington, Seattle, WA.

Wise MH, Linn IJ, Kennedy CR. 1981. A comparison of the feeding biology of mink *Mustela vison* and otter *Lutra lutra*. J Zool London 195:181-213.

Wobeser G, Nielsen NO, Schiefer B. 1976. Mercury and mink II. Experimental methyl mercury intoxication. Can J Comp Med 40:34-45.

Wofford HW, Wilsey CD, Neff GS, Giam CS, Neff JM. 1981. Bioaccumulation and metabolism of phthalate esters by oysters, brown shrimp, and sheepshead minnows. Ecotoxicol Environ Saf 5:202-210.

WR Consulting. 2004. Final sediment characterization report, Boyer Towing, Inc. dock replacement, Seattle, Washington. WR Consulting, Inc., Seattle, WA.

Wren CD, Hunter DB, Leatherland JF, Stokes PM. 1987. The effects of polychlorinated biphenyls and methylmercury, singly and in combination on mink. II: Reproduction and kit development. Arch Environ Contam Toxicol 16:449-454.

Wu RSS, Pollino CA, Au DWT, Zheng GJ, Yuen BBH, Lam PKS. 2003. Evaluation of biomarkers of exposure and effect in juvenile areolated grouper (*Epinephelus areolatus*) on foodborne exposure to benzo(a)pyrene. Environ Toxicol Chem 22(7):1568-1573.

Wydoski RS, Whitney RR. 2003. Inland fishes of Washington. 2nd ed. University of Washington Press, Seattle, WA.

Yates S. 1998. Marine wildlife from Puget Sound through the Inside Passage. Sasquatch Books, Seattle, WA.

You L, Casanova M, Bartolucci EJ, Fryczynski MW, Dorman DC, Everitt JI, Gaido KW, Ross SM, Heck HD. 2002. Combined effects of dietary phytoestrogen and synthetic endocrine-active compound on reproductive development in Sprague-Dawley rats: genistein and methoxychlor. Toxicol Sci 66:91-104.

MAPS

Wind Ward environmental LLC

0 0.25 0.5 Miles

0 0.25 0.5 Kilometers

N

Map A.1-1. Lower Duwamish Waterway study area

Prepared by CEJ 06/16/06, mod CEH 08/22/06, 08/03/07; Map 2387, \data\gis\ERA







Benthic invertebrate community surveys[a]

- Leon (1977-1978)
- West et al. (1989-1999)
- Williams (1989)
- Cordell et al. (1993-1999)
- King County WQA (1997)
- Ecology (2000)
- LDWG (2004)

Crab, clam, or shrimp surveys[a]

- ESG crab survey transect (1999)
- LDWG clam survey transect (2003)
- LDWG crab and shrimp survey location (2004)
- Navigation channel
- River mile

[a] Surveys were conducted for various purposes, which are discussed in Section A.2.2.2.

Windward environmental LLC

0 0.1 0.2 Miles

0 0.1 0.2 Kilometers

Scale is the same for each inset map

Map A.2-1. Locations of various LDW benthic invertebrate surveys conducted from 1977 to 2004

Sources: Leon 1980; Williams 1990; Cordell et al. 1996, 1997, 1999, 2001; King County 1999b; Ecology 2000a; ESG 1999; West et al. 2001; Windward 2004a, 2004b, 2005b, 2005c, 2005f, 2006a.

Prepared by CEH 05/26/06; mod 08/30/06, 12/07/06, 02/05/07, 05/15/07, 08/03/07; MAP 2330; W:\Projects\00-08-06_Duwamish_RI\data\gis\ERA\Phase 2



Map A.2-2. Tissue sampling locations in the Lower Duwamish Waterway

Sources: King County 1999d; Windward 2004c, 2005b, 2005c, 2006a.

Prepared by CEH, 06/13/06, mod CEH 08/30/06, 12/05/06, 12/19/06, 02-05-07, 08/03/07; MAP 2379; \data\gis\ERA\Phase 2\



Wind Ward environmental LLC

0 0.2 0.4 Miles

0 0.2 0.4 Kilometers

Scale is the same for each inset map

Map A.2-3. Baseline surface sediment sampling locations and porewater sampling areas

Sources: Ecology 2003; Exponent 1998; Floyd Snider McCarthy 2004; Hart Crowser 1993; Herrera 1997; Integral 2004; King County 1996, 1999d,1999e, 2000a, 2000b, 2000c, 2000d, 2001a, 2001b, 2002; NOAA 1997, 1998; Rhône-Poulenc 1995; Weston 1993, 1998, 1999, 2000; Windward 2004a, 2004b, 2005a, 2005b, 2005c, 2005d, 2005e; WR Consulting 2004.

Prepared by CEH 08/10/06, Mod CEH 8/30/06, 12/20/06, 02/05/07, 08/03/07, Map 2352, \data\gis\ERA\Phase 2\

0 25 50 Feet
0 10 20 Meters

N

Map A.3-1. Porewater sampling locations at GWI

Sources: Windward 2005a; Terra Vac and Floyd/Snider 2000.

Prepared by CEH, 08/30/06, mod 02/05/07, 08/03/07; MAP 2023; W:\Projects\00-08-06_Duwamish_RI\data\gis\Porewater\ERA



0 25 50 Feet

0 10 20 Meters

N

Map A.3-2. Porewater sampling locations at Boeing Plant 2 / Jorgensen Forge

Sources: Windward 2005a; Weston 1998.

Prepared by CEH, 08/30/06, mod 02/05/07, 08/03/07; MAP 2023; W:\Projects\05-08-06_Duwamish_RI\data\gis\Porewater\ERA





Detected concentration of TBT as ion (µg/kg dw) in surface sediment

- > 200
- > 100 and ≤ 200
- > 50 and ≤ 100
- > 20 and ≤ 50
- ≤ 20
- TBT not detected (RLs 1.0 - 5.3 µg/kg dw)
- Subtidal gastropod pilot survey sampling location
- Subtidal gastropod imposex analysis sampling location
- Intertidal zone
- Subtidal zone
- Navigation channel
- River mile
- **G1b** Gastropod sampling location ID*

* Only gastropod sampling locations have location IDs.

Windward environmental LLC

0 0.2 0.4 Miles

0 0.2 0.4 Kilometers

Scale is the same for each inset map

Map A.3-3. Gastropod sampling locations and TBT concentrations in baseline surface sediment samples

Sources: Windward 2004e, 2005b, d, e, 2006b; sources for baseline surface sediment data provided on Map A.2-2.

Prepared by STS 06/16/06, mod CEH 08/22/06, 12/20/06, 02/05/07, 08/03/07, Map 2336, \data\gis\ERA\Phase 2







Detected concentration of TBT as ion (µg/kg dw) in surface sediment*

- > 200
- > 100 and ≤ 200
- > 50 and ≤ 100
- > 20 and ≤ 50
- ≤ 20
- TBT not detected (RLs 1.0 - 5.3 µg/kg dw)
- Intertidal zone
- 0
- Navigation channel
- River mile

* Only benthic invertebrate tissue and co-located sediment sampling locations have location IDs. B1a through B10a are intertidal locations; B1b through B10b are subtidal locations.

Windward environmental LLC

0 0.1 0.2 Miles

0 0.1 0.2 Kilometers

Scale is the same for each inset map

Map A.3-4. TBT concentrations in baseline surface sediment samples and benthic invertebrate tissue and co-located sediment sampling locations

Sources: Windward 2005 b, d, e; Sources for baseline sediment data provided on Map A.2-2.

Prepared by STS 06/16/06, mod CEH 08/30/06, 12/20/06, 02/05/07, 08/03/07; Map 2323, W:\Projects\00-08-06_Duwamish_RI\data\gis\ERA\Phase 2\







LDWG toxicity test results (2005)*

bivalve amphipod

polychaete

> CSL

> SQS and ≤ CSL

≤ SQS

King County (2000)
DUD206 > SQS and ≤ CSL for echinoderm and polychaete toxicity tests; DUD200 through DUD205 no exceedances

River mile

Navigation channel

* Symbol oriented along the north-south axis in each frame.

0 0.1 0.2 Miles

0 0.1 0.2 Kilometers

Scale is the same for each inset map

Map A.3-5. Results of baseline toxicity testing conducted with LDW surface sediment samples

Sources: Ecology 2000; King County 2000; Windward 2005 d, e

Prepared by STS 06/16/06, mod CEH 08/30/06, 12/20/06, 02/05/07; Map 2320 \data\gis\ERA\Phase 2\





Exposure area

- Area 1 (RM 0.0 - RM 1.8)
- Area 2 (RM 1.1 - RM 3.1)
- Area 3 (RM 3.3 - RM 5.0)

Sandpiper foraging habitat quality

Habitat defined for intertidal zone only

- High
- Poor
- No habitat present
- Intertidal area not surveyed for sandpiper foraging habitat

Sampling location

- Surface sediment chemistry for all parameters for intertidal areas with foraging habitat
- Benthic invertebrate tissue identified by location ID

King County water quality sampling locations for metals and BEHP:

- Brandon CSO
- Norfolk CSO
- South and West Michigan CSOs

King County water quality sampling locations for PCBs:

- LTKE03
- LTUM03

- Navigation channel
- River mile

Windward environmental LLC

0 0.2 0.4 Miles

0 0.2 0.4 Kilometers

Scale is the same for each inset map

Map A.5-1. Exposure areas and sampling locations of data used in the spotted sandpiper exposure assessment

Sources: King County 1999a; Mickelson and Williston 2006; Windward 2004f, 2005b; sources for baseline surface sediment data provided on Map A.2-2.

Prepared by STS 05-16-06, mod CEH 08-10-06, STS 12-20-06, CEH 02-05-07, 08/03/07; Map 2283; \data\gis\ERA\Phase 2

Analyte	Exceedance Factor	
	SQS	CSL
PCBs	9.8	1.8

Chemicals in red exceeded CSL. The EF has no regulatory relevance and is presented here to provide an indication of the general magnitude of the concentration. Sampling locations represented by circles or squares were analyzed for all SMS chemicals. Surface sediment samples were collected at depths ≤15 cm below mudline.

Only locations with detected exceedances outside the Duwamish/ Diagonal dredged and capped or thin-layer placement areas are shown with exceedance factors. Colored symbols within these areas represent samples collected prior to dredging, capping or thin-layer placement. Ongoing monitoring data will be incorporated into the Phase 2 RI as available.

SQS/CSL categories for all surface sediment locations*

- > CSL, detected
- > SQS and ≤ CSL, detected
- > CSL, non-detect
- > SQS and ≤ CSL, non-detect
- ≤ SQS, detected
- ≤ SQS, non-detect

Locations where only PCBs were analyzed

- > CSL, detected
- > SQS and ≤ CSL, detected
- ≤ SQS, detected
- Dredged area
- Dredged and capped area
- Thin-layer placement
- EOF, CSO, or CSO/SD
- Pipe of unknown use
- Private outfall
- Publicly owned storm drain
- Seep
- Stream, channel, or swale
- Navigation channel
- River mile

Toxicity test results**: bivalve, amphipod, polychaete

* Symbols for locations sampled by LDWG are enlarged and identified by the location ID. Symbols for locations with detected SQS or CSL exceedances are identified by the location ID. Where more than one sample was collected at a location, symbol priority is as shown above.
** Toxicity test symbol oriented along north-south axis



Analyte	Abbreviation
1,2,4-Trichlorobenzene	124-TCB
1,2-Dichlorobenzene	12-DCB
2,4-Dimethylphenol	24-DMP
Benzofluoranthenes (total-calc'd)	Benzofluoranthenes
Bis(2-ethylhexyl)phthalate	BEHP
Butyl benzyl phthalate	BBP
Hexachlorobenzene	HCB
Hexachlorobutadiene	HCBD
PCBs (total calc'd)	PCBs
Pentachlorophenol	PCP
Total HPAH (calc'd)	Total HPAH
Total LPAH (calc'd)	Total LPAH

DR005	BEHP	1.6	
	BBP	1.1	
LDW-SS17	BEHP	1.4	
EST231	PCBs	1.1	
DUD200	4-Methylphenol	1.1	1.1
	BEHP	1.5	
	PCBs	1.1	
	BBP	1.1	
DUD_12C	PCBs	1.7	
	BEHP	1.6	
DUD043	BEHP	2.6	1.5
	PCBs	2.3	
	BBP	1.3	
DUD_10C	PCBs	3.1	
DUD_11C	BEHP	2.5	1.5
	PCBs	2.3	
	BBP	1.4	
DR006	BEHP	2.1	1.3
	BBP	1.5	
DUD044	PCBs	28	5.1
	BEHP	4.5	2.7
	HCB	1.4	
	BBP	1.2	
K-03	Phenol	1.3	
	PCBs	1.1	
DUD_9C	BEHP	1.3	
	PCBs	1.1	
DUD045	PCBs	2.3	
	BBP	1.3	
	BEHP	1.2	
DR081	PCBs	6.9	1.3
	BEHP	1.8	1.1
DUD032	Mercury	1.5	1.0
	PCBs	3.0	
	BEHP	1.3	
DUD_8C	PCBs	21	3.9
	BEHP	2.8	1.7
	Mercury	2.1	1.5
	Silver	1.0	1.0
	Cadmium	1.0	
CH1038	PCBs	6.8	1.2
DUD033	PCBs	1.5	
	BEHP	1.3	
DUD204	BEHP	2.1	1.3
	4-Methylphenol	1.1	1.1
	BBP	2.7	
DR059	BEHP	2.0	1.2
	BBP	1.1	
DUD_1C	BEHP	3.8	2.3
	PCBs	1.5	
DUD009	BEHP	2.3	1.4
	Fluoranthene	2.3	
	BBP	2.2	
	Total HPAH	1.1	
DUD205	4-Methylphenol	2.1	2.1
	BEHP	1.8	1.1
	PCBs	1.3	
	BBP	1.2	
DUD_2C	BEHP	2.3	1.4
	PCBs	1.4	
DUD010	N-Nitrosodiphenylamine	3.9	2.8
	BEHP	3.7	2.5
	Fluoranthene	1.8	1.2
	PCBs	6.6	
	BBP	4.9	
	Chrysene	1.4	
	Benzo(a)anthracene	1.2	
	Total HPAH	1.2	
	Benzofluroanthenes	1.0	
DR010	BEHP	1.1	
DUD014	BBP	3.7	

Map A.6-1a. Phase 2 sediment toxicity test results compared to SMS biological effects criteria and detected chemical concentrations at baseline surface sediment locations compared to SMS chemical criteria (RM 0.0-1.0)

Wind Ward environmental LLC

Prepared by STS 06/30/06, mod CEH 08/30/06; Map 2293

Source information provided by EPA and Ecology based on a preliminary file search. Tax parcel information provided by Seattle Public Utilities, May 2002. Some tax parcel polygons were edited by Windward to conform to the LDW shoreline for the purpose of map presentation. The locations of outfalls and other pipes shown on this figure were identified during a City of Seattle survey conducted during May-June 2003 (Herrera 2004). As part of the survey, the locations of permitted outfalls were first identified using available drainage and outfall maps for waterfront properties obtained from the Washington Department of Ecology National Pollutant Discharge Elimination System (NPDES) permit files. Outfalls and pipes that were observed in the field during low tides were then surveyed in the field to establish their locations. The status of permitted outfalls is currently being verified by the Lower Duwamish Waterway Group (LDWG) through interviews with agency personnel and individual LDWG members' staff, as appropriate.
In the future, known outfalls will be designated as either "combined sewer overflow, combined sewer overflow/storm drains, or emergency overflows;" "publicly owned storm drains;" or "private outfalls." Private outfalls will include two categories: 1) NPDES-permitted outfalls (e.g., storm drains, non-contact cooling water, process wastewater), and 2) other outfalls that are not included under an active NPDES permit. Outfalls whose discharge has been terminated and that are no longer included under an active NPDES permit will be identified as "formerly permitted outfalls." Pipes that cannot be identified as an outfall through agency permit file records review will be identified as "pipes of unknown use." A comprehensive survey of property owners will not be conducted.

Analyte	Abbreviation
1,2,4-Trichlorobenzene	124-TCB
1,2-Dichlorobenzene	12-DCB
2,4-Dimethylphenol	24-DMP
Benzofluoranthenes (total-calc'd)	Benzofluoranthenes
Bis(2-ethylhexyl)phthalate	BEHP
Butyl benzyl phthalate	BBP
Hexachlorobenzene	HCB
Hexachlorobutadiene	HCBD
PCBs (total calc'd)	PCBs
Pentachlorophenol	PCP
Total HPAH (calc'd)	Total HPAH
Total LPAH (calc'd)	Total LPAH



PCBs	18	3.4
Mercury	1.7	1.2

Fluoranthene	3.9	2.7
Pyrene	1.6	1.3
Total HPAH	1.5	1.1
PCBs	1.8	
Chrysene	1.5	
Benzo(a)anthracene	1.2	

Acenaphthene	16	4.6
Fluorene	10	3.0
Dibenzofuran	11	2.9
2-Methylnaphthalene	4.2	2.5
Total LPAH	4.6	2.2
Phenanthrene	7.5	1.6
Naphthalene	2.6	1.5
Fluoranthene	5.3	
PCBs	2.7	
Total HPAH	2.2	
Chrysene	1.6	
Benzo(a)anthracene	1.5	
Mercury	1.1	
Benzofluroanthenes	1.1	
Benzo(a)pyrene	1.0	

Mercury	2.7	1.8

PCBs	2.3	

PCBs	1.2	
BEHP	1.0	

Copper	3.2	3.2
Zinc	1.9	
BEHP	1.6	
Arsenic	1.2	

Copper	3.4	3.4
Arsenic	2.8	1.7
Zinc	2.1	

BEHP	1.1	
BBP	1.0	

Arsenic	14	8.7
Copper	3.6	3.6
Zinc	6.9	2.9
Lead	1.7	1.5
Mercury	1.9	1.3
Chrysene	1.3	
Fluoranthene	1.3	
Phenanthrene	1.3	
BEHP	1.2	
Acenaphthene	1.1	
BBP	1.1	

Arsenic	3.0	1.8
Copper	1.6	1.6
Zinc	1.9	

Fluoranthene	3.1	
Phenanthrene	1.4	
Chrysene	1.1	
Total HPAH	1.0	

PCBs	5.5	1.0
Arsenic	2.8	1.7
Zinc	1.5	

Benzyl alcohol	1.2	
PCBs	1.2	

PCBs	2.5	

PCBs	1.7	

PCBs	1.4	

Chrysene	2.5	
Zinc	3.7	1.6
PCBs	1.9	
Phenol	1.2	

PCBs	5.7	1.0

PCBs	1.3	

PCBs	3.2	

HCB	9.7	1.6

BEHP	1.2	

BEHP	2.0	1.2
BBP	1.8	

PCBs	1.1	

PCBs	1.3	

PCBs	1.1	

PCBs	1.9	

BEHP	2.0	1.2

PCBs	2.5	

PCBs	2.1	

PCBs	1.2	

PCBs	1.3	

PCBs	3.6	

BBP	3.1	

LDW-SSB4a
(Surface sediment sampling event)

PCBs	3.8	
PCP	1.1	

B4a (Benthic sampling event)

Fluoranthene	2.9	
Benzo(g,h,i)perylene	1.4	
PCBs	1.4	
Total HPAH	1.4	
Indeno(1,2,3-cd)pyrene	1.3	
Chrysene	1.2	
Phenanthrene	1.2	

Analyte	Exceedance Factor	
	SQS	CSL
PCBs	9.8	1.8

Chemicals in red exceeded CSL. The EF has no regulatory relevance and is presented here to provide an indication of the general magnitude of the concentration. Sampling locations represented by circles or squares were analyzed for all SMS chemicals. Surface sediment samples were collected at depths ≤15 cm below mudline.

Only locations with detected exceedances are shown with exceedance factors.

SQS/CSL categories for all surface sediment locations*

- > CSL, detected
- > SQS and ≤ CSL, detected
- > CSL, non-detect
- > SQS and ≤ CSL, non-detect
- ≤ SQS, detected
- ≤ SQS, non-detect

Locations where only PCBs were analyzed

- > CSL, detected
- > SQS and ≤ CSL, detected
- ≤ SQS, detected
- ≤ SQS, nondetect

- Dredged area
- Dredged area with thin-layer placement
- EOF, CSO, or CSO/SD
- Pipe of unknown use
- Private outfall
- Publicly owned storm drain
- Seep
- Stream, channel, or swale
- Navigation channel
- River mile

Toxicity test results**
bivalve, amphipod, polychaete

* Symbols for locations sampled by LDWG are enlarged and identified by the location ID. Symbols for locations with detected SQS or CSL exceedances are identified by the location ID. Where more than one sample was collected at a location, symbol priority is as shown above.
** Toxicity test symbol oriented along the north-south axis

Map A.6-1b. Phase 2 sediment toxicity test results compared to SMS biological effects criteria and detected chemical concentrations at baseline surface sediment locations compared to SMS chemical criteria (RM 1.0-2.0)

0 250 500 Feet
75 150 Meters
N

Wind Ward environmental LLC

Prepared by STS 06/30/06, mod CEH 08/30/06; Map 2293

Source information provided by EPA and Ecology based on a preliminary file search. Tax parcel information provided by Seattle Public Utilities, May 2002. Some tax parcel polygons were edited by Windward to conform to the LDW shoreline for the purpose of map presentation. The locations of outfalls and other pipes shown on this figure were identified during a City of Seattle survey conducted during May-June 2003 (Herrera 2004). As part of the survey, the locations of permitted outfalls were first identified using available drainage and outfall maps for waterfront properties obtained from the Washington Department of Ecology National Pollutant Discharge Elimination System (NPDES) permit files. Outfalls and pipes that were observed in the field during low tides were then surveyed in the field to establish their locations. The status of permitted outfalls is currently being verified by the Lower Duwamish Waterway Group (LDWG) through interviews with agency personnel and individual LDWG members' staff, as appropriate.
In the future, known outfalls will be designated as either "combined sewer overflow, combined sewer overflow/storm drains, or emergency overflows;" "publicly owned storm drains;" or "private outfalls." Private outfalls will include two categories: 1) NPDES-permitted outfalls (e.g., storm drains, non-contact cooling water, process wastewater), and 2) other outfalls that are not included under an active NPDES permit. Outfalls whose discharge has been terminated and that are no longer included under an active NPDES permit will be identified as "formerly permitted outfalls." Pipes that cannot be identified as an outfall through agency permit file records review will be identified as "pipes of unknown use." A comprehensive survey of property owners will not be conducted.



Map A.6-1c. Phase 2 sediment toxicity test results compared to SMS biological effects criteria and detected chemical concentrations at baseline surface sediment locations compared to SMS chemical criteria (RM 2.0-3.0)

0 250 500 Feet

0 75 150 Meters

N

Wind Ward environmental LLC

Prepared by STS 06/30/06, mod CEH 08/30/06, 03/08/07; Map 2293

Source information provided by EPA and Ecology based on a preliminary file search. Tax parcel information provided by Seattle Public Utilities, May 2002. Some tax parcel polygons were edited by Windward to conform to the LDW shoreline for the purpose of map presentation. The locations of outfalls and other pipes shown on this figure were identified during a City of Seattle survey conducted during May-June 2003 (Herrera 2004). As part of the survey, the locations of permitted outfalls were first identified using available drainage and outfall maps for waterfront properties obtained from the Washington Department of Ecology National Pollutant Discharge Elimination System (NPDES) permit files. Outfalls and pipes that were observed in the field during low tides were then surveyed in the field to establish their locations. The status of permitted outfalls is currently being verified by the Lower Duwamish Waterway Group (LDWG) through interviews with agency personnel and individual LDWG members' staff, as appropriate.
In the future, known outfalls will be designated as either "combined sewer overflow, combined sewer overflow/storm drains, or emergency overflows;" "publicly owned storm drains;" or "private outfalls." Private outfalls will include two categories: 1) NPDES-permitted outfalls (e.g., storm drains, non-contact cooling water, process wastewater), and 2) other outfalls that are not included under an active NPDES permit. Outfalls whose discharge has been terminated and that are no longer included under an active NPDES permit will be identified as "formerly permitted outfalls." Pipes that cannot be identified as an outfall through agency permit file records review will be identified as "pipes of unknown use." A comprehensive survey of property owners will not be conducted.



Analyte	Abbreviation
1,2,4-Trichlorobenzene	124-TCB
1,2-Dichlorobenzene	12-DCB
2,4-Dimethylphenol	24-DMP
Benzofluoranthenes (total-calc'd)	Benzofluoranthenes
Bis(2-ethylhexyl)phthalate	BEHP
Butyl benzyl phthalate	BBP
Hexachlorobenzene	HCB
Hexachlorobutadiene	HCBD
PCBs (total calc'd)	PCBs
Pentachlorophenol	PCP
Total HPAH (calc'd)	Total HPAH
Total LPAH (calc'd)	Total LPAH

Analyte	Exceedance Factor	
	SQS/SL	CSL/ML
PCBs	26	4.7

Toxicity test results**
bivalve
amphipod
polychaete

Chemicals in red exceeded CSL. The EF has no regulatory relevance and is presented here to provide an indication of the general magnitude of the concentration. Sampling locations represented by circles or squares were analyzed for all SMS chemicals. Surface sediment samples were collected at depths ≤15 cm below mudline.

Only locations with detected exceedances are shown with exceedance factors. Exceedance factors are not shown for locations within proposed removal areas.

SQS/CSL categories for all surface sediment locations*
- > CSL, detected
- > SQS and ≤ CSL, detected
- > CSL, non-detect
- > SQS and ≤ CSL, non-detect
- ≤ SQS, detected
- ≤ SQS, non-detect

Locations where only PCBs were analyzed
- > CSL, detected
- > SQS and ≤ CSL, detected
- ≤ SQS, detected

- Proposed removal area
- Dredged area
- EOF, CSO, or CSO/SD
- Pipe of unknown use
- Private outfall
- Publicly-owned storm drain
- Seep
- Stream, channel, or swale
- Navigation channel
- River mile

* Symbols for locations sampled by LDWG are enlarged and identified by the location ID. Symbols for locations with detected SQS or CSL exceedances are identified by the location ID. Where more than one sample was collected at a location, symbol priority is as shown above.
** Toxicity test symbol oriented along the north-south axis

R22	Benzo(g,h,i)perylene	3.2	1.3
	Indeno(1,2,3-cd)pyrene	3.2	1.3
	Dibenzo(a,h)anthracene	3.0	1.1
	Fluoranthene	2.5	
	Phenanthrene	2.1	
	Total HPAH	1.9	
	Chrysene	1.8	
	Benzo(a)pyrene	1.7	
	Arsenic	1.4	
	Benzofluroanthenes	1.4	
	Benzo(a)anthracene	1.4	
	PCBs	1.1	
	BEHP	1.0	
EST148	PCBs	2.5	
LDW-SS158	PCBs	1.7	
LDW-SS115	Fluoranthene	1.7	
	Phenanthrene	1.3	
	Chrysene	1.2	
	Dibenzo(a,h)anthracene	1.0	
	Total HPAH	1.0	
R23	Benzo(g,h,i)perylene	5.8	2.3
	Dibenzo(a,h)anthracene	5.9	2.2
	Indeno(1,2,3-cd)pyrene	5.6	2.2
	Benzo(a)pyrene	2.6	1.2
	Benzofluroanthenes	2.3	1.2
	BEHP	1.7	1.1
	PCBs	4.3	
	Fluoranthene	4.1	
	Phenanthrene	3.9	
	Total HPAH	3.1	
	Chrysene	2.8	
	BBP	2.4	
	Benzo(a)anthracene	2.1	
	Total LPAH	1.4	
	Acenaphthene	1.4	
	Fluorene	1.3	
	Dibenzofuran	1.2	
LDW-SS157	Benzoic acid	1.2	1.2
	BBP	1.3	
R26	BBP	2.0	
	PCBs	1.3	
R27	PCBs	1.9	
	BBP	1.2	
EIT060	PCBs	1.6	

Map A.6-1d. Phase 2 sediment toxicity test results compared to SMS biological effects criteria and detected chemical concentrations at baseline surface sediment locations compared to SMS chemical criteria (RM 3.0-4.0)

Prepared by STS 06/30/06, CEH 08/30/06; Map 2293

Source information provided by EPA and Ecology based on a preliminary file search. Tax parcel information provided by Seattle Public Utilities, May 2002. Some tax parcel polygons were edited by Windward to conform to the LDW shoreline for the purpose of map presentation. The locations of outfalls and other pipes shown on this figure were identified during a City of Seattle survey conducted during May-June 2003 (Herrera 2004). As part of the survey, the locations of permitted outfalls were first identified using available drainage and outfall maps for waterfront properties obtained from the Washington Department of Ecology National Pollutant Discharge Elimination System (NPDES) permit files. Outfalls and pipes that were observed in the field during low tides were then surveyed in the field to establish their locations. The status of permitted outfalls is currently being verified by the Lower Duwamish Waterway Group (LDWG) through interviews with agency personnel and individual LDWG members' staff, as appropriate.
In the future, known outfalls will be designated as either "combined sewer overflow, combined sewer overflow/storm drains, or emergency overflows;" "publicly owned storm drains;" or "private outfalls." Private outfalls will include two categories: 1) NPDES-permitted outfalls (e.g., storm drains, non-contact cooling water, process wastewater), and 2) other outfalls that are not included under an active NPDES permit. Outfalls whose discharge has been terminated and that are no longer included under an active NPDES permit will be identified as "formerly permitted outfalls." Pipes that cannot be identified as an outfall through agency permit file records review will be identified as "pipes of unknown use." A comprehensive survey of property owners will not be conducted.



Analyte	Abbreviation
1,2,4-Trichlorobenzene	124-TCB
1,2-Dichlorobenzene	12-DCB
2,4-Dimethylphenol	24-DMP
Benzofluoranthenes (total-calc'd)	Benzofluoranthenes
Bis(2-ethylhexyl)phthalate	BEHP
Butyl benzyl phthalate	BBP
Hexachlorobenzene	HCB
Hexachlorobutadiene	HCBD
PCBs (total calc'd)	PCBs
Pentachlorophenol	PCP
Total HPAH (calc'd)	Total HPAH
Total LPAH (calc'd)	Total LPAH

SD-SWY19	PCBs	9.2	1.7
SD-344-S	PCBs	36	6.6
SD-DUW89	Zinc	8.5	3.6
	PCBs	10	1.8
SD-DUW73	PCBs	1.6	
SD-307-S	PCBs	10.0	1.8
	Zinc	3.4	1.4
	Fluorene	3.0	
	Fluoranthene	3.1	
	Chrysene	2.7	
	Phenanthrene	2.2	
	Total HPAH	2.1	
	Benzo(a)anthracene	1.9	
	Benzofluroanthenes	1.4	
	Total LPAH	1.4	
	Phenol	1.0	
SD-SWY16	PCBs	3.7	
SD-341-S	PCBs	4.4	
SD-316-S	PCBs	3.5	
SD-330-S	PCBs	2.7	
SD-317-S	Phenol	2.6	
	PCBs	2.5	
SD-SWY17	PCBs	1.3	
SD-333-S	Copper	1.2	1.2
	PCBs	2.5	
SD-213-S	PCBs	2.3	
SD-309-S	PCBs	2.2	
	Fluorene	1.6	
	Phenanthrene	1.5	
	Dibenzofuran	1.3	
SD-332-S	PCBs	1.3	
SD-318-S	PCBs	3.8	
	Phenol	1.9	
SD-343-S	PCBs	1.1	
SD-208-S	PCBs	1.9	
SD-310-S	PCBs	2.3	
SD-215-S	PCBs	4.5	
SD-334-S	PCBs	1.6	
SD-335-S	PCBs	1.7	
SD-311-S	PCBs	15	2.8
	Phenol	2.9	1.0
SD-319-S	PCBs	15	2.8
SD-312-S	PCBs	3.9	
	Fluorene	1.6	
	Phenol	1.5	
	Phenanthrene	1.5	
	Dibenzofuran	1.2	
SD-342-S	PCBs	21	3.9
	Lead	1.9	1.6
SD-320-S	PCBs	46	8.5
SD-340-S	PCBs	1.2	
LDW-SS110	PCBs	48	8.8
	Lead	1.9	1.6
EST155	PCBs	1.6	
SD-211-S	PCBs	2.3	
SD-313-S	Chromium	2.2	2.2
	Lead	1.4	1.2
	PCBs	4.3	
	Phenol	1.5	
	Zinc	1.3	
SD-338-S	PCBs	3.3	
LDW-SS111	PCBs	12	2.2
	Chromium	1.8	1.7
	Lead	1.4	1.2
	Phenanthrene	1.4	
	Fluorene	1.2	
	Zinc	1.1	
SD-337-S	PCBs	5.4	
SD-314-S	PCBs	3.8	
EST156	PCBs	1.2	
SD-DUW93	PCBs	2.8	
SD-339-S	PCBs	2.3	
SD-216-S	PCBs	1.5	
SD-315-S	BBP	1.8	
	PCBs	1.4	
SD-217-S	PCBs	1.3	
SD-345-S	PCBs	1.2	
EST161	PCBs	1.6	
LDW-SS112	Arsenic	8.4	5.2
	BBP	2.4	
	PCBs	2.2	
	Fluoranthene	1.2	
SD-336-S	PCBs	1.3	
LDW-SS114	Arsenic	19	12
	BEHP	1.7	
	PCBs	4.5	
	Fluoranthene	1.3	
	Indeno(1,2,3-cd)pyrene	1.1	
	Chrysene	1.1	
EST162	PCBs	1.3	
EST147	PCBs	4.4	

Analyte	Exceedance Factor	
	SQS	CSL
PCBs	9.8	1.8

Chemicals in red exceeded CSL. The EF has no regulatory relevance and is presented here to provide an indication of the general magnitude of the concentration. Sampling locations represented by circles or squares were analyzed for all SMS chemicals. Surface sediment samples were collected at depths ≤15 cm below mudline.

Only locations with detected exceedances are shown with exceedance factors. Exceedance factors are not shown for proposed removal areas.

SQS/CSL categories for all surface sediment locations*

- > CSL, detected
- > SQS and ≤ CSL, detected
- > CSL, non-detect
- > SQS and ≤ CSL, non-detect
- ≤ SQS, detected
- ≤ SQS, non-detect

Locations where only PCBs were analyzed

- > CSL, detected
- > SQS and ≤ CSL, detected
- ≤ SQS, detected

- Dredged area
- Proposed removal area
- EOF, CSO, or CSO/SD
- Pipe of unknown use
- Private outfall
- Publicly owned storm drain
- Seep
- Stream, channel, or swale
- Navigation channel
- River mile

Toxicity test results**
bivalve, amphipod, polychaete

* Symbols for locations sampled by LDWG are enlarged and identified by the location ID. Symbols for other locations with detected SQS or CSL exceedances are identified by the location ID. Where more than one sample was collected at a location, symbol priority is as shown above.
** Toxicity test symbol oriented along the north-south axis

Map A.6-1e. Phase 2 sediment toxicity test results compared to SMS biological effects criteria and detected chemical concentrations at baseline surface sediment locations compared to SMS chemical criteria (RM 3.5-3.8)

0 10

0 50 100 Feet
20 Meters

N

Wind Ward environmental LLC

Prepared by STS 06/30/06, mod CEH 08/30/06; Map 2293

Source information provided by EPA and Ecology based on a preliminary file search. Tax parcel information provided by Seattle Public Utilities, May 2002. Some tax parcel polygons were edited by Windward to conform to the LDW shoreline for the purpose of map presentation. The locations of outfalls and other pipes shown on this figure were identified during a City of Seattle survey conducted during May-June 2003 (Herrera 2004). As part of the survey, the locations of permitted outfalls were first identified using available drainage and outfall maps for waterfront properties obtained from the Washington Department of Ecology National Pollutant Discharge Elimination System (NPDES) permit files. Outfalls and pipes that were observed in the field during low tides were then surveyed in the field to establish their locations. The status of permitted outfalls is currently being verified by the Lower Duwamish Waterway Group (LDWG) through interviews with agency personnel and individual LDWG members' staff, as appropriate.
In the future, known outfalls will be designated as either "combined sewer overflow, combined sewer overflow/storm drains, or emergency overflows;" "publicly owned storm drains;" or "private outfalls." Private outfalls will include two categories: 1) NPDES-permitted outfalls (e.g., storm drains, non-contact cooling water, process wastewater), and 2) other outfalls that are not included under an active NPDES permit. Outfalls whose discharge has been terminated and that are no longer included under an active NPDES permit will be identified as "formerly permitted outfalls." Pipes that cannot be identified as an outfall through agency permit file records review will be identified as "pipes of unknown use." A comprehensive survey of property owners will not be conducted.



Analyte	Abbreviation
1,2,4-Trichlorobenzene	124-TCB
1,2-Dichlorobenzene	12-DCB
2,4-Dimethylphenol	24-DMP
Benzofluoranthenes (total-calc'd)	Benzofluoranthenes
Bis(2-ethylhexyl)phthalate	BEHP
Butyl benzyl phthalate	BBP
Hexachlorobenzene	HCB
Hexachlorobutadiene	HCBD
PCBs (total calc'd)	PCBs
Pentachlorophenol	PCP
Total HPAH (calc'd)	Total HPAH
Total LPAH (calc'd)	Total LPAH

Analyte	Exceedance Factor	
	SQS	CSL
PCBs	9.8	1.8

Chemicals in red exceeded CSL. The EF has no regulatory relevance and is presented here to provide an indication of the general magnitude of the concentration. Sampling locations represented by circles or squares were analyzed for all SMS chemicals. Surface sediment samples were collected at depths ≤15 cm below mudline.

Only locations with detected exceedances are shown with exceedance factors.

SQS/CSL categories for all surface sediment locations*

- > CSL, detected
- > SQS and ≤ CSL, detected
- > CSL, non-detect
- > SQS and ≤ CSL, non-detect
- ≤ SQS, detected
- ≤ SQS, non-detect

Locations where only PCBs were analyzed

- > CSL, detected
- > SQS and ≤ CSL, detected
- ≤ SQS, detected

- Dredged area
- Dredged and capped area
- EOF, CSO, or CSO/SD
- Pipe of unknown use
- Private outfall
- Publicly owned storm drain
- Seep
- Stream, channel, or swale
- Navigation channel
- River mile

Toxicity test results**
bivalve, amphipod, polychaete

* Symbols for locations sampled by LDWG are enlarged and identified by the location ID. Symbols for locations with detected SQS or CSL exceedances are identified by the location ID. Where more than one sample was collected at a location, symbol priority is as shown above.
** Toxicity test symbol oriented along the north-south axis

Benzoic acid	1.4	1.4	
Benzoic acid	2.0	2.0	
Dibenzo(a,h)anthracene	2.2		
Fluoranthene	2.1		
Total HPAH	1.0		
PCBs	9.6	1.3	
Benzoic acid	1.3	1.3	
PCBs	1.6		
Mercury	1.6	1.1	
Benzoic acid	2.9	2.9	
Phenol	3.3	1.2	
BEHP	1.3		
Dibenzo(a,h)anthracene	1.3		
Phenol	3.3	1.2	
Dibenzo(a,h)anthracene	1.9		
BEHP	1.6		
Indeno(1,2,3-cd)pyrene	1.0		
Dibenzo(a,h)anthracene	2.2		
BEHP	1.3		
Benzo(g,h,i)perylene	1.3		
Indeno(1,2,3-cd)pyrene	1.3		
Fluoranthene	1.1		
Acenaphthene	1.6		
Dibenzo(a,h)anthracene	2.0		
PCBs	1.2		
Benzoic acid	2.6	2.6	
Dibenzo(a,h)anthracene	1.3		
Acenaphthene	2.9		
Dibenzofuran	1.7		
Fluorene	1.7		
Phenanthrene	1.5		
PCBs	1.8		
Benzo(g,h,i)perylene	1.6		
Dibenzo(a,h)anthracene	1.6		
Indeno(1,2,3-cd)pyrene	1.5		
Fluoranthene	1.1		
PCBs	1.3		
PCBs	1.1		
PCBs	1.2		
PCBs	1.9		
Acenaphthene	1.3		
Lead	1.4	1.2	
Phenol	2.6		
PCBs	1.7		
Acenaphthene	1.1		

NFK304	PCBs	1.5	
NFK305	PCBs	830	150
	Benzoic acid	6.9	6.9
NFK306	PCBs	3.8	
LDW-SS143	PCBs	33	6.0
NFK307	PCBs	4.2	
NFK308	PCBs	2.2	
NFK309	PCBs	1.1	
NFK310	PCBs	2.9	
NFK311	PCBs	1.7	
LDW-SS144	PCBs	2.1	
4	PCBs	28	5.1
5	PCBs	1.8	
6	PCBs	1.3	
7	PCBs	16	2.9
NFK507	PCBs	33	6.0
	BEHP	1.9	1.2
	BBP	1.1	
NFK508	PCBs	33	6.2
11	PCBs	1.5	
NFK503-Apr-04	PCBs	1.5	
NFK503-Apr-02	PCBs	2.5	
NFK501-Apr-04	PCBs	3.6	
NFK501-Apr-01	BBP	1.4	
R88	1,4-Dichlorobenzene	21	7.2
NFK004A	BEHP	2.0	1.4
	PCBs	2.9	
	BBP	2.2	
	Fluoranthene	1.3	
NFK005	BEHP	1.4	
NFK006	Dibenzo(a,h)anthracene	1.3	
	Phenanthrene	1.4	
	Indeno(1,2,3-cd)pyrene	1.2	

Map A.6-1f. Phase 2 sediment toxicity test results compared to SMS biological effects criteria and detected chemical concentrations at baseline surface sediment locations compared to SMS chemical criteria (RM 4.0-5.0)

0 210 420 Feet
0 75 150 Meters
N

Wind Ward environmental LLC

Prepared by STS 06/30/06, mod 08/30/06; Map 2293

Source information provided by EPA and Ecology based on a preliminary file search. Tax parcel information provided by Seattle Public Utilities, May 2002. Some tax parcel polygons were edited by Windward to conform to the LDW shoreline for the purpose of map presentation. The locations of outfalls and other pipes shown on this figure were identified during a City of Seattle survey conducted during May-June 2003 (Herrera 2004). As part of the survey, the locations of permitted outfalls were first identified using available drainage and outfall maps for waterfront properties obtained from the Washington Department of Ecology National Pollutant Discharge Elimination System (NPDES) permit files. Outfalls and pipes that were observed in the field during low tides were then surveyed in the field to establish their locations. The status of permitted outfalls is currently being verified by the Lower Duwamish Waterway Group (LDWG) through interviews with agency personnel and individual LDWG members' staff, as appropriate.
In the future, known outfalls will be designated as either "combined sewer overflow, combined sewer overflow/storm drains, or emergency overflows;" "publicly owned storm drains;" or "private outfalls." Private outfalls will include two categories: 1) NPDES-permitted outfalls (e.g., storm drains, non-contact cooling water, process wastewater), and 2) other outfalls that are not included under an active NPDES permit. Outfalls whose discharge has been terminated and that are no longer included under an active NPDES permit will be identified as "formerly permitted outfalls." Pipes that cannot be identified as an outfall through agency permit file records review will be identified as "pipes of unknown use." A comprehensive survey of property owners will not be conducted.







Nickel
- > ML, detect
- > SL and ≤ ML, detect
- ≤ SL, detect

Total DDTs
- > LOAEL, detect
- ≤ NOAEL, detect

Total chlordane
- > LOAEL, detect
- > NOAEL and ≤ LOAEL, detect
- ≤ NOAEL, detect

Intertidal zone
Subtidal zone
Navigation channel
River mile

Note: Multpile symbols are shown at a single location if more than one of these chemicals was detected.

Windward environmental LLC

0 0.1 0.2 Miles
0 0.1 0.2 Kilometers
Scale is the same for each inset map

Map A.6-2. SL/ML categories for nickel and NOAEL/LOAEL categories for total DDTs and total chlordane at LDW baseline surface sediment sampling locations

Sources: Windward 2005b, 2005d, 2005e; sources for baseline surface sediment data provided on Map A.2-2.

Prepared by CEH 05/26/06, mod CEH 08/28/06, STS 12/19/06, CEH 02/05/07, 08/03/07, Map 2341; \data\gis\ERA\Phase 2







> CSL toxicity or
> CSL chemistry if no toxicity data

> SQS and ≤ CSL toxicity or
> SQS and ≤ CSL chemistry if no toxicity data

≤ SQS toxicity or
≤ SQS chemistry if no toxicity data

Navigation channel

River mile

Exceedances of the SQS and CSL chemical criteria are based on detected concentrations.
For chemicals whose SQS and CSL are on an organic carbon normalized basis, if the TOC was < 0.5% or > 4.0%, the dry weight concentration of that chemical was compared to the lowest AET and second lowest AET, and exceedances of the lowest AET and second lowest AET were equated with exceedances of the SQS and CSL, respectively.

Windward environmental LLC

0 0.1 0.2 Miles
0 0.1 0.2 Kilometers

Scale is the same for each inset map

Map A.6-3. Exceedances of SQS and CSL (chemical criteria and toxicity combined) using Thiessen polygons for the LDW baseline surface sediment dataset

Sources: Windward 2005b, 2005d, 2005e; sources for baseline surface sediment data provided on Map A.2-2.

Prepared by CEH 06/16/06, mod CEH 08/28/06, 08/03/07; Map 2383; \data\gis\ERA\Phase 2







PCBs
SQS = 12 mg/kg OC
CSL = 65 mg/kg OC

For locations with TOC < 0.5% or > 4.0%
Lowest AET = 130 µg/kg dw
Second lowest AET = 1,000 µg/kg dw

- > CSL, detected
- > SQS and ≤ CSL, detected
- ≤ SQS, detected
- Non-detect
- Navigation channel
- River mile

Exceedances of the SQS or CSL chemical criteria are shown even if toxicity tests for that sample demonstrated no toxicity, which would override any exceedances of the SQS or CSL chemical criteria.

Windward environmental LLC

0 0.1 0.2 Miles
0 0.1 0.2 Kilometers

Scale is the same for each inset map

Map A.6-4. Exceedances of SQS and CSL chemical criteria using Thiessen polygons for PCBs in the LDW baseline surface sediment dataset

Sources: Windward 2005b, 2005d, 2005e; sources for baseline surface sediment data provided on Map A.2-2.

Prepared by CEH 06/16/06, mod CEH 08/28/06, 08/03/07; Map 2324; \data\gis\ERA\Phase 2







Bis(2-ethylhexyl) phthalate
SQS = 47 mg/kg OC
CSL = 78 mg/kg OC

For locations with TOC < 0.5% or > 4.0%
Lowest AET = 1,300 µg/kg dw
Second lowest AET = 1,900 µg/kg dw

- > CSL, detected
- > SQS and ≤ CSL, detected
- ≤ SQS, detected
- Non-detect
- Navigation channel
- River mile

Exceedances of the SQS or CSL chemical criteria are shown even if toxicity tests for that sample demonstrated no toxicity, which would override any exceedances of the SQS or CSL chemical criteria.

Windward environmental LLC

0 0.1 0.2 Miles
0 0.1 0.2 Kilometers

Scale is the same for each inset map

Map A.6-5. Exceedances of SQS and CSL chemical criteria using Thiessen polygons for bis(2-ethylhexyl) phthalate in the LDW baseline surface sediment dataset

Sources: Windward 2005b, 2005d, 2005e; sources for baseline surface sediment data provided on Map A.2-2.

Prepared by CEH 06/16/06; mod CEH 08/28/06; 08/03/07; Map 2324; \data\gis\ERA\Phase 2





SQS and CSL categories for baseline surface sediment sampling locations

1,2,4-Trichlorobenzene

- > CSL, detect
- > CSL, non-detect
- > SQS and ≤ CSL, non-detect
- ≤ SQS, detect
- ≤ SQS, non-detect
- Navigation channel
- River mile

Wind Ward environmental LLC

0 0.2 0.4 Miles

0 0.2 0.4 Kilometers

Scale is the same for each inset map

Map A.6-6. 1,2,4-Trichlorobenzene detected concentrations and RLs compared to SQS and CSL chemical criteria at baseline surface sediment sampling locations

Prepared by AMW, 06-27-06, mod CEH 08-29-06, STS 12-19-06, CEH 02-05-07, 08/03/07; MAP 2402; \data\gis\ERA\Phase 2



SQS and CSL categories for baseline surface sediment sampling locations (SQS = CSL for 2,4-dimethylphenol)

2,4-Dimethylphenol

- > CSL, detect
- > CSL, non-detect
- ≤ SQS, non-detect
- Navigation channel
- River mile



Wind Ward environmental LLC

0 0.2 0.4 Miles

0 0.2 0.4 Kilometers

Scale is the same for each inset map

Map A.6-7. 2,4-Dimethylphenol detected concentrations and RLs compared to SQS and CSL chemical criteria at baseline surface sediment sampling locations

Prepared by AMW, 06-27-06, mod CEH 08-29-06, STS 12-19-06, CEH 02-05-07, 08/03/07; MAP 2403; \data\gis\ERA\Phase 2





SQS and CSL categories for baseline surface sediment sampling locations

Hexachlorobenzene

- > CSL, detect
- > SQS and ≤ CSL, detect
- > CSL, non-detect
- > SQS and ≤ CSL, non-detect
- ≤ SQS, detect
- ≤ SQS, non-detect
- Navigation channel
- River mile location

Wind Ward environmental LLC

0 0.2 0.4 Miles

0 0.2 0.4 Kilometers

Scale is the same for each inset map

Map A.6-8. Hexachlorobenzene detected concentrations and RLs compared to SQS and CSL chemical criteria at baseline surface sediment sampling locations

Prepared by AMW, 06-27-06, mod CEH 08-29-06, STS 12-19-06, CEH 02-05-07, 08/03/07; MAP 2405; \data\gis\ERA\Phase 2



Wind Ward environmental LLC

0 0.1 0.2 Miles
0 0.1 0.2 Kilometers
Scale is the same for each inset map

Map A.6-9. Baseline surface sediment sampling locations with only non-detect RLs (no detected chemicals) greater than SQS or CSL chemical criteria

Prepared by CEH 07/11/06, mod CEH 08/30/06, STS 12-19-06, CEH 02-05-07, 08/03/67; Map 2417; \data\gis\ERA\Phase 2





Wind Ward LLC environmental

N

0 100 200 Feet

0 20 40 Meters

Scale is the same for each inset map

Map A.6-10. Areas with total PCB concentration greater than SQS and CSL, as represented by Thiessen polygons versus IDW interpolation

Prepared by CEH, 12/18/2006, mod 02/05/07, MAP 2719, W:\Projects\00-08-06_Duwamish_RI\data\gis\ERA\Phase 2

ATTACHMENT 1
BENTHIC INVERTEBRATE SPECIES IDENTIFIED IN THE LOWER DUWAMISH WATERWAY

Attachment 1 Benthic Invertebrate Species Identified in the Lower Duwamish Waterway

Note: The species names listed in the table below are as reported in the references. Because of revisions to the benthic invertebrate nomenclature over the time span of the references (1978 to 2005), the same species may be listed under different names.

Phylum	Class	Order	Family	Genus and Species
Cnidaria				
	Hydrozoa			
		Hydroida		
			Tubulariidae	
				Euphysa sp.
			Sertulariidae	
				Hydrallmania distans
		Nynantheae spp.		
	Anthozoa (sea anemones)			
		Actiniaria		
			Edwardsiidae	
				Edwardsia sp.
				Edwardsia californica
				Edwardsia callimorpha
				Edwardsia leidya
				Edwardsia sipunculoids
			Metridiidae	
				Metridium senile
		Pennatulacea		
			Pennatulidae	
				Ptilosarcus gurneyi
Platyhelminthes				
	Turbellaria (flatworms)			
		Polycladida		
			Leptoplanidae	
			Stylochidae	
				Kaburakia excelsa
				Rhabdocoela sp.
Nemertea (proboscis worms)				
	Anopla			
		Heteronemertea		
			Lineidae	
				Cerebratulus californiensis
				Cerebratulus sp.
		Palaeonemertea		
			Carinomidae	
				Carinoma mutabilis
			Tubulanidae	
				Tubulanus cingulatus
				Tubulanus pellucidus/polymorphus
				Tubulanus sp.
	Enopla			
		Hoplonemertea		
			Emplectonematidae	
				Paranemertes californica
				Paranemertes peregrina
Nematoda				
Annelida (segmented worms)				
	Archianellida			
	Oligochaeta			
			Megascolecidae	
				Enchytraeus sp.
			Naididae	
				Paranais sp.
	Polychaeta			
			Ampharetidae	
				Ampharete lobrops
				Ampharete cf crassiseta
				Amphicteis sp.
				Amphicteis scaphobranchiata
				Asabellides sibirica
				Asabellides lineata
				Pseudoamphicteis sp.
				Hobsonia florida
			Arabellidae	
			Arenicolidae	
				Abarenicola pacifica
			Autolytinae	
			Capitellidae	
				Barantolla nr americana
				Capitella capitata
				Heteromastus filiformis
				Heteromastus filobranchus
				Heteromastus sp.
				Mediomastus sp
				Notomastus hemipodus

Phylum	Class	Order	Family	Genus and Species
				Nodomastus sp.
			Chaetopteridae	
				Mesochaetopterus taylori
				Phyllochaetopterus prolifica
				Spiochaetopterus pottsi
			Chrysopetalidae	
				Paleanotus bellis
			Cirratulidae	
				Aphelochaeta sp.
				Aphelochaeta cf glandaria
				Aphelochaeta monilaris
				Caulleriella pacifica
				Chaetozone acuta
				Chaetozone setosa
				Chaetozone nr setosa
				Chaetozone sp. N1
				Chaetozone sp.
				Cirratulus sp.
				Monticellina secunda
				Monticellina serratiseta
				Monticellina sp.
				Tharyx multifilis
			Cossuridae	
				Cossura sp.
				Cossura pygodactylata
			Dorvilleidae	
				Dorvillea (Schistomeringos) annulata
				Protodorvillea gracilis
			Eunicidae	
			Glyceridae	
				Glycera americana
				Glycera nana
				Glycera capitata
			Goniadidae	
				Glycinde armigera
				Glycinde picta
				Glycinde polygnatha
				Goniada sp.
				Goniada maculate
			Hesionidae	
				Micropodarke dubia
				Ophiodromus pugettensis
				Podarkeopsis glabra
			Lumbrineridae	
				Lumbrineris californiensis
				Lumbrineris luti
				Scoletoma luti
			Maldanidae	

Phylum	Class	Order	Family	Genus and Species
				Euclymene zonalis
				Euclymeninae sp.
				Praxillella gracilis
				Rhodine bitorquata
			Nephtyidae	
				Nephtys sp.
				Nephtys cornuta
				Nephtys ferruginea
			Nereidae	
				Neanthes limnicola
				Neanthes sp.
				Nereis procera
				Nereis vexillosa
				Nereis sp.
				Platyneris bicanaliculata
			Onuphidae	
				Onuphis iridescens
			Opheliidae	
				Ammotrypane sp.
				Ammotrypane aulogaster
				Armandia brevis
				Ophelina acuminata
			Orbiniidae	
				Leitoscoloplos pugettensis
				Levinsenia gracilis
				Phylo felix
				Scoloplos acmeceps
				Scoloplos sp.
			Paraonidae	
				Aricidea lopezi
			Pectinariidae	
				Pectinaria californiensis
				Pectinaria granulata
			Pholoidae	
				Pholoides asperus
			Phyllodocidae	
				Anaitides sp.
				Eteone californica
				Eteone longa
				Eteone spilotus
				Eteone sp.
				Eumida longicornuta
				Phyllodoce groenlandica
				Phyllodoce hartmanae
				Phyllodoce longipes
				Phyllodoce sp.
			Pilargiidae	
				Pilargus maculata

Phylum	Class	Order	Family	Genus and Species
			Polynoidae	
				Gatyana treadwelli
				Harmothoe imbricata
				Harmothoe sp.
				Malmgreniella macginitiei
				Malmgreniella sp.
				Tenonia priops
			Sabellariidae	
				Neosabellaria cementarium
			Sabellidae	
				Sabella sp.
				Manayunkia aestuarina
				Euchone incolor
				Euchone limnicola
				Fabricia pacifica
				Fabricia sp.
			Sigalionidae	
				Pholoe minuta
				Pholoe sp. N1
				Pholoe sp
				Sthenelais berkeleyi
			Sphaerodoridae	
				Sphaerodoropsis sphaerulifer
			Spionidae	
				Dipolydora caulleryi
				Dipolydora socialis
				Dipolydora sp.
				Laonice sp.
				Polydora uncata
				Polydora cornuta
				Polydora cardilia
				Polydora quadrilobata
				Polydora sp.
				Prionospio lighti
				Prionospio multibranchiata
				Prionospio steenstrupi
				Prionospio sp.
				Prionospio jubata
				Paraprionospio pinnata
				Pseudopolydora kempi japonica
				Pseudopolydora kempi
				Pseudopolydora paucibranchiata
				Pygospio elegans
				Pygospio sp.
				Spio filicornis
				Spiophanes berkeleyorum
			Syllidae	
				Exogone lourei
				Exogone molestra
				Sphaerosyllis ranunculus
				Typosyllis cornuta
				Typosyllis heterochaeta
				Typosyllis hyperioni
			Terebellidae	
				Amphitrite cirrata
				Artacama coniferi
				Eupolymnia heterobranchia
				Lanassa venusta
				Lanassa venusta venustai
				Pista nr brevibranchiata
				Polycirrus sp. 1
				Polycirrus sp.
				Streblosoma bairdi
				Terebellides californica
				Terebellides sp.
Mollusca				
	Bivalvia			
		Myoida		
			Hiatellidae	
				Hiatella arctica
			Cuspidariidae	
				Cardiomya pectinata
			Myidae	
				Cryptomya californica
				Mya arenaria
		Mytiloida		
			Mytilidae	
				Megacrenella Columbiana
				Modiolus modiolus
				Mytilus edulis
				Mytilus sp.
				Solamen columbianum
		Nuculoidea		
			Nuculidae	
				Ennucula tenius
				Nucula tenuis
			Nuculanidae	
				Nuculana minuta

Phylum	Class	Order	Family	Genus and Species
		Ostreoida		
			Anomiidae	
				Pododesmus cepio
				Pododesmus macrochisma
			Pectinidae	
				Chlamys hastata
		Pholadomyoida		
			Lyonsiidae	
				Lyonsia californica
			Pandoridae	
				Pandora filosa
				Pandora sp.
			Thraciidae	
				Thracia trapezoides
		Veneroida		
			Cardiidae	
				Clinocardium sp.
				Clinocardium nuttali
			Kelliidae	
				Odontogena borealis
			Lasaeidae	
				Rochefortia tumida
			Lucinidae	
				Lucinoma acutlineata
				Lucinoma annulatum
				Parvilucina tenuisculpta
			Mactridae	
				Simomactra falcata
			Montacutidae	
				Mysella tumida
				Mysella sp.
			Semelidae	
				Semele rubropicta
			Solenidae	
				Solen sicarius
			Tellinidae	
				Macoma balthica
				Macoma carlottensis
				Macoma elimata
				Macoma expansa
				Macoma golikovi
				Macoma incongrua
				Macoma inquinata
				Macoma nasuta
				Macoma yoldiformis
				Macoma sp.
				Tellina modesta
				Tellina sp.

Phylum	Class	Order	Family	Genus and Species
			Thyasirdae	
				Axinopsida serricata
				Thyrasira flexuosa
			Veneridae	
				Compsomyax subdiaphana
				Nutricola lordi
				Protothaca staminea
				Psephidia lordii
				Saxidomus giganteus
				Transennella tantilla
	Gastropoda (snails)			
		Heterostropha		
			Pyramidellidae	
				Turbonilla sp.
		Mesogastropoda		
			Epitoniidae	
				Epitonium sp.
			Melanellidae	
				Melanella sp.
			Rissoidae	
				Alvania compacta
				Barleeia sp.
			Turritellidae	
				Tachyrhynchus sp.
		Neogastropoda		
			Nassariidae	
				Nassarius mendicus
				Nassarius sp.
			Columbellidae	
				Alia carinata
				Mitrella gouldii alias *Astyris gausapata*
				Nitidella gouldi
			Conidae	
				Kurtzia arteaga
			Muricidae	
				Nucella lamellosa
			Olividae	
				Olivella baetica
		Neotaenioglassa		
			Eulimidae	
				Polygireulima rutila
			Littorinidae	
				Lacuna vincta
			Naticidae	
				Polinices lewisii
		Systellommatopora		
			Onchidiidae	

Phylum	Class	Order	Family	Genus and Species
				Onchidella sp.
	Opisthobranchia (subclass)			
			Cylichnidae	
				Cylichna attonsa
			Pyramidellidae	
				Odostomia sp.
		Nudibranchia		
			Aeolidacea	
			Arminidae	
				Armina californica
		Cephalaspidea		
			Gastropteridae	
				Gastropteron pacificum
			Doridiidae	
				Melanochlamys diomedea
		Pteropoda		
	Aplacaphora			
			Chaetodermatidae	
				Chaetoderma sp.
Arthropoda				
	Arachnida			
		Acari		
			Halacaridae	
	Crustacea			
		Amphipoda		
			Protellidae	
				Tritella pilimana
			Pleustidae	
				Incisocalliope sp.
				Eochelidium miraculum
				Eochelidium sp.
				Chromopleustes oculatus
			Aoridae	
				Aoroides inermis
				Aoroides sp.
				Grandidierella japonica
			Ampeliscidae	
				Byblis millsi
			Ampithoidae	
				Ampithoe lacertosa
				Ampithoe valida
				Ampithoe sp.
			Anisogammaridae	
				Eogammarus confervicolus
				Anisogammarus confervicolus
				Anisogammarus pugettensis
				Anisogammarus sp.
			Caprellidae	

Phylum	Class	Order	Family	Genus and Species
			Corophiidae	
				Corophium acherrusicum
				Corophium salmonis
				Corophium spinicorne
				Corophium insidiosum
				Corophium sp.
			Eusiridae	
				Paramoera sp.
			Isaeidae	
				Photis brevipes
				Photis sp.
			Ischyroceridae	
				Protomedeia prudens
				Protomedeia sp.
			Lysianassidae	
				Hippomedon caecus
				Orchomene decipiens
			Melitidae	
				Desdimelita desdichada
				Melita desdichada
				Melita sp.
			Oedicerotidae	
				Americhelidium shoemakeri
				Monoculoides sp.
				Westwoodilla caecula
			Podoceridae	
				Dyopedos sp.
		Cladocera		
			Podonidae	
				Podon leuckarti
		Euphausiacea		
			Euphausid	
		Isopoda		
			Idoteidae	
				Idotea sp.
			Paramunnidae	
				Munnogonium sp.
				Munnogonium tillerae
			Pleurogoniidae	
				Pleurogonium rubricundum
			Sphaeromatidae	
				Gnorimosphaeroma insulare
				Gnorimosphaeroma oregonesis
			Epicaridea	
		Cumacea		
			Diastylidae	
				Diastylis santamariensis
			Lampropidae	

Phylum	Class	Order	Family	Genus and Species
				Lamprops quadriplicata
			Nannastacidae	
				Cumella vulgaris
			Leuconidae	
				Eudorella pacifica
				Nippoleucon hinumensis
		Tanaidacea		
			Leptocheliidae	
				Leptochelia sp.
				Leptochelia dubia
				Leptochelia savignyi
			Tanaidae	
				Sinelobus stanfordi
				Tanais sp.
		Mysidacea		
			Mysidae	
				Neomysis mercedis
				Alienacanthomysis macropsis
		Decapoda		
			Cancridae	
				Cancer gracilis
				Cancer magister
				Cancer oregonensis
				Cancer productus
			Callianassidae	
				Neotrypaea sp.
			Crangonidae	
				Crangon sp.
				Crangon alaskensis
				Crangon franciscorum
				Mesocrangon munitella
			Epialtidae	
				Pugettia producta
			Hippolytidae	
				Eualus pusiolus
				Eualus sp.
				Spirontocaris prionata
			Paguridae	
				Elassochirus sp.
				Paragus caurinus
				Paragus sp.
			Pandalidae	
				Pandalus danae
			Panopeidae	
				Lophopanopeus bellus
			Pinnotheridae	
				Pinnixa schmitti
				Pinnixa sp.

Phylum	Class	Order	Family	Genus and Species
			Pisidae	
				Loxorhynchus crispatus
		Thoracica		
			Balanomorpha (suborder)	
			Balanidae	
				Balanus crenatus
				Balanus sp.
	Copepoda (subclass)			
		Harpacticoida		
			Ancorabolidae	
			Ameiridae	
				Ameira sp.
				Nitocra sp. A
				Nitocra sp.
			Canthocamptidae	
				Leimia vaga
				Cletocamptus sp.
				Mesochra sp.
				Mesochra rapines
			Canuellidae	
				Coullana canadensis
			Cletodidae	
				Acrenhydrosoma sp.
				Enhydrosoma sp.
			Cylindropsyllidae	
			Darcythompsoniidae	
			Diosaccidae	
				Amphiascopsis cinctus
				Amphiascopsis sp.
				Amphiascoides sp.
				Amonardia perturbata
				Amonardia normani
				Diosaccus sp.
				Diosaccus spinatus
				Bulbamphiascus sp.
				Robertsonia sp.
				Typhlamphiascus pectinifer
				Typhlamphiascus sp.
				Stenhelia asetosa
				Stenhelia peniculata
				Stenhelia sp.
				Schizopera knabi
				Schizopera sp.
			Ectinosomatidae	
				Pseudobradya sp.
			Harpacticidae	
				Harpacticus uniremis
				Harpacticus sp.

Phylum	Class	Order	Family	Genus and Species
				Harpacticus compressus
				Harpacticus obscurus
				Harpacticus spinulosus
				Harpacticus arcticus
				Zaus sp.
			Huntemanniidea	
				Nannopus palustris
				Huntemannia jadensis
			Laophontidae	
				Heterolaophonte discophora
				Heterolaophonte longisetigera
				Heterolaophonte hamondi
				Laophonte sp.
				Laophonte cornuta
				Laophonte elongata
				Echinolaophontes sp.
				Onychocamptus mohammed
				Paralaophonte sp.
				Paralaophonte pacifica
				Paralaophonte perplexa
				Pseudonychocamptus sp.
			Longipediidae	
				Longipedia sp.
			Normanellidae	
				Normanella sp.
			Orthopsyllidae	
				Orthopsyllus illgi
			Paramesochridae	
				Apodopsyllus sp.
			Parastenheliidae	
				Parastenhelia hornelli
				Parastenhelia spinosa
			Peltidiidae	
			Tachidiidae	
				Microarthridion littorale
				Tachidius disciples
				Tachidius triangularis
			Tegastidae	
			Thalestridae	
				Dactylopodia crassipes
				Dactylopodia vulgaris
				Dactylopodia tisboides
				Dactylopodia paratisboides
				Dactylopodia glacialis
				Diarthrodes sp.
				Idomene sp.
				Paradactylopodia sp.
				Parathalestris sp.

Phylum	Class	Order	Family	Genus and Species
				Rhynchothalestris helgolandica
			Tisbidae	
				Scutellidium sp.
				Tisbe sp.
		Cyclopoida		
			Cyclopoidae	
				Halicyclops sp.
			Oithonidae	
				Oithona similis
				Oithona longirastris
		Calanoida		
			Temoridae	
				Eurytemora sp.
				Eurytemora americana
			Centropagidae	
				Centropages abdominalis
			Pseudodiaptomidae	
				Pseudodiaptomus marinus
			Stephidae	
				Stephos sp.
			Calanidae	
				Calanus sp.
			Paracalanidae	
				Paracalanidae sp.
			Clausocalanidae	
				Microcalanus sp.
				Pseudocalanus sp.
			Acartiidae	
				Acartia sp.
				Acartia longiremis
		Poecilostomatoida		
			Corycaeidae	
				Corycaeus anglicus
			Clausidiidae	
				Hemicyclops sp.
			Ergasilidae	
			Oncaeidae	
				Oncaea sp.
	Ostracoda			
		Myodocopida		
			Cylindroleberididae	
			Philomedidae	
				Euphilomedes carcharodonta
				Euphilomedes producta
		Podocopida		
	Insecta (larvae)			
		Acarina		
		Ceratopogonidae		

Phylum	Class	Order	Family	Genus and Species
		Coleoptera		
		Diptera (pupa)		
			Dolichopodidae (larvae)	
			Chironomidae (larvae)	
		Empididae		
		Collembola		
		Trichoptera		
		Thysanoptera		
Sipuncula				
			Golfingiidae	
				Thysanocardia nigra
Bryozoa				
		Cheilostomata		
			Bugulidae	
			Hippothoidae	
				Celleporella hyalina
		Ctenostomata		
			Alcyonidiidae	
				Alcyonidium sp.
		Pedicellinida		
			Pedicellinidae	
				Barentsia benedeni
Echinodermata				
	Asteroidea			
		Spinulosida		
			Solasteridae	
				Solaster stimpsoni
		Forcipulatida		
			Asteriidae	
				Evasterias troschelii
				Pycnopodia helianthoides

Phylum	Class	Order	Family	Genus and Species
				Pisaster sp.
			Luidiidae	
				Luidia sp.
	Stelleroidea			
		Ophiurida		
			Amphiuridae	
				Amphiodia sp.
				Amphiodia digitata
				Amphiodia (Amphispina) urtica
				Amphipholis squamata
	Holothuroidea			
		Dendrochirotida		
			Cucumariidae	
				Pentamera lissoplaca
				Pentamera trachyplaca
				Pentamera populifera
				Pentamera sp.
Cephalorhyncha				
	Priapulida			
			Priapuloidae	
				Priapulus caudatus
Rhizopoda				
	Rhizopodea			
		Foraminiferida		
Rotifera				

Sources: Bingham (1978); Leon (1980); Williams (1990); Cordell et al. (1996; Cordell et al. 1997); Taylor et al. (1999); Striplin Environmental Associates (1998); Windward (2005)

REFERENCES

Bingham CR. 1978. Aquatic disposal field investigations, Duwamish Waterway, disposal site Puget Sound, Washington. Appendix G: Benthic community structural changes resulting from dredged material disposal, Elliott Bay disposal site. Technical report D-77-24. U. S. Army Engineer Waterways Experiment Station, Vicksburg, MS.

Cordell JR, Tear LM, Simenstad CA, Hood WG. 1996. Duwamish river coastal America restoration and reference sites: Results from 1995 monitoring studies. Fish Research Institute, University of Washington, Seattle, WA.

Cordell JR, Tear LM, Jensen K, Luiting V. 1997. Duwamish river coastal America restoration and reference sites: Results from 1996 monitoring studies. Fisheries Research Institute, University of Washington, Seattle, WA.

Leon H. 1980. Final Report: Terminal 107 environmental studies. Benthic community impact study for terminal 107 (Kellogg Island) and vicinity. Prepared for Port of Seattle Planning and Research Department. Pacific Rim Planners, Inc., Seattle, WA.

SEA. 1998. Benthic infaunal communities in the vicinity of the Duwamish Diagonal combined sewer overflow. Final report. Prepared for King County Water Quality Division, Seattle, WA. Striplin Environmental Associates, Inc., Olympia, WA.

Taylor WJ, Shreffler DK, Cordell JR. 1999. Duwamish East Waterway channel deepening project: alternative dredge disposal sites juvenile salmonid and epibenthic prey assessment. Technical report. Preliminary draft. Prepared for Port of Seattle. Taylor Associates, Seattle, WA.

Williams MS. 1990. Port of Seattle Terminal 107 (Kellogg Island), biological assessment - 1989. Parametrix, Inc, Bellevue, WA.

Windward. 2005. Lower Duwamish Waterway remedial investigation. Data report: Taxonomic identifications of benthic invertebrate communities. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

ATTACHMENT 2
FISH SPECIES IDENTIFIED IN THE LOWER DUWAMISH WATERWAY

Attachment 2 Fish Species Identified in the Lower Duwamish Waterway

Common Name	Scientific Name	Family	Abundance[a]	Abundance Citation	Environment	Habitat	E/H Citation	Diet	Diet Citation
American shad	*Alosa sapidissima*	Engraulidae	rare	9, 10, 11, 12, 41	anadromous	bays, estuaries, freshwater	32	plankton, copepods, mysids, small fish	33
Bay goby	*Lepidogobius lepidus*	Gobiidae	rare	2, 3, 6	marine (estuary)	benthic (mud bottom)	13	benthic organisms	28
Bay pipefish	*Syngnathus grisiolineatum*	Syngnathidae	common	11	marine	demersal (associated with eel grass in the intertidal areas)	15	isopods, amphipods	14
			rare	6, 10					
Big skate	*Raja binoculata*	Rajidae	rare	7, 11	marine	benthic (sandy and gravelly bottoms)	16	crustaceans, fish	14
Blackbelly eelpout	*Lycodopsis pacifica*	Zoarcidae	rare	11	marine	over soft bottoms	32	worms, crustaceans, small bivalves, brittle stars	34
Bluegill	*Lepomis macrochirus*	Centrarchidae	rare	41	freshwater	weed beds	21	snails, worms, small crayfish, insects	21
Brown rockfish	*Sebastes auriculatus*	Scorpaenidae	rare	11, 12	marine	shallow, low-profile, rocky reefs	32	finfish, benthic crustaceans, fish eggs, larvae	35
Buffalo sculpin	*Enophrys bison*	Cottidae	rare	1, 2, 3, 4, 7, 11, 12	marine (estuary)	benthic (inshore rocky and sandy areas)	13	mainly algae, also amphipods, small fishes, crabs, polychaetes, nudibranchs, isopods	13, 29
Bull trout	*Salvelinus confluentes*	Salmonidae	rare	6, 9	anadromous	benthopelagic (near shore)	21	mainly fish, plus zooplankton	31
Butter sole	*Isopsetta isolepis*	Pleuronectidae	common	6	marine (estuary)	benthic (sandy bottom)	13	worms, fish, shrimps	14
			rare	7					
Chinook salmon[b]	*Oncorhynchus tshawytscha*	Salmonidae	abundant	1, 4, 5, 6, 9, 10, 41	anadromous	benthopelagic	27	juveniles: insects, epibenthic crustaceans, pelagic organisms	30
			rare	2					
Chum salmon	*Oncorhynchus keta*	Salmonidae	abundant	5, 6, 9, 41	anadromous	benthopelagic	27	juveniles: copepods, amphipods, cumaceans, euphausiids	29
			common	10					
			rare	1, 4					

Common Name	Scientific Name	Family	Abundance[a]	Abundance Citation	Environment	Habitat	E/H Citation	Diet	Diet Citation
C-O sole	*Pleuronichthys coenosus*	Pleuronectidae	rare	7, 11	marine	benthic (flat bottoms, rocky areas)	13	isopods, fish, polychaetes, amphipods, turbellarians, bivalves	29
Coho salmon[b]	*Oncorhynchus kisutch*	Salmonidae	abundant	6, 9, 10	anadromous	benthopelagic	27	juveniles: insects, epibenthic crustaceans, pelagic organisms, small fish	29
			common	4, 10, 41					
			rare	1, 2					
Crescent gunnel	*Pholis laeta*	Pholidae	rare	6, 9, 11, 41	marine (estuary)	demersal (intertidal areas, under rocks)	13	gammarid amphipods, copepods, tanaids, isopods	29
Cutthroat trout	*Oncorhynchus clarki*	Salmonidae	rare	1, 4, 5, 6, 9, 10, 41	anadromous	benthopelagic	22	fish, epibenthic crustaceans, pelagic organisms, insects	18
Dolly Varden	*Salvelinus malma*	Salmonidae	rare	1, 4	freshwater	benthopelagic	21	fish, epibenthic crustaceans, pelagic organisms, insects	14
Dover sole	*Microstomus pacificus*	Pleuronectidae	common	2, 11	marine	benthic (mud bottom)	13	benthic invertebrates, echinoderms, mollusks, polychaetes	24
			rare	3					
English sole	*Parophrys vetulus*	Pleuronectidae	abundant	2, 3, 4, 7, 11, 12	marine (estuary)	benthic (sand and mud bottoms)	18	cumaceans, gammarid amphipods, polychaetes, tanaids, crabs, bivalves	29
			rare	1, 6					
Eulachon	*Thaleichthys pacificus*	Osmeridae	rare	3	anadromous	pelagic	13	plankton (only feeds while at sea)	20
Flathead sole	*Hippoglossoides elassodon*	Pleuronectidae	rare	2, 11, 12	marine	benthic (soft mud bottom, adults below 180 m)	13	polychaetes, cumaceans, gammarid amphipods, isopods, bivalves	29
Gunnel sp.	*Apodichthys sp.*	Pholidae	rare	10	marine	intertidal zone among rocks and shallow eelgrass beds	32	small crustaceans, mollusks	13
Great sculpin	*Myoxocephalus polyacanthocephalus*	Cottidae	rare	11	marine	intertidal areas, sand and mud bottoms	13	small fish	13

Common Name	Scientific Name	Family	Abundance[a]	Abundance Citation	Environment	Habitat	E/H Citation	Diet	Diet Citation
Hybrid sole	*Inopsetta Isopsetta ischyra*	Pleuronectidae	rare	1, 12	marine (estuary)	benthic	13	benthic organisms	14
Kelp perch	*Brachyistius frenatus*	Embiotocidae	rare	9	marine	among fronds in kelp beds from near surface to depths of about 30 m	32	small crustaceans, parasites	13
Largescale sucker	*Catostomus macrocheilus*	Catostomidae	rare	1, 2, 4, 6, 41	freshwater	demersal	21	algae, diatoms, insects, amphipods, and mollusks	20
Longfin sculpin	*Jordania zonope*	Cottidae	rare	11	marine	demersal, intertidal areas, rocky areas and kelp	13	amphipods, benthic copepods, crabs, shrimp, gastropods, polychaetes	38
Longfin smelt	*Spirinchus thaleichthys*	Osmeridae	abundant	1, 2, 11	anadromous	benthopelagic (close to shore, in bays and estuaries)	21	crab larvae, copepods, mysid shrimp	29
			common	12					
			rare	7, 9					
Longnose dace	*Rhinichthys cataractae*	Cyprinidae	rare	6	freshwater	demersal	21	mayflies, blackflies, and midges	20
Longnose skate	*Raja rhina*	Rajidae	rare	11	marine	partially or entirely buried in sand or silt bottoms	36	small fish, crustaceans, worms, mollusks	36
Mountain whitefish	*Prosopium williamsoni*	Salmonidae	rare	1, 6, 9	freshwater	benthopelagic	14	insects, invertebrates, eggs, small fish	14
Northern pikeminnow	*Ptychocheilus oregonensus*	Cyprinidae	rare	1, 6	freshwater	benthopelagic	20	insects, fish	20
Northern ronquil	*Ronquilus jordani*	Bathymasteridae	rare	11	marine	demersal	13	polychaetes, plankton, invertebrates, cladocerans, copepods	14
Northern sculpin	*Icelinus borealis*	Cottidae	rare	6	marine	demersal	13	benthic crustaceans, shrimps/prawns	14, 29
Pacific cod	*Gadus macrocephalus*	Gadidae	rare	2, 3, 4	marine	(demersal, continental shelf and upper slopes)	23	fish, octopi, large crustaceans, worms, amphipods	26, 29
Pacific herring	*Clupea pallasi*	Clupeidae	abundant	4, 9, 11	marine	benthopelagic (coastal, first year in bays)	14	planktonic crustaceans, fish larvae	14, 29
			common	1, 2, 7, 12					
			rare	6, 10					

Common Name	Scientific Name	Family	Abundance[a]	Abundance Citation	Environment	Habitat	E/H Citation	Diet	Diet Citation
Pacific sand dab	*Citharichthys sordidus*	Paralichthyidae	common	11	marine	over soft sand bottoms	13	benthic crustaceans, worms	24
			rare	12					
Pacific sandlance	*Ammodytes hexapterus*	Ammodytidae	abundant	6, 9, 41	marine (brackish)	benthopelagic (surface or burrowed in sand)	13	zooplankton	17, 29
			common	4					
			rare	1, 10, 11					
Pacific staghorn sculpin	*Leptocottus armatus*	Cottidae	abundant	1, 2, 3, 4, 6, 9, 10, 11, 12	marine (lower estuary, offshore)	benthic (sandy bottom)	13	isopods, bivalve siphons, polychaetes, crabs, fish, tanaids, shrimp	19
			common	7					
Pacific tomcod	*Microgadus proximus*	Gadidae	abundant (juveniles)	7, 11	marine (brackish)	benthic (over sand)	23	shrimps, amphipods, isopods, gastropods, mussels, fishes	24
			common	2, 3, 12					
			rare	1, 4					
Padded sculpin	*Artedius fenestralis*	Cottidae	common	2, 3	marine	benthic	13	gammarid amphipods, isopods, tanaids, shrimp, copepods, small fish	18, 29
			rare	7, 12					
Peamouth chub	*Mylocheilus caurinus*	Cyprinidae	rare	9, 41	freshwater	demersal (brackish)	21	aquatic insects, larvae, terrestrial insects, crustaceans, mollusks, small fish	21
Penpoint gunnel	*Apodichthys flavidus*	Pholidae	rare	5, 6, 9	marine (estuary)	demersal (intertidal tide pools)	13	isopods, amphipods, shrimp, gastropods, other epibenthic crustaceans	29
Pile perch	*Rhacochilus vacca*	Embiotocidae	abundant	12	marine	demersal (rocky shores; near kelp, pilings, underwater structures)	13	isopods, bivalves, crabs, amphipods	29
			common	4, 7, 11					
			rare	1, 2, 3, 6, 9, 41					
Pink salmon[b]	*Oncorhynchus gorbuscha*	Salmonidae	rare	6, 41	anadromous	benthopelagic	27	juveniles: copepods, amphipods, barnacle larvae, cumaceans	27, 28
Plainfin midshipman	*Porichthys notatus*	Batrachoididae	common	11	marine	benthic (nearshore shelf, sand/mud bottom)	18	crustaceans, fish	14
			rare	2					

Common Name	Scientific Name	Family	Abundance[a]	Abundance Citation	Environment	Habitat	E/H Citation	Diet	Diet Citation
Prickly sculpin	*Cottus asper*	Cottidae	common	12, 41	marine	benthic	13	benthic organisms	20
			rare	1, 2, 3, 4, 6, 9, 11					
Pygmy poacher	*Odontopyxis trispinosa*	Agonidae	rare	2, 3, 7, 11	marine	demersal (soft bottoms)	13	epibenthic invertebrates	14
Ratfish	*Hydrolagus colliei*	Chimeridae	rare	2, 7, 11	marine	demersal (sandy bottom)	13	worms, bivalves, crustaceans, fishes	17, 29
Redsided shiner	*Richardsonius balteatus*	Cyprinidae	common	6	freshwater	demersal	20	zooplankton, algae, insects	20
Rex sole	*Errex zachirus*	Pleuronectidae	rare	11	marine	demersal	37	worms, benthic crustaceans, mollusks	24
River lamprey	*Lampetra ayresi*	Petromyzontidae	rare	1, 4, 6, 9	anadromous	demersal	14	adult: fish juveniles: detritus, algae	20
Rock sole	*Lepidopsetta bilineata*	Pleuronectidae	abundant	7,11	marine (estuary)	benthic (more pebbly bottom than most other flatfish)	13	isopods, gammarid amphipods, polychaetes, cumaceans, bivalves, crabs, fish	29
			common	2, 3, 12					
Rockfish	*Sebastes* spp.	Scorpaenidae	rare	1, 8	marine	demersal (near structure)	25	crabs, gammarid amphipods, mysids, shrimp, fish	26
Roughback sculpin	*Chitonotus pugeteneis*	Cottidae	common	11,12	marine	benthic (sand/mud bottom)	13	shrimps and other crustaceans	18
			rare	2, 3, 7					
Saddleback gunnel	*Pholis ornata*	Pholidae	rare	3, 5, 6, 9, 11, 12, 41	marine (estuary)	demersal (sandy bottom)	13	amphipods, isopods, polychaetes, copepods, cumaceans	29
Sand sole	*Psettichthys melanostictus*	Pleuronectidae	common	1, 2, 3, 7, 11, 12	marine, estuary	benthic (sandy bottom)	14	fishes, worms, crustaceans, and mollusks	14, 29
			rare	1					
Sailfin sculpin	*Nautichthys oculofasciatus*	Hemitripteridae	rare	11	marine	over rocks from inshore to depths of 110 m, often with algae	32	finfish, benthic crustaceans	19
Sharpnose sculpin	*Clinocottus acuticeps*	Cottidae	rare	6	marine	benthic (sand/vegetation)	13	benthic organisms	22

Common Name	Scientific Name	Family	Abundance[a]	Abundance Citation	Environment	Habitat	E/H Citation	Diet	Diet Citation
Shiner surfperch	*Cymatogaster aggregata*	Embiotocidae	abundant	1, 4, 5, 6, 7, 9, 10, 11, 12, 41	marine (estuary)	demersal (in shallow water, around eelgrass beds, piers and pilings commonly in bays and quiet back waters)	13	amphipods, cumaceans, polychaetes, copepods, isopods, algae	22, 29
			common	2, 3					
Slender sole	*Lyopsetta exilis*	Pleuronectidae	rare	3, 11	marine	benthic (> 200 m depth)	13	carnivore	24
Snake prickleback	*Lumpenus saggita*	Stichaeidae	abundant	1, 2, 3, 4, 6	marine	benthopelagic (shallow bays and offshore waters)	13	bivalves, marine worms, amphipods	29
			common	9, 10, 11, 12					
			rare	7, 41					
Sockeye salmon[b]	*Oncorhynchus nerka*	Salmonidae	rare	40, 41	anadromous	benthopelagic	27	juveniles: insects, epibenthic crustaceans, pelagic organisms	28
Soft sculpin	*Gilbertidia sigalutes*	Cottidae	rare	4	marine	demersal	13	epibenthic crustaceans, phytoplankton, fish eggs/larvae	14
Speckled sanddab	*Citharichthys stigmaeus*	Bothidae	rare	7, 9, 11	marine	benthic (sandy bottom)	13	crustaceans, fish	19
Spiny dogfish	*Squalus acanthias*	Squalidae	rare	2, 11	marine	benthopelagic	26	primarily fish	27
Starry flounder	*Platichthys stellatus*	Pleuronectidae	abundant	1, 2, 3, 4, 6, 7, 9, 10, 11, 12, 41	marine (estuary, brackish)	benthic	22	isopods, fish, gammarid amphipods, polychaetes, gastropods, worms	14
			common	5					
Steelhead[b]	*Oncorhynchus mykiss*	Salmonidae	common	9, 10	anadromous	benthopelagic	39	juveniles: insects, epibenthic crustaceans, pelagic organisms	29
			rare	1, 4, 5, 6, 11, 41					
Striped seaperch	*Embiotoca lateralis*	Embiotocidae	common	1, 4, 12	marine	demersal	13	amphipods, isopods, crabs, shrimp	29
			rare	2, 3, 5, 6, 7, 9, 10					
Sturgeon poacher	*Podothecus acipenserinus*	Agonidae	rare	3, 11	marine	demersal (soft bottom)	13	cumaceans, gammarid amphipods, shrimp, copepods, polychaetes, tanaids	29
Surf smelt	*Hypomesus pretiosus*	Osmeridae	abundant	9	marine (brackish)	benthopelagic	22	isopods, cumaceans, larvaceans, copepods,	29
			common	1, 4, 6, 7, 41					

Common Name	Scientific Name	Family	Abundance[a]	Abundance Citation	Environment	Habitat	E/H Citation	Diet	Diet Citation
			rare	11				amphipods	
Three-spine stickleback	*Gasterosteus aculeatus*	Gasterosteidae	abundant	41	marine, anadromous	benthopelagic (in/near vegetation)	21	worms, crustaceans, insects/larvae, small fish	20, 29
			common	1, 5, 6,10, 11					
			rare	4, 12					
Tidepool sculpin	*Oligocottus maculosus*	Cottidae	rare	41	marine	demersal	13	unknown	
Torrent sculpin	*Cottus rhotheus*	Cottidae	rare	11	freshwater	demersal	21	crustaceans, midges and mayflies larvae, minnows	21
Tubesnout poacher	*Pallasina barbata*	Agonidae	rare	3, 11	marine	demersal (eelgrass & seaweeds)	13	amphipods, polychaetes, copepods, mysids	29
Walleye pollock	*Theragra chalcogramma*	Gadidae	rare	1, 2, 4	freshwater	benthopelagic	23	insects, midge larvae, fish	14
Whitespotted greenling	*Hexagrammos stelleri*	Hexagrammidae	common	7	marine (intertidal)	demersal (nearshore, near rocks, pilings and eelgrass beds)	23	gammarid amphipods, shrimp, crabs, fish, polychaetes	29
			rare	2, 11					

[a] Abundance: abundant (numerically dominant); common (occurs in most samples); rare (occurs in few samples). Abundance characterizations reflect data collected by authors in the cited study. These data may reflect sampling gear bias for the species identified.

[b] Adults are found in the LDW only as they migrate to spawning ground upstream of the LDW and include wild and hatchery species.

E/H – environment/habitat

Citations

1. Matsuda et al. (1968)
2. Miller et al. (1975)
3. Miller et al. (1977a)
4. Weitkamp and Campbell (1980)
5. Taylor et al. (1999)
6. Warner and Fritz (1995)
7. West et al. (2001)
8. Malins et al. (1980)
9. Shannon (2006)
10. Windward (2004)
11. Windward (2005)
12. Windward (2006)
13. Eschmeyer et al. (1983)
14. Hart (1973)
15. Dawson (1985)
16. McEachran and Dunn (1998)
17. Armstrong (1996)
18. Clemens and Wilbey (1961)
19. Fitch and Lavenberg (1975)
20. Scott and Crossman (1973)
21. Page and Burr (1991)
22. Morrow (1980)
23. Cohen et al. (1985)
24. Pearcy and Hancock (1978)
25. Lamb and Edgel (1986)
26. Cox and Francis (1997)
27. Groot and Margolis (1998)
28. Grossman (1979)
29. Miller et al. (1977b)
30. Cordell et al. (2001)
31. Rieman and McIntyre (1993)
32. Gilbert and Williams (2002)
33. Whitehead (1985)
34. Anderson (1994)
35. Hobson (2000)
36. Florida Museum of Natural History (2005)
37. Cooper and Chapleau (1998)
38. Demetropoulos et al. (1990)
39. Gall and Crandell (1992)
40. Kerwin and Nelson (2000)
41. Cordell et al. (2006)

REFERENCES

Anderson ME. 1994. Systematics and osteology of the Zoarcidae (Teleostei: Perciformes). Ichthyol Bull JLB Smith Inst Ichthyol 60:1-120.

Armstrong RH. 1996. Alaska's fish. A guide to selected species. Alaska Northwest Books, Portland, OR.

Clemens WA, Wilby GV. 1961. Fishes of the Pacific coast of Canada. 2nd edition. Publication no. 68. Fisheries Resources Board of Canada.

Cooper JA, Chapleau F. 1998. Monophyly and intrarelationships of the family Pleuronectidae (Pleuronectiformes), with a revised classification. Fish Bull 96(4):686-726.

Cordell J, Toft J, Cooksey M, Gray A. 2006. Fish assemblages and patterns of chinook salmon abundance, diet, and growth at restored sites in the Duwamish River. King Conservation District and Salmon Recovery Funding Board, Seattle, WA.

Cordell JR, Tear LM, Jensen K. 2001. Biological monitoring at Duwamish River coastal America restoration and reference sites: A seven-year retrospective. SAFS-UW-0108. Wetlands Ecosystem Team, School of Aquatic and Fisheries Sciences, University of Washington, Seattle, WA.

Cox G, Francis M. 1997. Sharks and rays of New Zealand. Canterbury University Press, Christchurch, New Zealand.

Dawson CE. 1985. Indo-Pacific pipefishes (Red Sea to the Americas). The Gulf Coast Research Laboratory, Ocean Springs, MS.

Demetropoulos CL, Braithwaite LF, Maurer BA, Whiting D. 1990. Foraging and dietary strategies of two sublittoral cottids, *Jordania zonope* and *Artedius harringtoni*. J Fish Biol 37(1):19-32.

Eschmeyer WN, Herald ES, Hammann H. 1983. Pacific coast fishes. Peterson Field Guide Series. Houghton Mifflin, Boston, MA.

Fitch JE, Lavenberg RJ. 1975. Tidepool and nearshore fishes of California. California Natural History Guides no. 38. University of California Press, Berkeley and Los Angeles, CA.

Florida Museum of Natural History. 2005. Biological profiles: longnose skate. Ichthyology at the Florida Museum of Natural History: Education-Biological Profiles [online]. Florida Museum of Natural History, University of Florida, Gainesville, FL. [Cited 26 August 2005]. Available from: www.flmnh.ufl.edu/fish/Gallery/Descript/LongnoseSkate/LongnoseSkate.html.

Gall GAE, Crandell PA. 1992. The rainbow trout. Aquaculture 100:1-10.

Gilbert CR, Williams JD. 2002. National Audobon Society field guide to fishes. Alfred A. Knopf, Inc., New York, NY.

Groot C, Margolis L, eds. 1998. Pacific salmon life histories. UBC Press, Vancouver, BC.

Grossman GD. 1979. Demographic characteristics of an intertidal bay goby (*Lepidogobius lepidus*). Environ Biol Fish 4(3):207-218.

Hart JL. 1973. Pacific fishes of Canada. Fisheries Research Board of Canada, Ottawa, ON.

Hobson ES. 2000. Interannual variation in predation on first-year *Sebastes* spp. by three northern California predators. Fish Bull 99(2):292-302.

Kerwin J, Nelson TS, eds. 2000. Habitat limiting factors and reconnaissance assessment report, Green/Duwamish and Central Puget Sound watersheds (WRIA 9 and Vashon Island). Washington Conservation Commission, Lacey, WA, and King County Department of Natural Resources, Seattle, WA.

Lamb A, Edgell P. 1986. Coastal fishes of the Pacific Northwest. Harbour Publishing Co. Ltd., Madeira Park, BC.

Malins DC, McCain BB, Brown DW, Sparks AK, Hodgins HO. 1980. Chemical contaminants and biological abnormalities in central and southern Puget Sound. NOAA Technical Memorandum OMPA-2. Environmental Conservation Division, National Marine Fisheries Service, Seattle, WA.

Matsuda RI, Isaac GW, Dalseg RD. 1968. Fishes of the Green-Duwamish River. Water Quality Series No. 4. Municipality of Metropolitan Seattle, Seattle, WA.

McEachran JD, Dunn CA. 1998. Phylogenetic analysis of skates, a morphologically conservative clade of elasmobranchs (Chondrichthyes: Rajidae). Copeia 1998(2):271-290.

Miller BS, Wingert RC, Borton SF. 1975. Ecological survey of demersal fishes in the Duwamish River and at West Point 1974. Prepared for Municipality of Metropolitan Seattle. Report no. FRI-UW-7509. Fisheries Research Institute, University of Washington, Seattle, WA.

Miller BS, McCain BB, Wingert RC, Borton SF, Pierce KV, Griggs DT. 1977a. Ecological and disease studies of demersal fishes in Puget Sound near METRO-operated sewage treatment plants and in the Duwamish River. FRI-UW-7721. Fisheries Research Institute, University of Washington, Seattle, WA.

Miller BS, Simenstad CA, Moulton LL, Fresh KL, Funk FC, Karp WA, Borton SF. 1977b. Puget Sound baseline program nearshore fish survey. Final report, July 1974-June 1977. Prepared for Washington Department of Ecology. Fisheries Research Institute, University of Washington, Seattle, WA.

Morrow JE. 1980. The freshwater fishes of Alaska. Animal Resources Ecology Library, University of British Columbia, Vancouver, BC.

Page LM, Burr BM. 1991. A field guide to freshwater fishes of North America north of Mexico. Peterson Field Guides, Houghton Mifflin Company, Boston, MA.

Pearcy WG, Hancock D. 1978. Feeding habits of Dover sole, *Microstomus pacificus*; rex sole, *Glyptocephalus zachirus*; slender sole, *Lyopsetta exilis*; and Pacific sanddab, *Citharichthys sordidus*, in a region of diverse sediments and bathymetry off Oregon. Fish Bull 76(3):641-651.

Rieman BE, McIntyre JD. 1993. Demographic and habitat requirements for conservation of bull trout. USFS Publication IN-GTR-302. US Fish and Wildlife Service, Washington, DC.

Scott WB, Crossman EJ. 1973. Freshwater fishes of Canada. Bulletin 184. Fisheries Research Board of Canada, Ottawa, ON, Canada.

Shannon J. 2006. Personal communication (e-mails to Matt Luxon, Windward Environmental, regarding Duwamish-Elliot beach seine data collected by Taylor Associates in 1998, 2000, and 2002-2003. Biologist, Taylor Associates, Seattle, WA. March 30, 2006.

Taylor WJ, Shreffler DK, Cordell JR. 1999. Duwamish East Waterway channel deepening project: alternative dredge disposal sites juvenile salmonid and epibenthic prey assessment. Technical report. Preliminary draft. Prepared for Port of Seattle. Taylor Associates, Seattle, WA.

Warner EJ, Fritz RL. 1995. The distribution and growth of Green River chinook salmon (*Oncorhynchus tshawytscha*) and chum salmon (*Oncorhynchus keta*) outmigrants in the Duwamish estuary as a function of water quality and substrate. Water Resources Division, Muckleshoot Indian Tribe, Auburn, WA.

Weitkamp DE, Campbell RF. 1980. Port of Seattle Terminal 107 fisheries study. Parametrix, Inc., Bellevue, WA.

West JE, O'Neill SM, Lippert G, Quinnell S. 2001. Toxic contaminants in marine and anadromous fishes from Puget Sound, Washington. Results of the Puget Sound ambient monitoring program fish component 1989-1999. Washington Department of Fish and Wildlife, Olympia, WA.

Whitehead PJP. 1985. Clupeoid fishes of the world (suborder Clupeioidei). An annotated and illustrated catalogue of the herrings, sardines, pilchards, sprats, shads, anchovies and wolf-herrings. Part 1 - Chirocentridae, Clupeidae and Pristigasteridae. FAO species catalogue. FAO Fish Synopses 125, vol 7. Food and Agriculture Organization of the United Nations, Rome, Italy.

Windward. 2004. Lower Duwamish Waterway remedial investigation. Juvenile chinook salmon data report. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005. Lower Duwamish Waterway remedial investigation. Data report: Fish and crab tissue collection and chemical analyses. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2006. Lower Duwamish Waterway remedial investigation. Data report: 2005 fish and crab tissue collection and chemical analyses. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

ATTACHMENT 3
DATA MANAGEMENT RULES

Attachment 3 Data Management Rules

LABORATORY REPLICATES

Chemical concentrations obtained from the analysis of laboratory duplicates or replicates (two or more analyses on the same sample) are averaged for a closer representation of the "true" concentration as compared to the results of a single analysis. Averaging rules are dependent on whether the individual results are detected concentrations or reporting limits (RLs) for undetected analytes. If all concentrations are detects for a given parameter, the values are simply averaged arithmetically. If all concentrations are undetected for a given parameter, the minimum RL is reported. If the concentrations are a mixture of detected concentrations and RLs, any two or more detected concentrations are averaged arithmetically and RLs are ignored. If there is a single detected concentration and one or more RLs, the detected concentration is reported. The latter two rules are applied regardless of whether the RLs are higher or lower than the detected concentration.

LOCATION AVERAGING

The baseline surface sediment dataset contains averaged results of chemical concentrations of discrete samples collected at a single sampling location that were submitted to the laboratory as individual samples and analyzed separately. The averaging rules used for location averaging are the same as for laboratory replicates described above. A sampling location with averaged chemical concentrations is presented as a single "sample" in the ERA text and data tables. This type of averaging is performed in the following instances.

- The chemical concentrations obtained from the analyses of a surface sediment sample, and its field duplicate or replicate, are averaged to obtain a single concentration of the chemical for the sampling location.
- Surface sediment data have been collected repeatedly at certain locations within a six-month period.[1] For these locations that have multiple samples collected at different times, the results of these individual samples are averaged to a single chemical concentration for that location.

[1] An assumption was made in the memorandum describing the baseline surface sediment dataset (Windward 2006) that two or more samples collected from the same location within a six-month period reflect spatial variability, rather than temporal variability. Therefore, these results were averaged together. For multiple samples collected more than six months apart from the same location, such as data collected within monitoring programs, any differences in chemical concentrations also reflect temporal variability. Therefore, only the most recent sample from those locations was included in the baseline surface sediment dataset; no averaging occurred in these situations.

- The baseline surface sediment dataset contains two locations that were re-sampled in a different sampling event, outside of a monitoring program. These sample results were also averaged to obtain a single representative result for that location.

SIGNIFICANT FIGURES AND ROUNDING

The laboratories reported results with different numbers of significant figures depending on the instrument, parameter, and the concentration relative to the reporting limit (RL). The reported (or assessed) precision of each observation is explicitly stored in the project database as a record of the number of significant figures assigned by the laboratory. The tracking of significant figures becomes important when calculating averages and performing other data summaries.

When a calculation involves addition, such as totaling polychlorinated biphenyls (PCBs) or polycyclic aromatic hydrocarbons (PAHs), the calculation can only be as precise as the least precise number that went into the calculation. For example (assuming two significant figures):

> 210 + 19 = 229, but this would be reported as 230 because the trailing zero in the number 210 is not significant.

When a calculation involves multiplication or division, such as when carbon normalizing is used, all significant figures are carried through the calculation, and then the total result is rounded at the end of the calculation to reflect the value used in the calculation with the fewest significant figures. For example:

> 59.9 x 1.2 = 71.88, to be reported as 72 because there are two significant figures in the number 1.2.

When rounding, if the number following the last significant figure is less than 5, the digit is left unchanged. If the number following the last significant figure is equal to or greater than 5, the digit is increased by 1.

DILUTIONS

All analyte concentrations within the calibration range of the instrument in the lowest analytical dilution are selected as the final result. Any analyte concentrations that exceed the calibration range and are qualified as estimated by the laboratory as an exceedance (E-qualified) are rejected by the data validator. The values for these analytes are selected from the analysis of the sample dilution in which the analyte concentration is within the calibration range of the instrument. In cases where the result from the lowest analytical dilution is qualified by the laboratory or the validator, the validator uses best professional judgment to determine whether or not the qualification warrants the selection of the result from another analytical dilution as the final result.

MULTIPLE RESULTS FOR THE SAME ANALYTE USING ONE ANALYTICAL METHOD

Multiple analyses of a sample for a group of analytes can occur as a result of laboratory quality assurance (QA) issues that may only affect a subset of the analyte group. In these cases, there may be multiple results for certain analytes. The data validator uses the following rules to select a single value when multiple results are reported by the laboratory for a single analyte in a single sample using the same method.

- If all results are detected without qualification as an estimated value (i.e., J- or E-qualifier), then the result from the lowest analytical dilution is selected. If multiple, unqualified results from the same analytical dilution are available, the highest concentration is selected as a health-protective approach.
- If a mixture of estimated (i.e., J-qualified) and unqualified detected results are reported, then the unqualified detected result is selected.
- If all results are reported as detected with estimated qualification, the "best result" is selected using best professional, technical judgment.
- If both undetected and detected results are reported, then the detected result is selected.
- If all results are reported as undetected, then the lowest RL is selected.

MULTIPLE RESULTS FOR AN ANALYTE DETERMINED BY DIFFERENT ANALYTICAL METHODS

In cases where a single analyte is reported by more than one method, the preferred method is identified in the quality assurance project plan (QAPP). The results of this method are selected as the final value by the data validator unless the validator identifies a QA issue that warrants the selection of the results from an alternative method. These instances and the justification for decisions are documented in the data validation report. In cases where the results are generated in two separate analytical groups that are not submitted to the validator together, the QA manager is responsible for evaluating the results and determining the most appropriate final result.

CALCULATING TOTALS

Concentrations for analyte sums are calculated as follows:

- **Total PCBs** are calculated, in accordance with the methods of the Washington State Sediment Management Standards (SMS), using only detected values for seven Aroclor mixtures.[2] For individual samples in which none of the seven Aroclor mixtures is detected, total PCBs are given a value equal to the highest RL of the seven Aroclors and assigned a U-qualifier indicating the lack of detected concentrations.

[2] Aroclors 1016, 1221, 1232, 1242, 1248, 1254, and 1260.

- **Total low-molecular-weight PAHs (LPAHs), high-molecular-weight PAHs (HPAHs), PAHs, and benzofluoranthenes** are also calculated in accordance with the methods of the SMS. Total LPAHs are the sum of detected concentrations for naphthalene, acenaphthylene, acenaphthene, fluorene, phenanthrene, and anthracene. Total HPAHs are the sum of detected concentrations for fluoranthene, pyrene, benzo(a)anthracene, chrysene, total benzofluoranthenes, benzo(a)pyrene, indeno(1,2,3,-c,d)pyrene, dibenzo(a,h)anthracene, and benzo(g,h,i)perylene. Total benzofluoranthenes are the sum of the b (i.e., benzo(b)fluoranthene), j, and k isomers. Because the j isomer is rarely quantified, this sum is typically calculated with only the b and k isomers. For samples in which all individual compounds within any of the three groups described above are undetected, the single highest RL for that sample represents the sum.
- **Total DDTs** are calculated using only detected values for the six DDT isomers: 2,4′-DDD; 4,4′-DDD; 2,4′-DDE; 4,4′-DDE; 2,4′-DDT; and 4,4′-DDT. For individual samples in which none of the isomers are detected, total DDTs are given a value equal to the highest RL of the six isomers and assigned a U-qualifier, indicating the lack of detected concentrations.
- **Total chlordane** is calculated using only detected values for the following compounds: alpha-chlordane, gamma-chlordane, oxychlordane, cis-nonachlor, and trans-nonachlor. For individual samples in which none of these compounds is detected, total chlordane is given a value equal to the highest RL of the five compounds listed above and assigned a U-qualifier, indicating the lack of detected concentrations.

CALCULATION OF PCB CONGENER TEQS

PCB congener toxic equivalents (TEQs) are calculated using the World Health Organization (WHO) consensus toxic equivalency factor (TEF) values for fish, birds (Van den Berg et al. 1998) and mammals (Van den Berg et al. 2006) as presented in Table 3-1. The TEQ is calculated as the sum of each congener concentration multiplied by the corresponding TEF value. When the congener concentration is reported as non-detected, then the TEF is multiplied by zero, half the RL or the full RL, depending on the calculation method specified.

Table 1. PCB congener TEF values

PCB CONGENER NUMBER	TEF VALUE FOR FISH (unitless)	TEF VALUE FOR BIRDS (unitless)	TEF VALUE FOR MAMMALS (unitless)
77	0.0001	0.05	0.0001
81	0.0005	0.1	0.0003
105	<0.000005	0.0001	0.00003
114	<0.000005	0.0001	0.00003
118	<0.000005	0.00001	0.00003

PCB Congener Number	TEF Value for Fish (unitless)	TEF Value for Birds (unitless)	TEF Value for Mammals (unitless)
123	<0.000005	0.00001	0.00003
126	0.005	0.1	0.1
156	<0.000005	0.0001	0.00003
157	<0.000005	0.0001	0.00003
167	<0.000005	0.00001	0.00003
169	0.00005	0.001	0.03
189	<0.000005	0.00001	0.00003

References

Van den Berg M, Birnbaum L, Bosveld ATC, Brunström B, Cook P, Feeley M, Giesy JP, Hanberg A, Hasegawa R, Kennedy S, Kubiak T, Larsen JC, van Leeuwen FXR, Djien Liem AK, Nolt C, Peterson RE, Poellinger L, Safe S, Schrenk D, Tillitt D, Tysklind M, Younes M, Waern F, Zacharewski T. 1998. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. Environ Health Perspec 106(12):775-792.

Van den Berg M, Birnbaum LS, Denison M, De Vito M, Farland W, Feeley M, Fiedler H, Hakansson H, Hanberg A, Haws L, Rose M, Safe S, Schrenk D, Tohyama C, Tritscher A, Tuomisto J, Tysklind M, Walker N, Peterson RE. 2006. The 2005 World Health Organization reevaluation of human and mammalian toxic equivalency factors for dioxins and dioxin-like compounds. Tox Sci 93(2):223-241.

Windward. 2006. Lower Duwamish Waterway remedial investigation. Technical memorandum: Criteria for defining the baseline surface sediment dataset for use in the Lower Duwamish Waterway Phase 2 RI/FS. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

ATTACHMENT 4
RESULTS OF
CHEMICALS OF INTEREST SCREEN

Attachment 4 Results of Chemicals of Interest Screen

Chemical	Bio-accumulative Chemical? (Y/N)	Sediment		Intertidal Sediment		Tissue			Selected as COPC? (Y/N)	Selected as Sandpiper COPC? (Y/N)	Selected as Crab COPC? (Y/N)
		Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Any Tissue (No. of detects/No. of samples)	Crab[a] (No. of detects/ No. of samples)	Benthic Invertebrate (No. of detects/No. of samples)			
Metals and Trace Elements											
Aluminum	N	453/453	Y	154/154	Y	na	na	na	N	N	N
Antimony	N	139/552	Y	57/181	Y	130/164	26/29	63/67	Y	Y	Y
Arsenic	Y	754/814	Y	307/357	Y	164/164	29/29	67/67	Y	Y	Y
Arsenic (inorganic)	N	na	na	na	na	36/36	12/12	20/20	N	N	N
Barium	N	418/418	Y	142/142	Y	na	na	na	N	N	N
Beryllium	N	451/463	Y	146/156	Y	na	na	na	N	N	N
Cadmium	Y	565/797	Y	218/348	Y	164/164	29/29	67/67	Y	Y	Y
Calcium	N	418/418	Y	142/142	Y	na	na	na	N	N	N
Chromium	N	811/811	Y	357/357	Y	121/164	9/29	47/67	Y	Y	Y
Chromium VI	Y	1/7	Y	na	na	na	na	na	Y	N	N
Cobalt	N	556/556	Y	232/232	Y	143/143	26/26	60/60	Y	Y	Y
Copper	Y	814/814	Y	357/357	Y	164/164	29/29	67/67	Y	Y	Y
Iron	N	451/451	Y	152/152	Y	na	na	na	N	N	N
Lead	Y	814/814	Y	357/357	Y	164/164	29/29	67/67	Y	Y	Y
Magnesium	N	428/428	Y	143/143	Y	na	na	na	N	N	N
Manganese	N	448/448	Y	154/154	Y	na	na	na	N	N	N
Mercury	N	715/831	Y	274/352	Y	167/169	29/29	65/67	Y	Y	Y
Methylmercury	Y	20/20	Y	4/4	Y	na	na	na	Y	Y	N
Molybdenum	N	199/257	Y	105/119	Y	142/142	26/26	60/60	Y	Y	Y
Nickel	Y	771/773	Y	323/324	Y	163/164	29/29	67/67	Y	Y	Y
Potassium	N	429/429	Y	145/145	Y	na	na	na	N	N	N
Selenium	Y	277/629	Y	71/249	Y	132/132	26/26	60/60	Y	Y	Y
Silver	Y	481/782	Y	174/345	Y	139/164	29/29	67/67	Y	Y	Y
Sodium	N	418/418	Y	140/140	Y	na	na	na	N	N	N

Chemical	Bio-accumulative Chemical? (Y/N)	Sediment		Intertidal Sediment		Tissue			Selected as COPC? (Y/N)	Selected as Sandpiper COPC? (Y/N)	Selected as Crab COPC? (Y/N)
		Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Any Tissue (No. of detects/No. of samples)	Crab[a] (No. of detects/ No. of samples)	Benthic Invertebrate (No. of detects/No. of samples)			
Thallium	N	325/635	Y	90/249	Y	103/134	22/26	56/62	Y	Y	Y
Tin	N	163/251	Y	15/43	Y	na	na	na	N	N	N
Vanadium	N	556/556	Y	232/232	Y	114/140	16/26	50/60	Y	Y	Y
Zinc	Y	810/811	Y	354/355	Y	164/164	29/29	67/67	Y	Y	Y
Organometals											
Monobutyltin as ion	N	89/117	Y	36/42	Y	92/144	24/26	49/62	Y	Y	Y
Dibutyltin as ion	N	106/147	Y	43/52	Y	136/144	26/26	56/62	Y	Y	Y
Tributyltin as ion	Y	143/159	Y	47/56	Y	158/181	21/29	58/67	Y	Y	Y
Tetrabutyltin as ion	N	15/119	Y	3/43	Y	1/133	0/26	0/62	Y	N	N
Butyltin (total)	N	29/37	Y	0/4	N	na	na	na	N	N	N
Alkylated PAHs											
C1-Chrysenes	N	20/20	Y	13/13	Y	5/20	na	5/20	Y	N	N
C1-Dibenzothiophenes	N	14/20	Y	11/13	Y	1/20	na	1/20	Y	Y	N
C1-Fluoranthene/pyrene	N	19/20	Y	12/13	Y	12/20	na	12/20	Y	Y	N
C1-Fluorenes	N	10/20	Y	5/13	Y	2/20	na	2/20	Y	Y	N
C1-Phenanthrenes/ anthracenes	N	20/20	Y	13/13	Y	7/20	na	7/20	Y	Y	N
C2-Chrysenes	N	20/20	Y	13/13	Y	1/20	na	1/20	Y	N	N
C2-Dibenzothiophenes	N	15/20	Y	10/13	Y	1/20	na	1/20	Y	Y	N
C2-Fluorenes	N	16/20	Y	9/13	Y	4/20	na	4/20	Y	Y	N
C2-Naphthalenes	N	20/20	Y	13/13	Y	1/20	na	1/20	Y	Y	N
C2-Phenanthrenes/ anthracenes	N	20/20	Y	13/13	Y	6/20	na	6/20	Y	Y	N
C3-Chrysenes	N	20/20	Y	13/13	Y	0/20	na	0/20	N	N	N
C3-Dibenzothiophenes	N	16/20	Y	11/13	Y	0/20	na	0/20	N	N	N
C3-Fluorenes	N	18/20	Y	11/13	Y	4/20	na	4/20	Y	Y	N
C3-Naphthalenes	N	20/20	Y	13/13	Y	1/20	na	1/20	Y	Y	N
C3-Phenanthrenes/ anthracenes	N	20/20	Y	13/13	Y	7/20	na	7/20	Y	Y	N
C4-Chrysenes	N	17/20	Y	10/13	Y	0/20	na	0/20	N	N	N

Chemical	Bio-accumulative Chemical? (Y/N)	Sediment		Intertidal Sediment		Tissue			Selected as COPC? (Y/N)	Selected as Sandpiper COPC? (Y/N)	Selected as Crab COPC? (Y/N)
		Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Any Tissue (No. of detects/No. of samples)	Crab[a] (No. of detects/ No. of samples)	Benthic Invertebrate (No. of detects/No. of samples)			
C4-Naphthalenes	N	20/20	Y	13/13	Y	1/20	na	1/20	Y	Y	N
C4-Phenanthrenes/ anthracenes	N	20/20	Y	13/13	Y	4/20	na	4/20	Y	Y	N
PAHs											
1-Methylnaphthalene	N	20/20	Y	13/13	Y	19/20	na	19/20	Y	Y	N
2-Chloronaphthalene	N	0/743	N	0/310	N	0/164	0/29	0/67	N	N	N
2-Methylnaphthalene	N	139/780	Y	69/341	Y	100/164	18/29	52/67	Y	Y	Y
Acenaphthene	Y	301/790	Y	113/345	Y	109/164	26/29	60/67	Y	Y	Y
Acenaphthylene	Y	121/780	Y	59/341	Y	96/164	15/29	48/67	Y	Y	Y
Anthracene	Y	552/790	Y	181/345	Y	106/164	25/29	59/67	Y	Y	Y
Benzo(a)anthracene	Y	717/790	Y	292/345	Y	111/164	23/29	57/67	Y	Y	Y
Benzo(a)pyrene	Y	716/784	Y	294/341	Y	70/164	3/29	34/67	Y	Y	Y
Benzo(b)fluoranthene	Y	723/784	Y	302/340	Y	88/164	9/29	43/67	Y	Y	Y
Benzo(e)pyrene	N	20/20	Y	13/13	Y	20/20	na	20/20	Y	Y	N
Benzo(g,h,i)perylene	Y	648/785	Y	254/343	Y	71/164	9/29	43/67	Y	Y	Y
Benzo(k)fluoranthene	Y	696/784	Y	284/340	Y	86/164	9/29	43/67	Y	Y	Y
Chrysene	Y	739/790	Y	311/345	Y	113/164	23/29	57/67	Y	Y	Y
Dibenzo(a,h)anthracene	Y	400/790	Y	139/345	Y	40/164	2/29	31/67	Y	Y	Y
Dibenzofuran	N	246/789	Y	90/344	Y	107/164	26/29	59/67	Y	Y	Y
Fluoranthene	Y	759/790	Y	327/345	Y	128/164	24/29	59/67	Y	Y	Y
Fluorene	Y	371/790	Y	117/345	Y	110/164	26/29	60/67	Y	Y	Y
Indeno(1,2,3-cd)pyrene	Y	692/785	Y	280/343	Y	69/164	9/29	42/67	Y	Y	Y
Naphthalene	N	148/780	Y	71/341	Y	54/164	9/29	30/67	Y	Y	Y
Perylene	N	20/20	Y	13/13	Y	13/20	na	13/20	Y	Y	N
Phenanthrene	Y	724/790	Y	294/345	Y	108/164	26/29	60/67	Y	Y	Y
Pyrene	Y	750/790	Y	321/345	Y	120/164	24/29	59/67	Y	Y	Y
Total PAH (calc'd)	N	769/790	Y	335/345	Y	134/164	26/29	61/67	Y	Y	N

Chemical	Bio-accumulative Chemical? (Y/N)	Sediment		Intertidal Sediment		Tissue			Selected as COPC? (Y/N)	Selected as Sandpiper COPC? (Y/N)	Selected as Crab COPC? (Y/N)
		Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Any Tissue (No. of detects/No. of samples)	Crab[a] (No. of detects/ No. of samples)	Benthic Invertebrate (No. of detects/No. of samples)			
Phthalates											
Bis(2-ethylhexyl)phthalate	N	635/794	Y	275/345	Y	28/164	4/29	21/67	Y	Y	Y
Butyl benzyl phthalate	N	390/784	Y	122/341	Y	25/159	14/29	15/65	Y	N	Y
Diethyl phthalate	N	42/794	Y	23/345	Y	39/164	15/29	17/67	Y	Y	Y
Dimethyl phthalate	N	136/784	Y	49/341	Y	5/164	5/29	5/67	Y	N	Y
Di-n-butyl phthalate	N	181/784	Y	93/341	Y	8/164	2/29	6/67	Y	Y	Y
Di-n-octyl phthalate	N	48/794	Y	23/345	Y	0/164	0/29	0/67	N	N	N
Other SVOCs											
1,2,4-Trichlorobenzene	N	5/778	N	0/339	N	0/164	0/29	0/67	N	N	N
1,2-Dichlorobenzene	Y	18/778	N	1/339	N	0/164	0/29	0/67	N	N	N
1,2-Diphenylhydrazine	N	0/109	N	0/31	N	0/32	0/3	0/7	N	N	N
1,3-Dichlorobenzene	Y	3/767	N	1/330	N	0/164	0/29	0/67	N	N	N
1,4-Dichlorobenzene	Y	35/778	N	10/339	N	0/164	0/29	0/67	N	N	N
2,4,5-Trichlorophenol	N	0/733	N	0/306	N	2/164	0/29	1/67	N	N	N
2,4,6-Trichlorophenol	N	0/733	N	0/306	N	2/164	0/29	1/67	N	N	N
2,4-Dichlorophenol	N	0/733	N	0/306	N	1/164	0/29	0/67	N	N	N
2,4-Dimethylphenol	N	1/773	N	1/339	N	2/164	0/29	1/67	N	N	N
2,4-Dinitrophenol	N	0/721	N	0/306	N	0/145	0/29	0/67	N	N	N
2,4-Dinitrotoluene	N	0/733	N	0/306	N	0/164	0/29	0/67	N	N	N
2,6-Dinitrotoluene	N	0/733	N	0/306	N	1/164	0/29	0/67	N	N	N
2-Chlorophenol	N	0/733	N	0/306	N	0/164	0/29	0/67	N	N	N
2-Methylphenol	N	3/783	N	1/344	N	19/164	0/29	0/67	N	N	N
2-Nitroaniline	N	0/721	N	0/306	N	0/164	0/29	0/67	N	N	N
2-Nitrophenol	N	0/733	N	0/306	N	0/164	0/29	0/67	N	N	N
3,3'-Dichlorobenzidine	N	0/692	N	0/294	N	0/136	0/26	0/64	N	N	N
3-Nitroaniline	N	0/709	N	0/301	N	0/144	0/26	0/64	N	N	N

Chemical	Bio-accumulative Chemical? (Y/N)	Sediment		Intertidal Sediment		Tissue			Selected as COPC? (Y/N)	Selected as Sandpiper COPC? (Y/N)	Selected as Crab COPC? (Y/N)
		Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Any Tissue (No. of detects/No. of samples)	Crab[a] (No. of detects/ No. of samples)	Benthic Invertebrate (No. of detects/No. of samples)			
4,6-Dinitro-o-cresol	N	0/721	N	0/306	N	0/164	0/29	0/67	N	N	N
4-Bromophenyl phenyl ether	Y	1/733	N	0/306	N	0/162	0/29	0/67	N	N	N
4-Chloro-3-methylphenol	N	1/721	N	1/306	N	1/162	0/29	0/65	N	N	N
4-Chloroaniline	N	0/686	N	0/301	N	2/136	0/27	1/64	N	N	N
4-Chlorophenyl phenyl ether	Y	0/733	N	0/306	N	1/164	0/29	0/67	Y	N	N
4-Methylphenol	N	78/793	Y	37/348	Y	18/164	0/29	13/67	Y	Y	N
4-Nitroaniline	N	0/709	N	0/301	N	0/154	0/27	0/65	N	N	N
4-Nitrophenol	N	0/721	N	0/306	N	5/162	0/29	2/65	N	N	N
Aniline	N	1/250	N	0/121	N	0/153	0/26	0/62	N	N	N
Benzaldehyde	N	6/10	Y	0/4	N	na	na	na	N	N	N
Benzidine	N	0/7	N	0/5	N	0/111	0/18	0/52	N	N	N
Benzoic acid	N	69/781	Y	40/345	Y	112/164	0/29	33/67	Y	Y	N
Benzyl alcohol	N	14/771	N	9/341	N	37/164	4/29	12/67	N	N	Y
Biphenyl	N	20/20	Y	13/13	Y	20/20	na	20/20	Y	Y	N
Bis(2-chloroethoxy)methane	N	1/733	N	0/306	N	1/164	0/29	0/67	N	N	N
Bis(2-chloroethyl)ether	N	0/733	N	0/306	N	0/164	0/29	0/67	N	N	N
Bis(2-chloroisopropyl)ether	N	0/733	N	0/306	N	0/164	0/29	0/67	N	N	N
Caffeine	N	1/31	N	0/11	N	0/32	0/3	0/7	N	N	N
Caprolactam	N	1/10	Y	0/4	N	na	na	na	N	N	N
Carbazole	N	385/743	Y	124/310	Y	2/164	0/29	0/67	Y	N	N
Coprostanol	N	43/107	Y	8/27	Y	0/32	0/3	0/7	N	N	N
Dibenzothiophene	N	20/20	Y	13/13	Y	15/20	na	15/20	Y	Y	N
Hexachlorobenzene	Y	46/781	Y	20/342	Y	21/164	5/29	14/67	Y	Y	Y
Hexachlorobutadiene	Y	0/780	N	0/341	N	0/164	0/29	0/67	N	N	N
Hexachlorocyclopentadiene	Y	1/682	N	0/299	N	0/160	0/29	0/63	N	N	N
Hexachloroethane	Y	0/761	N	0/325	N	0/163	0/29	0/67	N	N	N
Isophorone	N	2/743	N	1/310	N	1/164	0/29	0/67	N	N	N

Chemical	Bio-accumulative Chemical? (Y/N)	Sediment		Intertidal Sediment		Tissue			Selected as COPC? (Y/N)	Selected as Sandpiper COPC? (Y/N)	Selected as Crab COPC? (Y/N)
		Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Any Tissue (No. of detects/No. of samples)	Crab[a] (No. of detects/ No. of samples)	Benthic Invertebrate (No. of detects/No. of samples)			
Methyl isobutyl ketone	N	0/42	N	0/10	N	na	na	na	N	N	N
Nitrobenzene	N	0/733	N	0/306	N	0/164	0/29	0/67	N	N	N
N-Nitrosodimethylamine	N	0/292	N	0/130	N	0/164	0/29	0/67	N	N	N
N-Nitroso-di-n-propylamine	N	0/733	N	0/306	N	3/162	0/29	1/65	N	N	N
N-Nitrosodiphenylamine	N	23/780	N	11/341	N	1/164	0/29	1/67	N	N	N
Pentachlorophenol	N	12/747	N	8/318	N	16/164	0/29	5/67	N	N	N
Phenol	N	254/793	Y	84/348	Y	39/164	1/29	24/67	Y	Y	Y
Pyridine	N	0/12	N	na	na	na	na	na	N	N	N
Retene	N	10/19	Y	2/4	Y	na	na	na	N	N	N
PCBs											
Aroclor-1016	N	0/981	N	0/443	N	0/237	0/37	0/75	N	N	N
Aroclor-1016/1242	Y	na	na	na	na	6/6	na	na	N	N	N
Aroclor-1221	Y	0/855	N	0/334	N	0/243	0/37	0/75	N	N	N
Aroclor-1232	Y	0/855	N	0/334	N	0/243	0/37	0/75	N	N	N
Aroclor-1242	Y	103/982	Y	27/442	Y	0/237	0/37	0/75	Y	Y	N
Aroclor-1248	Y	203/991	Y	76/447	Y	107/243	29/37	36/75	Y	Y	Y
Aroclor-1254	Y	776/983	Y	296/445	Y	237/243	37/37	73/75	Y	Y	Y
Aroclor-1254/1260	Y	8/8	Y	2/2	Y	na	na	na	Y	Y	N
Aroclor-1260	Y	783/983	Y	323/445	Y	180/243	37/37	45/75	Y	Y	Y
Aroclor-1262	Y	2/12	Y	2/6	Y	na	na	na	Y	Y	N
Aroclor-1268	Y	1/11	Y	1/5	Y	na	na	na	Y	Y	N
PCBs (total calc'd)	Y	1203/1288	Y	509/552	Y	238/243	37/37	74/75	Y	Y	Y
Pesticides											
2,4'-DDD	Y	5/93	Y	4/53	Y	36/150	11/26	26/60	Y	Y	Y
2,4'-DDE	Y	2/93	N	2/53	N	14/150	0/26	3/60	Y	Y	N
2,4'-DDT	Y	29/93	Y	23/53	Y	134/150	26/26	49/60	Y	Y	Y
4,4'-DDD	Y	67/197	Y	28/75	Y	112/161	12/26	31/60	Y	Y	Y
4,4'-DDE	Y	30/197	Y	19/75	Y	122/161	26/26	49/60	Y	Y	Y

Chemical	Bio-accumulative Chemical? (Y/N)	Sediment		Intertidal Sediment		Tissue			Selected as COPC? (Y/N)	Selected as Sandpiper COPC? (Y/N)	Selected as Crab COPC? (Y/N)
		Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Any Tissue (No. of detects/No. of samples)	Crab[a] (No. of detects/ No. of samples)	Benthic Invertebrate (No. of detects/No. of samples)			
4,4'-DDT	Y	41/197	Y	27/75	Y	136/161	26/26	56/60	Y	Y	Y
DDTs (total-calc'd)	Y	78/197	Y	34/75	Y	150/161	26/26	60/60	Y	Y	Y
Aldrin	Y	4/197	N	3/75	N	6/161	0/26	3/60	Y	Y	N
Dieldrin	Y	9/197	N	6/75	Y	11/161	1/26	6/60	Y	Y	Y
alpha-BHC	Y	3/197	N	3/75	N	12/161	4/26	6/60	Y	Y	Y
beta-BHC	Y	4/197	N	4/75	Y	53/161	0/26	14/60	Y	Y	N
delta-BHC	Y	3/158	N	3/66	N	3/161	0/26	3/60	Y	Y	N
gamma-BHC	Y	11/197	Y	7/75	Y	16/161	1/26	4/60	Y	Y	Y
alpha-Chlordane	Y	13/153	Y	10/62	Y	38/150	2/26	3/60	Y	Y	Y
gamma-Chlordane	Y	25/153	Y	18/62	Y	130/150	26/26	50/60	Y	Y	Y
Chlordane	Y	5/44	Y	2/13	Y	0/11	na	na	Y	Y	Y
alpha-Endosulfan	Y	10/151	Y	9/60	Y	50/161	4/26	10/60	Y	Y	Y
beta-Endosulfan	Y	4/153	N	3/62	N	32/161	0/26	8/60	Y	Y	N
Endosulfan	N	1/46	N	1/15	Y	na	na	na	N	N	N
Endosulfan sulfate	N	3/195	N	3/73	N	4/161	0/26	0/60	N	N	N
Endrin	Y	4/197	N	4/75	Y	31/161	0/26	11/60	Y	Y	N
Endrin aldehyde	N	6/186	N	6/69	Y	25/161	5/26	9/60	N	Y	Y
Endrin ketone	N	7/141	N	5/62	Y	4/150	0/26	2/60	N	N	N
Heptachlor	Y	8/197	N	2/75	N	10/161	0/26	1/60	Y	Y	N
Heptachlor epoxide	Y	5/197	N	2/75	N	54/161	16/26	24/60	Y	Y	Y
Methoxychlor	Y	11/197	Y	8/75	Y	8/161	3/26	7/60	Y	Y	Y
Mirex	Y	3/93	N	1/53	N	0/150	0/26	0/60	N	N	N
cis-Nonachlor	Y	0/58	N	0/25	N	0/18	na	na	N	N	N
Oxychlordane	Y	0/58	N	0/25	N	6/18	na	na	Y	N	N
Toxaphene	Y	2/195	N	2/73	N	0/161	0/26	0/60	N	N	N
trans-Nonachlor	Y	0/58	N	0/25	N	6/18	na	na	Y	N	N
Total chlordane (calc'd)	Y	28/153	Y	20/62	Y	133/150	26/26	50/60	Y	Y	Y
VOCs											
1,1,1,2-Tetrachloroethane	N	0/37	N	0/7	N	na	na	na	N	N	N

Chemical	Bio-accumulative Chemical? (Y/N)	Sediment		Intertidal Sediment		Tissue			Selected as COPC? (Y/N)	Selected as Sandpiper COPC? (Y/N)	Selected as Crab COPC? (Y/N)
		Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Any Tissue (No. of detects/No. of samples)	Crab[a] (No. of detects/ No. of samples)	Benthic Invertebrate (No. of detects/No. of samples)			
1,1,1-Trichloroethane	N	0/42	N	0/10	N	na	na	na	N	N	N
1,1,2,2-Tetrachloroethane	N	0/42	N	0/10	N	na	na	na	N	N	N
1,1,2-Trichloroethane	N	0/42	N	0/10	N	na	na	na	N	N	N
1,1,2-Trichlorotrifluoroethane	N	0/40	N	0/10	N	na	na	na	N	N	N
1,1-Dichloroacetone	N	0/35	N	0/6	N	na	na	na	N	N	N
1,1-Dichloroethane	N	0/42	N	0/10	N	na	na	na	N	N	N
1,1-Dichloroethene	N	0/42	N	0/10	N	na	na	na	N	N	N
1,1-Dichloropropene	N	0/37	N	0/7	N	na	na	na	N	N	N
1,2,3-Trichlorobenzene	N	0/37	N	0/7	N	na	na	na	N	N	N
1,2,3-Trichloropropane	N	0/37	N	0/7	N	na	na	na	N	N	N
1,2,4-Trimethylbenzene	Y	1/37	N	0/7	N	na	na	na	N	N	N
1,2-Dibromo-3-chloropropane	N	0/37	N	0/7	N	na	na	na	N	N	N
1,2-Dibromoethane (EDB)	N	0/37	N	0/7	N	na	na	na	N	N	N
1,2-Dichloroethane	N	0/42	N	0/10	N	na	na	na	N	N	N
1,2-Dichloroethene (total)	N	0/2	N	na	na	na	na	na	N	N	N
1,2-Dichloropropane	N	0/42	N	0/10	N	na	na	na	N	N	N
1,3,5-Trimethylbenzene	N	0/37	N	0/7	N	na	na	na	N	N	N
1,3-Dichloropropane	N	0/37	N	0/7	N	na	na	na	N	N	N
1-Chlorobutane	N	0/37	N	0/7	N	na	na	na	N	N	N
2,2-Dichloropropane	N	0/37	N	0/7	N	na	na	na	N	N	N
2-Chloroethyl vinyl ether	N	0/3	N	0/3	N	na	na	na	N	N	N
2-Chlorotoluene	N	0/37	N	0/7	N	na	na	na	N	N	N
2-Hexanone	N	0/42	N	0/10	N	na	na	na	N	N	N
2-Nitropropane	N	0/37	N	0/7	N	na	na	na	N	N	N
4-Chlorotoluene	N	0/37	N	0/7	N	na	na	na	N	N	N
Acetone	N	3/42	Y	3/10	Y	na	na	na	N	N	N
Allyl chloride	N	0/37	N	0/7	N	na	na	na	N	N	N
Benzene	N	0/42	N	0/10	N	na	na	na	N	N	N
Bromobenzene	N	0/37	N	0/7	N	na	na	na	N	N	N

Chemical	Bio-accumulative Chemical? (Y/N)	Sediment		Intertidal Sediment		Tissue			Selected as COPC? (Y/N)	Selected as Sandpiper COPC? (Y/N)	Selected as Crab COPC? (Y/N)
		Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Any Tissue (No. of detects/No. of samples)	Crab[a] (No. of detects/ No. of samples)	Benthic Invertebrate (No. of detects/No. of samples)			
Bromochloromethane	N	0/37	N	0/7	N	na	na	na	N	N	N
Bromodichloromethane	N	0/42	N	0/10	N	na	na	na	N	N	N
Bromoform	N	0/42	N	0/10	N	na	na	na	N	N	N
Bromomethane	N	0/42	N	0/10	N	na	na	na	N	N	N
Carbon disulfide	N	13/42	Y	1/10	Y	na	na	na	N	N	N
Carbon tetrachloride	N	0/42	N	0/10	N	na	na	na	N	N	N
Chloroacetonitrile	N	0/2	N	na	na	na	na	na	N	N	N
Chlorobenzene	N	0/42	N	0/10	N	na	na	na	N	N	N
Chloroethane	N	0/42	N	0/10	N	na	na	na	N	N	N
Chloroform	Y	0/42	N	0/10	N	na	na	na	N	N	N
Chloromethane	N	0/42	N	0/10	N	na	na	na	N	N	N
cis-1,2-Dichloroethene	N	0/40	N	0/10	N	na	na	na	N	N	N
cis-1,3-Dichloropropene	N	0/42	N	0/10	N	na	na	na	N	N	N
p-Cymene	N	3/37	Y	0/7	N	na	na	na	N	N	N
Dibromochloromethane	N	0/42	N	0/10	N	na	na	na	N	N	N
Dibromomethane	N	0/37	N	0/7	N	na	na	na	N	N	N
Dichlorodifluoromethane	N	0/7	N	0/2	N	na	na	na	N	N	N
Dichloromethane	N	1/42	N	0/10	N	na	na	na	N	N	N
Diethyl ether	N	0/37	N	0/7	N	na	na	na	N	N	N
Ethyl methacrylate	N	0/37	N	0/7	N	na	na	na	N	N	N
Ethylbenzene	N	0/42	N	0/10	N	na	na	na	N	N	N
Iodomethane	N	0/37	N	0/7	N	na	na	na	N	N	N
Isopropylbenzene	N	0/37	N	0/7	N	na	na	na	N	N	N
Methacrylonitrile	N	0/37	N	0/7	N	na	na	na	N	N	N
Methyl Acrylate	N	0/37	N	0/7	N	na	na	na	N	N	N
Methyl ethyl ketone	N	15/42	Y	4/10	Y	na	na	na	N	N	N
Methyl methacrylate	N	0/37	N	0/7	N	na	na	na	N	N	N
n-Butylbenzene	N	0/37	N	0/7	N	na	na	na	N	N	N
n-Propylbenzene	N	0/37	N	0/7	N	na	na	na	N	N	N

Chemical	Bio-accumulative Chemical? (Y/N)	Sediment		Intertidal Sediment		Tissue			Selected as COPC? (Y/N)	Selected as Sandpiper COPC? (Y/N)	Selected as Crab COPC? (Y/N)
		Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Any Tissue (No. of detects/No. of samples)	Crab[a] (No. of detects/ No. of samples)	Benthic Invertebrate (No. of detects/No. of samples)			
Pentachloroethane	N	0/37	N	0/7	N	na	na	na	N	N	N
sec-Butylbenzene	N	0/37	N	0/7	N	na	na	na	N	N	N
Styrene	N	0/42	N	0/10	N	na	na	na	N	N	N
tert-Butyl methyl ether	N	0/37	N	0/7	N	na	na	na	N	N	N
tert-Butylbenzene	N	0/37	N	0/7	N	na	na	na	N	N	N
Tetrachloroethene	N	2/42	N	0/10	N	na	na	na	N	N	N
Toluene	N	4/42	Y	0/10	N	na	na	na	N	N	N
trans-1,2-Dichloroethene	N	0/40	N	0/10	N	na	na	na	N	N	N
trans-1,3-Dichloropropene	N	0/42	N	0/10	N	na	na	na	N	N	N
trans-1,4-Dichloro-2-butene	N	0/35	N	0/6	N	na	na	na	N	N	N
Trichloroethene	N	0/42	N	0/10	N	na	na	na	N	N	N
Trichlorofluoromethane	N	0/40	N	0/10	N	na	na	na	N	N	N
Vinyl acetate	N	0/3	N	0/3	N	na	na	na	N	N	N
Vinyl chloride	N	0/42	N	0/10	N	na	na	na	N	N	N
Xylene (ortho)	N	0/40	N	0/10	N	na	na	na	N	N	N
Xylene (meta and para)	N	0/40	N	0/10	N	na	na	na	N	N	N
Xylene (total)	N	0/2	N	na	na	na	na	na	N	N	N
Total xylenes (calc'd)	N	0/40	N	0/10	N	na	na	na	N	N	N
Dioxins/Furans[b]											
2,3,7,8-TCDD	Y	17/43	Y	11/17	Y	na	na	na	Y	Y	N
1,2,3,7,8-PeCDD	Y	19/43	Y	12/17	Y	na	na	na	Y	Y	N
1,2,3,4,7,8-HxCDD	Y	19/43	Y	12/17	Y	na	na	na	Y	Y	N
1,2,3,6,7,8-HxCDD	Y	35/43	Y	15/17	Y	na	na	na	Y	Y	N
1,2,3,7,8,9-HxCDD	Y	31/43	Y	15/17	Y	na	na	na	Y	Y	N
1,2,3,4,6,7,8-HpCDD	Y	41/43	Y	17/17	Y	na	na	na	Y	Y	N
OCDD	Y	43/43	Y	17/17	Y	na	na	na	Y	Y	N
2,3,7,8-TCDF	Y	34/43	Y	15/17	Y	na	na	na	Y	Y	N
1,2,3,7,8-PeCDF	Y	18/43	Y	12/17	Y	na	na	na	Y	Y	N
2,3,4,7,8-PeCDF	Y	19/43	Y	12/17	Y	na	na	na	Y	Y	N

Chemical	Bio-accumulative Chemical? (Y/N)	Sediment		Intertidal Sediment		Tissue			Selected as COPC? (Y/N)	Selected as Sandpiper COPC? (Y/N)	Selected as Crab COPC? (Y/N)
		Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Detection Frequency (No. of detects/No. of samples)	Detected in ≥ 5% of Sediment Samples? (Y/N)	Any Tissue (No. of detects/No. of samples)	Crab[a] (No. of detects/ No. of samples)	Benthic Invertebrate (No. of detects/No. of samples)			
1,2,3,4,7,8-HxCDF	Y	30/43	Y	14/17	Y	na	na	na	Y	Y	N
1,2,3,6,7,8-HxCDF	Y	19/43	Y	12/17	Y	na	na	na	Y	Y	N
1,2,3,7,8,9-HxCDF	Y	18/43	Y	12/17	Y	na	na	na	Y	Y	N
2,3,4,6,7,8-HxCDF	Y	19/43	Y	12/17	Y	na	na	na	Y	Y	N
1,2,3,4,6,7,8-HpCDF	Y	40/43	Y	16/17	Y	na	na	na	Y	Y	N
1,2,3,4,7,8,9-HpCDF	Y	27/43	Y	14/17	Y	na	na	na	Y	Y	N
OCDF	Y	42/43	Y	17/17	Y	na	na	na	Y	Y	N
Total TCDD	Y	20/25	Y	4/5	Y	na	na	na	Y	Y	N
Total PeCDD	Y	1/25	N	0/5	N	na	na	na	N	N	N
Total HxCDD	Y	23/25	Y	5/5	Y	na	na	na	Y	Y	N
Total HpCDD	Y	23/25	Y	5/5	Y	na	na	na	Y	Y	N
Total TCDF	Y	23/25	Y	5/5	Y	na	na	na	Y	Y	N
Total PeCDF	Y	22/25	Y	4/5	Y	na	na	na	Y	Y	N
Total HxCDF	Y	23/25	Y	5/5	Y	na	na	na	Y	Y	N
Total HpCDF	Y	23/25	Y	5/5	Y	na	na	na	Y	Y	N

[a] Eight hepatopancreas samples and twenty-one calculated whole body samples. The LDWRI 2005 crab samples were analyzed for PCB Aroclors only and include an additional four hepatopancreas samples and four calculated whole body samples.

[b] No quantitative LDW risk estimates were calculated for ecological risks from dioxins/furans. LDWG, EPA, and Ecology agreed that such data were not needed for remedial decision-making, because remedial decisions to address dioxins/furans will be based on concentrations of dioxins/furans in LDW sediment relative to their concentrations in urban background sediment from the greater Seattle metropolitan area

COPC – chemical of potential concern

HpCDD – heptachlorodibenzo-*p*-dioxin

HpCDF – heptachlorodibenzofuran

HxCDD – hexachlorodibenzo-*p*-dioxin

HxCDF – hexachlorodibenzofuran

N – no

na – not analyzed

OCDD – octachlorodibenzo-*p*-dioxin

OCDF – octachlorodibenzofuran

PAH – polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

PeCDD – pentachlorodibenzo-*p*-dioxin

PeCDF – pentachlorodibenzofuran

SVOC – semivolatile organic compound

TCDD – tetrachlorodibenzo-*p*-dioxin

TCDF – tetrachlorodibenzofuran

VOC – volatile organic compound

Y – yes

ATTACHMENT 5
ALL ACCEPTABLE DECAPOD STUDIES FOR CHEMICALS OF INTEREST FOR CRAB

Attachment 5 All Acceptable Decapod Studies for Chemicals of Interest for Crab

Analyte	Test Species	Marine or Fresh-water	Tissue Type	NOAEL	LOAEL	Unit	Exposure Duration	Exposure Route	Endpoint	Source	Notes
Metals and Trace Elements											
Arsenic	grass shrimp	marine	whole body	1.28	na	mg/kg ww	28 days	water	growth	Lindsay and Sanders (1990)	accumulation test where no mortality observed (not a toxicity test)
Arsenic	grass shrimp	marine	whole body	1.15	na	mg/kg ww	28 days	water	growth	Lindsay and Sanders (1990)	accumulation test where no mortality observed (not a toxicity test)
Cadmium	red swamp crayfish	fresh-water	not measured–injection study	0.5	na	mg/kg ww	21 days	single injection	reproduction	Reddy et al. (1997)	no significant effect on ovarian maturation was observed; NOAEL is injected dose (whole body tissues were not measured)
Cadmium	grass shrimp	marine	whole body	0.6	na	mg/kg ww	14 days	exposed sediment	mortality	Rule and Alden (1996)	NOAEL estimated assuming 80% moisture in organism; no significant mortalities observed in accumulation study where shrimp were exposed to cadmium only (up to 5 mg/kg in sediment); estimated from figure
Cadmium	grass shrimp	marine	whole body	na	2.6	mg/kg ww	21 days	water	mortality	Vernberg et al. (1977)	at LOAEL – approximately 20% increase in mortality after 7 days at 5 ppt salinity; no statistics; little increase in mortality after day 7; only one cadmium exposure dose at varying salinities.
Cadmium	shore crab	marine	muscle	4.9	9.5	mg/kg ww	40 days	1 and 10 ppm in water for 40 days	survival	Jennings and Rainbow (1979)	measured in muscle; NOAEL based on 1 ppm exposure; LOAEL based on 10 ppm exposure. Concentrations lowest in five tissue types (midgut gland, gills, muscle tissue, remaining tissue
Cadmium	adult Norway lobster	marine	hepato-pancreas	5.7	na	mg/kg ww	50 days	13 mg/kg ww in diet	survival	Canli and Furness (1995)	measured in hepatopancreas; no mortality
Cadmium	adult Norway lobster	marine	gills	2.7	na	mg/kg ww	50 days	13 mg/kg ww in diet	survival	Canli and Furness (1995)	measured in gills; no mortality

Analyte	Test Species	Marine or Fresh-water	Tissue Type	NOAEL	LOAEL	Unit	Exposure Duration	Exposure Route	Endpoint	Source	Notes
Cadmium	adult Norway lobster	marine	tail muscle	0.13	na	mg/kg ww	50 days	13 mg/kg ww in diet	survival	Canli and Furness (1995)	measured in tail muscle; no mortality
Cadmium	adult Norway lobster	marine	hepato-pancreas	46	na	mg/kg ww	30 days	100 µg/L in water	survival	Canli and Furness (1995)	measured in hepatopancreas; no mortality
Cadmium	adult Norway lobster	marine	gills	32	na	mg/kg ww	30 days	100 µg/L in water	survival	Canli and Furness (1995)	measured in gills; no mortality
Cadmium	adult Norway lobster	marine	tail muscle	0.58	na	mg/kg ww	30 days	100 µg/L in water	survival	Canli and Furness (1995)	measured in tail muscle; no mortality
Cadmium	crustaceans	fresh-water	whole body	na	3.5-10.3	mg/kg ww	22 days	water	prenatal mortality	Marshall (1978) in Dillon (1984)	reduced longevity and increased prenatal mortality
Cadmium	crayfish		whole body	14.9	22	mg/kg ww	5 months	water	mortality	Thorp et al. (1979)	significant increase in mortality at LOAEL; water contaminated with 10 µg/L cadmium; no effect on growth at LOAEL; NOAEL is whole-body concentration where no significant mortality was observed where crayfish were exposed to 5 µg/L cadmium in water
Cadmium	crayfish	fresh-water	whole body	na	28.4	mg/kg ww	2 weeks	water	mortality	Mirenda (1986b)	at LOAEL – 25% mortality after 2 weeks in lowest dose group
Cadmium	crayfish	fresh-water	whole body	534	na	mg/kg ww	approx. 8 days	water	mortality	Gillespie et al. (1977)	no significant difference in mortality (1/15 died) in crayfish exposed to 1 ppm cadmium in water; NOAEL is average tissue concentration
Chromium	juvenile (2nd instar) sand crab	marine	whole body	1	3.2	mg/kg ww	30 days	0.1 and 0.3 mg/L in water	growth	Mortimer and Miller (1994)	NOAEL and LOAEL values read of Figure 2. Paper provide equations that would slightly lower LOAEL (2.67) and increase NOAEL (1.2)

Analyte	Test Species	Marine or Fresh-water	Tissue Type	NOAEL	LOAEL	Unit	Exposure Duration	Exposure Route	Endpoint	Source	Notes
Copper	grass shrimp	marine	whole body	40	na	mg/kg ww	14 days	exposed sediment	mortality	Rule et al. (1996)	NOAEL estimated assuming 80% moisture in organism; no significant mortalities observed in accumulation study where shrimp were exposed to copper only (up to 25 mg/kg in sediment); estimated from figure (a little less than 200 mg/kg); relationship of copper uptake to body burden was weak
Copper	crayfish	fresh-water	whole body	50	na	mg/kg ww	48 hours	water	mortality	Evans (1980)	accumulation test where no mortality was observed (not a toxicity test); NOAEL estimated from study assuming 80% moisture in organism
Copper	crayfish	fresh-water	thorax and abdomen	34	na	mg/kg ww	48 hours	water	mortality	Evans (1980)	accumulation test where no mortality was observed (not a toxicity test); NOAEL estimated from study assuming 80% moisture in organism
Mercury	red swamp crayfish	fresh-water	not measured - injection study	na	0.5	mg/kg ww	21 days	single injection	reproduction	Reddy et al. (1997)	ovarian maturation was significantly reduced at LOAEL; LOAEL is injected dose (whole body tissues were not measured); reduced ovarian maturation may cause a reduction in egg laying
Mercury	adult male shore crab	fresh and salt water	hepato-pancreas	na	1	mg/kg ww	32 hours	1 mg/L in water	survival	Bianchini and Gilles (1996)	concentration is lowest of three crab species tested (*Carcinus maenas, Eriocheir sinensis*, and *Cancer pagurus*)
Mercury	adult Norway lobster	marine	hepato-pancreas	0.99	na	mg/kg ww	50 days	164 mg/kg ww in diet	survival	Canli and Furness (1995)	measured in hepatopancreas; no mortality
Mercury	Adult Norway lobster	marine	gills	0.77	na	mg/kg ww	50 days	164 mg/kg ww in diet	survival	Canli and Furness (1995)	measured in gills; no mortality
Mercury	adult Norway lobster	marine	tail muscle	0.23	na	mg/kg ww	50 days	164 mg/kg ww in diet	survival	Canli and Furness (1995)	measured in tail muscle; no mortality

Analyte	Test Species	Marine or Fresh-water	Tissue Type	NOAEL	LOAEL	Unit	Exposure Duration	Exposure Route	Endpoint	Source	Notes
Mercury	adult Norway lobster	marine	hepato-pancreas	1.43	na	mg/kg ww	30 days	10 µg/L in water	survival	Canli and Furness (1995)	measured in hepatopancreas; no mortality
Mercury	adult Norway lobster	marine	gills	46.5	na	mg/kg ww	30 days	10 µg/L in water	survival	Canli and Furness (1995)	measured in gills; no mortality
Mercury	adult Norway lobster	marine	tail muscle	0.34	na	mg/kg ww	30 days	10 µg/L in water	survival	Canli and Furness (1995)	measured in tail muscle; no mortality
Vanadium	shrimp	marine	whole body	0.6	na	mg/kg ww	21 days	100 µg/L in water	mortality	Miramand et al. (1981)	no mortality observed
Zinc	crayfish	fresh-water	whole body	12.7	35.2	mg/kg ww	2 weeks	water	mortality	Mirenda (1986a)	at LOAEL = 23% mortality; at NOAEL = 6% mortality; NOAEL and LOAEL estimated from study assuming 80% moisture in organism. Other tissue types analyzed in the study: gills, abdominal muscle, carapace, and intestine.
Zinc	crayfish	fresh-water	hepato-pancreas	42.5	85.6	mg/kg ww	2 weeks	water	mortality	Mirenda (1986a)	at LOAEL = 23% mortality; at NOAEL = 6% mortality; NOAEL and LOAEL estimated from study assuming 80% moisture in organism. Other tissue types analyzed in the study: gills, abdominal muscle, carapace, and intestine.

Analyte	Test Species	Marine or Fresh-water	Tissue Type	NOAEL	LOAEL	Unit	Exposure Duration	Exposure Route	Endpoint	Source	Notes
SVOCs											
1,4-dichloro-benzene	sand crab	marine	whole body	na	1,600	µg/kg ww	90 hours	concentration range 0.1-10 ppm in water	mortality	Mortimer and Connell (1994)	values derived from Fig.3. Read tissue conc. as 1/16 of 100, 4/16 of 100, and 14/16 of 100 = 6.3, 25, and 87.5 mmol/kg lipid. 1,4-dichlorobenzene = 147 g/mol, which then gives 926, 3675, and 12,863 mg/kg lipid. Assume 0.9% lipid and 80% moisture, which gives: 1.6 mg/kg ww, 6.6 mg/kg ww, and 23 mg/kg ww. ERED from the same paper comes up with four values: 5.88, 14.7, 33.8, and 85.27 mg/kg for LC100 – does not state ww or dw.
1,4-dichloro-benzene	sand crab	marine	whole body	na	6,600	µg/kg ww	60 hours	concentration range 0.1-10 ppm in water	mortality	Mortimer and Connell (1994)	
1,4-dichloro-benzene	sand crab	marine	whole body	na	23,000	µg/kg ww	40 hours	concentration range 0.1-10 ppm in water	mortality	Mortimer and Connell (1994)	
Naphthalene	spot shrimp	marine	whole body	na	50	µg/kg ww	24 hours	concentration range 8-12 ppb in water	mortality	Sanborn and Malins (1977)	LOAEL read of Fig. 1. Bovine serum albumin was added to the seawater without telling why
Phthalates											
1,2-di-2-ethylhexyl phthalate	penaied shrimp	marine	whole body	18,300	na	µg/kg ww	14 days	diet	mortality	Hobson et al (1984)	no mortality at the highest exposure concentration
Phthalates	crustaceans	fresh-water	whole body	320-26,800	na	µg/kg ww	21 days	water	reproduction	Brown and Thompson (1982) s cited in Dillon (1984)	no effects on reproduction after one-generation exposure
PCBs											
PCBs (Aroclor 1254)	crayfish	fresh-water	whole body	1,220	na	µg/kg ww	21 days	water	mortality	Sanders and Chandler (1972)	NOAEL based on dose given for 21 days in biomagnification study; based on magnification factor and water concentration with no effects data; NOAEL concentration reported in study = 6.1 mg/kg dw
PCBs (Aroclor 1254)	pink shrimp	marine	whole body	1,300	3,900	µg/kg ww	48 hours	water	mortality	Duke et al. (1970)	100% mortality at LOAEL; 0% mortality at NOAEL; no control was reported

Analyte	Test Species	Marine or Fresh-water	Tissue Type	NOAEL	LOAEL	Unit	Exposure Duration	Exposure Route	Endpoint	Source	Notes
PCBs (Aroclor 1254)	pink shrimp	marine	whole body	na	16,000	µg/kg ww	20 days	water	mortality	Duke et al. (1970)	72% mortality at LOAEL (0% mortality in control); LOAEL is average concentration in dead shrimp, average concentration is higher in surviving shrimp subject to same dose-33 ppm
PCBs (Aroclor 1254)	blue crab	marine	whole body	23,000	na	µg/kg ww	20 days	water	mortality	Duke et al. (1970)	equal mortality in control and dose group (5 µg/L) – 1/20 crabs died; NOAEL is average of 5 crabs (range of 18 to 27 ppm)
PCBs (Aroclor 1254)	grass shrimp	marine	whole body	18,000	27,000	µg/kg ww	16 days	water	mortality	Nimmo et al. (1974)	45% mortality at LOAEL – significantly higher than control group (25% mortality); NOAEL is 40% mortality, which paper states is not significantly higher than control ($p < 0.05$)
PCBs (Aroclor 1016)	grass shrimp	marine	whole body	na	1,100	µg/kg ww	96 hours	water	mortality	Hansen et al. (1974)	33% mortality at LOAEL; control is next lowest dose with 8% mortality; no statistics; shrimp collected at Gulf Breeze lab
PCBs (Aroclor 1016)	brown shrimp	marine	whole body	3,800	42,000	µg/kg ww	96 hours	water	mortality	Hansen et al. (1974)	43% mortality at LOAEL; 8% mortality at NOAEL; 0% mortality in control; no statistics
Organochlorine Pesticides											
Chlordane	pink shrimp	marine	whole body	710	1,700	µg/kg ww	96 hours	water	mortality	Parrish et al (1976)	10% mortality at NOAEL 55% mortality at LOAEL (two replicates no statistics on mortality)
Chlordane	grass shrimp	marine	whole body	4,500	9,100	µg/kg ww	96 hours	water	mortality	Parrish et al (1976)	15% mortality at NOAEL 45% mortality at LOAEL (two reps no stats on mortality). Tissue concentration lower at NOAEL than at a lower concentration with 0 mortality (tissue concentration 4.8 µg/g ww)
DDTs	crayfish (adult)	fresh-water	whole body	46	na	µg/kg ww	3 days	water	mortality	Johnson et al. (1971)	not a toxicity test; accumulation test with 0% mortality

Analyte	Test Species	Marine or Fresh-water	Tissue Type	NOAEL	LOAEL	Unit	Exposure Duration	Exposure Route	Endpoint	Source	Notes
DDTs	pink shrimp	marine	whole body	na	60	µg/kg ww	56 days	water	mortality	Nimmo et al. (1970)	30% mortality (control: 17%), includes DDE, DDD, and DDT; shrimp collected from Escambia and Bay counties, Florida; uncertainty with high mortality in control and field collected shrimp
DDTs	blue crab (juvenile)	marine	whole body	26	200	µg/kg ww	5 weeks	treated food	metabolic rate	Leffler (1975)	dosed with 4,4'-DDT, but total DDT metabolites measured in residues; mortality reported as "chlorinated hydrocarbon poisoning;" crabs collected from estuarine Cedar Key, Florida; metabolic rate measured – not relevant endpoint
Heptachlor	pink shrimp	marine	whole body	10	30	µg/kg ww	96 hours	water	mortality	Schimmel et al. (1976)	5% mortality at NOAEL, 80% at LOAEL
Heptachlor	grass shrimp	marine	whole body	200	970	µg/kg ww	96 hours	water	mortality	Schimmel et al. (1976)	13% mortality at NOAEL, 70% at LOAEL
Heptachlor epoxide	pink shrimp	marine	whole body	54	180	µg/kg ww	96 hours	water	mortality	Schimmel et al. (1976)	5% mortality at NOAEL, 80% at LOAEL
Heptachlor epoxide	grass shrimp	marine	whole body	550	2500	µg/kg ww	96 hours	water	mortality	Schimmel et al. (1976)	13% mortality at NOAEL, 70% at LOAEL
Methoxychlor	blue crab	marine	whole body	na	340	µg/kg ww	mean duration 63 days		mortality	Bookhout et al (1976)	states that concentrations > 1.0 ppb larval mortality was nearly complete after third stage. 1.3 ppb first concentration above 1.0 ppb. Tissue concentration at LOAEL is lowest measured value out of 3.
Methoxychlor	mud-crab	marine	whole body	na	230	µg/kg ww	mean duration 20 days	water	mortality	Bookhout et al (1976)	states that all concentrations > 1.0 ppb gave significant first stage larval mortality. 2.5 ppb first concentration above 1.0 ppb and had 72 to 86% survival. Tissue concentration at LOAEL is lowest measured value out of 4.

Analyte	Test Species	Marine or Fresh-water	Tissue Type	NOAEL	LOAEL	Unit	Exposure Duration	Exposure Route	Endpoint	Source	Notes
Methoxychlor	Dungeness crab (adults)	marine	whole body	na	570	µg/kg ww	up to 15 days	water	mortality	Armstrong et al. (1976)	LOAEL concentration 4.0 ug/L measured body concentration exposed to a water concentration of 7.5 µg/L. No tissue measurement at LOAEL of 4.0 µg/L
Methoxychlor	Dungeness crab (juveniles)	marine	whole body	na	150	µg/kg ww	18 days	water	mortality	Armstrong et al. (1976)	LOAEL concentration 0.04 µg/L. Figure 8 measured tissue concentration after 18 days at LOAEL concentration.

COI – chemical of interest

ERED – environmental residue effects database

LOAEL – lowest-observed-adverse-effect level

na – not available

NOAEL – no-observed-adverse-effect level

PAH – polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

ppb – parts per billion

ppm – parts per million

ppt – parts per thousand

SVOC – semivolatile organic compound

ww – wet weight

REFERENCES

Armstrong DA, Buchanan DV, Mallon MH, Caldwell RS, Millemann RE. 1976. Toxicity of the insecticide methoxychlor to the dungeness crab *Cancer magister*. Mar Biol 38:239-252.

Bianchini A, Gilles R. 1996. Toxicity and accumulation of mercury in three species of crabs with different osmoregulatory capacities. Bull Environ Contam Toxicol 57:91-98.

Bookhout CG, Costlow Jr JD, Monroe R. 1976. Effects of methoxychlor on larval development of mud-crab and blue crab. Wat Air Soil Pollut 5:349-365.

Brown D, Thompson RS. 1982. Phthalates and the aquatic environment: Part I. The effect of di-2ethylhexyl phthalate (DEHP) and di-isodecyl phthalate (DIDP) on the reproduction of *Daphnia magna* and observations on their bioconcentration. Chemosphere 11:417-426.

Canli M, Furness RW. 1995. Mercury and cadmium uptake from seawater and from food by the Norway lobster *Nephrops norvegicus*. Environ Toxicol Chem 14(5):819-828.

Dillon TM. 1984. Biological consequences of bioaccumulation in aquatics animals: an assessment of the current literature. Waterways Experiment Station, US Army Corps of Engineers, Vicksburg, MS.

Duke TW, Lowe JL, Wilson AJ, Jr. 1970. A polychlorinated biphenyl (Aroclor 1254) in the water, sediment, and biota of Escambia Bay, Florida. Bull Environ Contam Toxicol 5:171-180.

Evans ML. 1980. Copper accumulation in the crayfish (*Orconectes rusticus*). Bull Environ Contam Toxicol 24:916-920.

Gillespie RB, Reisine T, Massaro EJ. 1977. Cadmium uptake by the crayfish, *Orconectes propinquus propinquus* (Girard). Environ Res 13:364-368.

Hansen DJ, Parrish PR, Forester J. 1974. Aroclor 1016: Toxicity to and uptake by estuarine animals. Environ Res 7:363-373.

Hobson JF, Carter DE, Lightner DV. 1984. Toxicity of a phthalate ester in the diet of a penaied shrimp. J Toxicol Env Health 13:959-968.

Jennings JR, Rainbow PS. 1979. Studies on the uptake of cadmium by the crab *Carcinus maenas* in the laboratory. I. Accumulation from seawater and a food source. Mar Biol 50:131-139.

Johnson BT, Saunders CR, Sanders HO. 1971. Biological magnification and degradation of DDT and aldrin by freshwater invertebrates. J Fish Res Board Can 28:705-709.

Leffler CW. 1975. Effects of ingested mirex and DDT on juvenile *Callinectes sapidus* Rathbun. Environ Pollut 8:283-300.

Lindsay DM, Sanders JG. 1990. Arsenic uptake and transfer in a simplified estuarine food chain. Environ Toxicol Chem 9:391-395.

Marshall JS. 1978. Population dynamics of *Daphnia galeata mendotae* as modified by chronic cadmium stress. J Fish Res Board Can 35:461-469.

Miramand P, Guary JC, Fowler SW. 1981. Uptake, assimilation, and excretion of vanadium in the shrimp, *Lysmata seticaudata* (Risso), and the crab, *Carcinus maenas* (L.). J Exp Mar Biol Ecol 49:267-287.

Mirenda RJ. 1986a. Acute toxicity and accumulation of zinc in the crayfish, *Orconectes virilis* (Hagen). Bull Environ Contam Toxicol 37:387-394.

Mirenda RJ. 1986b. Toxicity and accumulation of cadmium in the crayfish, *Orconectes virilis* (Hagen). Arch Environ Contam Toxicol 15:401-407.

Mortimer MR, Connell DW. 1994. Critical internal and aqueous lethal concentrations of chlorobenzenes with the crab *Portunus pelagicus* (L). Ecotoxicol Environ Saf 28:298-312.

Mortimer MR, Miller GJ. 1994. Susceptibility of larval and juvenile instars of the sand crab, *Portunus pelagicus* (L.) to sea water contaminated by chromium, nickel or copper. Aust J Mar Freshwater Res 45:1107-1121.

Nimmo DR, Wilson Jr AJ, Blackman RR. 1970. Localization of DDT in the body organs of pink and white shrimp. Bull Environ Contam Toxicol 5(4):333-341.

Nimmo DR, Forester J, Heitmuller PT, Cook GH. 1974. Accumulation of Aroclor 1254 in grass shrimp (*Palaemonetes pugio*) in laboratory and field exposures. Bull Environ Contam Toxicol 11(4):303-308.

Parrish PR, Schimmel SC, Hansen DJ, Patrick Jr JM, Forester J. 1976. Chlordane: effects on several estuarine organisms. J Toxicol Environ Health 1:485-494.

Reddy PS, Tuberty SR, Fingerman M. 1997. Effects of cadmium and mercury on ovarian maturation in the red swamp crayfish, *Procambarus clarkii*. Ecotoxicol Environ Saf 37:62-65.

Rule JH, Alden RW. 1996. Interactions of Cd and Cu in anaerobic estuarine sediments. II. Bioavailability, body burdens and respiration effects as related to geochemical partitioning. Environ Toxicol Chem 15(4):466-471.

Sanborn HR, Malins DC. 1977. Toxicity and metabolism of naphthalene: a study with marine larval invertebrates. Proc Soc Exper Biol Med 154:151-155.

Sanders HO, Chandler JH. 1972. Biological magnification of a polychlorinated biphenyl (Aroclor 1254) from water by aquatic invertebrates. Environ Contam Toxicol 7(5):257-263.

Schimmel SC, Patrick Jr JM, Forester J. 1976. Heptachlor: toxicity to and uptake by several estuarine organisms. J Toxicol Environ Health 1:955-965.

Thorp JH, Giesy JP, Wineriter SA. 1979. Effects of chronic cadmium exposure on crayfish survival, growth, and tolerance to elevated temperatures. Arch Environ Contam Toxicol 8:449-456.

Vernberg WB, DeCoursey PJ, Kelly M, Johns DM. 1977. Effects of sublethal concentrations of cadmium on adult *Palaemonetes pugio* under static and flow-through conditions. Bull Environ Contam Toxicol 17(1):16-24.

ATTACHMENT 6
SEDIMENT TOXICITY REFERENCE VALUES FOR CHLORDANE AND DDTS

Attachment 6 Sediment Toxicity Reference Values for Chlordane and DDTs

Analyte	Test Species	Marine or Fresh-water	NOAEL (µg/kg dw)	LOAEL (µg/kg dw)	Exposure Duration	Endpoint	Source	Notes
Chlordane	amphipod AET	marine	2.8	na	10 days	AET	Gries and Waldow (1996)	AET values for chlordane amphipod 2.8, echinoderm >4.5 µg/kg dw or 0.16 >0.26 mg/kg OC.
Chlordane	all components of the aquatic ecosystem (e.g., bacteria, algae, macrophytes, invertebrates, fish) were considered, if data were available	freshwater	4.5	8.87	na	ISQG for NOAEL PEL for LOAEL	Canadian environmental quality guidelines (CCME 2002)	Sediment guidelines published by Environment Canada. A comprehensive review of the scientific literature performed for each substance. Each toxicological study retrieved from the scientific literature was evaluated for its overall acceptability to ensure that high-quality data were used in developing SQGs.
Chlordane	all components of the aquatic ecosystem (e.g., bacteria, algae, macrophytes, invertebrates, fish) were considered, if data were available	marine	2.26	4.79	na	ISQG for NOAEL PEL for LOAEL	Canadian environmental quality guidelines (CCME 2002)	Sediment guidelines published by Environment Canada. A comprehensive review of the scientific literature performed for each substance. Each toxicological study retrieved from the scientific literature was evaluated for its overall acceptability to ensure that high-quality data were used in developing SQGs.
Chlordane	varied	freshwater	na	17.6	na	consensus-based PEC	Ingersoll et al. (2000)	Consensus-based PEC based on PEL 8.9, SEL 60, TEL 30, and ERM 6 µg/kg dw.
p,p-DDE	benthic AET	marine	9	na	na	AET	Gries and Waldow (1996)	AET values for p,p-DDE amphipod 62, echinoderm 9.3, benthic 9.0 µg/kg dw or 6.0, >7.3, 0.31 mg/kg OC
DDE (sum of p,p and o,p isomers)	all components of the aquatic ecosystem (e.g., bacteria, algae, macrophytes, invertebrates, fish) were considered, if data were available	freshwater	1.42	6.75	na	ISQG for NOAEL PEL for LOAEL	Canadian environmental quality guidelines (CCME 2002)	Sediment guidelines published by Environment Canada. A comprehensive review of the scientific literature performed for each substance. Each toxicological study retrieved from the scientific literature was evaluated for its overall acceptability to ensure that high-quality data were used in developing SQGs.
DDE (sum of p,p and o,p isomers)	all components of the aquatic ecosystem (e.g., bacteria, algae, macrophytes, invertebrates, fish) were considered, if data were available	marine	2.07	374	na	ISQG for NOAEL PEL for LOAEL	Canadian environmental quality guidelines (CCME 2002)	Sediment guidelines published by Environment Canada. A comprehensive review of the scientific literature performed for each substance. Each toxicological study retrieved from the scientific literature was evaluated for its overall acceptability to ensure that high-quality data were used in developing SQGs.
sum DDE	varied	freshwater	na	31.3	na	consensus-based PEC	Ingersoll et al. (2000)	Consensus-based PEC based on PEL 6.75, SEL 190, TEL 50, and ERM 15 µg/kg dw.

Analyte	Test Species	Marine or Fresh-water	NOAEL (µg/kg dw)	LOAEL (µg/kg dw)	Exposure Duration	Endpoint	Source	Notes
4,4- DDE	varied	marine and estuary	2.2	27	na	ERL for NOAEL ERM for LOAEL	Long et al. (1995)	Marine and estuarine data compilation BEDS from lab spike sediment bioassays; field studies, and equilibrium partitioning studies; NOTE- relationship between incidence of effects and the concentration of total DDT and 4,4'-DDE was poor (only 50% incidence of effects were observed above the median concentration; for other chemicals incidence was 80 to 100% above the median concentration).
p,p,-DDD	benthic AET	marine	16	na	na	AET	Gries and Waldow (1996)	AET values for p,p-DDD amphipod 63, echinoderm 28, benthic 16 µg/kg dw or 3.1, 1.6, 1.0 mg/kg OC.
DDD (sum of p,p and o,p isomers)	all components of the aquatic ecosystem (e.g., bacteria, algae, macrophytes, invertebrates, fish) were considered, if data were available	freshwater	3.54	8.51	na	ISQG for NOAEL PEL for LOAEL	Canadian environmental quality guidelines (CCME 2002)	Sediment guidelines published by Environment Canada. A comprehensive review of the scientific literature performed for each substance. Each toxicological study retrieved from the scientific literature was evaluated for its overall acceptability to ensure that high-quality data were used in developing SQGs.
DDD (sum of p,p and o,p isomers)	all components of the aquatic ecosystem (e.g., bacteria, algae, macrophytes, invertebrates, fish) were considered, if data were available	marine	1.22	7.81	na	ISQG for NOAEL PEL for LOAEL	Canadian environmental quality guidelines (CCME 2002)	Sediment guidelines published by Environment Canada. A comprehensive review of the scientific literature performed for each substance. Each toxicological study retrieved from the scientific literature was evaluated for its overall acceptability to ensure that high-quality data were used in developing SQGs.
sum DDD	varied	freshwater	na	28	na	consensus-based PEC	Ingersoll et al. (2000)	Consensus-based PEC based on PEL 8.51, SEL 60, TEL 60, and ERM 20 µg/kg dw
DDD	*Hyalella azteca*	freshwater	1,200	4,000	42 days	survival	Ingersoll et al. (2005)	Sediment concentrations reported as µg/g OC. TOC = 1%.
DDD	*Hyalella azteca*	freshwater	na	30	42 days	growth	Ingersoll et al. (2005)	Sediment concentrations reported as µg/g OC. TOC = 1%.
DDD	*Hyalella azteca*	freshwater	1,200	4,000	42 days	reproduction	Ingersoll et al. (2005)	Sediment concentrations reported as µg/g OC. TOC = 1%.
DDT	*Rhepoxynius abronius*	marine	na	6,180	10 days	survival (LC50)	Murdoch et al. (1997)	Reported as LC50 6.18 mg/kg dw.
DDT (sum of p,p and o,p isomers)	all components of the aquatic ecosystem (e.g., bacteria, algae, macrophytes, invertebrates, fish) were considered, if data were available	marine	1.19	4.77	na	ISQG for NOAEL PEL for LOAEL	Canadian environmental quality guidelines (CCME 2002)	Sediment guidelines published by Environment Canada. A comprehensive review of the scientific literature performed for each substance. Each toxicological study retrieved from the scientific literature was evaluated for its overall acceptability to ensure that high-quality data were used in developing SQGs.

Analyte	Test Species	Marine or Fresh-water	NOAEL (µg/kg dw)	LOAEL (µg/kg dw)	Exposure Duration	Endpoint	Source	Notes
DDT (sum of p,p and o,p isomers)	all components of the aquatic ecosystem (e.g., bacteria, algae, macrophytes, invertebrates, fish) were considered, if data were available	freshwater	1.19	4.77	na	ISQG for NOAEL PEL for LOAEL	Canadian environmental quality guidelines (CCME 2002)	Sediment guidelines published by Environment Canada. A comprehensive review of the scientific literature performed for each substance. Each toxicological study retrieved from the scientific literature was evaluated for its overall acceptability to ensure that high-quality data were used in developing SQGs.
sum DDT	varied	freshwater	na	62.9	na	consensus-based PEC	Ingersoll et al. (2000)	Consensus-based PEC based on SEL 170, TET 50, and ERM 7 µg/kg dw.
sum DDT	varied	freshwater	na	710	na	SEL	Ingersoll et al. (2000)	Compilation of matching sediment chemistry and biological effects data from numerous studies.
sum DDT	varied	freshwater	na	50	na	TET	Ingersoll et al. (2000)	Compilation of matching sediment chemistry and biological effects data from numerous studies.
sum DDT	varied	freshwater	na	7	na	ERM	Ingersoll et al. (2000)	Compilation of matching sediment chemistry and biological effects data from numerous studies.
DDTs	*Neanthes arenaceodentata*	marine	na	308,000	28 days	growth	Lotufo et al. (2000)	Target concentration 18.5 mg/g OC. Measured at test start 13.83 mg/g OC and end 11.59 mg/g OC. TOC 2.66%. Used test end sediment concentration to calculate LOAEL.
DDTs	*Hyalella azteca*	freshwater	na	3,510	28 days	survival (LC50)	Lotufo et al. (2001b)	Reported as 9.9 nmol DDT equivalents/g dw. Molecular weight for DDT 354.51.
DDTs	*Leptocheirus plumulosus*	marine	na	1,985	10 days	survival (LC50)	Lotufo et al. (2001a)	Reported as 5.6 nmol/g dw (5.5 to 5.7 95% confidence interval).
DDTs	*Hyalella azteca*	freshwater	567	1,063	28 days	survival	Lotufo et al. (2001b)	Significant and non-significant difference in survival from control (LOAEL and NOAEL). Reported as 3.0 and 1.6 nmol DDT equivalents/g dw, respectively. Molecular weight for DDT 354.51.
DDTs	*Diporeia* spp.	freshwater	na	2,910	28 days	survival (LC50)	Lotufo et al. (2001b)	Reported as 8.2 nmol DDT equivalents/g dw. Molecular weight for DDT 354.51.
Total DDTs	amphipod AET	marine	24	na	10 days	AET	Gries and Waldow (1996)	AET values for total DDT amphipod 24, echinoderm 37µg/kg dw or 1.4, 8.8 mg/kg OC.
Total DDTs	varied	freshwater	na	572	na	consensus-based PEC	Ingersoll et al. (2000)	Consensus-based PEC based on PEL 4450, SEL 120, TET na, and ERM 350 µg/kg dw.
Total DDT	*Neanthes arenaceodentata*	marine	8510	na	120 days	Full life-cycle toxicity test (survival, growth, fecundity, reproduction)	Chapman (1996)	Based on field collected sediment; alternative predictive approach estimated that above 8.51 mg/kg dw total DDT, adverse effects would be expected to occur; OC normalized concentrations are also provided.

Analyte	Test Species	Marine or Fresh-water	NOAEL (µg/kg dw)	LOAEL (µg/kg dw)	Exposure Duration	Endpoint	Source	Notes
Total DDT	varied	marine and estuary	1.58	46.1	na	ERL for NOAEL ERM for LOAEL	Long et al. (1995)	Marine and estuarine data compilation BEDS (biological effects database for sediment) from lab spike sediment bioassays; field studies, and equilibrium partitioning studies. Note: relationship between incidence of effects and the concentration of total DDT and 4,4'-DDE was poor (only 50% incidence of effects were observed above the median concentration; for other chemicals incidence was 80 to 100% above the median concentration).
Total DDT	marine polychaete (*Heteromastus filiformis*)	marine	na	7,500	28 days	significant reduced feeding rate	Mulsow & Landrum (1995)	Laboratory 28-day bioaccumulation study using spiked sediment; no effect on feeding rate in polychaetes exposed to 4.0 mg/kg dw total DDT.
Total DDT	*Hyalella azteca*	freshwater	na	11,000	10 days	LC50	Nebeker et al. (1989)	Spiked sediment toxicity tests with ranging TOC (3-11%); at 5 mg/kg dw (TOC ranging from 3 to 7.2%) mortality is less than 10%.

AET – apparent effects threshold (highest no-hit value)

BEDS – biological effects database for sediment

ERL – effects range – low

ERM – effects range – medium

ISQG – interim sediment quality guideline

LC50 – concentration that causes the death of 50% of a group of test animals

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

na – not available

PEC – probable effect concentration

PEL – probable effects level

SEL – severe effects level

SQG – sediment quality guideline

TEL – threshold effects level

TET – toxic effect threshold

REFERENCES

CCME. 2002. Canadian sediment quality guidelines for the protection of aquatic life: summary tables. Canadian Environmental Quality Guidelines, 1999, updated 2001, updated 2002 [online]. Canadian Council of Ministers of the Environment, Winnipeg, MB. Updated 2002. Available from: http://www.ccme.ca/assets/pdf/e7_002.pdf.

Chapman PM. 1996. A test of sediment effects concentrations: DDT and PCB in the Southern California Bight. Environ Toxicol Chem 15(7):1197-1198.

Gries TH, Waldow KH. 1996. Progress re-evaluating Puget Sound apparent effects thresholds (AETs). Volume I: 1994 amphipod and echinoderm larval AETs. Draft report. Prepared for Puget Sound Dredged Disposal Analysis (PSDDA) agencies. Washington Department of Ecology, Olympia, WA.

Ingersoll CG, MacDonald DD, Wang N, Crane JL, Field LJ, Haverland PS, Kemble NE, Lindskoog RA, Severn C, Smorong DE. 2000. Prediction of sediment toxicity using consensus-based freshwater sediment quality guidelines. US Environmental Protection Agency, Chicago, IL.

Ingersoll CG, Wang N, Hayward JMR, Jones JR, Jones SB, Ireland DS. 2005. A field assessment of long-term laboratory sediment toxicity tests with the amphipod *Hyalella azteca*. Environ Toxicol Chem 24(11):2853-2870.

Long ER, MacDonald DD, Smith SL, Calder FD. 1995. Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments. Environ Manage 19:81-97.

Lotufo GR, Farrar JD, Bridges TS. 2000. Effects of exposure source, worm density, and sex on DDT bioaccumulation and toxicity in the marine polychaete *Neanthes arenaceodentata*. Environ Toxicol Chem 19(2):472-484.

Lotufo GR, Farrar JD, Duke BM, Bridges TS. 2001a. DDT toxicity and critical body residue in the amphipod *Leptocheirus plumulosus* in exposures to spiked sediment. Arch Environ Contam Toxicol 41:142-150.

Lotufo GR, Landrum PF, Gedeon ML. 2001b. Toxicity and bioaccumulation of DDT in freshwater amphipods in exposures to spiked sediments. Environ Toxicol Chem 20(4):810-825.

Mulsow SG, Landrum P. 1995. Bioaccumulation of DDT in a marine polychaete, the conveyor-belt deposit feeder *Heteromastus filiformis* (Claparede). Chemosphere 31(4):3141-3152.

Murdoch MH, Chapman PM, Norman DM, Quintino VM. 1997. Spiking sediment with organochlorines for toxicity testing. Environ Toxicol Chem 16(7):1504-1509.

Nebeker AV, Schuytema GS, Griffis WL, Barbitta JA, Carey LA. 1989. Effect of sediment organic carbon on survival of *Hyalella azteca* exposed to DDT and endrin. Environ Toxicol Chem 8:705-18.

ATTACHMENT 7
SUMMARY OF TOXICITY DATA FOR VOLATILE ORGANIC COMPOUNDS IN POREWATER

Attachment 7 Summary of Toxicity Data for Volatile Organic Compounds in Porewater

A literature search for relevant toxicity studies for the 16 volatile organic compounds (VOCs) detected in porewater samples was conducted using two databases, ECOTOX and BIOSIS. The following sections describe each database, the methods used to search the databases, and the search results. This information was first presented in the porewater data and analysis report (Windward 2006).

DATABASE DESCRIPTIONS

ECOTOX includes three formerly independent databases, AQUIRE, TERRETOX, and PHYTOTOX. The AQUIRE database is the most relevant source available for aquatic toxicology data. AQUIRE includes lethal, sublethal, and residue effects data for saltwater and freshwater aquatic species, with coverage spanning from 1915 to the present. AQUIRE includes more than 227,800 records for more than 7,300 chemicals and 3,700 freshwater and marine organisms. Sources for studies in AQUIRE include journal articles, published and unpublished reports, miscellaneous government databases and files, gray literature, and independent laboratory test results.

For studies to be included in ECOTOX, the author(s) must report the species, chemical information, and exposure duration. Only single chemical studies are included (i.e., studies testing chemical mixtures are excluded). The majority of the literature is from 1972 to the present. ECOTOX is continually updated, but there may be a time lag of up to six months between the time the literature search is conducted and entry of new studies into ECOTOX. Therefore, EPA recommends searching other sources for literature published within the last year.

Therefore, BIOSIS was searched to supplement the ECOTOX search results. BIOSIS is the most comprehensive database available based on the number of records, the number of records added annually, and the number of journals reviewed (6,000 journal titles). BIOSIS contains references to primary journal literature in the biological sciences (botany, ecology, and zoology) as well as medical research. BIOSIS coverage is from 1969 to the present, and it currently includes over 15,000,000 records, with 600,000 new records added each year. BIOSIS does not include unpublished reports, government studies, or independent laboratory results.

DATABASE SEARCH METHODS

The ECOTOX database was searched first for relevant toxicity studies involving invertebrate species with growth, mortality (including immobilization), reproductive, or developmental endpoints. Studies with invertebrates were preferred because, as discussed in the Work Plan (Windward 2004), the purpose of the porewater study is to evaluate risk to benthic invertebrates. However, if there were no invertebrate data in

the ECOTOX database, then fish studies with growth, mortality (including immobilization), reproductive, or developmental endpoints were included in the search. Toxicity data for both freshwater and marine species were included because of the wide salinity range in the Lower Duwamish Waterway (LDW).

After searching ECOTOX, BIOSIS was searched to identify articles published in 2003 to 2005 for each of the 16 VOCs of interest. To make the search comprehensive, each chemical name, its Chemical Abstracts Service (CAS) number, and any chemical name synonyms were searched on BIOSIS for the years 2003 to 2005. In addition, for three VOCs with very few or no data in ECOTOX (1,1-dichloroethane, 1,2-dichloroethene, and vinyl chloride), a more extensive search was conducted using BIOSIS with no restrictions on publication date.

STUDY SELECTION

The studies identified in the literature search were reviewed to assess the most appropriate toxicity data for comparison with the porewater concentrations of individual VOCs. In many cases, very few toxicity data were available. In particular, relatively few data were available for certain chemicals for a range of families, as required for derivation of ambient water quality criteria (AWQC) (Stephan et al. 1985), and very few lowest-observed-adverse-effects levels (LOAELs) were available. Instead, many studies typically reported lethal concentrations for 50% of a test population (i.e., LC50 values) and no-observed-adverse-effects levels (NOAELs) for a range of different species. In addition, only one or two studies were identified for some of the VOCs. Therefore, because of the limitations of the available dataset, the rules for study selection had to be modified from those generally used to select toxicity reference values (TRVs). Ideally, TRVs would be selected by identifying the lowest LOAEL and the highest NOAEL below that LOAEL for the same relevant test species. However, because the available data generally did not support this approach, best professional judgment was required to select the most appropriate toxicity values for each VOC of interest. The following general rules were followed:

- If available, the study identified with the lowest effects level (preferably a LOAEL) for each VOC was selected and reviewed
- If available, the study with the lowest NOAEL for each VOC was selected and reviewed, provided that more one NOAEL was not available for a given test species and endpoint. When there were multiple NOAELs for the same test species/endpoint for a given VOC, the study with the highest NOAEL for that test species/endpoint was selected and reviewed.

In many cases, the available toxicity data were uncertain because LOAELs were rarely reported, NOAELs were dependent on the chosen test dilution series, and very few studies reported both effect and no-effect concentrations for a single species and

endpoint.[1] However, rather than assigning uncertainty factors to the toxicity data in this memorandum, the actual NOAELs, LOAELs, and LC50s are reported.

STUDY ACCEPTABILITY

Criteria were established to determine the acceptability of each study. These criteria are based on Stephan et al. (1985), which presents guidelines used to develop Environmental Protection Agency's (EPA's) AWQC, and on Suter and Tsao (1996), which presents methods used to derive Tier II values.[2] Two types of criteria were developed in which some criteria were required, while others were preferred, but not required. Studies had to meet the following required criteria to be accepted:

- Negative control tests must be used.
- For a no-effect result, the exposure period must be no less than 48 hours for daphnids, and no less than 96 hours for fish.
- The salinity must be no greater than 35 parts per thousand in tests using *Artemia salina* (brine shrimp), to represent conditions relevant to those found in the LDW.

The following additional criteria were preferred, but not required, for a study to be accepted:

- Test containers should be covered to minimize volatilization.[3]
- Standard test methods should be used for those tests/organisms/endpoints that have standardized protocols.
- Reported effects in tests should be compared to controls using statistical methods.
- Control media and test media should be identical in all respects except for the treatment.

---

[1] The actual effect threshold is uncertain for a NOAEL without a corresponding LOAEL.

[2] Tier II values are secondary acute and chronic values derived for chemicals with some toxicity data, but not enough to meet the data requirements for development of AWQC.

[3] Most of the studies reviewed noted that test vessels were covered to minimize volatilization during toxicity testing. However, a few studies did not comment on whether test vessels were covered. If test vessels were not covered during testing, but the VOC concentration was measured throughout the study, coverage should not impact the result as long as the concentration remained stable. On the other hand, if the concentration was not measured or the concentration declined over the course of the study, then the effects concentration could be an overestimate (i.e., effects could be occurring at a lower concentration than the nominal concentration). Therefore, although the lack of coverage documentation leads to additional uncertainty, rejection of a study (without another study available at a lower effects concentration) would lead to the selection of study with a higher effects concentration which could be argued to be less protective. Therefore, this criterion was included in the preferred category rather than the required category.

- Flow-through tests are preferred, followed by static renewal and then static tests.
- Organisms should be less than 24 hours old for acute and chronic daphnid tests and up to third instar for midge acute tests; juvenile or larval life stages are preferred for acute tests for other organisms.
- Chemical concentrations should be measured in the test rather than presented as nominal concentrations.
- The chemistry of the test environment (e.g., dissolved oxygen (DO), pH, temperature, salinity) should be reported.
- DO should not fall below 40% saturation in static tests and should not fall below 60% saturation in flow-through tests.
- Test organisms should not be fed during acute tests, with the exception of saltwater annelids and mysids.
- Methods used to prepare treatment solutions should be described in detail.
- The study should report dose-related toxicity information, and preferably report both a NOAEL and LOAEL.
- The chemical source and purity should be noted.

LITERATURE SEARCH RESULTS

Table 1 summarizes the results of the literature search. NOAELs and LOAELs found in the literature search are presented. Because a large number of the available values for some chemicals represented LC50, only the lowest LC50 for a given test organism is shown in Table 1.

In the course of the literature search, several papers were identified that established NOAELs for specific VOCs (e.g., De Rooij et al. 2004a, b; Van Wijk et al. 2004). These papers contained summaries of aquatic toxicity studies, which the authors classified as either: 1) valid without restriction, 2) valid with restrictions to be considered with care, 3) invalid, or 4) not assignable. Only those studies classified as either 1 or 2 are included in Table 1.

Table 1. Aquatic toxicity data obtained from the literature search

VOC	Reported Concentration (µg/L)	Effect Level	Test Organism	Test Duration	Endpoint	Source[a]
1,1-Dichloroethane	202,000	LC50	*Poecilia reticulate* (guppy)	14d	mortality	Könemann (1981)
	7,800	FCV	Based on narcosis model	na	na	DiToro et al. (2000)
	39,600	FAV	Based on narcosis model	na	na	DiToro et al. (2000)
1,1-Dichloroethene	**2,400**	NOAEL	*Daphnia magna* (water flea)	48h	mortality	LeBlanc (1980)
	11,600	LC50	*Daphnia magna* (water flea)	48h	mortality	Dill et al. (1980)
	79,000	LC50	*Daphnia magna* (water flea)	48h	mortality	LeBlanc (1980)
	224,000	LC50	*Americamysis bahia* (opossum shrimp)	96h	mortality	ECOTOX, Reference No. 9607
1,2-dichlorobenzene	1970	LC50	*Americamysis bahia* (opossum shrimp)	96h	mortality	ECOTOX, Reference No. 9607
	26,000	LD50	*Cloeon dipterum* (may fly)	24h	mortality	ECOTOX, Reference No. 6954
	26,000	LD50	*Cloeon dipterum* (may fly)	48h	mortality	ECOTOX, Reference No. 6954
	1,000	LOEL	*Crassostrea virginica* (American oyster)	24h	growth	ECOTOX, Reference No. 3708
	550	EC50	*Daphnia magna* (water flea)	14d	reproduction	ECOTOX, Reference No. 15526
	2,400	LC50	*Daphnia magna* (water flea)	24h	mortality	ECOTOX, Reference No. 5184
	68,000	LC50	*Daphnia magna* (water flea)	24h	mortality	ECOTOX, Reference No. 5718
	2,400	LC50	*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 5184
	2,200	LC50	*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 6629
	630	NOAEL	*Daphnia magna* (water flea)	21d	reproduction	ECOTOX, Reference No. 847
	> 100,000	EC50	*Mercenaria mercenaria* (hard clam)	48h	develop	ECOTOX, Reference No. 2400
	> 100,000	EC50	*Mercenaria mercenaria* (hard clam)	12d	mortality	ECOTOX, Reference No. 2400
	14,300	LC50	*Palaemonetes pugio* (grass shrimp)	24h	mortality	ECOTOX, Reference No. 875
	10,300	LC50	*Palaemonetes pugio* (grass shrimp)	48h	mortality	ECOTOX, Reference No. 875
	9,400	LC50	*Palaemonetes pugio* (grass shrimp)	96h	mortality	ECOTOX, Reference No. 875
	10,000	LC50	*Palaemonetes pugio* (grass shrimp)	96h	mortality	ECOTOX, Reference No. 2965
	2,300	EC50	*Tanytarsus dissimilis* (midge)	48h	mortality	ECOTOX, Reference No. 16044
	19,900	LC50	*Tanytarsus dissimilis* (midge)	24h	mortality	ECOTOX, Reference No. 10579
	12,000	LC50	*Tanytarsus dissimilis* (midge)	48h	mortality	ECOTOX, Reference No. 10579
	11,800	LC50	*Tanytarsus dissimilis* (midge)	48h	mortality	ECOTOX, Reference No. 16044
	51,000	EC50	*Tetrahymena pyriformis* (ciliate)	24h	growth	ECOTOX, Reference No. 11258
1,2-Dichloroethane	**6,927**	LC50	*Artemia salina* (brine shrimp)	72h	mortality	Sanchez-Fortun et al. (1997)
	11,000	NOAEL	*Daphnia magna* (water flea)	28d	reproduction	Richter et al. (1983)
	42,000	NOAEL	*Daphnia magna* (water flea)	28d	growth	Richter et al. (1983)
	< 68,000	NOAEL	*Daphnia magna* (water flea)	28d	mortality	Richter et al. (1983)

VOC	Reported Concentration (µg/L)	Effect Level	Test Organism	Test Duration	Endpoint	Source[a]
	> 100,000	LC50	*Gammarus fasciatus* (scud)	96h	mortality	ECOTOX, Reference No. 6797
	> 100,000	LC50	*Pteronarcys californicus* (stonefly)	96h	mortality	ECOTOX, Reference No. 6797
	113,000	LC50	*Americamysis bahia* (opossum shrimp)	96h	mortality	ECOTOX, Reference No. 9607
	186,000	LC50	*Elminius modestus* (Australian barnacle)	48h	mortality	ECOTOX, Reference No. 13535
	220,000	LC50	*Daphnia magna* (water flea)	48h	mortality	Richter et al. (1983)
	900,000	LC50	*Ophryotrocha labronica* (polychaete)	96h	mortality	ECOTOX, Reference No. 5902
1,2-Dichloropropane	**42,000**	LC50	*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 5184
	> 100,000	LC50	*Crangon crangon* (sand shrimp)	48h	mortality	ECOTOX, Reference No. 906
	58,000	LC50	*Daphnia magna* (water flea)	24h	mortality	ECOTOX, Reference No. 5184
	53,000	LC50	*Elminius modestus* (Australian barnacle)	48h	mortality	ECOTOX, Reference No. 19535
	< 22,000		*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 5184
1,4-Dichlorobenzene	1,990	LC50	*Americamysis bahia* (opossum shrimp)	96h	mortality	ECOTOX, Reference No. 9607
	93	LC50	*Artemia salina* (brine shrimp)	24h	mortality	ECOTOX, Reference No. 11926
	16,500	EC50	*Brachionus calycilorus* (rotifer)	48h	reproduction	ECOTOX, Reference No. 20484
	3125	NOAEL	*Brachionus calycilorus* (rotifer)	48h	reproduction	ECOTOX, Reference No. 20484
	9,400	LC50	*Chironomus thummi* (midge)	48h	mortality	ECOTOX, Reference No. 4072
	940	NOAEL	*Chironomus thummi* (midge)	48h	mortality	ECOTOX, Reference No. 4072
	930	EC50	*Daphnia magna* (water flea)	14d	reproduction	ECOTOX, Reference No. 15526
	17,000	LC50	*Daphnia magna* (water flea)	24h	mortality	ECOTOX, Reference No. 5184
	9,800	LC50	*Daphnia magna* (water flea)	24h	mortality	ECOTOX, Reference No. 12055
	19,400	LC50	*Daphnia magna* (water flea)	24h	mortality	ECOTOX, Reference No. 12055
	14,800	LC50	*Daphnia magna* (water flea)	24h	mortality	ECOTOX, Reference No. 12055
	6,600	LC50	*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 5184
	9,900	LC50	*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 12055
	11,500	LC50	*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 12055
	9,500	LC50	*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 12055
	2,200	LC50	*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 6629
	300	NOAEL	*Daphnia magna* (water flea)	21d	reproduction	ECOTOX, Reference No. 847
	400	NOAEL	*Daphnia magna* (water flea)	21d	reproduction	ECOTOX, Reference No. 20484
	100,000	LC50	*Palaemonetes pugio* (grass shrimp)	48h	mortality	ECOTOX, Reference No. 875
	60,500	LC50	*Palaemonetes pugio* (grass shrimp)	96h	mortality	ECOTOX, Reference No. 875
	36,000	LC50	*Palaemonetes pugio* (grass shrimp)	96h	mortality	ECOTOX, Reference No. 2965
	147	LC50	*Portunus pelagicus* (crab)	10h	mortality	ECOTOX, Reference No. 4745
	14,700	LC50	*Portunus pelagicus* (crab)	1000h	mortality	ECOTOX, Reference No. 4745

VOC	Reported Concentration (µg/L)	Effect Level	Test Organism	Test Duration	Endpoint	Source[a]
	3,000	EC50	*Tanytarsus dissimilis* (midge)	2h	mortality	ECOTOX, Reference No. 16044
	19,200	EC50	*Tanytarsus dissimilis* (midge)	24h	mortality	ECOTOX, Reference No. 10579
	10,900	EC50	*Tanytarsus dissimilis* (midge)	48h	mortality	ECOTOX, Reference No. 10579
	13,000	EC50	*Tanytarsus dissimilis* (midge)	48h	mortality	ECOTOX, Reference No. 16044
	32,000		*Chironomus thummi* (midge)	48h	mortality	ECOTOX, Reference No. 4072
	680		*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 5184
	400		*Daphnia magna* (water flea)	28f	reproduction	ECOTOX, Reference No. 10712
1,2-Dichloroethene (cis or trans)	**6,785**	LC50	*Artemia salina* (brine shrimp)	72h	mortality	Sanchez-Fortun et al. (1997)
	140,000	LC50	*Lepomis macrochirus* (bluegill)	24-96h	mortality	ECOTOX, Reference No. 590
Benzene	180	NOAEL	*Cancer magister* (Dungeness crab)	24d	mortality, growth, development	Caldwell et al. (1976)
	1,100	LC100	*Cancer magister* (Dungeness crab)	24d	mortality	Caldwell et al. (1976)
	2,968	NOAEL	*Ceriodaphnia dubia* (water flea)	7d	reproduction	Niederlehner et al. (1998)
	6,000	LC50	*Gammarus* sp. (scud)	96h	mortality	ECOTOX, Reference No. 13419
	10,000	LC50	*Ischnura elegans* (damselfly)	48h	mortality	ECOTOX, Reference No. 15788
	12,419	LC50	*Ceriodaphnia dubia* (water flea)	7d	mortality	Niederlehner et al. (1998)
	15,000	LC50	*Daphnia pulex* (water flea)	96h	mortality	ECOTOX, Reference No. 15337
	16,796	LC50	*Crangon franciscorum* (shrimp)	96h	mortality	ECOTOX, Reference No. 558
	21,000	LC50	*Artemia salina* (brine shrimp)	48h	mortality	ECOTOX, Reference No. 2408
	27,000	LC50	*Palaemonetes pugio* (shrimp)	96h	mortality	ECOTOX, Reference No. 420
	34,000	LC50	*Cloeon dipterum* (mayfly)	48h	mortality	ECOTOX, Reference No. 15788
	48,000	LC50	*Corixa* sp. (water boatman)	48h	mortality	ECOTOX, Reference No. 15788
	59,600	LC50	*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 7069
	71,000	LC50	*Culex pipiens* (northern house mosquito)	48h	mortality	ECOTOX, Reference No. 10574
	74,000	LC50	*Dugesia lugubris* (vortex worm)	48h	mortality	ECOTOX, Reference No. 15788
	82,000	LC50	*Diaptomus forbesi* (copepod)	≤96h	mortality	ECOTOX, Reference No. 7800
	100,000	LC50	*Chironomus thummi* (midge)	48h	mortality	ECOTOX, Reference No. 15788
	111,800	LC100	*Homarus americanus* (American lobster)	1h	mortality	ECOTOX, Reference No. 5353
	120,000	LC50	*Asellus aquaticus* (aquatic sowbug)	48h	mortality	ECOTOX, Reference No. 15788
	130,000	LC50	*Nemoura cinerea* (stonefly)	48h	mortality	ECOTOX, Reference No. 15788
	190,000	LC50	*Katelysia opima* (marine bivalve)	96h	mortality	ECOTOX, Reference No. 9017
	200,000	LC50	*Aedes aegypti* (yellow fever mosquito)	48h	mortality	ECOTOX, Reference No. 14863
	230,000	LC50	*Lymnaea stagnalis* (great pond snail)	48h	mortality	ECOTOX, Reference No. 15788
	> 320,000	LC50	*Erpobdella octoculata* (leech)	48h	mortality	ECOTOX, Reference No. 15788
	> 320,000	LC50	*Tubificidae* (oligochaete)	48h	mortality	ECOTOX, Reference No. 15788

VOC	Reported Concentration (µg/L)	Effect Level	Test Organism	Test Duration	Endpoint	Source[a]
	356,000	LC50	*Daphnia cucullata* (water flea)	48h	mortality	ECOTOX, Reference No. 2017
	377,000	LC50	*Crassostrea gigas* (Pacific oyster)	48h	mortality	ECOTOX, Reference No. 8621
	550,000[b]	LC50	*Lymnaea stagnalis* (great pond snail)	120h	mortality	ECOTOX, Reference No. 13419
Carbon disulfide	**1,900**	LC50	*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 11455
Chlorobenzene	320	NOAEL	*Daphnia magna* (water flea)	16d	reproduction	Hermens et al. (1984)
	< 1,400	NOAEL	*Daphnia magna* (water flea)	10d	mortality	Cowgill and Milazzo (1991)
	2,500	EC50	*Daphnia magna* (water flea)	14d	reproduction	Calamari et al. (1983)
	3,890	NOAEL	*Ceriodaphnia dubia* (water flea)	7-10d	mortality	ECOTOX, Reference No. 212
	12,000	NOAEL	*Ceriodaphnia dubia* (water flea)	7-10d	reproduction	ECOTOX, Reference No. 212
	16,400	LC50	*Americamysis bahia* (opossum shrimp)	96h	mortality	ECOTOX, Reference No. 9607

VOC	Reported Concentration (µg/L)	Effect Level	Test Organism	Test Duration	Endpoint	Source[a]
Tetrachloroethene	**331**	NOAEL	*Ceriodaphnia dubia* (water flea)	7d	reproduction	Niederlehner et al. (1998)
	332	LC50	*Artemia salina* (brine shrimp)	72h	mortality	Sanchez-Fortun et al. (1997)
	400	NOAEL	*Daphnia magna* (water flea)	21d	reproduction	ECOTOX, Reference No. 56345
	829	LC50	*Ceriodaphnia dubia* (water flea)	7d	mortality	Niederlehner et al. (1998)
	900	EC50	*Dugesia japonica* (flatworm)	7d	growth	ECOTOX, Reference No. 12513
	1,300	LC50	*Palaemonetes pugio* (grass shrimp)	96h	mortality	ECOTOX, Reference No. 14563
	1,300	LC50	*Nereis arenaceodentata* (polychaete)	96h	mortality	ECOTOX, Reference No. 14563
	1,400	LC50	*Dugesia japonica* (flatworm)	7d	mortality	ECOTOX, Reference No. 12513
	1,800	LC50	*Moina macrocopa* (water flea)	3h	mortality	ECOTOX, Reference No. 12513
	3,500	LC50	*Elminius modestus* (Australian barnacle)	48h	mortality	ECOTOX, Reference No. 19535
	3,600	LC50	*Tallaperia maria* (stonefly)	96h	mortality	ECOTOX, Reference No. 14563
	7,000	LC50	*Tanytarsus dissimilis* (midge)	48h	mortality	ECOTOX, Reference No. 16044
	9,100	LC50	*Gammarus annulata* (scud)	96h	mortality	ECOTOX, Reference No. 14563
	9,100	LC50	*Daphnia magna* (water flea)	48h	mortality	ECOTOX, Reference No. 15981
	10,200	LC50	*Americamysis bahia* (opossum shrimp)	96h	mortality	ECOTOX, Reference No. 9607
	13,200	LC50	*Acartia tonsa* (calanoid copepod)	96h	mortality	ECOTOX, Reference No. 14563
	17,400	LC50	*Crangon septemspinosa* (bay shrimp)	96h	mortality	ECOTOX, Reference No. 14563
	21,600	NOAEL	*Gammarus minus* (scud)	96h	mortality	ECOTOX, Reference No. 14563
	93,400	LC50	*Physa heterostropha* (pond snail)	96h	mortality	ECOTOX, Reference No. 14563

VOC	Reported Concentration (µg/L)	Effect Level	Test Organism	Test Duration	Endpoint	Source[a]
Toluene (methylbenzene)	**737**	NOAEL	*Ceriodaphnia dubia* (water flea)	7d	reproduction	Niederlehner et al. (1998)
	1,000	NOAEL	*Daphnia magna* (water flea)	21d	reproduction	ECOTOX, Reference No. 847
	5,600	NOAEL	*Chironomus thummi*	48h	mortality	ECOTOX, Reference No. 4072
	9,500	LC50	*Palaemonetes pugio* (daggerblade shrimp)	96h	mortality	Tatem and Johnson (1978)
	14,700	LC50	*Eualus* sp. (shrimp)	96h	mortality	Korn et al. (1979)
	23,500	LC50	*Hemigrapsus nudus* (shore crab)	8d	mortality	ECOTOX, Reference No. 12879
	24,200	LC50	*Nitocra spinipes* (harpacticoid copepod)	96h	mortality	ECOTOX, Reference No. 7800
	28,000	LC50	*Cancer magister* (Dungeness crab)	96h	mortality	ECOTOX, Reference No. 5035
	33,000	LC50	*Artemia* sp. (brine shrimp)	24h	mortality	ECOTOX, Reference No. 2408
	38,900	NOAEL	*Physa heterostropha* (pond snail)	96h	mortality	ECOTOX, Reference No. 14396
	58,000	LC50	*Gammarus minus* (scud)	96h	mortality	ECOTOX, Reference No. 14396
	113,000	LC50	*Brachionus sp.* (rotifer)	24h	mortality	ECOTOX, Reference No. 9385
	172,000	LC50	*Crassostrea gigas* (Pacific oyster)	48h	mortality	ECOTOX, Reference No. 8621
	225,000	LC50	*Katelysia opima* (marine bivalve)	96h	mortality	ECOTOX, Reference No. 9017
	447,000	LC50	*Diaptomus forbesi* (calanoid copepod)	96h	mortality	ECOTOX, Reference No. 11282
	1,047,000	EC50	*Crassostrea gigas* (Pacific oyster)	48h	development	ECOTOX, Reference No. 8621
	1,100,000	LC50	*Melanoides tuberculata* (snail)	96h	mortality	ECOTOX, Reference No. 18198
Trichloroethene	1,700	LC50	*Platyhelminthes* sp. (flatworm)	7d	mortality	Yoshioka et al. (1986)
	2,200	NOAEL	*Daphnia magna* (water flea)	48h	mortality	LeBlanc (1980)
	2,300	LC50	*Daphnia* sp. (water flea)	3h	mortality	Yoshioka et al. (1986)
	2,300	NOAEL	*Daphnia magna* (water flea)	21d	reproduction	ECOTOX, Reference No. 56378
	7,095	NOAEL	*Ceriodaphnia dubia* (water flea)	7d	reproduction	Niederlehner et al. (1998)
	8,000	LOAEL	*Daphnia magna* (water flea)	21d	reproduction	ECOTOX, Reference No. 56378
	14,000	LC50	*Mysidopsis bahia* (mysid shrimp)	96h	mortality	Ward et al. (1986)
	16,951	LC50	*Ceriodaphnia dubia* (water flea)	7d	mortality	Niederlehner et al. (1998)
	18,000	LC50	*Daphnia magna* (water flea)	48h	mortality	LeBlanc (1980)
	20,000	LC50	*Elminius modestus* (Australian barnacle)	48h	mortality	ECOTOX, Reference No. 19535
	24,000	LC50	*Gammarus pulex* (scud)	48h	mortality	ECOTOX, Reference No. 15788
	30,000	LC50	*Asellus aquaticus* (aquatic sowbug)	48h	mortality	ECOTOX, Reference No. 15788
	39,000	LC50	*Daphnia pulex* (water flea)	48h	mortality	ECOTOX, Reference No. 2017
	42,000	LC50	*Cloeon dipterum* (mayfly)	48h	mortality	ECOTOX, Reference No. 15788
	48,000	LC50	*Aedes aegypti* (yellow fever mosquito)	48h	mortality	ECOTOX, Reference No. 14863
	49,000	LC50	*Ischnura elegans* (damselfly)	48h	mortality	ECOTOX, Reference No. 15788
	55,000	LC50	*Culex pipiens* (northern house mosquito)	48h	mortality	ECOTOX, Reference No. 10574

VOC	Reported Concentration (µg/L)	Effect Level	Test Organism	Test Duration	Endpoint	Source[a]
	56,000	LC50	*Lymnaea stagnalis* (great pond snail)	24 to 48h	mortality	ECOTOX, Reference No. 10574
	56,000	LC50	*Daphnia cucullata* (water flea)	48h	mortality	ECOTOX, Reference No. 2017
	64,000	LC50	*Chironomus* (midge)	48h	mortality	ECOTOX, Reference No. 15788
	70,000	LC50	*Nemoura cinerea* (stonefly)	48h	mortality	ECOTOX, Reference No. 15788
	75,000	LC50	*Hydra oligactix* (hydra)	48h	mortality	ECOTOX, Reference No. 10574
	75,000	LC50	*Erpobdella octoculata* (leech)	48h	mortality	ECOTOX, Reference No. 15788
	132,000	LC50	*Tubificidae* (oligochaete)	48h	mortality	ECOTOX, Reference No. 15788
Vinyl chloride (chloroethene)	**12,800**	FCV	Based on narcosis model	na	na	DiToro et al. (2000)
	65,300	FAV	Based on narcosis model	na	na	DiToro et al. (2000)
	128,000	NOAEL	*Brachydanio rerio* (zebrafish)	96h	mortality	Groeneveld et al. (1993), as cited in de Rooij et al. (2004b)
	388,000	LC100	*Esox lucius* (northern pike)	10d	mortality	Brown et al. (1977)

[a] References for ECOTOX sources can be found at: http://cfpub.epa.gov/ecotox/advanced_query.htm.

[b] Additional higher concentrations were reported in ECOTOX

d – days

EC50 – effects concentration for 50% of a test population

FAV – final acute value

FCV – final chronic value

h – hours

LC50 – lethal concentration for 50% of a test population

LC100 - lethal concentration for 100% of a test population

LOAEL – lowest-observed-effect concentration

na – not available

NOAEL – no-observed-effect concentration

VOC – volatile organic compound

Bold identifies concentrations that were selected as toxicity reference values.

SUMMARY OF SELECTED TOXICOLOGICAL STUDIES

1,1-Dichloroethane and vinyl chloride

There were no invertebrate and very few fish toxicity data available for 1,1-dichloroethane and vinyl chloride. Therefore, the toxicity values for these two chemicals were derived using the narcosis toxicity model for aquatic organisms presented in DiToro et al. (2000). The narcosis model is based on the calculation of chemical concentrations in the target lipid phase of the organism associated with narcotic mortality. This model was calibrated by DiToro et al. (2000) using a database of LC50s for 156 organic chemicals and 33 species including fish, amphibians, arthropods, mollusks, polychaetes, coelenterates, and protozoans. The predicted chronic values (PCVs) for 1,1-dichloroethane and vinyl chloride were obtained from the final acute values (FAVs) by using an acute-to-chronic ratio of 5.09, which was calculated by DiToro et al. (2000) as the geometric mean of the available acute-to-chronic ratios.

1,1-Dichloroethene

The NOAEL and LOAEL were selected from two different studies. The lowest LOAEL for 1,1-dichloroethene was reported in a study using *Daphnia magna* (Dill et al. 1980). In this study, *Daphnia magna* (< 24 hours old) were exposed to 1,1-dichloroethene for 48 hours in a static environment. The mortality rate was determined at test termination, and an LC50 of 11, 600 µg/L was calculated. The NOAEL was selected from a similar study with *Daphnia magna* (LeBlanc 1980). *Daphnia magna* (< 24 hours old) were exposed to 1,1-dichloroethene for 48 hours in a static test. The NOAEL was 2,400 µg/L.

1,2-Dichlorobenzene

The lowest LOAEL was selected from a study using *Daphnia magna* (Calamari et al. 1983). In this study, *Daphnia magna* were exposed to 1,2-dichlorobenzene for 14 days. The LOAEL was based on a reproductive EC50 of 550 µg/L. Because the selected LOAEL was lower than any NOAEL reported in the literature, a NOAEL of 11 µg/L was derived by dividing the LOAEL by 50.

1,2-Dichloroethane

The LOAEL for 1,2-dichloroethane was selected from a study using *Artemia salina* (Sanchez-Fortun et al. 1997). In this study, *Artemia* shrimp larvae were exposed to 1,2-dichloroethane for 72 hours. The mortality rate was determined at test termination, and an LC50 of 6,927 µg/L was calculated. Because the selected LOAEL was lower than any NOAEL reported in the literature, a NOAEL of 139 µg/L was derived by dividing the LOAEL by 50.

1,2-Dichloropropane
The lowest LOAEL was selected from a study using *Daphnia magna* (LeBlanc 1980). In this study, *Daphnia magna* were exposed to 1,2-dichlorobenzene for 48 hours. The mortality rate was determined at test termination and an LC50 of 6,927 µg/L was calculated. Because no NOAEL was available in the literature, a NOAEL of 840 µg/L was derived by dividing the LOAEL by 50.

1,4-Dichlorobenzene
The lowest LOAEL was selected from a study using *Portunus pelagicus* (Mortimer and Connell 1994). In this study, *Portunus pelagicus* were exposed to 1,4-dichlorobenzene for 10 hours. The mortality rate was determined at test termination and an LC50 of 147 µg/L was calculated. Because no NOAEL was available in the literature, a NOAEL of 3 µg/L was derived by dividing the LOAEL by 50.

1,2-Dichloroethene (cis or trans)
The lowest LOAEL was selected from a study using *Artemia salina* (Sanchez-Fortun et al. 1997). In this study, *Artemia* shrimp larvae were exposed to 1,2-dichloroethane for 72 hours. The mortality rate was determined at test termination and an LC50 of 6,785 µg/L was calculated. The only other toxicity data available for this chemical is an LC50 of 140,000 g/L from a bluegill study. Therefore, a NOAEL of 136 µg/L was derived by dividing the LOAEL by 50.

Benzene
The LOAEL for benzene was selected from a study by Caldwell et al. (1976). This study was conducted with larval stages of the Dungeness crab, *Cancer magister*. Larvae were exposed to benzene solutions within a few hours after larvae were hatched and mortality, growth, and development were monitored for up to 50 days. At a benzene concentration of 1,100 µg/L, all larvae died after 24 days of exposure. No effects on mortality, growth, or development were observed at the next lowest concentration of 180 µg/L. Therefore, these concentrations were selected as the LOAEL and NOAEL, respectively, for benzene.

Carbon disulfide
The lowest LOAEL was selected from a study using *Daphnia magna* (Van Leeuwen et al. 1985). In this study, *Daphnia magna* were exposed to carbon disulfide for 48 hours. The mortality rate was determined at test termination and an LC50 of 1,900 µg/L was calculated. Because no NOAEL was available in the literature, a NOAEL of 38 µg/L was derived by dividing the LOAEL by 50.

Chlorobenzene
The NOAEL and LOAEL were selected from two different studies. The lowest LOAEL was selected from a study using *Daphnia magna* (Calamari et al. 1983). In this study, *Daphnia magna* were exposed to chlorobenzene for 14 days in a static renewal environment. The LOAEL was based on a reproductive EC50 of 2,500 µg/L. The

NOAEL was selected from a study by Cowgill and Milazzo (1991). In this study, *Daphnia magna* were exposed to chlorobenzene for 10 days in a renewed static environment. No effect on mortality was observed at a chlorobenzene concentration of <1,400 µg/L. Therefore, 1,400 µg/L was selected as the NOAEL for chlorobenzene.

Tetrachloroethene

The NOAEL and LOAEL were selected from two different studies. The lowest LOAEL for tetrachloroethene was selected from a study using *Artemia salina* (Sanchez-Fortun et al. 1997). In this study, *Artemia* shrimp larvae were exposed to tetrachloroethene for 72 hours. The mortality rate was determined at test termination and an LC50 of 332 µg/L was calculated. The NOAEL was selected from a study using *Ceriodaphnia dubia* (Niederlehner et al. 1998). In this study, *Ceriodaphnia dubia* were exposed to tetrachloroethene for 7 days in a static environment. At test termination the LC50 was reported as 829 µg/L and the reproductive NOAEL as 331 µg/L. Because the LC50 reported for *Artemia* was lower than the LC50 reported for *Ceriodaphnia dubia*, the *Artemia* LC50 was selected as the LOAEL.

Toluene

The NOAEL and LOAEL were selected from two different studies. The lowest LOAEL available in the literature was not selected because the authors did not indicate whether a negative control was used (Tatem et al. 1978). Instead, the LOAEL was selected from a study by Korn et al. (1979). In this study, adult shrimp (*Eualus* spp.) were exposed to toluene for 96 hours in a static environment. The mortality rate was determined at test termination and an LC50 of 14,700 µg/L was calculated. Methods used to reduce evaporative losses were not discussed in the study and comparison of measured concentrations at both test initiation and termination showed that toluene declined substantially during the test, thus indicating that the LC50 may be an overestimate (i.e., the actual LC50 may be lower). However, because this LC50 was the lowest concentration at which an effect was reported, it was selected as the LOAEL. The NOAEL for toluene was selected from a study using *Ceriodaphnia dubia* (Niederlehner et al. 1998). In this study, *Ceriodaphnia dubia* were exposed to toluene for 7 days in a static environment. A reproductive NOAEL of 737 µg/L was reported at test termination.

Trichloroethene

The two lowest LOAEL for trichloroethene were from two studies by Yoshioka et al. (1986). Both studies were considered unacceptable either because the test did not represent a relevant environmental stressor (heads were removed from flatworms and the endpoints were abnormal head regeneration and mortality) or very few study details were presented and the use of a negative control was not discussed. The third lowest concentration where effects were observed was a reproductive LOAEL of 8,000 µg/L, but the derivation of this value could not be reviewed because the paper was in German. The fourth lowest concentration with effects was from an acceptable study by Ward et al. (1986). In this study, *Mysidopsis bahia* (mysid shrimp) were exposed to

trichloroethene for 96 hours in a static environment. The mortality rate was determined at test termination and an LC50 of 14,000 µg/L was calculated. The lowest NOAEL for trichloroethene was selected from a study by LeBlanc (1980). In this study, *Daphnia magna* (< 24 hours old) were exposed to trichloroethene for 48 hours in a static environment. The NOAEL was 2,200 µg/L.

REFERENCES

Brown ER, Sinclair T, Keith L, Beamer P, Hazdra JJ, Nair V, Callaghan O. 1977. Chemical pollutants in relation to diseases in fish. Ann NY Acad Sci 298:535-546.

Calamari D, Galassi S, Setti F, Vighi M. 1983. Toxicity of selected chlorobenzenes to aquatic organisms. Chemosphere 12(2):253-262.

Caldwell RS, Calderone EM, Mallon MH. 1976. Effects of a seawater-soluble fraction of Cook Inlet crude oil and its major aromatic components on larval stages of the Dungeness crab, *Cancer magister* Dana. In: Wolfe DA, ed, Fate and effects of petroleum hydrocarbons in marine ecosystems and organisms: proceedings of a symposium November 10-12, 1976, Olympic Hotel, Seattle, Washington. Pergamon Press, New York, NY, pp 210-220.

Cowgill UM, Milazzo DP. 1991. The sensitivity of *Ceridaphnia dubia* and *Daphnia magna* to seven chemicals utilizing the three-brood test. Arch Environ Contam Toxicol 20:211-217.

De Rooij C, Defourny C, Thompson RS, Garny V, Lecloux A, Van Wijk D. 2004a. 1,1,1-trichloroethane marine risk assessment with special reference to the OSPARCOM region: North Sea. Environ Monit Assess 97(1-3):39-56.

De Rooij C, Defourny C, Thompson RS, Garny V, Lecloux A, Van Wijk D. 2004b. Vinyl chloride marine risk assessment with special reference to the OSPARCOM region: North Sea. Environ Monit Assess 97(1-3):57-67.

Dill DC, McCarty WM, Alexander HC, Bartlett EA. 1980. Toxicity of 1,1-dichloroethylene (vinylidene chloride) to aquatic organisms. EPA-600/3-80-057. US Environmental Protection Agency, Duluth, MN.

DiToro DM, McGrath JA, Hansen DE. 2000. Technical basis for narcotic chemicals and polycyclic aromatic hydrocarbon criteria. I. Water and tissue. Environ Toxicol Chem 19(8):1951-1970.

Groeneveld AHC, Zijlstra AG, Feenstra AFM, Berends AG. 1993. The acute toxicity of vinyl chloride to the zebra fish. As cited in De Rooij et al. 1994. Solvay-Duphar Report 56635, 64/92.

Hermens J, Canton H, Janssen P, De Jong R. 1984. Quantitative structure-activity relationships and toxicity studies of mixtures of chemicals with anaesthetic potency: acute lethal and sublethal toxicity to *Daphnia magna*. Aquat Toxicol 5:143-154.

Könemann H. 1981. Quantitative structure-activity relationships in fish toxicity studies. Part 1: relationship for 50 industrial pollutants. Toxicology 19:209-221.

Korn S, Moles DA, Rice SD. 1979. Effects of temperature on the median tolerance limit of pink salmon and shrimp exposed to toluene, naphthalene, and Cook Inlet crude oil. Bull Environ Contam Toxicol 21:521-525.

LeBlanc GA. 1980. Acute toxicity of priority pollutants to water flea (*Daphnia magna*). Bull Environ Contam Toxicol 24:684-691.

Mortimer MR, Connell DW. 1994. Critical internal and aqueous lethal concentrations of chlorobenzenes with the crab *Portunus pelagicus* (L). Ecotoxicol Environ Saf 28:298-312.

Niederlehner BR, Cairns JJ, Smith EP. 1998. Modeling acute and chronic toxicity of nonpolar narcotic chemicals and mixtures to *Ceriodaphnia dubia*. Ecotoxicol Environ Saf 39:136-146.

Richter JE, Peterson SF, Kleiner DF. 1983. Acute and chronic toxicity of some chlorinated benzenes, chlorinated ethanes, and tetrachloroethylene to *Daphnia magna*. Arch Environ Contam Toxicol 12:679-684.

Sanchez-Fortun S, Sanz, Santa-Maria A, Ros JM, De Vicente ML, Encinas MT, Vinagre E, Barahona MV. 1997. Acute sensitivity of three age classes of *Artemia salina* larvae to seven chlorinated solvents. Bull Environ Contam Toxicol 59:445-451.

Stephan CE, Mount DI, Hansen DJ, Gentile JH, Chapman GA, Brungs WA. 1985. Guidelines for deriving numerical national water quality criteria for the protection of aquatic organisms and their uses. PB85-227049. Office of Research and Development, US Environmental Protection Agency, Washington, DC.

Suter GW, Tsao CL. 1996. Toxicological benchmarks for screening potential contaminants of concern for effects on aquatic biota: 1996 revision. Prepared for U.S. Department of Energy Office of Environmental Management. Risk Assessment Program, Health Sciences Research Division.

Tatem HE, Johnson JE. 1978. Aquatic disposal field investigations, Duwamish Waterway disposal site, Puget Sound, Washington: evaluative summary. Dredged material research program technical report no. D-77-24. Waterway Experiment Station, US Army Corps of Engineers, Vicksburg, MS.

Tatem HE, Cox BA, Anderson JW. 1978. The toxicity of oils and petroleum hydrocarbons to estuarine crustaceans. Estuar Coast Mar Sci 6:365-373.

Van Leeuwen CJ, Maas-Diepeveen JL, Niebeek G, Vergouw WHA, Griffioen PS, Luijken MW. 1985. Aquatic toxological aspects of dithiocarbamates and related compounds. I. Short-term toxicity tests. Aquat Toxicol 7:145-164.

Van Wijk D, Thompson RS, De Rooij C, Garny V, Lecloux A, Kanne R. 2004. Monochlorobenzene marine risk assessment with special reference to the OSPARCOM region: North Sea. Environ Monit Assess 97(1-3):87-102.

Ward GS, Tolmsoff AJ, Petrocelli SR. 1986. Acute toxicity of trichloroethylene to saltwater organisms. Bull Environ Contam Toxicol 37:830-836.

Windward. 2004. Lower Duwamish Waterway remedial investigation. Task 8: Phase 2 RI work plan. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2006. Lower Duwamish Waterway remedial investigation. Data and analysis report: porewater sampling of Lower Duwamish Waterway. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Yoshioka Y, Ose Y, Sato T. 1986. Correlation of the five test methods to assess chemical toxicity and relation to physical properties. Ecotoxicol Environ Saf 12:15-21.

ATTACHMENT 8
ACCEPTABLE TOXICITY STUDIES FOR THE SELECTION OF FISH TOXICITY REFERENCE VALUES

Attachment 8 Acceptable Toxicity Studies for the Selection of Fish Toxicity Reference Values

Table 1. Acceptable toxicity studies for selection of fish dietary toxicity reference values

Chemical	Chemical Form	Test Species	Life Stage	NOAEL (mg/kg dw[a])	LOAEL (mg/kg dw[a])	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
Arsenic	sodium arsenite	rainbow trout	juvenile	20[b]	30[b]	food	6 wks	growth	body weight	Oladimeji et al. (1984)	concentrations in figure and text do not agree in study: 20 mg/kg is mentioned both as an effect level and a NOAEL in the text; however, it is shown in the figure to be not significant; 20 mg/kg = assumed NOAEL; food consumption not measured; fish fed prepared diet treated with chemical salts; only the nominal concentration was reported
Arsenic	disodium arsenate heptahydrate	rainbow trout	juvenile	8	44	food	16 wks	growth	body weight	Cockell et al. (1991)	feed refusal accompanied effects; fish fed prepared diet treated with chemical salts
Arsenic	disodium arsenate heptahydrate	rainbow trout	juvenile		49	food	24 wks	growth	body weight	Cockell et al. (1991)	feed refusal accompanied effects; effects attributed to feed refusal; fish fed prepared diet treated with chemical salts
Arsenic	disodium arsenate heptahydrate	rainbow trout	juvenile		55	food	8 days	growth	body weight	Cockell et al. (1992)	feed refusal accompanied effects; fish fed prepared diet treated with chemical salts
Arsenic	disodium arsenate	rainbow trout	juvenile		58	food	12 days	growth	body weight	Cockell and Bettger (1993)	feed refusal accompanied effects; fish fed prepared diet treated with chemical salts
Arsenic	disodium arsenate heptahydrate	rainbow trout	juvenile	32	60	food	12 days	growth	body weight	Cockell et al. (1992)	feed refusal accompanied effects; fish fed prepared diet treated with chemical salts
Arsenic	disodium arsenate heptahydrate	rainbow trout	juvenile	33	65	food	24 wks	growth	body weight	Cockell et al. (1991)	feed consumption did not differ from controls; body weight gain reduced at 12 weeks in fish fed 33 mg/kg arsenic but not at 24 weeks (body weight was recovered); fish fed prepared diet treated with chemical salts
Arsenic	disodium arsenate	rainbow trout	juvenile		137	food	8 days	growth	body weight	Cockell and Hilton (1988)	greater than 10% mortality at LOAEL; feed refusal accompanied effects; study also reported effects of organic arsenic which was not toxic at high levels; fish fed prepared diet treated with chemical salts
Arsenic	arsenic trioxide	rainbow trout	juvenile		180	food	8 days	growth	body weight	Cockell and Hilton (1988)	feed refusal accompanied effects; greater than 10% mortality at LOAEL; study also reported effects of organic arsenic which was not toxic at high levels; fish fed prepared diet treated with chemical salts
Arsenic	disodium arsenate heptahydrate	striped bass	juvenile	52.3	188.8	food	6 days	growth	body weight	Blazer et al. (1997)	feed refusal accompanied effects; fish fed prepared diet treated with chemical salts
Cadmium	$Cd(NO_3)_2$	rockfish	juvenile	0.1[c]	0.5	food	60 days	growth	body weight and length growth rate; condition factor	Kim et al. (2004); Kang et al. (2005)	body weight and body length growth rate were significantly affected in both studies at 25 and 125 mg/kg dw; significance of effect on body weight and body length growth rate at 0.5 and 5 mg/kg dw was not consistent between the two studies; condition factor was reduced at all diet concentrations (0.5, 5, 25, and 125 mg/kg), but was not significant at 25 mg/kg; a significant inverse relationship was observed between weight gain and the exposure concentration of dietary Cd at 25 and 125 mg/kg dw; fish fed prepared diet treated with chemical salts
Cadmium	cadmium chloride	rainbow trout	fry	55[b]		primarily food	60 days	survival, growth	100% survival, body weight and length	Mount et al. (1994)	fish exposed to copper, cadmium, lead, and zinc in water at 23.0, 0.97, 3.32, and 46.3 µg/L in addition to dietary exposure; only no-effect reported; fish fed *Artemia* exposed to cadmium chloride in water; dietary concentrations corrected for a theoretical 20% loss resulting from depuration of copper from *Artemia.*
Cadmium	$Cd(NO_3)_2$	rockfish	juvenile	125		food	60 days	survival		Kim et al. (2004)	fish fed prepared diet treated with chemical salts
Cadmium		guppy		171[b]		food	10-30 days	growth		Hatakeyama and Yasuno (1982)	fed exposed *Monia macrocopa,* significant growth effect at day 10 disappeared at day 20 in fish fed 126 and 171 µg/g dw; fish fed prepared diet treated with chemical salts
Cadmium	cadmium chloride	guppy	adult	210[b]		food	2 mo	reproduction	# live fry, fry mortality, premature embryos	Hatakeyama and Yasuno (1987)	fed exposed *Chironomus yoshimatsui* (midge larvae), no reproductive effect after 2 mo

Table 1, continued

Chemical	Chemical Form	Test Species	Life Stage	NOAEL (mg/kg dw[a])	LOAEL (mg/kg dw[a])	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
Cadmium		Atlantic salmon	juvenile	250		food	4 months	growth	growth rate (body weight)	Lundebye et al. (1999)	fish fed prepared diet treated with chemical salts
Cadmium	cadmium chloride	guppy	2 mo	274[b]		food	30 days	growth	body weight	Hatakeyama and Yasuno (1987)	fed exposed *Chironomus yoshimatsui* (midge larvae), no growth effect by day 30
Cadmium		rainbow trout	juvenile	294		food	15-30 days	growth, survival	specific growth rate, survival	Baldisserotto et al. (2005)	fish fed prepared diet treated with chemical salts
Cadmium		rainbow trout	juvenile	471		food	28 days	survival, growth	growth rate, survival	Franklin et al. (2005)	fish fed prepared diet treated with chemical salts
Cadmium	cadmium chloride	guppy	30 day old	500[b]	800[b]	food	7 mo	reproduction	cumulative number of fry produced	Hatakeyama and Yasuno (1987)	fed exposed *Chironomus yoshimatsui* (midge larvae); cumulative number of fry decreased to about 60% of the control in fish exposed to 80 and 160 µg/L; no discussion of statistics; LOAEL and NOAEL are estimated using figure, no replication
Cadmium	cadmium chloride	guppy	30 day old		1,250[b]	food	7 mo	growth, survival	female growth, female survival	Hatakeyama and Yasuno (1987)	fed exposed *Chironomus yoshimatsui* (midge larvae); female body weight decreased - 68% of control at the 48th day in fish exposed to 160 µg/L; 6 of 7 females died that were exposed to 160 µg/L; no discussion of statistics- unclear on growth effects at the other doses; LOAEL is estimated using figure; no effect on male growth, no replication
Cadmium	cadmium nitrate	rainbow trout	juvenile	786	1,395	food	30 days	survival		Szebedinsky et al. (2001)	57% survival observed in fish fed 1,395 mg/kg; in a separate experiment reported in this study, 92% survival was reported for juvenile rainbow trout exposed to dietary cadmium concentrations of 1,419 mg/kg dw over a 36-day exposure period; survival data were not statistically analyzed in either experiment
Cadmium	cadmium nitrate	rainbow trout	juvenile	1,395	2,265	food	30 days	growth	specific growth rate (weight)	Szebedinsky et al. (2001)	
Cadmium	cadmium sulfate	rainbow trout			10,000	food	28 days	survival		Handy (1993)	39% (14 of 36) mortality between day 3 and 23 of exposure period; fish fed prepared diet treated with chemical salts
Chromium	chromium (III)	grey mullet	2 yrs old	9.42		food and sediment	8 wks	growth		Walsh et al. (1994)	significant increase in growth of the chromium-exposed fish (not an adverse effect); gray mullet consume large amounts of sediment during feeding and fish were exposed to chromium in both diet and sediments (46mg/kg) simultaneously
Copper	copper	channel catfish	fingerling	8	16	food	16 wks	growth	body weight	Murai et al. (1981)	significant effects at 4 wks in 16- and 32-mg/kg treatments; growth gain per feed consumed was significantly lower for fish fed 8 mg/kg; fish fed prepared diet treated with chemical salts
Copper	copper sulfate pentahydrate	channel catfish	fingerling	40		food	13 wks	growth	body weight	Gatlin and Wilson (1986)	fish fed prepared diet treated with chemical salts
Copper	copper sulfate	rockfish	juvenile	50	100	food	60 days	growth	body weight (growth rate)	Kang et al. (2005)	fish fed prepared diet treated with chemical salts
Copper	copper sulfate pentahydrate	Atlantic salmon	parr	98		food	12 wks	growth, survival	survival, condition factor, body weight,	Lorentzen et al. (1998)	
Copper	copper sulfate	rainbow trout		200		food	32 days	survival		Handy (1992)	only no-effect level concentration reported; fish fed prepared diet treated with chemical salts.
Copper	copper sulfate pentahydrate	rainbow trout		664[b]		food	24 wks	growth		Lanno et al. (1985b)	significant effects at 16 wks; at 24 wks, growth effects no longer apparent; only no-effect level concentration reported; fish fed prepared diet treated with chemical salts
Copper	copper sulfate	rainbow trout		684		food	42 days	growth		Miller et al. (1993)	only no-effect level concentration reported; fish fed prepared diet treated with chemical salts
Copper	copper sulfate pentahydrate	Atlantic salmon	parr	691.3[b]		food	4 wks	growth	length, weight, or condition factor	Berntssen et al. (1999b)	only no-effect level concentration reported; fish fed prepared diet treated with chemical salts
Copper	copper sulfate pentahydrate	Atlantic salmon	fry	500	700	food	3 mos	growth		Lundebye et al. (1999)	fish fed prepared diet treated with chemical salts

Table 1, continued

Chemical	Chemical Form	Test Species	Life Stage	NOAEL (mg/kg dw[a])	LOAEL (mg/kg dw[a])	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
Copper	copper sulfate pentahydrate	rainbow trout		287[b]	730[b]	food	8 wks	growth		Lanno et al. (1985b)	feed refusal and 19% mortality was observed at 1,585 mg/kg-diet; fish fed prepared diet treated with chemical salts
Copper	copper sulfate pentahydrate	rainbow trout		730[b]		food	8 wks	survival		Lanno et al. (1985b)	study showed potential effects at higher concentrations- confounded by food avoidance; feed refusal and 19% mortality were observed at 1,585-mg/kg diet; only no-effect level concentration reported; fish fed prepared diet treated with chemical salts
Copper	copper sulphate pentahydrate	rainbow trout			796[b]	food	16 wks	growth		Lanno et al. (1985a)	fish were fed on control diet for 12 d after exposure period; fish fed prepared diet treated with chemical salts
Copper	copper chloride	rainbow trout	fry	352[b]		primarily food	60 days	survival	survival	Mount et al. (1994)	fish exposed to copper, cadmium, lead, and zinc in water at 23.0, 0.97, 3.32, and 46.3 µg/L in addition to dietary exposure; mortality observed in fish fed *Artemia* with Cu concentrations of 830 mg/kg was attributed to elevated copper concentrations in the water; fish fed live Artemia exposed to copper chloride in water; dietary concentrations corrected for a theoretical 20% loss resulting from depuration of copper from *Artemia*
Copper	copper sulfate	Atlantic salmon		638[b]	868[b]	food	3 mo	growth		Berntssen et al. (1999a)	fish fed prepared diet treated with chemical salts
Copper	copper chloride	rainbow trout	fry	800[b]		primarily food	60 days	growth	body weight and length	Mount et al. (1994)	fish exposed to copper, cadmium, lead, and zinc in water at 23.0, 0.97, 3.32, and 46.3 µg/L in addition to dietary exposure; only no-effect level concentration reported; fish fed live *Artemia* exposed to copper chloride in water; dietary concentrations corrected for a theoretical 20% loss resulting from depuration of copper from *Artemia*
Copper	copper sulfate pentahydrate	rainbow trout		1,042		food	28 days	survival/ growth		Kamunde et al. (2001)	non-significant growth inhibition at NOAEL; only no-effect level concentration reported; fish fed prepared diet
Copper	copper sulfate pentahydrate	grey mullet			2,397[b]	food	67 days	growth		Baker et al. (1998)	effects associated with reduced feeding
Copper	copper sulfate	rainbow trout		10,000		food	28 days	survival		Handy (1993)	only no-effect level concentration reported; fish fed prepared diet treated with chemical salts
Lead	lead nitrate	rainbow trout	fry	210[b]		primarily food	60 days	survival, growth	100% survival, body weight and length	Mount et al. (1994)	fish exposed to copper, cadmium, lead, and zinc in water at 23.0, 0.97, 3.32, and 46.3 µg/L in addition to dietary exposure; fish fed live *Artemia* exposed to lead nitrate in water
Lead		rainbow trout		7,040		food		growth		Goettl et al. (1976)	fish fed prepared diet treated with chemical salts
Silver		rainbow trout		3000		food		growth		Galvez and Wood (1999)	fish fed prepared diet treated with chemical salts
Vanadium	sodium orthovanadate	rainbow trout	juvenile	2.04[d]	10.2	food	12 wks	growth	body weight	Hilton and Bettger (1988)	reduced feeding at all doses but significant increase in food consumption/weight gain
Zinc	zinc chloride	rainbow trout	fry	1,900[b]		primarily food	60 days	growth, survival	body weight and length; 97% survival	Mount et al. (1994)	fish exposed to copper, cadmium, lead, and zinc in water at 23.0, 0.97, 3.32, and 46.3 µg/L in addition to dietary exposure. fish fed live *Artemia* exposed to zinc chloride in water
Zinc		rainbow trout	fingerling	1,000	2,000	food	6 wks	growth		Takeda and Shimma (1977)	fish fed at same dose zinc with 0.5% Ca experienced no adverse effects.
Benzo(a)pyrene		areolated grouper	juvenile	81		food	4 wks + 4 wks recovery	survival, growth	survival, body weight & length	Wu et al. (2003)	exposure was dietary force feeding - fish were force fed pellet into larynx
Benzo(a)pyrene		English sole	juvenile	47	116	food	10-12 days	growth		Rice et al. (2000)	diet of exposed polychaetes exposed for 28 d; exposure concentrations calculated from feeding rates
Benzo(a)pyrene		rainbow trout	juvenile	100	1,000	food	28 days	growth		Hart and Heddle (1991)	fish fed prepared diet treated with benzo(a)pyrene
Benzo(a)pyrene		rainbow trout	3 mo. old		1,000	food	18 mo	growth	reduced body weight	Hendricks et al. (1985)	fish fed prepared diet treated with benzo(a)pyrene; total mortality was higher in the control group
Benzo(a)pyrene		rainbow trout	juvenile (3 mos old)	1,000		food	18 mo	survival		Hendricks et al. (1985)	mortality was greater in control; NOAEL supported by additional 9-week study; fish fed prepared diet treated with benzo(a)pyrene

Table 1, continued

Chemical	Chemical Form	Test Species	Life Stage	NOAEL (mg/kg dw[a])	LOAEL (mg/kg dw[a])	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
PAHs	PAH mixture[a]	chinook salmon	juvenile	280[b]		food	7 wks	growth		Palm et al. (2003)	14 PAHs included in diet: acenaphthene, benzo(a)anthracene, benzo(b)fluoranthene, benzo(g,h,i)perylene, benzo(k)fluoranthene, chrysene, dibenzo(a,h)anthracene, fluoranthene, fluorene, indeno(1,2,3-cd)pyrene, naphthalene, phenanthrene, and pyrene
PAHs	PAH mixture[a]	chinook salmon	juvenile	280[b]		food	4 wks exposure, 2 wks immuno-challenge	disease resistance		Palm et al. (2003)	14 PAHs included in diet: acenaphthene, benzo(a)anthracene, benzo(b)fluoranthene, benzo(g,h,i)perylene, benzo(k)fluoranthene, chrysene, dibenzo(a,h)anthracene, fluoranthene, fluorene, indeno(1,2,3-cd)pyrene, naphthalene, phenanthrene, and pyrene; fish were exposed to *Listonella anguillarum* following PAH exposure. No difference was observed between PAH-exposed fish and controls in either fish that were vaccinated against the bacterium or those that weren't vaccinated
PAHs	PAH mixture	chinook salmon	juvenile	324	951	food	53 days	growth	reduced dry-weight body weight	Meador et al. (2006)	21 PAHs included in diet: naphthalene, 2-methylnaphthalene, dimethylnaphthalene, dibenzothiophene, acenaphthene, fluorene, 1,8-dimethyl(9H)fluorene, phenanthrene, 9-ethylphenanthrene, 9-ethyl-10-methylphenanthrene, 1-methyl-7-isopropylphenanthrene, anthracene, fluoranthene, pyrene, methyl pyrene, benzo(a)anthracene, chrysene, benz(a)pyrene, benzo(k)fluoranthene, benzo(g,h,i)perylene, and dibenzanthracene

[a] Concentrations of elemental metal or specific PAH chemical. Except where noted, concentrations were as administered, and were assumed to approximate a dry weight concentration.

[b] Converted to dry weight basis based on moisture content reported in paper.

[c] NOAEL estimated using uncertainty factor of 5 (chronic LOAEL to chronic NOAEL).

[d] NOAEL estimated using uncertainty factor of 10 (acute LOAEL to chronic NOAEL).

LOAEL – lowest-observed-adverse-effect level

NOAEL – no-observed-adverse-effect level

PAH – polycyclic aromatic hydrocarbon

Table 2. Acceptable toxicity studies for selection of fish whole-body tissue residue TRVs

Chemical	NOAEL (WB)	LOAEL (WB)	Conv. Factor	NOAEL (egg)	LOAEL (egg)	Unit (ww)	Test Species	Lifestage	Exposure Mode	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
2,3,7,8-TCDD	2.4[a]	12	28			pg/g	rainbow trout	adult	food	approx 300 days	survival	reduced survival	Giesy et al. (2002)	LOAEL of 0.44 in fillet tissue measured at day 200 converted to concentration at approximately day 300 when mortality was observed based on recommendation from the author (Giesy 2006)
2,3,7,8-TCDD	46	85				pg/g	lake whitefish	juvenile	food	30 days	growth	body weight	Fisk et al. (1997)	exposed for 30 days and effect measured for 180 days post-exposure; unclear when residues measured; assumed WB residues, however liver tissue possibly excluded; 29% mortality in control
2,3,7,8-TCDD	87	102	2.56	34	40	pg/g	lake trout	egg (converted to adult WB)	water		reproduction (egg exposure)	sac-fry survival	Walker et al. (1994)	study measured through routes of exposure for the egg (maternal transfer, water, injection)–egg residues not statistically different based on exposure route; significant mortality observed at LOAEL; adult concentration was estimated using egg:adult conversion factor of 2.56 based on Tietge et. al. (1998)
2,3,7,8-TCDD	38	102	2.56	15	40	pg/g	lake trout	egg (converted to adult WB)	water	48 hrs	reproduction (egg exposure)	early life-stage survival	Spitsbergen et al. (1991)	22.5% cumulative mortality (egg, hatching, sac-fry) at LOAEL statistically significant; mortality at NOAEL (15.8%) not statistically different from control (18%); egg:adult conversion factor of 2.56 based on Tietge et. al. (1998)
2,3,7,8-TCDD	90	108	2.56	35	42	pg/g	lake trout	egg (converted to adult WB)	water	48 hrs	reproduction (egg exposure)	early life-stage survival	Guiney et al. (1996)	NOAEL and LOAEL (LC50) based on hatchery fish eggs; LOAEL is LC50; NOAEL and LC50 also reported in field-collected eggs (from Lake Ontario) and reference eggs (from Lake Ontario); Lake Ontario NOAEL=30-45 pg/g and LC50=44-72 pg/g; Lake Superior NOAEL=34 pg/g; egg:adult conversion factor of 2.56 based on Tietge et. al. (1998)
2,3,7,8-TCDD	59	128	2.56	23	50	pg/g	lake trout	egg (converted to adult WB)	maternal transfer	11 wks prior to spawning	reproduction (egg exposure)	sac-fry survival	Walker et al. (1994)	parental fish exposed via diet; study measured through routes of exposure for the egg (maternal transfer, water, injection)–egg residues not statistically different based on exposure route; significant mortality observed at LOAEL; mortality in control=7.2%; egg:adult conversion factor of 2.56 based on Tietge et. al. (1998)
2,3,7,8-TCDD	87	141	2.56	34	55	pg/g	lake trout	egg (converted to adult WB)	water	48 hrs	reproduction (egg exposure)	early life-stage and sac-fry survival	Walker et al. (1991); Walker et al. (1992)	statistically significant effects observed at LOAEL, frequency of sac-fry edema also statistically significant; LC50=65 pg/g; egg:adult conversion factor of 2.56 based on Tietge et al. (1988)
2,3,7,8-TCDD	72	150				pg/g	rainbow trout	juvenile	food	30 days	growth	body weight	Fisk et al. (1997)	exposed for 30 days and effect measured for 180 days post-exposure; unclear when residues measured; assumed WB residues, however liver tissue possibly excluded
2,3,7,8-TCDD	215	225	2.56	84	88	pg/g	brook trout	egg (converted to adult WB)	maternal transfer	throughout egg devl'p	reproduction (egg exposure)	swim-up and juvenile survival	Johnson et al. (1998)	parental fish (1.5 yr old) fed TCDD until spawning; LOAEL is calculated LC10; LC10 is less than residue at which mortality was observed (at 156 pg/g ww); NOAEL is egg residue where no significant mortality was observed; egg residues are approximate; egg:adult conversion factor of 2.56 based on Tietge et al. (1988)
2,3,7,8-TCDD	346	474	2.56	135	185	pg/g	brook trout	egg (converted to adult WB)	water	48 hrs	reproduction (egg exposure)	sac-fry survival	Walker and Peterson (1994)	NOAEL and LOAEL as reported in study; at LOAEL 25% mortality; at NOAEL 10% mortality; egg:adult conversion factor of 2.56 based on Tietge et al. (1988);
2,3,7,8-TCDD	310	579	2.56	121	226	pg/g	lake trout	egg (converted to adult WB)	water	48 hrs	reproduction (egg exposure)	sac-fry growth (length), hatchability	Walker et al. (1991)	statistically significant effects observed at LOAEL; egg:adult conversion factor of 2.56 based on Tietge et al. (1988)

Table 2, continued

Chemical	NOAEL (WB)	LOAEL (WB)	Conv. Factor	NOAEL (egg)	LOAEL (egg)	Unit (ww)	Test Species	Lifestage	Exposure Mode	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
2,3,7,8-TCDD	448	691	2.56	175	270	pg/g	lake herring	egg (converted to adult WB)	water	100 days	reproduction (egg exposure)	early life-stage growth and survival	Elonen et al. (1998); Spehar et al. (1997)	statistically significant effects observed at LOAEL; six other fish species tested- lake herring was the most sensitive fish exposed to 2,3,7,8-TCDD; LC50=902 pg/g; LC10=509 pg/g; egg:adult conversion factor of 2.56 based on Tietge et al. (1988)
2,3,7,8-TCDD		714	2.56		279	pg/g	rainbow trout	egg (converted to adult WB)	water	48 hrs	reproduction (egg exposure)	sac-fry survival	Walker et al. (1992)	LC50s also reported in study; sac-fry mortality most sensitive endpoint; egg:adult conversion factor of 2.56 based on Tietge et al. (1988)
2,3,7,8-TCDD		780				pg/g	rainbow trout	10-15cm	water	2 and 6 hrs	growth	body weight	Branson et al. (1985)	no statistics; LOAEL presented is based on tissue residues measured 28d after end of exposure, LOAEL is misleading; higher LOAEL should be used (max WB for dose, WB at end of exp, etc); tissue burdens assoc with same dose but measured earlier in exp were higher
2,3,7,8-TCDD		980				pg/g	rainbow trout	fry	water	~28 days	survival		Mehrle et al. (1988)	LOAEL is based on tissue residue measured at the end of the exposure period
2,3,7,8-TCDD		990				pg/g	rainbow trout	fry	water	~28 days	growth		Mehrle et al. (1988)	
2,3,7,8-TCDD	602	1,114	2.56	235	435	pg/g	fathead minnow	egg (converted to adult WB)	water	100 days	reproduction (egg exposure)	early life-stage growth and survival	Elonen et al. (1998); Spehar et al. (1997)	statistically significant effects observed at LOAEL; LC50=539 pg/g; LC10=293 pg/g; egg:adult conversion factor of 2.56 based on Tietge et al. (1988)
2,3,7,8-TCDD	1,486					pg/g	brook trout	adult	food	182 days	survival, growth, gonadal development, egg production		Tietge et al. (1998)	NOAEL is initial concentration in female fish exposed to 1,200 pg TCDD/g-food; Giesy et al. (2002) report adverse effects in rainbow trout at similar exposure levels after 250 days of exposure
2,3,7,8-TCDD		1,626	2.56		635	pg/g	killifish (mummi-chog)	egg (converted to adult WB)	water	through hatch or 30 days post-hatch	reproduction (egg exposure)	hatchability	Prince and Cooper (1995)	hatchability (61%); study compared the effects of TCDD-treated water on field-collected "non-impacted" eggs (from clean site - Tuckerton, NJ) and "impacted" eggs (from contaminated site - Newark Bay, NJ); no dose response from "impacted" eggs; egg:adult conversion factor of 2.56 based on Tietge et al. (1988)
2,3,7,8-TCDD	125	2,170				pg/g	coho salmon	juvenile	water	96 hrs	growth		Miller et al. (1979)	body burdens measured after 114 days in clean water following exposure, thus not conservative
2,3,7,8-TCDD	125	2,170				pg/g	coho salmon	juvenile	water	96 hrs	survival		Miller et al. (1979)	body burdens measured after 114 days in clean water following exposure
2,3,7,8-TCDD	986	2,189	2.56	385	855	pg/g	channel catfish	egg (converted to adult WB)	water	100 days	reproduction (egg exposure)	early life-stage growth and survival	Elonen et al. (1998); Spehar et al. (1997)	statistically significant effects observed at LOAEL; LC50=644 pg/g; LC10=429 pg/g
2,3,7,8-TCDD		2,410				pg/g	medaka	juvenile	water	12 days	growth	body weight	Schmieder et al. (1995)	no statistics; growth increased 74% vs. 94% in control
2,3,7,8-TCDD	1,165	2,429	2.56	455	949	pg/g	medaka	egg (converted to adult WB)	water	100 days	reproduction (egg exposure)	early life-stage growth and survival	Elonen et al. (1998); Spehar et al. (1997)	statistically significant effects observed at LOAEL; LC50=1,110 pg/g; LC10=656 pg/g; egg:adult conversion factor of 2.56 based on Tietge et al. (1988)
2,3,7,8-TCDD	2,171	3,123	2.56	848	1,220	pg/g	white sucker	egg (converted to adult WB)	water	100 days	reproduction (egg exposure)	early life-stage growth and survival	Elonen et al. (1998); Spehar et al. (1997)	statistically significant effects observed at LOAEL; LC50=1,890 pg/g; LC10=1,590 pg/g; egg:adult conversion factor of 2.56 based on Tietge et al. (1988)
2,3,7,8-TCDD	3,046	4,608	2.56	1,190	1,800	pg/g	northern pike	egg (converted to adult WB)	water	100 days	reproduction (egg exposure)	early life-stage growth and survival	Elonen et al. (1998); Spehar et al. (1997)	statistically significant effects observed at LOAEL; LC50=2,460 pg/g; LC10=1,530 pg/g; egg:adult conversion factor of 2.56 based on Tietge et al. (1988)
2,3,7,8-TCDD	1,085	5,120	2.56	424	2,000	pg/g	zebra fish	egg (converted to adult WB)	water	100 days	reproduction (egg exposure)	early life-stage growth and survival	Elonen et al. (1998); Spehar et al. (1997)	statistically significant effects observed at LOAEL; LC50=2,610 pg/g; LC10=1,610 pg/g; egg:adult conversion factor of 2.56 based on Tietge et al. (1988)

Table 2, continued

Chemical	NOAEL (WB)	LOAEL (WB)	Conv. Factor	NOAEL (egg)	LOAEL (egg)	Unit (ww)	Test Species	Lifestage	Exposure Mode	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
2,3,7,8-TCDD		69,000				pg/g	fathead minnow	juvenile	water	28 day	survival		Adams et al. (1986)	no statistics—100% mort; LOAEL is average concentration in dead fish
2,3,7,8-TCDD	1,570	1,380,000				pg/g	rainbow trout	young	food	105 days	survival		Hawkes and Norris (1977)	
2,3,7,8-TCDD	1,570	1,380,000				pg/g	rainbow trout	young	food	105 days	growth	body weight	Hawkes and Norris (1977)	
PCBs (Aroclor 1260)		520				µg/kg	common barbel	adult	food	50 days	reproduction	fecundity	Hugla and Thome (1999)	Fecundity effect was not dose-responsive; number of fish tested unclear; uncertain statistical significance of fecundity endpoint; ww concentration converted from dw assuming 20% solids
PCBs (Aroclor mixture)					857	µg/kg	Atlantic salmon	embryo	water	48 hrs	reproduction (egg exposure)	live fry body weight	Fisher et al. (1994)	no effect on reproduction were observed
PCBs (Aroclor 1254)	980					µg/kg	chinook salmon	juvenile	food	4 wks	growth, survival		Powell et al. (2003)	whole body burdens ranged from 740 to 980 µg/kg ww over the 13-day period following treatment; only no-effect level reported; no effect on growth, survival, or survival following immunological challenge
PCBs (Aroclor 1254; egg)[a]					1,640	µg/kg	rainbow trout	egg	maternal transfer	60 days	reproduction (egg exposure)	fry growth	Hendricks et al. (1981)	eggs were exposed via maternal transfer from single gravid female fed 200 µg/g PCBs for 60 days
PCBs (Aroclor 1260)		2,640				µg/kg	common barbel	adult	food	75 days	reproduction	lack of spawning in first reproductive season; egg and larval mortality; fecundity	Hugla and Thome (1999)	fecundity effect was not dose-responsive; number of fish tested unclear; uncertain statistical significance of fecundity endpoint; ww concentration converted from dw assuming 20% solids
PCBs (Aroclor 1254)					3,200	µg/kg	Atlantic croaker	egg	maternal transfer	2 wks during reproduction (adults)	reproduction (egg exposure)	reduction in larval growth rate	McCarthy et al. (2003)	parental fish fed dietary PCBs, eggs exposed via maternal transfer; residues not clearly presented
PCBs (Aroclor 1254)	8,000					µg/kg	rainbow trout	14 wks	food	32 wks	growth, survival		Lieb et al. (1974)	only no-effect level reported
PCBs (Aroclor 1254)	1,900	9,300				µg/kg	sheepshead minnow	adult		28 days	reproduction	decreased fry survival	Hansen et al. (1974a)	elevated PCB concentrations in control fish; egg production was artificially stimulated by injecting the fish with human chorionic gonadotropic hormone
PCBs (Aroclor 1268)	15,000					µg/kg	mummichog	adult	food	~6 wks	reproduction	fertilization and hatching success, larval survival	Matta et al. (2001)	two generations of progeny observed; only no-effect level reported; offspring weight was significantly greater for PCB-exposed fish with parental tissue burdens equal to or greater than 1,300 µg/kg ww; however, this was not considered an adverse effect
PCBs (Aroclor 1254)	27,000	46,000				µg/kg	spot		water	20 days	survival		Hansen et al. (1971)	mortality did not appear directly related to body burden; bb increased with exposure duration; NOAEL (catfish) = 32
PCBs (Aroclor 1254)	31,000	71,000				µg/kg	brook trout	fry- exposure to eggs	water	10 days prior to hatch and 118 days after hatch	growth	reduced growth	Mauck et al. (1978)	residue measured at 118 days but was absent at 118 days.
PCBs (Aroclor 1254)					77,900	µg/kg	brook trout	egg	water	21 days	reproduction (egg exposure)	reduced hatchability	Freeman and Idler (1975)	75% hatching at LOAEL and 92% hatching in control; concentration in back muscle of dose fish with affected hatchability was 32.8 mg/kg ww
PCBs (Aroclor 1016)	110,000					µg/kg	sheepshead minnow	fry	water	4 wks	reproduction	fertilization and hatching success, larval survival	Hansen et al. (1975)	intermittent-flow toxicity test; no effect: fertilization success, survival of embryos to hatching, or survival of fry; only no-effect level reported; Juvenile tissue concentration LOAEL for reduced survival at the same exposure level was 220,000 µg/kg ww

Table 2, continued

Chemical	NOAEL (WB)	LOAEL (WB)	Conv. Factor	NOAEL (egg)	LOAEL (egg)	Unit (ww)	Test Species	Lifestage	Exposure Mode	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
PCBs (Aroclor 1016)		106,000				µg/kg	pinfish		water	33 days	survival, behavior	loss of equilibrium; erratic swimming	Hansen et al. (1974b)	significant reduction in survival (50% mortality relative to 6% in control)
PCBs (Aroclor 1254:1260 mixture)	120,000					µg/kg	rainbow trout	young	water	90 days	survival		Mayer et al. (1985)	mortality observed; not significantly different; dose was 1:2 ratio of Aroclor 1254:1260; only no-effect level reported
PCBs (Aroclor 1254:1260 mixture)	70,000	120,000				µg/kg	rainbow trout	young	water	90 days	growth		Mayer et al. (1985)	
PCBs (Aroclor 1254)	71,000	125,000				µg/kg	brook trout	fry- exposure to eggs	water	10days prior to hatch and 118days after hatch	survival	fry survival	Mauck et al. (1978)	reduced fry survival; 21 to 100% mortality; tissue residue measured at 118 days; Median hatching time and egg hatchability were not affected; larval growth was initially reduced, but not by the end of the test
PCBs (Aroclor 1016)	57,000	200,000				µg/kg	sheepshead minnow	fry	water		survival	fry survival	Hansen et al. (1975)	
PCBs (Clophen A50)		250,000				µg/kg	goldfish		water	5-21 days	survival	survival	Hattula and Karlog (1972)	LOAEL is concentration in individual dead fish
PCBs (Aroclor 1254)		429,000 (female)				µg/kg	fathead minnow		water		reproduction	reduced spawning	Nebeker et al. (1974)	mean terminal residue; egg hatchability and fry survival was not affected
PCBs (Aroclor 1242, 1254, or 1260)		1,860 – 749,000				µg/kg	fathead minnow		water	up to 300 hrs	survival	lethal body burden	van Wezel et al. (1995)	Tissue concentrations of individual fish that died in less than 20 hours ranged from 1,800 to 30,000 µg/kg ww; tissue concentrations of individual fish that died at 100 to 300 hours ranged from 120,000 to 749,000 µg/kg ww.
Mercury (mercuric chloride)	0.2					mg/kg	guppy	male adult	sediment and water	20 days	survival		Kudo and Mortimer (1979)	only no-effect level reported
Mercury					14	µg/kg	catfish	4 days post-hatch	water	spawning until 4 d post-hatch	survival	67% survival	Birge et al. (1979)	LOAEL is sac-fry concentration at 4 days post-hatch; control data was not reported; 17% survival was reported in catfish with WB residues of 1.15 µg/g ww
Mercury (inorganic Hg)					36	µg/kg	rainbow trout	alevins	exposed to contamin-ated sediment	20 days	reproduction (egg exposure)	egg/embryo survival reduced by 46%	Birge et al. (1979)	LOAEL is based on alevin (sac-fry) concentration of 0.036; adult concentration was estimated using sac-fry:adult conversion factor of 64.4 based on rainbow trout data from Niimi (1983) and assuming 3:2 ratio of concentration of sac-fry:egg
Mercury					41	µg/kg	rainbow trout	4 days post-hatch	water	spawning until 4 days post-hatch	survival	45% survival	Birge et al. (1979)	LOAEL is sac-fry concentration at 4 days post-hatch; 94% survival was reported in controls; 39 - 79% survival was reported in eggs with concentrations of 0.0682 µg/g ww; adult concentration was estimated using sac-fry:adult conversion factor of 64.4 based on rainbow trout data from Niimi (1983) and assuming 3:2 ratio of concentration of sac-fry:egg
Mercury (methylmercury)	230					µg/kg	golden shiner		food	90 days	survival	predator avoidance	Webber and Haines (2003)	Field collected fish (but from a manmade cement lined pond). Predator avoidance was reduced at this NOAEL; growth accompanied by reduced feeding was reduced at NOAEL but not dose-responsive, no effect on growth was observed in fish fed 959 ng/g ww where WB residue = 518 ug/g ww
Mercury (methyl-mercuric chloride)	200	470				µg/kg	mummichog	adult	water	42 days	survival		Matta et al. (2001)	effects observed in male fish but not females
Mercury (methylmercury)				990		µg/kg	walleye (field-collected)	egg	maternal transfer		reproduction (egg exposure)	fertilization and hatching success, larval length	Latif et al. (2001)	field-collected fish; adult concentration was estimated using egg:adult conversion factor of 36.05 based on average data reported in five species in Niimi (1983); mercury concentrations of diet not reported
Mercury (mercuric chloride)	0.8	1,310				µg/kg	fathead minnow	3 mos	food	60 days	growth		Snarski and Olson (1982)	
Mercury (methyl-mercuric chloride)	2,280					µg/kg	rainbow trout	fingerling	water	24 days	growth		Phillips and Buhler (1978)	only no-effect level reported

Table 2, continued

Chemical	NOAEL (WB)	LOAEL (WB)	Conv. Factor	NOAEL (egg)	LOAEL (egg)	Unit (ww)	Test Species	Lifestage	Exposure Mode	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
Mercury (methylmercury)	2,700	3,400				µg/kg	brook trout	embryo – adult	water	756 days	reproduction	reduced number of viable eggs produced	McKim et al. (1976)	residue measured in parental fish at 39 weeks; NOAEL is recommended no effect level presented in study (for abnormalities and multigenerational mortalities); NOAEL based on the data for no effect on the number of viable eggs produced was 1.1 µg/g ww
Mercury (methylmercury)					3,800	µg/kg	grayling	sac-fry (converted to adult WB)	water	1st 10 days of develop-ment	survival	increased fry mortality	Fjeld et al. (1998)	NOAEL and LOAEL reported as yolk-fry concentration; adult concentration was estimated using sac-fry:adult conversion factor of 64.4 based on rainbow trout data from Niimi (1983) and assuming 3:2 ratio of concentration of sac-fry:egg
Mercury (mercuric chloride)	2,750	4,180				µg/kg	fathead minnow	3 month old	water	60 days	survival		Snarski and Olson (1982)	
Mercury (mercuric chloride)	2,840	4,470				µg/kg	fathead minnow	larvae-adult	water	287 days	reproduction		Snarski and Olson (1982)	
Mercury (methylmercury)	5,000					µg/kg	rainbow trout	juvenile	water	84 days	growth, survival		Lock (1975)	only no-effect level reported
Mercury (mercuric chloride)		5,600				µg/kg	goldfish	4.5-6.5 cm	water	2 days	survival		Heisinger et al. (1979)	Concentration converted from dry weight to wet weight assuming 80% moisture in whole fish
Mercury (methyl-mercuric chloride)	5,670					µg/kg	rainbow trout	fingerling	water	24 days	growth		Phillips and Buhler (1978)	only no-effect level reported
Mercury (methyl-mercuric chloride)		6,500				µg/kg	bluegill	juvenile	water	12.5 days	survival		Cember et al. 1978	
Mercury (methyl-mercuric chloride)	8,630					µg/kg	rainbow trout	fingerling	water	24 days	growth		Phillips and Buhler (1978)	brook trout = 9.4 (NOAEL); only no-effect level reported
Mercury	9,400					µg/kg	brook trout	39 wks old	water	756 days	growth, survival	reduced juvenile weight	McKim et al. (1976)	residues measured in parental fish at 39 weeks; only no-effect level reported
Mercury (methylmercury)		10,000				µg/kg	rainbow trout	fingerling	food	84 days	growth	reduced growth and final body weight (after 84 days)	Rodgers and Beamish (1982)	reduced growth associated with feeding; approximated from graph (and text); at 14 days of treatment Hg concentrations in WB tissues were lower (approx. 5 µg/g ww); however, effects were observed (and significant) over the 84 days of exposure
Mercury (methylmercury)	10,900					µg/kg	fathead minnow	larvae-adult	water	336 days	growth, survival		Olson et al. (1975)	only no-effect level reported
Mercury (methyl-mercuric chloride)	1,100	11,000				µg/kg	mummichog		food	42 days	reproduction	F1 fertilization success	Matta et al. (2001)	residue measured in parental fish
Mercury (methyl-mercuric chloride)	12,000					µg/kg	rainbow trout	sub-adult	water	75 days	growth, survival	body weight	Niimi and Lowe-Jinde (1984)	only no-effect level reported
Mercury (methyl-mercuric chloride)	12,000					µg/kg	mummichog		food	42 days	reproduction	F1 hatchability, survival, fecundity, F2 larval survival	Matta et al. (2001)	residue measured in female parental fish; only no-effect level reported
Mercury (methyl-mercuric chloride)	12,000					µg/kg	mummichog		food	42 days	reproduction	fecundity, fertilization success, offspring weight, female survival	Matta et al. (2001)	offspring of medium and high doses were larger than offspring of control fish; only no-effect level reported; residue measured in female parental fish
Mercury	29,000					µg/kg	rainbow trout	fingerling		84 days	survival		Rodgers and Beamish (1982)	only no-effect level reported
Mercury (methylmercury)				16,000	29,000	µg/kg	Japanese medaka	egg (converted to adult WB)	water	16 days	reproduction (egg exposure)	hatchability	Heisinger and Green (1975)	at LOAEL - 20.8% hatchability; at NOAEL - 58.3% hatchability; at control - 46.7% hatchability; adult concentration was estimated using egg:adult conversion factor of 36.05 based on average data reported in five species in Niimi (1983)
Selenium	1,200	1,600				µg/kg	National criteria	na	na	na	protection of aquatic life	chronic criteria	EPA (2004)	concentration converted to ww assuming 80% moisture content;

Table 2, continued

Chemical	NOAEL (WB)	LOAEL (WB)	Conv. Factor	NOAEL (egg)	LOAEL (egg)	Unit (ww)	Test Species	Lifestage	Exposure Mode	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
TBT	18	159				µg/kg	Japanese flounder	larvae	food	approx 65 days	growth	body weight	Shimasaki et al. (2003)	all test fish were genetically XX fish by parents both XX but one parent phenotypically male and functional; No replication in study; survival was not significantly affected at any dose; mortality was observed in all groups, including the control group.
TBT		1,054				µg/kg	Japanese medaka	adult	food	3 wks	reproduction	reduced swim-up failure	Nakayama et al. (2005)	LOAEL is concentration in adult female fish using an adult:egg conversion factor of 8.57 based on Nirmala et al. (1999); effects were observed in offspring; concentration in egg tissues was 123 µg/kg
TBT		2,390				µg/kg	Japanese medaka	adult	food	3 wks	reproduction	reduced hatching success, swim-up success, and embryonic success	Nirmala et al. (1999)	LOAEL is concentration in adult female fish; effects were observed in offspring
Chlordane (alpha-chlordane and gamma-chlordane)	190 - 710					µg/kg	goldfish		food and water	96 hrs	survival	survival	Moore et al. (1977)	uncontrolled study; NOAEL is based on assumed 100% survival; NOAEL is based on the assumption that no mortality was observed because none was reported
Chlordane (cis-chlordane)		1,360				µg/kg	goldfish		water	24 hrs	survival	survival 3/9 fish died	Feroz and Khan (1979)	uncontrolled study; no clear residue burden and response to mortality effects; LOAEL is based on residues in carcasses of 3 dead fish; surviving fish tissue residues were 2.9 µg/g ww (at 10 days after exposure) and 2.1 µg/g (at 25 days after exposure)
Chlordane (tech chlordane)		16,600				µg/kg	pinfish	fry/ juvenile	water	96 hrs	survival	30% mortality	Parrish et al. (1976)	96-hr exposure; tissue residues in surviving fish exposed to 5.4 µg/L aqueous chlordane; seawater
Chlordane (tech chlordane)	87,000					µg/kg	sheepshead minnow	fry/juvenile	water	28 day	survival	3.7% mortality	Parrish et al. (1976)	28-day embryo/fry exposure; tissue residues in surviving fish exposed to 7.1 µg/L; tissue residues not reported in fish exposed to higher aqueous concentrations (100% mortality); seawater
Chlordane (tech chlordane)		281,000				µg/kg	sheepshead minnow	fry/juvenile	water	96 hrs	survival	25% mortality	Parrish et al. (1976)	96-hr exposure; tissue residues in surviving fish exposed to 15 µg/L; seawater; tissue residue concentration at lowest water treated concentration (unbounded)
DDT (mixture)	25					µg/kg	golden shiner		food	6-15 days	survival	survival	Courtney and Reed (1971)	converted from dry weight to wet weight using factor given in paper
DDT (mixture)		550				µg/kg	pinfish	4.8 g	diet	15 days	survival	44% mortality at 10 days	Butler (1969)	4% mortality in control; residues were much higher in live fish (LOAEL is residue of dead fish); at 7 days live fish WB residue = 2,700 µg/kg ww; residues were 640 µg/kg ww where morality was 10% (at 14 days); poorly reported study; no clear dose-response
DDT (mixture)	1,800	1,800				µg/kg	cutthroat trout	21 months	water	612 days (mortality observed at 111 days)	survival		Allison et al. (1964)	LOAEL is tissue concentration at 111 days (3.7 months) in fish exposed to 0.1 mg/kg DDT in water where mortality was significant after "4 months" (approximately 120d), note that tissue concentrations at this dose increased to 3.0 mg/kg at the next sampling
DDT (total)	1,920					µg/kg	brook trout	juvenile		120 days	survival		Macek and Korn (1970)	no effect level reported
DDT (total)	1,200	2,000				µg/kg	cutthroat trout	yearlings	water	20 mo (mortality observed at 4 mo)	survival	survival	Allison et al. (1963)	LOAEL is tissue concentration at 56 days (1.9 months) in fish exposed to 1.0 mg/kg DDT in water where mortality was significant after "4 months" (approximately 120d), note that tissue concentrations at this dose increased to 4,300 µg/kg at the next sampling.
DDT (total)		2,800 - 3,000				µg/kg	brook trout	sex mat yearlings	food	156 days	reproduction	offspring (sac-fry and embryo) mortality	Macek (1968b)	LOAEL is range of parental tissue residues; residues in offspring (fry) ranged from 2,620 to 3,090 µg/kg ww

Table 2, continued

Chemical	NOAEL (WB)	LOAEL (WB)	Conv. Factor	NOAEL (egg)	LOAEL (egg)	Unit (ww)	Test Species	Lifestage	Exposure Mode	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
DDT (total)	620	3,650				µg/kg	chinook salmon	0.61g	food	40 days	survival		Buhler et al. (1969)	tissue concentration at 4 and 7 days; growth increase attributed to size -selective mortality
DDT (total)	4,670					µg/kg	rainbow trout	15 g	food	140 days	growth, survival	survival, growth	Macek et al. (1970)	no effect level reported
DDT (total)		5,200				µg/kg	killifish		water	24 hrs	survival	25% mortality at 24 hrs	Crawford and Guarino (1976)	LOAEL is the weighted sum of residues in 10 different tissues of fish exposed to 0.1 ppm DDT for two 24hr DDT doses at 24 hrs after exposures; at 8 days after exposures, weighted sum of residues was 3,900 µg/kg ww, however, mortality started occurring at 24 hours.
DDT (mixture)	3,900	5,500				µg/kg	cutthroat trout	21 month old	food	612 days	survival	survival	Allison et al. (1964)	control diet had 760 µg/kg DDT; LOAEL tissue residue measured at day 166 in fish fed to 1,000 µg/kg body weight (bw) DDT and where mortality was observed; NOAEL is highest tissue concentration in fish fed 300 µg/kg bw DDT where significant mortality was not observed over entire exposure duration of 20 months; concentration is total chlorinated hydrocarbons
DDT (mixture)		5,600				µg/kg	pinfish	3.0 g	food	21 days	survival	35% mortality at 21 days	Butler (1969)	0% mortality in control; LOAEL is residue of dead fish; residues were 3,300 µg/kg ww where morality was 10% (at 14 days); poorly reported study; no clear dose-response of tissues with mortality across the experiments presented in the paper
DDT (mixture)		7,890				µg/kg	pinfish	3.0 g	food	2 days	survival	63.5% mortality	Butler (1969)	0% mortality in control; LOAEL is residue of dead fish; poorly reported study; no clear dose-response of tissues with mortality across the experiments presented in the paper
DDT (total)	7,600					µg/kg	brook trout fry	fry	food	156 days	growth, survival	survival, growth	Macek (1968b)	other fish species from study not reported because NOAEL/LOAELs are much higher; no effect level reported; length but not weight of male fish was significantly greater than controls
DDT (total)	11,200					µg/kg	brook trout	under yearling	food	31 wks	growth	increased growth	Macek (1968a)	NOAEL effect is an increase in growth (significant); sum of DDD, DDE, and DDT concentrations
DDT (total)	11,400	12,100				µg/kg	chinook salmon	1.1 g	food	40 days	survival	survival	Buhler et al. (1969)	tissues sampled at 4 and 7 days
DDT (total)		20,200 - 45,800				µg/kg	brook trout	under yearling	food	26 wks	survival	reduced survival during stress (starvation)	Macek (1968a)	sum of DDD, DDE, 2,4'-DDT and 4,4'-DDT in fish exposed to 2.0 mg/kg/wk DDT
DDT (total)		24,000				µg/kg	green sunfish/ pumpkin--seed		water	90 days	survival	survival	Hamelink et al. (1971)	
DDT (mixture)	24,000					µg/kg	Atlantic menhaden		food	48 (109)[a]	growth	growth	Warlen et al. (1977)	fish exposed for 48 days and observed for 109 days after exposure; no effect level reported
DDT (total)		26,500				µg/kg	mosquito fish		water	16 days	survival	survival	Pillai et al. (1977)	
DDT (total)	40,000					µg/kg	fathead minnow	juvenile-adult		266 days	survival	survival	Jarvinen et al. (1976; 1977)	no effect level reported
DDT (total)	51,400	92,700				µg/kg	sailfin molly		water	21 days	growth, survival	survival, growth	Benton et al. (1994)	converted from dry weight to wet weight based on reported 24% solids; sum of 2,4'-DDT, 4,4'-DDT, 2,4'-DDE, 4,4'-DDE, 2,4'-DDD, and 4,4'-DDD
DDT (total)	16,600	69,600				µg/kg	coho salmon	3.7 g	food	60 days	survival	survival	Buhler et al. (1969)	tissues sampled at 4 and 7 days
DDT (total)	130,000					µg/kg	goldfish	9.1 g	food and water	58 days	survival	survival	Rhead and Perkins (1984)	converted from dry weight to wet weight assuming 20% solids; no effect level reported

Table 2, continued

Chemical	NOAEL (WB)	LOAEL (WB)	Conv. Factor	NOAEL (egg)	LOAEL (egg)	Unit (ww)	Test Species	Lifestage	Exposure Mode	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
DDT (total)		200,000				µg/kg	goldfish	9.1 g	food and water	38 days	survival	survival	Rhead and Perkins (1984)	converted from dry weight to wet weight assuming 20% solids
Dieldrin	85 - 120					µg/kg	largemouth bass	adult	food	30-50 days	survival, growth	body weight, length, condition factor, survival	Muller et al. (2004)	estimated from graph; not clear if gonads included (measured at 1-3.5% of whole body concentration); blood not included
Dieldrin	120	200				µg/kg	rainbow trout	juvenile	food and water	16 wks	survival	33% mortality	Shubat and Curtis (1986)	5 of 15 fish died in high treated water (no maintenance diet treatment) and 4 of 15 fish died in high treated water and maintenance treated diet; no statistics; assumed no mortalities in control; no replication
Dieldrin	1,400					µg/kg	rainbow trout	juvenile		16 wks	growth		Shubat and Curtis (1986)	highest residue concentration in growth diet group (maintenance diet group with residues up to 0.36 ppm); no effect observed in body weight of treated fish groups
Endosulfan (tech endosulfan (sum of endosulfan I, II, and endosulfan sulfate)	3.1[b]	31				µg/kg	spot		water	96 hrs	survival	35% mortality; 10% mortality (control)	Schimmel et al. (1977a)	tissue residues of surviving fish
Endosulfan (tech endosulfan (sum of endosulfan I, II, and endosulfan sulfate)	195	272				µg/kg	pinfish		water	96 hrs	survival	35% mortality LOAEL; 5% NOAEL	Schimmel et al. (1977a)	tissue residues of surviving fish
Endosulfan (tech endosulfan (sum of endosulfan I, II, and endosulfan sulfate)		360				µg/kg	mullet		water	96 hrs	survival	40% mort; 0% mortality (control)	Schimmel et al. (1977a)	tissue residues of surviving fish
Endrin (technical endrin)	1.15[b]	11.5				µg/kg	largemouth bass	fingerlings	water	20 days	survival	40% fingerling mortality	Fabacher (1976)	one dose; no statistics; static tanks treated every 5 days; residues in dead fish
Endrin (analytical endrin)		240				µg/kg	fathead minnow	30d	food and water	300 days (also reprod period)	survival		Jarvinen and Tyo (1978)	statistically significant results at LOAEL
Endrin	307					µg/kg	channel catfish	fingerlings	food	198 days	survival		Argyle et al. (1973)	no statistics; only no-effect level reported
Endrin	307					µg/kg	channel catfish	fingerlings	food	198 days	growth		Argyle et al. (1973)	no statistics; only no-effect level reported; LOAEL is average residue from day 20 to day 198
Endrin	410	720				µg/kg	channel catfish	fingerlings	water	55 days	survival		Argyle et al. (1973)	no statistics - 40% mortality; LOAEL is concentration at day 26 when mortality began; NOAEL is mean concentration when residue levels were maximal from days 49-55
Endrin	110	880				µg/kg	sheepshead minnow	all	water	4 wks	survival	juvenile survival- 1 month survival of F2	Hansen et al. (1977)	saltwater species, embryos spawned from field-collected adults; tissue residues in juvenile fish- effects based on mortality effects in fry
Endrin	260	940				µg/kg	sheepshead minnow	all	water	2 generations	reproduction	female fertility	Hansen et al. (1977)	saltwater species, embryos spawned from field-collected adults
Endrin		1,660				µg/kg	golden shiner		water	8 hrs	survival	100% mortality	Ludke et al. (1968)	LOAEL is average residue at time of death; average residue over 8 hours = 0.97 mg/kg ww (resistant [i.e. pre-exposed] test fish not evaluated)
gamma-Hexachloro-cyclohexane (lindane)	1,580[c]	79,000				µg/kg	sheepshead minnow	17-21 mm			survival	LR50	Schimmel et al. (1977b)	LR50
Heptachlor (tech heptachlor: 65% heptachlor; 22% trans-chlordane; 2% cis-chlordane; 2% nonachlor)	150[b]	1,500				µg/kg	spot		water	96 hrs	survival	LOAEL- 25% mortality	Schimmel et al. (1976)	no statistical methods; saltwater study

Table 2, continued

Chemical	NOAEL (WB)	LOAEL (WB)	Conv. Factor	NOAEL (egg)	LOAEL (egg)	Unit (ww)	Test Species	Lifestage	Exposure Mode	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
Heptachlor (technical heptachlor)		1,900 - 2,600				µg/kg	bluegill		food	up to 140 days	growth	average body weight	Andrews et al. (1966)	fish were exposed to 5 ppm technical heptachlor; growth effects were not statistically evaluated; LOAEL is range of heptachlor residue concentrations (in two replicates) where average body weight was reduced (approx 90% of controls) after 28 days; growth e
Heptachlor	1,700	5,300				µg/kg	spot		water	96 hrs	survival	NOAEL-0% mortality; LOAEL 85% mortality	Schimmel et al. (1976)	no statistical methods; saltwater study
Heptachlor (technical heptachlor)		17,300 - 24,800				µg/kg			water	24 hrs	survival	90% mortality	Andrews et al. (1966)	fish were exposed to technical heptachlor; 90% mortality was observed after 72 hrs following exposure period in fish exposed to 0.0500 and 0.0375 ppm technical heptachlor; LOAEL is tissue residues at these dose levels at 72 hrs; heptachlor epoxide, other related compounds and DDT were also measured in fish tissue
Heptachlor		20,000				µg/kg	sheepshead minnow		water	96 hrs	survival	LOAEL- 35% mortality	Schimmel et al. (1976)	no statistical methods; saltwater study
Heptachlor (tech heptachlor: 65% heptachlor; 22% trans-chlordane; 2% cis-chlordane; 2% nonachlor)\	5,700	34,000				µg/kg	pinfish		water	96 hrs	survival	NOAEL-5% mortality; LOAEL-50% mortality	Schimmel et al. (1976)	no statistical methods; saltwater study
Heptachlor epoxide (tech heptachlor)	80[b]	800 - 900				µg/kg	bluegill		food	up to 140 days	growth	average body weight	Andrews et al. (1966)	fish were exposed to 5 ppm technical heptachlor; growth effects were not statistically evaluated; LOAEL is range of heptachlor residue concentrations (in two replicates) where average body weight was reduced (approx 90% of controls) after 28 days; growth effects were more pronounced at higher dose levels and after greater exposure duration; heptachlor, other related compounds and DDT were also measured in fish tissue
Heptachlor epoxide (tech heptachlor)		1,660 - 2,400				µg/kg	bluegill		water	24 hrs	survival	90% mortality	Andrews et al. (1966)	fish were exposed to technical heptachlor; 90% mortality was observed after 72 hrs following exposure period in fish exposed to 50 and 37.5 µg/kg technical heptachlor; LOAEL is tissue residues at these dose levels at 72 hrs; heptachlor, other related compounds and DDT were also measured in fish tissue
Hexachlorobenzene	46,500					µg/kg	fathead minnow		water	28-day exposure	survival		Nebeker et al. (1989)	
Hexachlorobenzene	46,500					µg/kg	fathead minnow		water	28-day exposure	growth		Nebeker et al. (1989)	
Hexachlorobenzene	97,000					µg/kg	fathead minnow	embryo - juvenile	water	32 days	growth, survival	final body weight, survival	Carlson and Kosian (1987)	no significant effect on growth or survival
Hexachlorobenzene	97,000					µg/kg	fathead minnow	embryo	water	32-day exposure	growth, survival		Carlson and Kosian (1987)	development/mortality; embryos 4-12 hrs old; exposure period during embryo-to-juvenile stage; NOAEL is mean tissue residue n=2
Hexachlorobenzene	282,000					µg/kg	fathead minnow		sediment and water	28-day exposure	survival		Schuytema et al. (1990)	exposure period followed by 15-28 day depuration; uptake/bioaccumulation study
Hexachlorobenzene	468,000					µg/kg	fathead minnow		water	28-day exposure	survival		Schuytema et al. (1990)	exposure period followed by 15-28 day depuration; uptake/bioaccumulation study
Methoxychlor	50	300				µg/kg	brook trout	yearlings	food	30 days	growth	body weight	Oladimeji and Leduc (1975)	fish were fed four different concentrations of methoxychlor at four different rates resulting in the same dose of 0.67 mg/kg bw/d; LOAEL is WB concentration in fish fed at 1% ingestion rate

Table 2, continued

Chemical	NOAEL (WB)	LOAEL (WB)	Conv. Factor	NOAEL (egg)	LOAEL (egg)	Unit (ww)	Test Species	Lifestage	Exposure Mode	Exposure Duration	Endpoint	Endpoint Effect	Source	Notes
4-methylphenol	1,530[c]	76,500				µg/kg	rainbow trout		injection	96 hrs	survival	LD50	Kaiser et al. (1984)	LOAEL is injected LD50 dose following 96-hr exposure; tissue residues not measured
4-methylphenol		78,900				µg/kg	rainbow trout		injection	96 hrs	survival	LD50	Hodson et al. (1988)	LOAEL is injected LD50 dose following 96-hr exposure
Phenol	1,470[c]	73,400				µg/kg	rainbow trout		water	13 hrs	survival	lethal body burden	McKim & Schmieder (1990)	LOAEL is reported lethal residue - limited detail on study methods
Phenol	76,000	114,000				µg/kg	goldfish		water	25 hrs	survival	LC50	Kobayashi et al. (1979)	LOAEL is LC50 body burden based on dead fish exposed to 60 µg/L; NOAEL is estimated body burden using BCF reported in study (1.9) and water concentration (40 µg/L) where 95% survival was reported over the duration of the experiment
Phenol		238,000				µg/kg	goldfish		water	5 hrs	survival	LC50	Kishino and Kobayashi (1995)	LOAEL is LC 50 from pH 8 study; LC50s were 273, 238, and 420 µg/g at pH 6, 8 and 10, respectively; mean residues of dead fish from all exposure levels from pH levels 6-10 were: 223 - 309 µg/g
Phenol		420,000				µg/kg	rainbow trout		injection	96 hrs	survival	LD50	Kaiser et al. (1984)	LOAEL is injected LD50 dose following 96-hr exposure; tissue residues not measured
Bis (2-ethylhexyl) phthalate	390					µg/kg	rainbow trout - sac fry			10 day prior to hatch and effects measured 24 days post hatch	reproduction (sac-fry survival)	13% mortality at NOAEL (not statistically different than control); 6% mortality in control	Mehrle and Mayer (1976)	fry tissue residues calculated from bioconcentration factor and water concentrations reported in paper.
Bis (2-ethylhexyl) phthalate	1,000 - 6,750					µg/kg	sheepshead minnow		water	24 hrs	survival	survival	Wofford et al. (1981)	NOAEL is based on assumed 100% survival; NOAEL is based on the assumption that no mortality was observed because none was reported
Butylbenzyl phthalate	6,450					µg/kg	bluegill		water	21 days	survival	survival	Barrows et al. (1980)	NOAEL is based on assumed 100% survival; NOAEL is based on the assumption that no mortality was observed because none was reported; study is a bioaccumulation study
Di(n)butyl phthalate	660					µg/kg	bluegill		water	42 days	survival	survival	Barrows et al. (1980)	NOAEL is based on assumed 100% survival; NOAEL is based on the assumption that no mortality was observed because none was reported; study is a bioaccumulation study
Di(n)butyl phthalate	1,170					µg/kg	sheepshead minnow		water	24 hrs	survival	survival	Wofford et al. (1981)	NOAEL is based on assumed 100% survival; NOAEL is based on the assumption that no mortality was observed because none was reported
Dimethyl phthalate	498					µg/kg	bluegill		water	21 days	survival	survival	Barrows et al. (1980)	NOAEL is based on assumed 100% survival; NOAEL is based on the assumption that no mortality was observed because none was reported; study is a bioaccumulation study
Diethyl phthalate	1,102					µg/kg	bluegill		water	21 days	survival	survival	Barrows et al. (1980)	NOAEL is based on assumed 100% survival; NOAEL is based on the assumption that no mortality was observed because none was reported; study is a bioaccumulation study
Benzoic acid	3,380					µg/kg	mosquito fish		water	24 hrs	survival	survival	Lu and Metcalf (1975)	highly uncertain study- exposure conditions was a mini-ecosystem with no mortality results discussed (bioaccumulation study); was assumed that no mortality was observed because none was reported

[a] NOAEL estimated using uncertainty factor of 5 (chronic LOAEL to chronic NOAEL).

[b] NOAEL estimated using uncertainty factor of 10 (acute LOAEL to chronic NOAEL).

[c] NOAEL estimated using uncertainty factor of 50 (LC50 to chronic NOAEL).

LC10 – concentration that causes the death of 10% of a group of test animals

LC50 – concentration that causes the death of 50% of a group of test animals

LOAEL – lowest-observed-adverse-effect level

NC – TRVs not reported in database because study only injection dose was reported (no WB tissue residues were reported)

NOAEL – no-observed-adverse-effect level

PCB – polychlorinated biphenyl

TCDD – tetrachlorodibenzo-*p*-dioxin

TBT – tributyltin

ww – concentration in fish tissue was reported on a wet weight basis

REFERENCES

Adams WJ, DeGraeve GM, Sabourin TD, Cooney JD, Mosher GM. 1986. Toxicity and bioconcentration of 2,3,7,8-TCDD to fathead minnows (*Pimephales promelas*). Chemosphere 15(9-12):1503-1511.

Allison D, Kallman BJ, Cope OB, Van Valin C. 1963. Insecticides: effects on cutthroat trout of repeated exposure to DDT. Science 142(3594):958-961.

Allison DT, Kollman BJ, Cope OB, Van Valin C. 1964. Some chronic effects of DDT on cutthroat trout. Research report 64. Bureau of Sport Fisheries and Wildlife, US Fish and Wildlife Service, Washington, DC.

Andrews AK, Van Valin CC, Stebbings BE. 1966. Some effects of heptachlor on bluegills. Trans Am Fish Soc 95:297-309.

Argyle RL, Williams GC, Dupree HK. 1973. Endrin uptake and release by fingerling channel catfish (*Ictalurus punctatus*). J Fish Res Board Can 30:1743-1744.

Baker RTM, Handy RD, Davies SJ, Snook JC. 1998. Chronic dietary exposure to copper affects growth, tissue lipid peroxidation, and metal composition of the grey mullet, *Chelon labrosus*. Mar Environ Res 45(4/5):357-365.

Baldisserotto B, Chowdhury MJ, Wood CM. 2005. Effects of dietary calcium and cadmium on cadmium accumulation, calcium and cadmium uptake from the water, and their interactions in juvenile rainbow trout. Aquat Toxicol 72:99-117.

Barrows ME, Petrocelli SR, Macek KJ, Carroll JR. 1980. Bioconcentration and elimination of selected water pollutants by bluegill sunfish (*Lepomis macrochirus*). In: Haque R, ed, Dynamics, exposure and hazard assessment of toxic chemicals. Ann Arbor Science Publishers, Ann Arbor, MI.

Benton MJ, Nimrod AC, Benson WH. 1994. Evaluation of growth and energy storage as biological markers of DDT exposure in sailfin mollies. Ecotoxicol Environ Saf 29:1-12.

Berntssen MHG, Lundebye AK, Maage A. 1999a. Effects of elevated dietary copper concentrations on growth, feed utilization and nutritional status of Atlantic salmon (*Salmo salar* L.) fry. Aquaculture 174:167-181.

Berntssen MHG, Hylland K, Bonga SEW, Maage A. 1999b. Toxic levels of dietary copper in Atlantic salmon (*Salmo salar* L.) parr. Aquat Toxicol 46:87-99.

Birge WJ, Black JA, Westerman AG, Hudson JE. 1979. The effects of mercury on reproduction of fish and amphibians. In: Nriagu JO, ed, The biogeochemistry of mercury in the environment. Elsevier/North-Holland Biomedical Press, New York, NY, pp 629-655.

Blazer VS, Fournie JW, Weeks-Perkins BA. 1997. Macrophage aggregates: biomarker for immune function in fishes? In: Dwyer FJ, Doane TR, Hinman ML, eds,

Environmental toxicology and risk assessment: Modeling and risk assessment. Vol 6. STP 1317. American Society for Testing and Materials, Philadelphia, PA, pp 360-375.

Branson DR, Takahashi IT, Parker WM, Blau GE. 1985. Bioconcentration kinetics of 2,3,7,8-tetrachlorodibenzo-p-dioxin in rainbow trout. Environ Toxicol Chem 4:779-788.

Buhler DR, Rasmusson ME, Shanks WE. 1969. Chronic oral DDT toxicity in juvenile coho and chinook salmon. Toxicol Appl Pharmacol 14:535-555.

Butler PA. 1969. Significance of DDT residues in estuarine fauna. In: Miller MW, Berg GG, eds, Chemical fallout: current research on persistent pesticides. Charles C. Thomas, Springfield, IL, pp 205-220.

Carlson AR, Kosian PA. 1987. Toxicity of chlorinated benzenes to fathead minnows (*Pimephales promelas*). Arch Environ Contam Toxicol 16:129-135.

Cockell KA, Bettger WJ. 1993. Investigations of the gallbladder pathology associated with dietary exposure to disodium arsenate heptahydrate in juvenile rainbow trout. Toxicology 77:233-248.

Cockell KA, Hilton JW. 1988. Preliminary investigation on the comparative chronic toxicity of four dietary arsenicals to juvenile rainbow trout (*Salmo gairdneri* R.). Aquat Toxicol 12:73-82.

Cockell KA, Hilton JW, Bettger WJ. 1991. Chronic toxicity of dietary disodium arsenate heptahydrate to juvenile rainbow trout (*Oncorhynchus mykiss*). Arch Environ Contam Toxicol 21:518-527.

Cockell KA, Hilton JW, Bettger WJ. 1992. Hepatobiliary and hematological effects of dietary disodium arsenate heptahydrate in juvenile rainbow trout (*Oncorhynchus mykiss*). Comp Biochem Physiol 103C(3):453-458.

Courtney CH, Reed JK. 1971. Accumulation of DDT from food and from water by golden shiner minnows, *Notemigonus crysoleucas*. 25th Annual Conference, Southeastern Association of Game and Fish Commissioners, Charleston SC. Southeastern Association of Game and Fish Commissioners, Frankfort, KY, pp 426-431.

Crawford RB, Guarino AM. 1976. Effects of DDT in *Fundulus*: studies on toxicity, fate, and reproduction. Arch Environ Contam Toxicol 4:334-348.

Elonen GE, Spehar RL, Holcombe GW, Johnson RD, Fernandez JD, Erickson RJ, Tietge JE, Cook PM. 1998. Comparative toxicity of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin to seven freshwater fish species during early life-stage development. Environ Toxicol Chem 17(3):472-483.

EPA. 2004. Draft aquatic life water quality criteria for selenium - 2004. EPA-822-D-04-001. US Environmental Protection Agency, Washington, DC.

Fabacher DL. 1976. Toxicity of endrin and an endrin-methyl parathion fomulation to largemouth bass fingerlings. Bull Environ Contam Toxicol 16(3):376-378.

Feroz M, Khan MAQ. 1979. Fate of ^{14}C-cis-chlordane in goldfish, *Carassius auratus* (L.). Bull Environ Contam Toxicol 23:64-69.

Fisher JP, Spitsbergen JM, Bush B, Jahan-Parwar B. 1994. Effect of embryonic PCB exposure on hatching success, survival, growth and developmental behavior in landlocked Atlantic salmon, *Salmo salar*. In: Gorsuch JW, Dwyer FJ, Ingersoll CG, La Point TW, eds, Environmental toxicology and risk assessment. Vol 2. ASTM STP 1216. American Society for Testing and Materials, Philadelphia, PA, pp 298-314.

Fisk AT, Yarechewski AL, Metner DA, Evans RE, Lockhart WL, Muir DCG. 1997. Accumulation, depuration and hepatic mixed-function oxidase enzyme induction in juvenile rainbow trout and lake whitefish exposed to dietary 2,3,7,8-tetrachlorodibenzo-p-dioxin. Aquat Toxicol 37:201-220.

Fjeld E, Haugen TO, Vollestad LA. 1998. Permanent impairment in the feeding behavior of grayling (*Thymallus thymallus*) exposed to methylmercury during embryogenesis. Sci Total Environ 213:247-254.

Franklin NM, Glover CN, Nicol JA, Wood CM. 2005. Calcium/cadmium interactions at uptake surfaces in rainbow trout: waterborne versus dietary routes of exposure. Environ Toxicol Chem 24(11):2954-2964.

Freeman HC, Idler DR. 1975. The effect of PCB on steroidogenesis and reproduction in the brook trout (*Salvelinus fontinalis*). Can J Biochem 53:666-670.

Galvez F, Wood CM. 1999. Physiological effects of dietary silver sulfide exposure in rainbow trout. Environ Toxicol Chem 18(1):84-88.

Gatlin DM, III, Wilson RP. 1986. Dietary copper requirement of fingerling channel catfish. Aquacult 54:277-285.

Giesy JP, Jones PD, Kannan K, Newsted JL, Tillitt DE, Williams LL. 2002. Effects of chronic dietary exposure to environmentally relevant concentrations if 2,3,7,8-tetrachlorodibenzo-p-dioxin on survival, growth, reproduction and biochemical responses of female rainbow trout (*Oncorhynchus mykiss*). Aquat Toxicol 59:35-53.

Goettl JP, Davies PH, Sinley JR. 1976. Water pollution studies. Colorado Fisheries Research Review 1972-1975. CO Div Wildlife Rev 8:68-75.

Guiney PD, Cook PM, Casselman JM, Fitzsimmons JD, Simonin HA, Zabel EW, Peterson RE. 1996. Assessment of 2,3,7,8-tetrachlorodibenzo-p-dioxin induced sac fry mortality in lake trout (*Salvelinus namaycush*) from different regions of the Great Lakes. Can J Fish Aquat Sci 53:2080-2092.

Hamelink JL, Waybrant RC, Ball RC. 1971. A proposal: exchange equilibria control the degree chlorinated hydrocarbons are biologically magnified in lentic environments. Trans Am Fish Soc 100(2):207-213.

Handy RD. 1992. The assessment of episodic metal pollution. II. The effects of cadmium and copper enriched diets on tissue contaminant analysis in rainbow trout (*Oncorhynchus mykiss*). Arch Environ Contam Toxicol 22:82-87.

Handy RD. 1993. The effect of acute exposure to dietary Cd and Cu on organ toxicant concentrations in rainbow trout, *Oncorhynchus mykiss*. Aquat Toxicol 27:1-14.

Hansen D, Schimmel SC, Forester J. 1974a. Aroclor 1254 in eggs of sheepshead minnows: effect on fertilization success and survival of embryos and fry. Proceedings of 27th Annual Conference, Southeastern Association of Game and Fish Commissioners, Hot Springs, AR, pp 420-426.

Hansen DJ, Parrish PR, Lowe JL, Wilson AJ, Jr, Wilson PD. 1971. Chronic toxicity, uptake, and retention of Aroclor 1254 in two estuarine fishes. Bull Environ Contam Toxicol 6:113-119.

Hansen DJ, Parrish PR, Forester J. 1974b. Aroclor 1016: Toxicity to and uptake by estuarine animals. Environ Res 7:363-373.

Hansen DJ, Schimmel SC, Forester J. 1975. Effects of Aroclor 1016 on embryos, fry, juveniles, and adults of sheepshead minnows (*Cyprinodon variegates*). Trans Am Fish Soc 104:584-588.

Hansen DJ, Schimmel SC, Forester J. 1977. Endrin: effects on the entire life cycle of a saltwater fish, *Cyprinodon variegatus*. J Toxicol Environ Health 3:721-733.

Hart DR, Heddle JA. 1991. Micronucleus assays in peripheral blood of rainbow trout: Timing of response and chemical mutagen sensitivity. Can Tech Rep Fish Aquat Sci 1774(2):993-1010.

Hatakeyama S, Yasuno M. 1982. Accumulation and effects of cadmium on guppy (*Poecilia reticulata*) fed cadmium-dosed Cladocera (*Moina macrocopa*). Bull Environ Contam Toxicol 29:159-166.

Hatakeyama S, Yasuno M. 1987. Chronic effects of Cd on the reproduction of the guppy (*Poecilia reticulata*) through Cd-accumulated midge larvae (*Chironomus yoshimatsui*). Ecotoxicol Environ Saf 14:191-207.

Hattula ML, Karlog O. 1972. Toxicity of polychlorinated biphenyls (PCBs) to goldfish. Acta Pharmacol Toxicol 31:238-240.

Hawkes CL, Norris LA. 1977. Chronic oral toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) to rainbow trout. Trans Am Fish Soc 106(6):641-645.

Heisinger JF, Green W. 1975. Mercuric chloride uptake by eggs of the ricefish and resulting teratogenic effects. Bull Environ Contam Toxicol 14(6):665-673.

Heisinger JF, Hansen CD, Kim JH. 1979. Effect of selenium dioxide on the accumulation and acute toxicity of mercuric chloride in goldfish. Arch Environ Contam Toxicol 8:279-283.

Hendricks JD, Scott WT, Putnam TP, Sinnhuber RO. 1981. Enhancement of aflatoxin B1 hepatocarcinogenesis in rainbow trout (*Salmo gairdneri*) embryos by prior exposure of gravid females to dietary Aroclor 1254. In: Branson DR, Dickson KL, eds, Aquat Toxicol and Hazard Assessment. Fourth Conference. ASTM STP 737. American Society for Testing and Materials, Philadelphia, PA, pp 203-214.

Hendricks JD, Meyers TR, Shelton DW, Casteel JL, Bailey GS. 1985. Hepatocarcinogenicity of benzo[a]pyrene to rainbow trout by dietary exposure and intraperitoneal injection. J Nat Cancer Inst 74:839-851.

Hilton JW, Bettger WJ. 1988. Dietary vanadium toxicity in juvenile rainbow trout: a preliminary study. Aquat Toxicol 12:63-71.

Hodson PV, Dixon DG, Kaiser KLE. 1988. Estimating the acute toxicity of waterborne chemicals in trout from measurements of median lethal dose and the octanol-water partition coefficient. Environ Toxicol Chem 7:443-454.

Hugla JL, Thome JP. 1999. Effects of polychlorinated biphenyls on liver ultrastructure, hepatic monooxygenases, and reproductive success in the barbel. Ecotoxicol Environ Saf 42:265-273.

Jarvinen AW, Tyo RM. 1978. Toxicity to fathead minnows of endrin in food and water. Arch Environ Contam Toxicol 7:409-421.

Jarvinen AW, Hoffman MJ, Thorslund TW. 1976. Toxicity of DDT food and water exposure to fathead minnows. EPA-600/3-76/114. US Environmental Protection Agency, Duluth, MN.

Jarvinen AW, Hoffman MJ, Thorslund TW. 1977. Long-term toxic effects of DDT food and water exposure on fathead minnows (*Pimephales promelas*). J Fish Res Board Can 34:2089-2103.

Johnson RD, Tietge JE, Jensen KM, Fernandez JD, Linnum AL, Lothenback DB, Holcombe GW, Cook PM, Christ SA, Lattier DL, Gordon DA. 1998. Toxicity of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin to early life stage brook trout (*Salvelinus fontinalis*) following parental dietary exposure. Environ Toxicol Chem 17(12):2408-2421.

Kaiser KLE, Dixon DG, Hodson PV. 1984. QSAR studies on chlorophenols, chlorobenzenes and para-substituted phenols. In: Kaiser KLE, ed, QSAR in environmental toxicology. D. Reidel Publishing Co., Dordrecht, the Netherlands, pp 189-206.

Kamunde CN, Grosell M, Lott JNA, Wood CM. 2001. Copper metabolism and gut morphology in rainbow trout (*Oncorhynchus mykiss*) during chronic sublethal dietary copper exposure. Can J Fish Aquat Sci 58:293-305.

Kang J-C, Kim S-G, Jang S-W. 2005. Growth and hematological changes of rockfish, *Sebastes schlegeli* (Hilgendorf) exposed to dietary Cu and Cd. J World Aquacult Soc 36(2):188-195.

Kim S-G, Kim J-W, Kang J-C. 2004. Effect of dietary cadmium on growth and haematological parameters of juvenile rockfish, *Sebastes schlegeli* (Hilgendorf). Aquacult Res 35:80-86.

Kishino T, Kobayashi K. 1995. Relation between toxicity and accumulation of chlorophenols at various pH, and their absorption mechanism in fish. Wat Res 29(2):431-442.

Kobayashi K, Akitake H, Manabe K. 1979. Relation between toxicity and accumulation of various chlorophenols in goldfish. Bull Jap Soc Sci Fish 45(2):173-175.

Kudo A, Mortimer DC. 1979. Pathways for mercury uptake by fish from bed sediments. Environ Pollut 13(1979):239-245.

Lanno RP, Slinger SJ, Hilton JW. 1985a. Effect of ascorbic acid on dietary copper toxicity in rainbow trout (*Salmo gairdneri* Richardson). Aquaculture 49:269-287.

Lanno RP, Slinger SJ, Hilton JW. 1985b. Maximum tolerable and toxicity levels of dietary copper in rainbow trout (*Salmo gairdneri* Richardson). Aquaculture 49:257-268.

Latif MA, Bodaly RA, Johnston TA, Fudge RJP. 2001. Effects of environmental and maternally derived methylmercury on the embryonic and larval stages of walleye (*Stizostedion vitreum*). Environ Pollut 111(2001):139-148.

Lieb AJ, Bills DD, Sinnhuber RO. 1974. Accumulation of dietary polychlorinated biphenyls (Aroclor 1254) by rainbow trout (*Salmo gairdneri*). J Agr Food Chem 22:638-642.

Lock RAC. 1975. Uptake of methylmercury by aquatic organisms from water and food. In: Koeman JH, Strik JJTWA, eds, Sublethal effects of toxic chemicals on aquatic organisms. Elsevier Press, Amsterdam, pp 61-79.

Lorentzen M, Maage A, Julshamn K. 1998. Supplementing copper to a fish meal based diet fed to Atlantic salmon parr affects liver copper and selenium concentrations. Aquacult Nutrit 4:67-72.

Lu P-Y, Metcalf RL. 1975. Environmental fate and biodegradability of benzene derivatives as studied in a model aquatic ecosystem. Environ Health Persp 10:268-284.

Ludke JL, Ferguson DE, Burke WD. 1968. Some endrin relationships in resistant and susceptible populations of golden shiners, *Notemigonus crysoleucas*. Trans Am Fish Soc 97:260-263.

Lundebye AK, Berntssen MHG, Bonga WSE, Maage A. 1999. Biochemical and physiological responses in Atlantic salmon (*Salmo salar*) following dietary exposure to copper and cadmium. Mar Pollut Bull 39:137-144.

Macek KJ. 1968a. Growth and resistance to stress in brook trout fed sublethal levels of DDT. J Fish Res Bd Can 25(11):2443-2451.

Macek KJ. 1968b. Reproduction in brook trout (*Salvelinus fontinalis*) fed sublethal concentrations of DDT. J Fish Res Board Can 25(9):1787-1796.

Macek KJ, Korn S. 1970. Significance of the food chain in DDT accumulation by fish. J Fisheries Res Board Can 27(8):1496-1498.

Macek KJ, Rodgers CR, Stalling DL, Korn S. 1970. The uptake, distribution and elimination of dietary ^{14}C-DDT and ^{14}C-Dieldrin in rainbow trout. Trans Am Fish Soc 99:689-695.

Matta MB, Linse J, Cairncross C, Francendese L, Kocan RM. 2001. Reproductive and transgenerational effects of methylmercury or Aroclor 1268 on *Fundulus heteroclitus*. Environ Toxicol Chem 20(2):327-335.

Mauck WL, Mehrle PM, Mayer FL. 1978. Effects of the polychlorinated biphenyl Aroclor 1254 on growth, survival, and bone development in brook trout (*Salvelinus fontinalis*). J Fish Res Board Can 35:1084-1088.

Mayer KS, Mayer FL, Witt A, Jr. 1985. Waste transformer oil and PCB toxicity to rainbow trout. Trans Am Fish Soc 114(6):869-886.

McCarthy ID, Fuiman LA, Alvarez MC. 2003. Aroclor 1254 affects growth and survival skills of Atlantic croaker *Micropogonias undulatus* larvae. Mar Ecol Prog Ser 252:295-301.

McKim JM, Schmeider PK. 1990. Bioaccumulation: does it reflect toxicity? In: Nagel R, Loskill R, eds, Bioaccumulation in aquatic systems: Contributions to the assessment. Proceedings, International Workshop, Berlin, 1990. VCG Publishers, New York, NY, pp 161-188.

McKim JM, Olson GF, Holcombe GW, Hunt EP. 1976. Long-term effects of methylmercuric chloride on three generations of brook trout (*Salvelinus fontinalis*): Toxicity, accumulation, distribution, and elimination. J Fish Res Board Can 33:2726-2739.

Meador JP, Sommers FC, Ylitalo GM, Sloan CA. 2006. Altered growth and related physiological responses in juvenile chinook salmon (*Oncorhynchus tshawytscha*) from dietary exposure to polycyclic aromatic hydrocarbons (PAHs). Can J Fish Aquat Sci 63:2364-2376.

Mehrle PM, Mayer FL. 1976. Di-2-ethylhexyl phthalate: residue dynamics and biological effects in rainbow trout and fathead minnows. Proceedings of

University of Missouri's 10th Annual Conference on Trace Substances in Environmental Health, June 8-10, Columbia, MO, pp 519-524.

Mehrle PM, Buckler DR, Little EE, Smith LM, Petty JD, Peterman PH, Stalling DL, De Graeve GM, Coyle JJ, Adams WJ. 1988. Toxicity and bioconcentration of 2,3,7,8-tetrachlorodibenzodioxin and 2,3,7,8-tetrachlorodibenzofuran in rainbow trout. Environ Toxicol Chem 7:47-62.

Miller PA, Lanno RP, McMaster ME, Dixon DG. 1993. Relative contributions of dietary and waterborne copper to tissue copper burdens and waterborne-copper tolerance in rainbow trout (*Oncorhynchus mykiss*). Can J Fish Aquat Sci 5(8):1683-1689.

Miller RA, Norris LA, Loper BR. 1979. The response of coho salmon and guppies to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in water. Trans Am Fish Soc 108:401-407.

Moore R, Toro E, Stanton M, Khan MAQ. 1977. Absorption and elimination of ^{14}C-*alpha*- and *gamma*-chlordane by a freshwater alga, daphnid, and goldfish. Arch Environ Contam Toxicol 6:411-420.

Mount DR, Barth AK, Garrison TD, Barten KA, Hockett JR. 1994. Dietary and waterborne exposure of rainbow trout (*Oncorhynchus mykiss*) to copper, cadmium, lead and zinc using a live diet. Environ Toxicol Chem 13(12):2031-41.

Muller JK, Johnson KG, Sepulveda MS, Borgert CJ, Gross TS. 2004. Accumulation of dietary DDE and dieldrin by largemouth bass, *Micropterus salmoides floridanus*. Bull Environ Contam Toxicol 73:1078-1085.

Murai T, Andrews JW, Smith RG, Jr. 1981. Effects of dietary copper on channel catfish. Aquaculture 22:353-357.

Nakayama K, Oshima Y, Nagafuchi K, Hano T, Shimasaki Y, Honjo T. 2005. Early-life-stage toxicity in offspring from exposed parent medaka, *Oryzias latipes*, to mixtures of tributyltin and polychlorinated biphenyls. Environ Toxicol Chem 24(3):591-596.

Nebeker AV, Puglisi FA, DeFoe DL. 1974. Effect of polychlorinated biphenyl compounds on survival and reproduction of the fathead minnow and flagfish. Trans Am Fish Soc 103:562-568.

Nebeker AV, Schuytema GS, Griffis WL, Barbitta JA, Carey LA. 1989. Effect of sediment organic carbon on survival of *Hyalella azteca* exposed to DDT and endrin. Environ Toxicol Chem 8:705-18.

Niimi AJ, Lowe-Jinde L. 1984. Differential blood cell ratios of rainbow trout (*Salmo gairdneri*) exposed to methylmercury and chlorobenzenes. Arch Environ Contam Toxicol 13:303-311.

Nirmala K, Oshima Y, Lee R, Imada N, Honjo T, Kobayashi K. 1999. Transgenerational toxicity of tributyltin and its combined effects with polychlorinated biphenyls on reproductive processes in Japanese medaka (*Oryzias latipes*). Environ Toxicol Chem 18(4):717-721.

Oladimeji AA, Leduc G. 1975. Effects of dietary methoxychlor on the food maintenance requirements of brook trout. Prog Wat Tech 7(3/4):587-598.

Oladimeji AA, Qadri SU, DeFreitas ASW. 1984. Long-term effects of arsenic accumulation in rainbow trout, *Salmo gairdneri*. Bull Environ Contam Toxicol 32:732-741.

Olson GF, Mount DI, Snarski VM, Thorslund TW. 1975. Mercury residues in fathead minnows, *Pimephales promelas* Rafinesque, chronically exposed to methylmercury in water. Bull Environ Contam Toxicol 14(2):129-134.

Palm RC, Jr, Powell DG, Skillman A, Godtfredsen K. 2003. Immunocompetence of juvenile chinook salmon against *Listonella anquillarum* following dietary exposure to polycyclic aromatic hydrocarbons. Environ Toxicol Chem 22(12):2986-2994.

Parrish PR, Schimmel SC, Hansen DJ, Patrick Jr JM, Forester J. 1976. Chlordane: effects on several estuarine organisms. J Toxicol Environ Health 1:485-494.

Phillips GR, Buhler DR. 1978. The relative contributions of methylmercury from food or water to rainbow trout (*Salmo gairdneri*) in a controlled laboratory environment. Trans Am Fish Soc 107(6):853-861.

Pillai MKK, Agarwal HC, Yadav DV. 1977. Tolerance, uptake and metabolism of DDT in *Gambusia affinis*. Ind J Exper Biol 15:40-41.

Powell DB, Palm RC, Jr, Skillman A, Godtfredsen K. 2003. Immunocompetence of juvenile chinook salmon against *Listonella anguillarum* following dietary exposure to Aroclor 1254. Environ Toxicol Chem 22(2):285-295.

Prince R, Cooper KR. 1995. Comparisons of the effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin on chemically impacted and nonimpacted subpopulations of *Fundulus heraclitus*: I. TCDD toxicity. Environ Toxicol Chem 14(4):579-587.

Rhead MM, Perkins JM. 1984. An evaluation of the relative importance of food and water as sources of p,p´-DDT to the goldfish, *Carassius auratus* (L.). Wat Res 18(6):719-725.

Rice CA, Myers MS, Willis ML, French BL, Casillas E. 2000. From sediment bioassay to fish biomarker – connecting the dots using simple trophic relationships. Mar Environ Res 50:527-533.

Rodgers DW, Beamish FWH. 1982. Dynamics of dietary methylmercury in rainbow trout, *Salmo gairdneri*. Aquat Toxicol 2(1982):271-290.

Schimmel SC, Patrick Jr JM, Forester J. 1976. Heptachlor: toxicity to and uptake by several estuarine organisms. J Toxicol Environ Health 1:955-965.

Schimmel SC, Patrick Jr JM, Wilson Jr AJ. 1977a. Acute toxicity to and bioconcentration of endosulfan by estuarine animals. In: Mayer FL, Hamelink JL, eds, Aquatic toxicology and hazard evaluation, ASTM STP 634. American Society for Testing and Materials, Philadelphia, PA, pp 241-252.

Schimmel SC, Patrick Jr JM, Forester J. 1977b. Toxicity and bioconcentration of BHC and lindane in selected estuarine animals. Arch Environ Contam Toxicol 6:355-363.

Schmieder P, Lothenbach D, Tietge J, Erickson R, Johnson RF. 1995. [^{3}H]-2,3,7,8-TCDD uptake and elimination kinetics of medaka (*Oryzias latipes*). Environ Toxicol Chem 14(10):1735-1743.

Schuytema GS, Krawczyk DF, Griffis WL, Nebeker AV, Robideaux ML. 1990. Hexachlorobenzene uptake by fathead minnows and macroinvertebrates in recirculating sediment/water systems. Arch Environ Contam Toxicol 19:1-9.

Shimasaki Y, Kitano T, Oshima Y, Inoue S, Imada N, Honjo T. 2003. Tributyltin causes masculinization in fish. Environ Toxicol Chem 22(1):141-144.

Shubat PJ, Curtis LR. 1986. Ration and toxicant preexposure influence dieldrin accumulation by rainbow trout (*Salmo gairdneri*). Environ Toxicol Chem 5:69-77.

Snarski VM, Olson GF. 1982. Chronic toxicity and bioaccumulation of mercuric chloride in the fathead minnow (*Pimephales promelas*). Aquat Toxicol 2:143-156.

Spehar RL, Cook PM, Elonen GE, Holcombe GW, Johnson RD, Fernandez JD, Tietge JE. 1997. Comparison of early-life-stage sensitivity of freshwater fish exposed as eggs to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). Submitted to Dioxin 97: 17th Symposium on Chlorinated Dioxins and Related Compounds, August 25-29, 1997, Indianapolis, IN. US Environmental Protection Agency, Duluth, MN.

Spitsbergen JM, Walker MK, Olson JR, Peterson RE. 1991. Pathologic alterations in early life stages of lake trout, *Salvelinus namaycush*, exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin as fertilized eggs. Aquat Toxicol 19:41-72.

Szebedinsky C, McGeer JC, McDonald DG, Wood CM. 2001. Effects of chronic Cd exposure via the diet or water on internal organ-specific distribution and subsequent gill Cd uptake kinetics in juvenile rainbow trout (*Oncorhynchus mykiss*). Environ Toxicol Chem 20(3):597-607.

Takeda H, Shimma Y. 1977. Effects of toxic amounts of dietary zinc on the growth and body components of rainbow trout at two levels of calcium. Bull Freshw Fish Res 27:103-109.

Tietge JE, Johnson RD, Jensen KM, Cook PM, Elonen GE, Fernandez JD, Holcombe GW, Lothenbach DB, Nichols JW. 1998. Reproductive toxicity and disposition

of 2,3,7,8-tetrachlorodibenzo-p-dioxin in adult brook trout (*Salvelinus fontinalis*) following a dietary exposure. Environ Toxicol Chem 17(12):2395-2407.

van Wezel AP, Sijm DTHM, Seinen W, Opperhuizen A. 1995. Use of lethal body burdens to indicate species differences in susceptibility to narcotic toxicants. Chemosphere 31(5):3201-3209.

Walker MK, Peterson RE. 1994. Toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin to brook trout (*Salvelinus fontinalis)* during early development. Environ Toxicol Chem 13:817-820.

Walker MK, Spitsbergen JM, Olson JR, Peterson RE. 1991. 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) toxicity during early life stage development of lake trout (*Salvelinus namaycush*). Can J Fish Aquat Sci 48:875-883.

Walker MK, Hufnagle LC, Clayton MK, Peterson RE. 1992. An egg injection method for assessing early life stage mortality of polychlorinated dibenzo-p-dioxins, dibenzofurans, and biphenyls in rainbow trout (*Oncorhynchus mykiss*). Aquat Toxicol 22:15-38.

Walker MK, Cook PM, Batterman AR, Butterworth BC, Berini C, Libal JJ, Hufnagle LC, Peterson RE. 1994. Translocation of 2,3,7,8-tetrachlorodibenzo-p-dioxin from adult female lake trout (*Salvelinus namaycush*) to oocytes: effects on early life stage development and sac fry survival. Can J Fish Aquat Sci 51:1410-1419.

Walsh AR, O'Halloran J, Gower AM. 1994. Some effects of elevated levels of chromium (III) in sediments to the mullet *Chelon labrosus* (R). Ecotoxicol Environ Saf 27:168-176.

Warlen SM, Wolfe DA, Lewis CW, Colby DR. 1977. Accumulation and retention of dietary ^{14}C-DDT by Atlantic menhaden. Trans Am Fish Soc 106(1):95-104.

Webber HM, Haines TA. 2003. Mercury effects on predator avoidance behavior of a forage fish, golden shiner (*Notemigonus crysoleucas*). Environ Toxicol Chem 22(7):1556-1561.

Wofford HW, Wilsey CD, Neff GS, Giam CS, Neff JM. 1981. Bioaccumulation and metabolism of phthalate esters by oysters, brown shrimp, and sheepshead minnows. Ecotoxicol Environ Saf 5:202-210.

Wu RSS, Pollino CA, Au DWT, Zheng GJ, Yuen BBH, Lam PKS. 2003. Evaluation of biomarkers of exposure and effect in juvenile areolated grouper (*Epinephelus areolatus*) on foodborne exposure to benzo(a)pyrene. Environ Toxicol Chem 22(7):1568-1573.

ATTACHMENT 9
ACCEPTABLE TOXICITY STUDIES FOR THE SELECTION OF BIRD DIETARY TOXICITY REFERENCE VALUES

Attachment 9 Acceptable Toxicity Studies for the Selection of Bird Dietary Toxicity Reference Values

Chemical	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source
Metals and Trace Elements									
Arsenic	sodium arsenate	mallard	6.1	na	food	10 wks	growth, survival	female	Camardese et al. (1990)
Arsenic	sodium arsenate	mallard	10	40	food	115-128 days	reproduction	delayed egg laying; depressed egg weight and shell thinning; decrease offspring body weight and production	Stanley et al. (1994)
Arsenic	sodium arsenite	mallard - young	25	50	food	154 days	survival	survival	USFWS (1964)
Cadmium	cadmium chloride	mallard young (females)	1.5	na	food	12 wks	growth	body weight	Cain et al. (1983)
Cadmium	$CdSO_4*8H_2O$	leghorn hen	0.73	2.9	food	48 wks	reproduction	egg production, shell thickness	Leach et al. (1979)
Cadmium	cadmium chloride	Japanese quail (chicks)	na	4.0	food	6 wks	growth	male body weight	Richardson et al. (1974)
Cadmium	cadmium chloride	mallard	19	na	food	90 days	survival		White and Finley (1978a)
Cadmium	cadmium chloride	mallard	20	na	food	30-90 days	survival, growth	body weight, adult survival	White and Finley (1978b)
Cadmium	cadmium chloride	mallard	1.5	20	food	30-90 days	reproduction	egg production	White and Finley (1978b)
Cadmium	cadmium chloride	leghorn chicks	na	24	food	21 days	growth	male body weight	Freeland and Cousins (1973)
Cadmium	cadmium chloride	leghorn chicks	na	40	food	20 days	growth	male body weight	Pritzl et al. (1974)
Cadmium	cadmium chloride	mallard	16	na	food	42 days	growth	body weight	DiGiulio and Scanlon (1984)
Cadmium	cadmium chloride	mallard	na	47	food	42 days	growth	body weight	DiGiulio and Scanlon (1984)
Chromium	copper sulfate	white leghorn layers (hens)	0.10	na	food	28 days	reproduction	egg weight and shell thickness	Lien et al. (2004)

Chemical	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source
Chromium	chromium 3+ as $CrK(SO_4)_2$	black duck	1.0	5.0	food	10 mo (and critical lifestage)	reproduction	duckling survival	Haseltine et al. (unpublished)
Chromium	Na_2CrO_4	Nochols chicks	7.7	na	food	22 days	growth, survival	male adult survival, male body weight	Romoser et al. (1961)
Cobalt	cobalt chloride	broiler chicks	2.31[b]	23.1	food	14 days	growth	body weight, survival	Diaz et al. (1994)
Copper	copper sulfate, copper amino acid complex	broiler chicks	2.1	na	food	17 days	growth, survival	body weight, survival	Dozier et al. (2003)
Copper	copper sulfate (hydrous)	hisex-brown hens	11.2	na	food	90 days	reproduction, survival	damaged egg ratio, egg weight and survival	Balevi & Coskun (2004)
Copper	copper sulfate	chicks, day-old	16	29	food	25 days	growth	growth	Smith (1969)
Copper	copper sulfate	chicks	21	41	food	4 wks	growth	growth and gizzard erosion	Poupoulis and Jensen (1976)
Copper	copper chloride	chicks	na	66	food	8-22 days	growth	body weight	Persia et al. (2004)
Copper	copper oxide	chicks	47	62	food	10 wks	growth/ survival	growth, survival	Mehring et al. (1960)
Copper	copper sulfate	white leghorn layers (hens)	15	na	food	28 days	reproduction	egg weight and shell thickness	Lien et al. (2004)
Lead	lead nitrate	mallards, first-year	2.5	na	food	12 wks	survival	survival	Finley et al. (1976)
Lead	metallic lead powder	American kestrel	5.82	na	food	5-7 mo	survival/ reproduction	survival, fertility, egg production, eggshell thinning	Pattee (1984)
Lead	lead acetate	Japanese quail	2.0	20	food	12 wks	reproduction	egg hatchability	Edens et al. (1976)
Lead	lead acetate	Japanese quail (chicks)	5.5	28	food	6 wks	growth	body weight	Morgan et al. (1975)
Mercury	methylmercury chloride	great egret, one day old	0.0091[b]	0.091	food	14 wks	growth	Growth	Spalding et al. (2000)
Mercury	methylmercury chloride	mallard	0.50	na	food	>60 days	reproduction	eggshell thickness	Heinz (1980)
Mercury	methylmercury chloride	Japanese quail (chicks at hatching)	na	0.9	food	5 days	survival	hatchling survival (16%)	Hill and Soares (1987)
Mercury	methylmercury chloride	zebra finch	0.72	1.4	food	76 days	survival	Survival	Scheuhammer (1988)

Chemical	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source
Mercury	methylmercury chloride	northern bobwhite, 12 day old	0.43	1.6	food	6 wks	survival	Survival	Spann et al. (1986a)
Mercury	mercuric chloride	Japanese quail, one day old	0.80	1.6	food	10 wks	reproduction	eggshell thickness	Stoewsand et al. (1971)
Mercury	dimethyl mercury	American kestrel	5.24	na	food	3 mo	survival	eggshell thickness	Peakall and Lincer (1972)
Mercury	mercuric chloride	Japanese quail (chicks at hatching)	na	62	food	5 days	survival	hatchling survival (12%)	Hill and Soares (1987)
Molybdenum	sodium molybdate	chicken	6.0[a]	30	food	21 days	reproduction	embryonic viability	Lepore and Miller (1965)
Nickel	nickel sulfate	broiler chicks	15	na	food	4 wks	growth	body weight gain	Weber and Reid (1968)
Nickel	nickel sulfate	broiler chicks	na	33	food	4 wks	growth	reduced body weight	Weber and Reid (1968)
Nickel	nickel acetate	broiler chicks	17	na	food	4 wks	growth	body weight gain	Weber and Reid (1968)
Nickel	nickel acetate	broiler chicks	na	38	food	4 wks	growth	reduced body weight	Weber and Reid (1968)
Nickel	nickel sulfate	mallard	na	107	food	90 days	survival, growth	survival, body weight	Cain and Pafford (1981)
Nickel	nickel sulfate	mallard	77	na	food	90 days	survival, growth	survival, body weight	Cain and Pafford (1981)
Nickel	nickel sulfate	mallard	132	na	food	90 days	survival, growth, reproduction	adult survival; body weight; hatchling weight	Eastin and O'Shea (1981)
Selenium	sodium selenite	broiler chicks	0.025	na	food	~40 days	growth, survival	body weight	Choct et al. (2004)
Selenium	sel-plex 50	broiler chicks	0.025	na	food	~40 days	growth, survival	body weight	Choct et al. (2004)
Selenium	seleno-methionine	mallard	0.42	na	food	~100 days	reproduction	offspring growth/survival	Heinz et al. (1989)
Selenium	seleno-methionine	mallard	na	0.82	food	~100 days	reproduction	offspring growth/survival	Heinz et al. (1989)
Selenium	Na_2SeO_3	mallard	0.50	na	food	4 wks before laying to 3 wks after hatching	reproduction	embryo abnormalities	Heinz et al. (1987)

Chemical	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source
Selenium	Na_2SeO_3	mallard	na	1.0	food	4 wks before laying to 3 wks after hatching	reproduction	embryo abnormalities	Heinz et al. (1987)
Selenium	Na_2SeO_3	mallard	1.0	na	food	4 wks before laying to 3 wks after hatching	growth	adult growth	Heinz et al. (1987)
Selenium	seleno-methionine	mallard	1.6	na	food	~100 days	survival, growth	body weight; adult survival	Heinz et al. (1989)
Selenium	Na_2SeO_3	mallard	na	2.5	food	4 wks before laying to 3 wks after hatching	growth	adult growth	Heinz et al. (1987)
Selenium	seleno-methionine	screech owl	1.0	3.2	food	~ 3 mo	growth, reproduction	body weight, hatching success, 5 days survival, clutch size, egg size and mass	Wiemeyer and Hoffman (1996)
Selenium	Na_2SeO_3	mallard	4.6	na	food	42 days	survival		Heinz et al. (1988)
Selenium	Na_2SeO_3	mallard		4.6	food	42 days	growth	body weight	Heinz et al. (1988)
Selenium	Na_2SeO_3	mallard	2.1		food	42 days	growth	body weight	Heinz et al. (1988)
Selenium	Na_2SeO_3	mallard	na	10	food	4 wks before laying to 3 wks after hatching	survival	adult survival	Heinz et al. (1987)
Selenium	Na_2SeO_3	mallard	2.5	na	food	4 wks before laying to 3 wks after hatching	survival	adult survival	Heinz et al. (1987)
Selenium	Na_2SeO_3	mallard	na	10	food	42 days	survival		Heinz et al. (1988)
Thallium	thallium sulfate	pheasant	2.4[b]	24	oral gavage	max 14 days	survival	LD50	Hudson et al. (1984)
Thallium	thallium sulfate	mallard	na	37	oral gavage	max 14 days	survival	LD50	Hudson et al. (1984)
Vanadium	ammonium metavanadate	white leghorn hens	2.0	na	food	6 wks	reproduction	egg weight, egg quality	Davis et al. (2002)
Vanadium	ammonium metavanadate	white leghorn hens	1.2	2.3	food	4 wks	growth	body weight	Ousterhout and Berg (1981)
Vanadium	vanadium sulfate	mallard	11.4	na	food	12 wks	growth, survival	body weight, survival	White and Dieter (1978)
Zinc	copper sulfate, copper amino acid complex	broiler chicks	17	na	food	17 days	growth, survival	body weight, survival	Dozier et al. (2003)

Chemical	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source
Zinc	zinc oxide, zinc sulfate, or zinc carbonate	white rock chicks	82	124	food	5 wks	growth	growth	Roberson and Schaible (1960)
Zinc	zinc sulfate	white leghorn hens	133	na	food and supplements	44 wks	reproduction	egg hatchability	Stahl et al. (1990)
Zinc	zinc carbonate	mallard (7 wks old)	na	300	food	60 days	survival	survival, leg paralysis	Gasaway and Buss (1972)
Zinc	zinc acetate	Hubbard broiler chicks	330	659	food	5 wks	growth/ survival	survival, reduced growth	Oh et al. (1979)
Zinc	zinc chloride	chicks	na	344	food	8-22 days	growth	body weight	Persia et al. (2004)
Organometals									
Tributyltin	TBTO	Japanese quail	22.5	na	food	6 wks	growth	body weight	Schlatterer et al. (1993)
Tributyltin	TBTO	Japanese quail	1.4	3.6	food	6 wks	reproduction	no hatched eggs per pair	Schlatterer et al. (1993)
Tributyltin	TBTO	Japanese quail	1.4	3.6	food	6 wks	reproduction	embryo survival in shell, hatchability	Coenen et al. (1992)
PAHs									
Benzo(a)pyrene	benzo(a)pyrene	pigeons	0.28[a]	1.4	weekly intramuscular injection	5 mo	reproduction	fertility, ovarian appearance	Hough et al. (1993)
Benzo(a)pyrene	benzo(a)pyrene	white rock chicken	33	na	food	30 days	growth	body weight gain	Rigdon and Neal (1963)
PAHs	aromatic hydrocarbon mixture including individual PAHs	mallard	8	40	food	7 mo	growth	little change in body weight	Patton and Dieter (1980)
PAHs	petroleum hydrocarbon mixture including PAHs)	mallard	400	na	food	7 mo	survival	survival	Patton and Dieter (1980)
Phthalates									
Bis(ethylhexyl) phthalate	BEHP	ringed turtle-dove	1.45	na	food	4 wks	reproduction	eggshell thickness	Peakall (1974)
Bis(ethylhexyl) phthalate	BEHP	European starling	67.8	na	food	30 days	growth	growth, food consumption	O'Shea and Stafford (1980)
Bis(ethylhexyl) phthalate	BEHP	chicken	na	329	food	230 days	reproduction	cessation of egg laying, abnormal ovaries	Ishida et al. (1982)
PCBs									

Chemical	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source
PCBs (Aroclor 1248)	Aroclor 1248	American kestrel	na	0.35	food	5.5 mo	reproduction	eggshell weight and thickness	Lowe and Stendell (1991)
PCBs (Aroclor 1248)	Aroclor 1248	screech owl	0.49	na	food	2 generations	reproduction	eggshell thickness, egg production, hatching success, fledging success	McLane and Hughes (1980)
PCBs (Aroclor 1242)	Aroclor 1242	Japanese quail	na	0.60	food	45 days	reproduction	eggshell thinning	Hill et al. (1976)
PCBs (Aroclor 1254)	Aroclor 1254	ringed turtle-dove	na	1.4	food	2 generations	reproduction	hatching success in second generation	Peakall et al. (1972); Peakall and Peakall (1973)
PCBs (Aroclor 1254)	Aroclor 1254	mallard	2.5	na	food	~ 1 mo	reproduction	Reproductive success	Custer and Heinz (1980)
PCBs (Aroclor 1254)	Aroclor 1254	mallard	3.9	na	food	4 mo	reproduction	egg production, eggshell thinning	Risebrough and Anderson (1975)
PCBs (Aroclor 1248: 1254:1260 mixture)	1:1:1 ratio of Aroclor 1248:1254:1260	American kestrel	na	5 to 7	food	100 days until eggs hatched	reproduction	egg laying in second generation (exposed in ovo); also some effect on clutch size and fledgling success	Fernie et al. (2000; 2001)
PCBs (Aroclor 1248: 1254:1260 mixture)	1:1:1 ratio of Aroclor 1248:1254:1260	American kestrel	na	5 to 7	food	1 mo prior to pairing until anticipated egg hatching	reproduction	cracked eggs, embryo abnormalities;	Fernie et al. (2003b)
PCBs (Aroclor 1248: 1254:1260 mixture)	1:1:1 ratio of Aroclor 1248:1254:1260	American kestrel	na	7	food	100 days	reproduction	offspring survival and offspring body weight	Fernie et al. (2003a)
PCBs (Aroclor 1242)	Aroclor 1242	mallard	na	15	food	12 wks	reproduction	hatchability, embryo survival, egg viability, embryo abnormalities	Haseltine and Prouty (1980)
Pesticides									
Aldrin	aldrin	quail	0.008[a]	0.04	food	5 mo	survival	97.5% survival in 127 days	DeWitt (1956)
Aldrin	aldrin	pheasant	na	0.7	food	16-20 wks	survival	100% survival in 46 days	DeWitt (1956)
Aldrin	aldrin	pen-reared pheasant chicks	10	20	gelatin capsule	7 wks (+ 9 wks untreated)	growth	growth	Hall et al. (1971)
Aldrin	aldrin	pen-reared pheasant chicks	na	92	gelatin capsule	7 wks (+ 9 wks untreated)	survival	growth	Hall et al. (1971)

Chemical	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source
Chlordane	tech chlordane	bobwhite quail	0.6	na	food	10 wks	growth, survival	body weight, adult survival	Ludke (1976)
Chlordane	exp chlordane (HCS 3260)	bobwhites (juvenile)	na	20	food	5 days	survival	50% survival	Hill et al. (1975); Heath et al. (1972)
Chlordane	exp chlordane (HCS 3260)	Japanese quail	na	21	food	5 days	survival	50% survival	Hill et al. (1975); Heath et al. (1972)
Chlordane	exp chlordane (HCS 3260)	ring-necked pheasant	na	25	food	5 days	survival	50% survival	Hill et al. (1975); Heath et al. (1972)
Chlordane	exp chlordane (HCS 3260)	mallard-young	na	115	food	5 days	survival	50% survival	Hill et al. (1975); Heath et al. (1972)
DDT	p,p'-DDT	quail	0.03	0.15	food	26 wks+12 days	reproduction	eggshell thickness	Stickel and Rhodes (1970)
Total DDT	tech DDT	mallard	0.18	na	food	11 mo	reproduction	eggshell weight and thickness	Davison and Sell (1974)
DDT	p,p'-DDT	mallard	0.19	na	food	11 mo	reproduction	eggshell weight and thickness	Davison and Sell (1974)
DDE	"DDE"	barn owls	na	0.32	food	2 yrs (2 nestings)	reproduction	eggshell breakage/ thickness; nestling survival	Mendenhall et al. (1983)
DDE	"DDE"	barn owls	0.32	na	food	2 yrs (2 nestings)	survival	adult survival	Mendenhall et al. (1983)
DDE	"DDE"	American kestrel	na	0.35	food	14 days	reproduction	eggshell thinning, egg permeability	Peakall et al. (1973)
DDD	tech DDD	mallard	na	0.90	food	2 yrs	reproduction	hatchling survival, production	Heath et al. (1969)
DDE	p,p'-DDE	mallard	na	0.90	food	2 yrs	reproduction	% cracked, hatchling survival/production, shell thickness, embryonation	Heath et al. (1969)
DDE	p,p-DDE	black duck	na	1.0	food	7 mo	reproduction	shell thickness, egg weight, hatchability, duckling survival	Longcore and Samson (1973)
DDE	DDE	mallard	na	1.0	food	30 days	reproduction	eggshell thinning	Kolaja (1977)
DDE	DDE	black duck	na	1.0	food	reproductive period (through incubation and hatching)	reproduction	eggshell thinning and cracking, embryo survival, duckling survival	Longcore et al. (1971)
DDT	DDT	mallard	na	1.0	food	30 days	reproduction	eggshell thinning	Kolaja (1977)

Chemical	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source
DDE	p,p' DDE	American kestrel	na	1.0	food	1 yr (2 clutches)	reproduction	eggshell thickness	Wiemeyer and Porter (1970); Porter and Wiemeyer (1972)
Dieldrin	dieldrin	Japanese quail	0.06	na	food	4 generation	survival, growth, reproduction	adult survival, body weight, fertility, hatchability, egg product	Shellenberger (1978)
Dieldrin	dieldrin	mallard	0.063	na	food	24 days	growth, survival	body weight, leg length, adult survival	Nebeker et al. (1992)
Dieldrin	dieldrin	barn owl	0.066	na	food	2 yrs (2 nestings)	survival, reproduction	adult survival, eggshell breakage/ thickness, nestling survival, hatching success, clutch size	Mendenhall et al. (1983)
Dieldrin	dieldrin	quail	0.080	0.12	food	5 mo	survival	17% survival vs. 9% in control	DeWitt (1956)
Dieldrin	dieldrin	mallard	na	0.16	food	> 1 year	reproduction	eggshell thinning	Lehner and Egbert (1969)
Dieldrin	dieldrin	bobwhite quail	0.24	0.47	food	2 mo	survival		Fergin and Schafer (1977)
Dieldrin	dieldrin	white leghorn chicken	0.3	na	food	16 wks	reproduction	egg hatch, 14-day chick survival	Graves et al. (1969)
Dieldrin	dieldrin	mallard	0.49	na	food	11 mo	reproduction	eggshell weight and thickness	Davison and Sell (1974)
Dieldrin	dieldrin	Japanese quail	0.54	na	food	75 days	reproduction	eggshell thinning	Hill et al. (1975)
Dieldrin	dieldrin	Japanese quail	na	0.60	food	up to 18 wks	reproduction	fertility	Walker et al. (1969)
Dieldrin	dieldrin	pheasant	na	0.60	food	16-20 wks	survival	100% survival by 68 days	DeWitt (1956)
Dieldrin	dieldrin	white leghorn males	na	0.65	food	20 wks	survival		Ahmed et al. (1978)
Dieldrin	dieldrin	quail	na	0.80	food	16-20 wks	survival	100% survival by 76 days; 24% in control	DeWitt (1956)
Dieldrin	dieldrin	red-light Sussex hybrids	na	0.83	food	13 mo adults, 6 wks chicks	growth	body weight	Brown et al. (1974)
Dieldrin	dieldrin	red-light Sussex hybrids	0.83	na	food	13 mo	reproduction	chick survival	Brown et al. (1974)

Chemical	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source
Dieldrin	dieldrin	ring-necked pheasant	na	0.86	food	8 wks	reproduction	offspring survival, egg production	Genelly and Rudd (1956)
Dieldrin	dieldrin	mallard	na	0.92	food	11 mo	reproduction	eggshell weight and thickness	Davison and Sell (1974)
Dieldrin	dieldrin	mallard	0.92	na	food	11 mo	survival, growth	body weight, adult survival	Davison and Sell (1974)
Endosulfan	endosulfan	gray partridge	10	na	food	4 wks- during critical lifestage	reproduction	number of eggs; egg fertility, embryo and chick survival; eggs hatched	Abiola (1992)
Endrin	endrin	quail	0.070	na	food	5 mo	survival		DeWitt (1956)
Endrin	endrin	quail	na	0.20	food	16-20 wks	survival		DeWitt (1956)
Endrin	endrin	mallard	0.29	na	food	7 mo	growth	body weight	Spann et al. (1986b)
Endrin	endrin	mallard	0.29	na	food	7 mo	reproduction	egg weight and fertility, shell thickness, duckling survival/ body weight, clutch size, hatch	Spann et al. (1986b)
Endrin	endrin	mallard	na	0.30	food	12 wks	growth, reproduction	body weight; onset egg production, embryo survival	Roylance et al. (1985)
Endrin	endrin	screech owl	0.33	na	food	mating-incubation	growth	body weight	Fleming et al. (1982)
Endrin	endrin	screech owl	na	0.33	food	mating-incubation (crit. lifestage)	reproduction	fledglings produced, egg production, number of hatched eggs	Fleming et al. (1982)
Endrin	endrin	pheasant	na	0.60	food	16-20 wks	survival	100% survival	DeWitt (1956)
gamma-Hexachloro-cyclohexane (Lindane)	gamma HCH	domestic mallard	1.6	3.6	oral intubation	8 wks	reproduction	eggshell thickness, size, quality; clutch size	Chakravarty and Lahiri (1986); Chakravarty et al. (1986)
gamma-Hexachloro-cyclohexane (Lindane)	gamma HCH	domestic mallard	20	na	oral intubation	8 wks	survival, growth	survival, body weight	Chakravarty et al. (1986)
Heptachlor	exp chlordane (HCS 3260)	bobwhites (juvenile)	0.5[b]	5	food	5 days	survival	50% survival	Hill et al. (1975); Heath et al. (1972)
Heptachlor	exp chlordane (HCS 3260)	Japanese quail	na	6	food	5 days	survival	50% survival	Hill et al. (1975); Heath et al. (1972)
Heptachlor	exp chlordane (HCS 3260)	ring-necked pheasant	na	13	food	5 days	survival	50% survival	Hill et al. (1975); Heath et al. (1972)

Chemical	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Endpoint Effect	Source
Heptachlor	exp chlordane (HCS 3260)	mallard-young	na	104	food	5 days	survival	50% survival	Hill et al. (1975); Heath et al. (1972)
Hexachlorobenzene	HCB	Japanese quail	1.1	na	food	90 days	reproduction	hatchability	Vos et al. (1971)
Hexachlorobenzene	HCB	Japanese quail	1.1	na	food	90 days	survival		Vos et al. (1971)
Hexachlorobenzene	HCB	Japanese quail	1.2	na	food	90 days	growth	body weight	Schwetz et al. (1974)
Hexachlorobenzene	HCB	Japanese quail	na	1.2	food	90 days	reproduction	chicks hatched and survival	Schwetz et al. (1974)
Hexachlorobenzene	HCB	Japanese quail	na	4.5	food	90 days	reproduction	hatchability	Vos et al. (1971)
Hexachlorobenzene	HCB	Japanese quail	na	5	food	90 days	survival		Vos et al. (1971)
Methoxychlor	technical methoxychlor	zebra finch (chicks at 5-11 days old)	34.6	346	oral gavage	1 wks	reproduction	number of eggs hatched, broken/missing eggs	Gee et al. (2004)
Methoxychlor	methoxychlor	zebra finch chicks	34.6	346	oral gavage	1 wks	survival	survival	Millam et al. (2002)
Methoxychlor	technical methoxychlor	bobwhites (juvenile), Japanese quail, ringed-necked pheasant, mallard (young)	831	na	food	5 days	survival	survival	Hill et al. (1975); Heath et al. (1972)

[a] NOAEL estimated using an uncertainty factor of 5 (chronic LOAEL to chronic NOAEL).

[b] NOAEL estimated using an uncertainty factor of 10 (acute/subchronic LOAEL to chronic NOAEL).

LOAEL – lowest-observed-effect concentration

na – not available

NOAEL – no-observed-effect concentration

PAH – polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

REFERENCES

Abiola FA. 1992. Ecotoxicity of organochloride insecticides: effects of endosulfan on bird reproduction and evaluation of its induction effect in partridge (*Perdix perdix* L.). Revue Med Vet 143(5):443-450.

Ahmed T, Arscott GH, Tinsley IJ. 1978. Effect of chlorinated hydrocarbons on reproductive performance of adult white leghorn male chickens. Poult Sci 57:1594-1598.

Balevi T, Coskun B. 2004. Effects of dietary copper on production and egg cholesterol content in laying hens. Brit Poult Sci 45(4):530-534.

Brown VKH, Robinson J, Thorpe E, Barrett JW. 1974. The toxicity of dieldrin (HEOD) to domestic fowl. Pestic Sci 5:567-586.

Cain BW, Pafford EA. 1981. Effects of dietary nickel on survival and growth of mallard ducklings. Arch Environ Contam Toxicol 10:737-745.

Cain BW, Sileo L, Franson JC, Moore J. 1983. Effects of dietary cadmium on mallard ducklings. Environ Res 32:286-297.

Camardese MB, Hoffman DJ, LeCaptain LJ, Pendleton GW. 1990. Effects of arsenate on growth and physiology in mallard ducklings. Environ Toxicol Chem 9:785-795.

Chakravarty S, Lahiri P. 1986. Effect of lindane on eggshell characteristics and calcium level in the domestic duck. Toxicology 42:245-258.

Chakravarty S, Mandal A, Lahiri P. 1986. Effect of lindane on clutch size and level of egg yolk protein in domestic duck. Toxicology 39:93-103.

Choct M, Naylor AJ, Reinke N. 2004. Selenium supplementation affects brioler growth performance, meat yield and feather coverage. Brit Poult Sci 45(5):677-683.

Coenen TMM, Brouwer A, Enninga IC, Koeman JH. 1992. Subchronic toxicity and reproduction effects if tri-n-butyltin oxide in Japanese quail. Arch Environ Contam Toxicol 23:457-463.

Custer TW, Heinz GH. 1980. Reproductive success and nest attentiveness of mallard ducks fed Aroclor 1254. Environ Pollut 21:313-318.

Davis EG, Miles RD, Butcher GD, Comer CS. 2002. Effects of dietary vanadium on performance and immune responses of commercial egg-type laying hens. J Appl Anim Res 22:113-124.

Davison KL, Sell JL. 1974. DDT thins shells of eggs from mallard ducks maintained on *ad libitum* or controlled-feeding regimens. Arch Environ Contam Toxicol 2(3):222-232.

DeWitt JB. 1956. Chronic toxicity to quail and pheasants of some chlorinated insecticides. Agric Food Chem 4(10):863-866.

Diaz GJ, Julian RJ, Squires EJ. 1994. Lesions in broiler chickens following experimental intoxication with cobalt. Avian Dis 38(2):308-316.

DiGiulio RT, Scanlon P. 1984. Sublethal effects of cadmium ingestion on mallard ducks. Arch Environ Contam Toxicol 13:765-771.

Dozier WA, III, Davis AJ, Freeman ME, Ward TL. 2003. Early growth and environmental implications of dietary zinc and copper concentrations and sources of broiler chicks. Brit Poult Sci 44(5):726-731.

Eastin WC, Jr, O'Shea TJ. 1981. Effects of dietary nickel on mallards. J Toxicol Environ Health 7:883-892.

Edens FW, Benton E, Bursian SJ, Morgan GW. 1976. Effect of dietary lead on reproductive performance in Japanese quail, *Coturnix coturnix japonica*. Toxicol Appl Pharmacol 38:307-314.

Fergin TJ, Schafer EC. 1977. Toxicity of dieldrin to bobwhite quail in relation to sex and reproductive status. Arch Environ Contam Toxicol 6:213-219.

Fernie K, Smiths J, Bortolotti G. 2003a. Developmental toxicity of in ovo exposure to polychlorinated biphenyls: I. Immediate and subsequent effects on first-generation nestling American kestrels (*Falco sparverius*). Environ Toxicol Chem 22(3):554-560.

Fernie K, Bortolotti G, Smits J. 2003b. Reproductive abnormalities, teratogenicity, and developmental problems in American kestrels (*Falco sparverius*) exposed to polychlorinated biphenyls. J toxicol Env Health Part A 66:2089-2103.

Fernie KJ, Bortolotti GR, Smits JE, Wilson J, Drouillard KG, Bird DM. 2000. Changes in egg composition of American kestrels exposed to dietary polychlorinated biphenyls. J Toxicol Environ Health Part A, 60:291-303.

Fernie KJ, Smits JE, Bortolotti GR, Bird DM. 2001. *In ovo* exposure to polychlorinated biphenyls: reproductive effects on second-generation American kestrels. Arch Environ Contam Toxicol 40:544-550.

Finley MT, Dieter MP, Locke LN. 1976. Sublethal effects of chronic lead ingestion in mallard ducks. J Toxicol Environ Health 1:929-937.

Fleming WJ, McLane MAR, Cromartie E. 1982. Endrin decreases screech owl productivity. J Wildl Manage 46(2):462-468.

Freeland JH, Cousins RJ. 1973. Effect of dietary cadmium on anemia, iron absorption, and cadmium binding protein in the chick. Nutr Rep Internat 8(5):337-347.

Gasaway WC, Buss IO. 1972. Zinc toxicity in the mallard duck. J Wildl Manage 36:1107-1117.

Gee JM, Craig-Veit CB, Millam JR. 2004. Posthatch methoxychlor exposure adversely affects reproduction of adult zebra finches, *Taeniopygia guttata*. Bull Environ Contam Toxicol 73:607-612.

Genelly RE, Rudd RL. 1956. Effects of DDT, toxaphene, and dieldrin on pheasant reproduction. Auk 73:529-539.

Graves JB, Bonner FL, McKnight WF, Watts AB, Epps EA. 1969. Residues in eggs, preening glands, liver, and muscle from feeding dieldrin-contaminated rice bran to hens and its effect on egg production, egg hatch, and chick survival. Bull Environ Contam Toxicol 4(6):375-383.

Hall JE, Greichus YA, Severson KE. 1971. Effects of aldrin on young pen-reared pheasants. J Wildl Manage 35(3):429-434.

Haseltine SD, Prouty RM. 1980. Aroclor 1242 and reproductive success of adult mallards (*Anas platyrhynchos*). Environ Res 23:29-34.

Haseltine SD, Sileo L, Hoffman DJ, Mulhern BD. unpublished. Effects of chromium on reproduction and growth in black ducks. As cited in Sample BE, Opresko DM, Suter GW. 1996. Toxicological benchmarks for wildlife. 1996 revision. ES/ERM-86/R3. Office of Environmental Management, US Department of Energy, Washington, DC.

Heath RG, Spann JW, Kreitzer JF. 1969. Marked DDE impairment of mallard reproduction in controlled studies. Nature 224:47-48.

Heath RG, Spann JW, Hill EF, Kreitzer JF. 1972. Comparative dietary toxicities of pesticides to birds. Wildlife no. 152. Patuxent Wildlife Research Center, Bureau of Sport Fisheries and Wildlife, Laurel, MD.

Heinz GH. 1980. Eggshell thickness in mallards fed methylmercury. Bull Environ Contam Toxicol 25:498-502.

Heinz GH, Hoffman DJ, Krynitsky AJ, Weller DMG. 1987. Reproduction in mallards fed selenium. Environ Toxicol Chem 6:423-433.

Heinz GH, Hoffman DJ, Gold LG. 1988. Toxicity of organic and inorganic selenium to mallard ducklings. Arch Environ Contam Toxicol 17:561-568.

Heinz GH, Hoffman DJ, Gold LG. 1989. Impaired reproduction of mallards fed an organic form of selenium. J Wildl Manage 53(2):418-428.

Hill EF, Soares JH, Jr. 1987. Oral and intramuscular toxicity of inorganic and organic mercury chloride to growing quail. J Toxicol Env Health 20:105-116.

Hill EF, Heath RG, Spann JW, Williams JD. 1975. Lethal dietary toxicities of environmental pollutants to birds. US Fish and Wildlife Service, Laurel, MD.

Hill EF, Heath RG, Williams JD. 1976. Effect of dieldrin and Aroclor 1242 on Japanese quail eggshell thickness. Bull Environ Contam Toxicol 16(4):445-453.

Hough JL, Baird MB, Sfeir GT, Pacini CS, Darrow D, Wheelock C. 1993. Benzo(a)pyrene enhances atherosclerosis in white carneau and show racer pigeons. Arterioscler Thromb 13:1721-1727.

Hudson RH, Tucker RK, Haegele MA. 1984. Handbook of toxicity of pesticides to wildlife. Resource publication 153. 2nd ed. US Fish and Wildlife Service, Washington, DC.

Ishida M, Suyama K, Adachi S, Hoshino T. 1982. Distribution of orally administered di-ethylhexyl phthalate in laying hens. Poult Sci 61:262-267.

Kolaja GJ. 1977. The effects of DDT, DDE and their sulfonated derivatives on eggshell formation in the mallard duck. Bull Environ Contam Toxicol 17:697-701.

Leach RM, Jr, Wang KW-L, Baker DE. 1979. Cadmium and the food chain: the effect of dietary cadmium on tissue composition in chicks and laying hens. J Nutr 109:437-443.

Lehner PN, Egbert A. 1969. Dieldrin and eggshell thickness in ducks. Nature 224:1218-1219.

Lepore PD, Miller RF. 1965. Embryonic viability as influenced by excess molybdenum in chicken breeder diets. Proc Soc Exper Biol Med 118:155-157.

Lien TF, Chen KL, Wu CP, Lu JJ. 2004. Effects of supplemental copper and chromium on the serum and egg traits of laying hens. Brit Poult Sci 45(4):535-539.

Longcore JR, Samson FB. 1973. Eggshell breakage by incubating black ducks fed DDE. J Wildl Manage 37(3):390-394.

Longcore JR, Samson FB, Whittendale Jr TW. 1971. DDE thins eggshells and lowers reproductive success of captive black ducks. Bull Environ Contam Toxicol 6(6):485-490.

Lowe TP, Stendell RC. 1991. Eggshell modifications in captive American kestrels resulting from Aroclor 1248 in the diet. Arch Environ Contam Toxicol 20:519-522.

Ludke JL. 1976. Organochlorine pesticide residues associated with survival: additivity of chlordane and endrin. Bull Environ Contam Toxicol 16:253-260.

McLane MAR, Hughes DL. 1980. Reproductive success of screech owls fed Aroclor 1248. Arch Environ Contam Toxicol 9:661-665.

Mehring AL, Brumbaugh JH, Sutherland AJ, Titus HW. 1960. The tolerance of growing chickens for dietary copper. Poult Sci 39:713-719.

Mendenhall VM, Klaas EE, McLane MAR. 1983. Breeding success of barn owls (*Tyto alba*) fed low levels of DDE and dieldrin. Arch Environ Contam Toxicol 12:235-240.

Millam JR, Craig-Veit CB, Batchelder ME, Viant MR, Herbeck TM, Woods LW. 2002. An avian bioassay for environmental estrogens: the growth response of zebra finch

(*Taeniopygia guttata*) chick oviduct to oral estrogens. Environ Toxicol Chem 21(12):2663-2668.

Morgan GW, Edens FW, Thaxton P, Parkhurst CR. 1975. Toxicity of dietary lead in Japanese quail. Poult Sci 54:1636-1642.

Nebeker AV, Griffis WL, Stutzman TW, Schuytema GS, Carey LA, Scherer SM. 1992. Effects of aqueous and dietary exposure of dieldrin on survival, growth and bioconcentration in mallard ducklings. Environ Toxicol Chem 11:687-699.

O'Shea TJ, Stafford CJ. 1980. Phthalate plasticizers: accumulation and effects on weight and food consumption in captive starlings. Bull Environ Contam Toxicol 25:345-352.

Oh SH, Nakaue H, Deagen JT, Whanger PD, Arscott GH. 1979. Accumulation and depletion of zinc in chick tissue metallothioneins. J Nutr 109: 1720-1729.

Ousterhout LE, Berg LR. 1981. Effects of diet composition on vanadium toxicity in laying hens. Poult Sci 60:1152-1159.

Pattee OH. 1984. Eggshell thickness and reproduction in American kestrels exposed to chronic dietary lead. Arch Environ Contam Toxicol 13:29-34.

Patton JF, Dieter MP. 1980. Effects of petroleum hydrocarbons on hepatic function in the duck. Comp Biochem Physiol 65C:33-36.

Peakall DB. 1974. Effects of di-n-butylphthalate and di-2-ethylhexylphthalate on the eggs of ring doves. Bull Environ Contam Toxicol 12:698-702.

Peakall DB, Lincer JL. 1972. Methyl mercury: its effect on eggshell thickness. Bull Environ Contam Toxicol 3(2):89-90.

Peakall DB, Peakall ML. 1973. Effect of a polychlorinated biphenyl on the reproduction of artificially and naturally incubated dove eggs. J Appl Ecol 10:863-868.

Peakall DB, Lincer JL, Bloom SE. 1972. Embryonic survival and chromosomal alterations caused by Aroclor 1254 in ring doves. Environ Health Perspect 1:103-104.

Peakall DB, Lincer JL, Risebrough RW, Pritchard JB, Kinter WB. 1973. DDE-induced eggshell thinning; structural and physiological effects in three species. Comp Gen Pharmac 4:305-313.

Persia ME, Baker DH, Parsons CM. 2004. Tolerance for excess basic zinc chloride and basic copper chloride in chicks. Brit Poult Sci 45(5):672-676.

Porter RD, Wiemeyer SN. 1972. DDE at low dietary levels kills captive American kestrels. Bull Environ Contam Toxicol 8(4):193-199.

Poupoulis C, Jensen LS. 1976. Effect of high dietary copper on gizzard integrity of the chick. Poult Sci 55:113-121.

Pritzl MC, Lie YH, Kienholz EW, Whiteman CE. 1974. The effect of dietary cadmium on development of young chickens. Poult Sci 53:2026-2029.

Richardson ME, Fox MRS, Bry BE, Jr. 1974. Pathological changes produced in Japanese quail by ingestion of cadmium. J Nutr 104:323-338.

Rigdon RH, Neal J. 1963. Fluorescence of chickens and eggs following the feeding of benzpyrene crystals. Tex Rep Biol Med 21(4):558-566.

Risebrough RW, Anderson DW. 1975. Some effects of DDE and PCB on mallards and their eggs. J Wildl Manage 39(3):508-513.

Roberson RH, Schaible PJ. 1960. The tolerance of growing chicks for high levels of different forms of zinc. Poult Sci 39:893-895.

Romoser GL, Dudley WA, Machlin LJ, Loveless L. 1961. Toxicity of vanadium and chromium for the growing chick. Poult Sci 40:1171-1173.

Roylance KJ, Jorgensen CD, Booth GM, Carter MW. 1985. Effects of dietary endrin on reproduction of mallard ducks. Arch Environ Contam Toxicol 14:705-711.

Scheuhammer AM. 1988. Chronic dietary toxicity of methylmercury in the zebra finch, *Poephila guttata*. Bull Environ Contam Toxicol 40:123-130.

Schlatterer B, Coenen TMM, Ebert E, Grau R, Hilbig V, Munk R. 1993. Effects of bis(tri-n-butyltin) oxide in Japanese quail exposed during egg laying period: an interlaboratory comparison study. Arch Environ Contam Toxicol 24:440-448.

Schwetz BA, Norris JM, Kociba RJ, Keeler PA, Cornier RF, Gehring PJ. 1974. Reproduction study in Japanese quail fed hexachlorobutadiene for 90 days. Toxicol Appl Pharmacol 30:255-265.

Shellenberger TE. 1978. A multi-generation toxicity evaluation of p,p'-DDT and dieldrin with Japanese quail. I. Effects of growth and reproduction. Drug Chem Toxicol 1(2):137-146.

Smith MS. 1969. Responses of chicks to dietary supplements of copper sulphate. Brit Poult Sci 10:97-108.

Spalding MG, Frederick PC, McGill HC, Bouton SN, McDowell LR. 2000. Methylmercury accumulation in tissues and its effects on growth and appetite in captive great egrets. J Wildl Dis 36(3):411-422.

Spann JW, Heinz GH, Camardese B, Hill EF, Moore JF, Murray HC. 1986a. Differences in survival among bobwhite fed methylmercury chloride dissolved in various carriers. Environ Toxicol Chem 5:721-724.

Spann JW, Heinz GH, Hulse CS. 1986b. Reproduction and health of mallards fed endrin. Environ Toxicol Chem 5:755-759.

Stahl JL, Greger JL, Cook ME. 1990. Breeding-hen and progeny performance when hens are fed excessive dietary zinc. Poult Sci 69:259-263.

Stanley TR, Jr, Spann JW, Smith GJ, Rosscoe R. 1994. Main and interactive effects of arsenic and selenium on mallard reproduction and duckling growth and survival. Arch Environ Contam Toxicol 26:441-451.

Stickel LF, Rhodes LI. 1970. The thin eggshell problem. In: Gillett JW, ed, Proceedings of the symposium, The Biological Impact of Pesticides in the Environment. Oregon State University, Corvallis, OR, pp 31-35.

Stoewsand GS, Anderson JA, Gutenmann WH, Bache CA, Lisk DJ. 1971. Eggshell thinning in Japanese quail fed mercuric chloride. Science 173:1030-1031.

USFWS. 1964. Pesticide-wildlife studies, 1963: a review of Fish and Wildlife Service investigations during the calendar year. Circular 199. Fish and Wildlife Service, US Department of the Interior, Washington, DC.

Vos JG, van der Maas HL, Musch A, Ram E. 1971. Toxicity of hexachlorobenzene in Japanese quail with special reference to porphyria, liver damage, reproduction, and tissue residues. Toxicol Appl Pharmacol 18:944-957.

Walker AIT, Neill CH, Stevenson DE, Robinson J. 1969. The toxicity of dieldrin (HEOD) to Japanese quail (*Coturnix coturnix japonica*). Toxicol Appl Pharmacol 15:69-73.

Weber CW, Reid BL. 1968. Nickel toxicity in growing chicks. J Anim Sci 95:612-616.

White DH, Dieter MP. 1978. Effects of dietary vanadium in mallard ducks. J Toxicol Environ Health 4:43-50.

White DH, Finley MT. 1978a. Effects of dietary cadmium in mallard ducks. In: Hemphill DD, ed, Trace substances in environmental health - XII. University of Missouri, Columbia, MO, pp 220-223.

White DH, Finley MT. 1978b. Uptake and retention of dietary cadmium in mallard ducks. Environ Res 17:53-59.

Wiemeyer SN, Hoffman DJ. 1996. Reproduction in eastern screech-owls fed selenium. J Wildl Manage 60(2):332-341.

Wiemeyer SN, Porter RD. 1970. DDE thins eggshells of captive American kestrels. Nature 227:737-738.

ATTACHMENT 10
ACCEPTABLE TOXICITY STUDIES FOR THE SELECTION OF MAMMAL DIETARY TOXICITY REFERENCE VALUES

Attachment 10 Acceptable Toxicity Studies for the Selection of Mammal Dietary Toxicity Reference Values

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
Metals									
Antimony	Sb_2O_3	rat	1,489	na	food	90 days	growth, survival	body weight	Hext et al. (1999)
Arsenic	Sodium arsenite-$NaAsO_2$	Osborne-Mendel rat	na	5.4	food	2 years	growth	female body weight	Byron et al. (1967)
Arsenic	Sodium arsenite-$NaAsO_2$	Osborne-Mendel rat	2.6	na	food	2 years	growth	female body weight	Byron et al. (1967)
Cadmium	$CdCl_2$	beagle dog	0.88	na	food	3 mos	growth	female body weight	Loeser and Lorke (1977b)
Cadmium	$CdCl_2$	Wistar rat	3.0	na	food	3 mos	survival, growth	adult survival, growth	Loeser and Lorke (1977a)
Cadmium	$CdCl_2$	Long-Evans rat	3.5	13	food	10 days (pregnancy)	growth	maternal body weight	Machemer and Lorke (1981)
Cadmium	$CdCl_2$	Long-Evans rat	13	na	food	10 days	survival, reproduction	adult survival; fertility, fetus weight/survival/ malformations	Machemer and Lorke (1981)
Cadmium	$CdCl_2$	shrew	115	na	food	12 wks	growth	female body weight	Dodds-Smith et al. (1992)
Cadmium	$CdCl_2$	shrew	na	115	food	12 wks	growth	male body weight	Dodds-Smith et al. (1992)
Cadmium	cadmium	Sprague-Dawley rat	na	189	food	12 wks	growth, reproduction	pup birth weight; adult body weight	Pond and Walker (1975)
Chromium	Cr picolinate	Sprague-Dawley rat	0.14	na	food	12 wks	growth	body weight and growth rate	Hasten et al. (1997)
Chromium	$CrCl_3$ or Cr picolinate	Sprague-Dawley rat	8.3	na	food	20 wks	growth, survival		Anderson et al. (1997)
Chromium	chromic oxide green	rat	1,292	na	food	90 days	growth, reproduction	body weight, litter size, pregnancy rate, pup malformation	Ivankovic and Preussman (1975)
Chromium	chromic oxide green	rat	1,466	na	food	2 yrs	growth, survival	body weight	Ivankovic and Preussman (1975)
Cobalt	cobaltous chloride	Sprague-Dawley rat	0.1[b]	1.0	diet	4 wks	growth	body weight	Chetty et al. (1979)

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
Cobalt	cobalt sulfate	guinea pigs	na	1.4	diet	5 wks	survival		Mohiuddin et al. (1970)
Cobalt	cobalt sulfate	guinea pigs	1.4	na	diet	5 wks	growth		Mohiuddin et al. (1970)
Cobalt	cobalt chloride	hooded rats	1.9	10	diet	3 days	growth	body weight	Wellman et al. (1984)
Copper	$CuSO_4$	mink	na	26	food	357 days	reproduction	kit survival, litter mass	Aulerich et al. (1982)
Copper	$CuSO_4$	mink	18	na	food	357 days	reproduction	kit survival, litter mass	Aulerich et al. (1982)
Copper	$CuSO_4$	mink	43	na	food	153-657 d	growth	body weight	Aulerich et al. (1982)
Copper	$CuSO_4*5H_2O$	rat	137	na	food	13 wks	survival		NTP (1993)
Copper	$CuSO_4*5H_2O$	rat	67	137	food	13 wks	growth	body weight	NTP (1993)
Copper	$CuSO_4*5H_2O$	rat	93	197	food	2 wks	growth	body weight	NTP (1993)
Copper	copper chloride	shrew	267	na	food	weanlings for 12 wks	survival, growth	body weight	Dodds-Smith et al. (1992)
Copper	$CuSO_4*5H_2O$	rat	305	na	food	2 wks	survival		NTP (1993)
Copper	$CuSO_4*5H_2O$	mouse	467	na	food	13 wks	survival		NTP (1993)
Copper	$CuSO_4*5H_2O$	mouse	227	467	food	13 wks	growth	body weight	NTP (1993)
Copper	$CuSO_4*5H_2O$	mouse	749	na	food	2 wks	survival, growth		NTP (1993)
Lead	Lead acetate	rat	11	90	food	2 years	growth	Offspring weight and kidney damage	Azar et al. (1973)
Mercury	Methylmercuric chloride	Wistar rat	0.0017[a]	0.0084	food	3 gen	growth		Verschuuren et al. (1976)
Mercury	Methylmercuric chloride	Wistar rat	0.19	na	food	3 gen	survival, reproduction		Verschuuren et al. (1976)
Mercury	Methylmercuric chloride	mink	0.16	0.25	food	93 days	growth, survival		Wobeser et al. (1976)
Mercury	methyl-mercury	mink	na	0.64	food	2 months	growth, survival		Aulerich et al. (1974)
Molybdenum	soluble arsenite	mouse	na	2.58	food and drink water	3 generation	reproduction, survival	decreased litter size	Schroeder and Mitchener (1971)
Nickel	NiCl2	Sprague-Dawley rat	20	na	food	14 days + 61 days untreated	growth	body weight	Nation et al. (1985)
Nickel	Ni	Wistar rat	na	20	food	3 generations	reproduction	increased number of stillborns in F1 generation	Ambrose et al. (1976)
Nickel	$Ni(SO)_4$	Wistar rat	na	87	food	2 yrs	growth	body weight	Ambrose et al. (1976)

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
Nickel	$Ni(SO)_4$	Wistar rat	8.4	na	food	2 yrs	growth	body weight	Ambrose et al. (1976)
Nickel	nickel acetate	Webster Swiss mouse	na	169	food	4 wks	growth	body weight	Weber and Reid (1969)
Nickel	nickel acetate	Webster Swiss mouse	210	na	food	weanling thru reprod	reproduction	number of pups weaned	Weber and Reid (1969)
Nickel	$Ni(SO)_4$	Wistar rat	230	na	food	2 yrs	survival		Ambrose et al. (1976)
Nickel	nickel acetate	Webster Swiss mouse	na	313	food	weanling thru reprod	reproduction	number of pups weaned	Weber and Reid (1969)
Nickel	$Ni(SO)_4$	beagle dog	2500	na	food	2 yrs	survival		Ambrose et al. (1976)
Selenium	$NaSeO_3$ or seleniferous wheat	Sprague-Dawley rat	na	0.080	food	6 wks	growth	body weight	Halverson et al. (1966)
Selenium	$NaSeO_3$ or seleniferous wheat	Sprague-Dawley rat	0.055	na	food	6 wks	growth	body weight	Halverson et al. (1966)
Selenium	$NaSeO_3$ or seleniferous wheat	Sprague-Dawley rat	na	0.14	food	6 wks	survival	survival	Halverson et al. (1966)
Selenium	$NaSeO_3$ or seleniferous wheat	Sprague-Dawley rat	0.13	na	food	6 wks	survival	survival	Halverson et al. (1966)
Selenium	L-selenomethionine	Wistar rat	na	0.16	food	110 days	growth	body weight	Behne et al. (1992)
Selenium	selenite	Wistar rat	0.16	na	food	110 days	growth	body weight	Behne et al. (1992)
Selenium	Na_2SeO_3, Nano-Se, or organic selenium	Sprague-Dawley rat	0.17	0.28	food	13 wks	growth	body weight	Jia et al. (2005)
Selenium	selenomethionine	hamster	0.36	na	food	21d	growth	body weight	Julius et al. (1983)
Selenium	selenomethionine	hamster	na	0.76	food	21d	growth	body weight	Julius et al. (1983)
Selenium	Na_2SeO_3	hamster	na	3.4	food	21d	growth	body weight	Julius et al. (1983)
Selenium	Na_2SeO_3	hamster	na	5.8	food	21 d	survival	female survival	Julius et al. (1983)

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
Thallium	thallium sulfate	rat	0.74	na	drink water	60 d	growth		Formigli et al. (1986)
Vanadium	vanadium sulfate	mouse	1.05	na	food and drink water	lifetime (1 yr)	growth	body weight	Schroeder and Balassa (1967)
Vanadium	sodium meta-vanadate	Wistar rat	na	2.7	food	10 weeks	growth	body weight	Adachi et al. (2000)
Vanadium	sodium meta-vanadate	Sprague-Dawley rat	na	6.5	food	reproduction period	growth, reproduction	maternal body weight, offspring body weight gain and survival	Elfant and Keen (1987)
Zinc	zinc oxide	Sprague-Dawley rat	160	320	food	gestation	reproduction	fetal growth, number of resorptions	Schlicker and Cox (1968)
Zinc	zinc oxide	ferret	149	433	food	2wks-6 mos	growth		Straube et al. (1980)
Zinc	zinc carbonate	rat	400	799	food	gestation	growth	body weight	Sutton and Nelson (1937)
Organometals									
TBT	tributyltin chloride	Wistar rat	0.4	2.0	food	multi-generational	reproduction	pup body weight	Omura et al. (2001)
TBT	tributyltin chloride	Wistar rat	na	2.0	food	reproduction period	reproduction	pup body weight	Makita et al. (2003); Makita et al. (2004)
TBT	bis (tri-n-butyltin) oxide	Wistar rat	0.21	2.1	food	106 wks	growth, survival	body weight (male), survival	Wester et al. (1990)
TBT	tributyltin chloride	Wistar rat	2.0	10	food	multi-generational	reproduction	pup body weight, percentage of live pups	Ogata et al. (2001)
DBT	dibutyltin dichloride	Wistar rat	3.8	7.6	gavage	3 days during pregnancy	growth	maternal body weight	Harazono and Ema (2003)
DBT	dibutyltin dichloride	Wistar rat	1.5	7.6	gavage	3 days during pregnancy	reproduction, growth	reduced no. implantations, no. of females pregnant, increased rate of postimplantation loss, reduced maternal body weight	Ema et al. (2003)
PAHs									
Benzo(a)pyrene	benzo(a)pyrene	mouse	2.0[a]	10	gavage	gestation (10 days)	reproduction	pup body weight, testes weight	MacKenzie and Angevine (1981)
Benzo(a)pyrene	benzo(a) pyrene	mouse	33.3	na	food	up to 115 days	survival	survival	Neal and Rigdon (1967)
Naphthalene	Naphthalene	mouse	133	na	gavage	90 d	survival, growth	body weight, adult survival	Shopp et al. (1984)
Naphthalene	Naphthalene	mouse	na	300	gavage	8 days of pregnancy	survival, reproduction	maternal survival; litter size	Plasterer et al. (1985)

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
1-methylnaphthalene	1-methylnapthalene	B6C3F1 mouse	150	na	food	81 weeks	growth	body weight	Murata et al. (1993)
2-methylnaphthalene	2-methylnapthalene	mouse	54	114	food	81 wks	growth	body weight	Murata et al. (1997)
Phthalates									
Bis (2-ethylhexyl) phthalate	BEHP: di(2-ethylhexyl) phthalate	mouse, CD-1	44	na	food	17 days (pregnancy)	reproduction	fetal abnormalities increased	Tyl et al. (1988)
Bis (2-ethylhexyl) phthalate	BEHP: di(2-ethylhexyl) phthalate	mouse, CD-1	na	91	food	17 days (pregnancy)	reproduction	fetal abnormalities increased	Tyl et al. (1988)
Bis (2-ethylhexyl) phthalate	BEHP: di(2-ethylhexyl) phthalate	guinea pig	93	na	food	1 yr	survival, growth	life expectancy, body weight	Carpenter et al. (1953)
Bis (2-ethylhexyl) phthalate	BEHP: di(2-ethylhexyl) phthalate	mouse, COBS	14	136	food	>24 wks (2 generations)	reproduction	number of litters, viability	Lamb et al. (1987)
Bis (2-ethylhexyl) phthalate	BEHP: di(2-ethylhexyl) phthalate	mouse, ICR-JCL	na	190	food	18 days (pregnancy)	reproduction	fetal death	Shiota et al. (1980)
Bis (2-ethylhexyl) phthalate	BEHP: di(2-ethylhexyl) phthalate	mouse, ICR-JCL	70	na	food	18 days (pregnancy)	reproduction	fetal death	Shiota et al. (1980)
Bis (2-ethylhexyl) phthalate	BEHP: di(2-ethylhexyl) phthalate	mouse, CD-1	na	191	food	17 days (pregnancy)	growth	maternal body weight	Tyl et al. (1988)
Bis (2-ethylhexyl) phthalate	BEHP: di(2-ethylhexyl) phthalate	mouse, CD-1	91	na	food	17 days (pregnancy)	growth	maternal body weight	Tyl et al. (1988)
Butyl benzyl phthalate	BBP: butyl benzyl phthalate	rat	250	750	food	multi-generational (2 generations)	growth, reproduction	body weight, reduced no. live pups/litter and no. implantations, reduced offspring body weight, delayed onset of puberty	Tyl et al. (2004)
Butyl benzyl phthalate	BBP: butyl benzyl phthalate	rat, Fisher 344	831	na	food	14 days	growth	male body weight	Agarwal et al. (1985)
Butyl benzyl phthalate	BBP: butyl benzyl phthalate	Wistar rat	na	845	food	11 days (pregnancy)	reproduction	maternal weight; post-implantation embryo loss	Ema et al. (1994)

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
Butyl benzyl phthalate	BBP: butyl benzyl phthalate	Wistar rat	991	na	food	20 days (pregnancy)	survival	maternal mortalities	Ema et al. (1992a; 1992b)
Butyl benzyl phthalate	BBP: butyl benzyl phthalate	Wistar rat	na	991	food	20 days (pregnancy)	reproduction	fetal death, fetal weight, malformations and resorptions	Ema et al. (1992a)
Butyl benzyl phthalate	BBP: butyl benzyl phthalate	rat, Fisher 344	na	1325	food	14 days	growth	male body weight	Agarwal et al. (1985)
Butyl benzyl phthalate	BBP: butyl benzyl phthalate	rat, Fisher 344	1570	na	food	14 days	survival	male	Agarwal et al. (1985)
Dibutyl phthalate	DBP: di-n-butyl phthalate	rat, Sprague-Dawley	16[a]	80	food	~24 wks (2 gen)	reproduction	fertility, pup weight, pup viability, mating	Wine et al. (1997)
Dibutyl phthalate	DBP: di-n-butyl phthalate	rat, Sprague-Dawley	641	na	food	~24 wks (2 gen)	survival		Wine et al. (1997)
Dibutyl phthalate	DBP: di-n-butyl phthalate	rat, Sprague-Dawley	320	641	food	~24 wks (2 gen)	growth	body weight	Wine et al. (1997)
Dibutyl phthalate	DBP: di-n-butyl phthalate	mouse, ICR-JCL	660	na	food	18 days (pregnancy)	growth	maternal weight	Shiota et al. (1980)
Dibutyl phthalate	DBP: di-n-butyl phthalate	mouse, ICR-JCL	na	660	food	18 days (pregnancy)	reproduction	decrease in male fetal weight	Shiota et al. (1980)
Dibutyl phthalate	DBP: di-n-butyl phthalate	mouse, ICR-JCL	350	na	food	18 days (pregnancy)	reproduction	decrease in male fetal weight	Shiota et al. (1980)
Dibutyl phthalate	DBP: di-n-butyl phthalate	rat, Sprague-Dawley	200	999	food	1 yr	survival		Smith (1953)
Diethyl phthalate	DEP: diethyl phthalate	mouse, COBS	3721	na	food	>24 wks (2 gen)	survival		Lamb et al. (1987)
Diethyl phthalate	DEP: diethyl phthalate	mouse, COBS	1860	3721	food	>24 wks (2 gen)	growth, reproduction	body weight; number of live pups per litter in F1 generation	Lamb et al. (1987)
Di-n-octyl phthalate	DNOP: di-n-octyl phthalate	rat, Sprague-Dawley	377	na	food	13 wks	growth, survival	body weight	Poon et al. (1997)
Di-n-octyl phthalate	DNOP: di-n-octyl phthalate	mouse, CD-1	7500	na	food	~ 26 wks	survival, growth, reproduction	survival; body weight; fertility, number of litters, viability, sex ratio, pup weights	Heindel et al. (1989)
Other SVOCs									
1,4-dichlorobenzene	1,4-DCB	Wistar rat	3.5	na	food	42 days	growth, reproduction	offspring body weight, average litter size, sex ratio, pup viability	Makita (2005)

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
1,4-dichlorobenzene	1,4-DCB	rat	54	107	gavage	1,4 or 13 wks	growth	body weight	Lake et al. (1997)
1,4-dichlorobenzene	1,4-DCB	mouse	429	na	gavage	1,4 or 13 wks	growth	body weight	Lake et al. (1997)
Dibenzothiophene	dibenzothio-pene	mouse	47[b]	470	gavage	<72 hrs	survival	LD50	Leighton (Leighton 1989)
Benzoic acid	Sodium benzoate	hamster	300	na	diet		survival, reproduction	no maternal toxicity or fetal toxicity	FDRL 1972 (as cited in IRIS (EPA 2006))
Benzoic acid		rat	80	na	diet	18 months	survival, growth	body weight, survival	Ignat'ev 1965 (as cited in IRIS (EPA 2006))
Benzoic acid		rat	50	750	diet	long-term	growth	body weight	Marquardt 1980 (as cited in IRIS (EPA 2006))
Phenol		rat	60	120	gavage	Gestational days 6 - 15	growth	maternal body weight	Argus Research Laboratories 1997 (as cited in IRIS (EPA 2006))
Phenol		rat	60	120	gavage		reproduction	decreased fetal body weight	Charles River Laboratories 1988 and NTP 1983a (as cited in IRIS (EPA 2006))
Phenol		mouse	140	280	gavage	Gestational days 6 - 15	survival, growth	survival and reduced body weight	NTP 1983b (as cited in IRIS (EPA 2006))
Biphenyl		rat	50	250	diet	3-generation	survival	longevity	Ambrose et al. 1960 (as cited in IRIS (EPA 2006))
PCBs									
PCBs (Clophen A50)	Clophen A50	mink	0.018[a]	0.089	food	18 months	reproduction	kit growth	Brunström et al. (2001)
PCBs (Aroclor 1254)		mink	na	0.13	food	6 mos	reproduction	reduced kit growth rate	Wren et al. (1987)
PCBs (Aroclor 1254)		mink	na	0.22	food	4 and 9 months prior to giving birth	reproduction	number of offspring per female, decrease in pup body weight	Ringer (1983)
PCBs (Aroclor 1254)		mink	0.13	0.26	food	4 months	reproduction	Number of kits born alive (0% at 4 wks)	Aulerich and Ringer (1977)
PCBs (Clophen A50)	Clophen A50	mink	0.27	na	food	18 months	growth	maternal body weight	Brunström et al. (2001)
PCBs (Aroclor 1254)		mink	na	0.39	food	88-102 days	reproduction	number of kits whelped and born alive (0%)	Aulerich et al. (1985)

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
PCB (mixture composition not reported)		mink	na	0.51	food	66 days	reproduction	number of kits born alive	Jensen et al. (1977)
PCBs (Aroclor 1242)		mink	na	0.65	food	8 months	reproduction	reproductive failure	Bleavins et al. (1980)
PCBs (Aroclor 1254)		mink	na	1.31	food	4 wks	weight gain in adults	weight gain in adults	Hornshaw et al. (1986)
PCBs (Aroclor 1254)		mink	na	1.64	food	3 months	reproduction	all whelps stillborn	Kihlstrom et al. (1992)
PCBs (Aroclor 1254)		mink	1.2	1.8	food	28 days	growth	female growth	Aulerich et al. (1986)
PCBs (Clophen A50)		mink	na	2.0	food	3 months	reproduction	all whelps stillborn	Kihlstrom et al. (1992)
PCBs (Aroclor 1254)		mink	1.5	2.4	food	28 days	growth	male and female growth	Aulerich et al. (1986)
PCBs (Aroclor 1016)		mink	na	2.6	food	8 months	reproduction/survival	birth weight and growth rate of kits, and 25 % adult female survival	Bleavins et al. (1980)
Organochlorine Pesticides									
Aldrin		Osborne-Mendel rat	0.8	4.1	diet	2 years	survival	increased survival	Reuber (1980)
beta-hexachloro cyclohexane (beta-BHC)	beta-HCH	Wistar rat	5.7	na	food	13 wks	growth, survival	body weight, survival	Van Velsen et al. (1986)
beta-hexachloro cyclohexane (beta-BHC)	beta-HCH	Wistar rat	na	31	food	13 wks	growth, survival	body weight, 50% survival	Van Velsen et al. (1986)
beta-hexachloro cyclohexane (beta-BHC)	beta-HCH	Wistar rat	64	na	food	2 wks	growth	body weight	Srinivasan et al. (1991)
Chlordane	tech chlordane	mouse	0.18	0.92	food	104 wk	growth	body weight-males	Khasawinah and Grutsch (1989)
Chlordane	tech chlordane	mouse	2.3	na	food	104 wk	survival		Khasawinah and Grutsch (1989)
Chlordane	tech chlordane	Osborne-Mendel rat	na	8.0	food	104 wk	growth, survival, reproduction	body weight, adult survival, viability of offspring	Ingle (1952)
Chlordane	tech chlordane	Osborne-Mendel rat	1.5	na	food	104 wk	growth, survival, reproduction	body weight, adult survival, viability of offspring	Ingle (1952)
Chlordane	tech chlordane	albino rat	na	11	food	407 d	growth	body weight-males	Ambrose et al. (1953)

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
Chlordane	tech chlordane	albino rat	4.9	na	food	407 d	growth	body weight-males	Ambrose et al. (1953)
Chlordane	tech chlordane	albino rat	11	22	food	407 d	survival		Ambrose et al. (1953)
trans-Nonachlor	trans-nonachlor	Sprague-Dawley rat	2.5	25	gavage	28 days	growth, survival	body weight- female	Bondy et al. (2000)
Chlordane	tech chlordane	Sprague-Dawley rat	25	na	food	28 days	growth	body weight	Bondy et al. (2000)
Chlordane	tech chlordane	Sprague-Dawley rat	40	na	food	reproduction period	growth	body weight	Cassidy et al. (1994)
DDT	o,p'-DDT	Sprague-Dawley rat	0.24	na	food	2 generation	reproduction	litter size and weight and uterine involution	Duby et al. (1971)
DDT	p,p'-DDT	mouse	0.6	na	food	5 generation	survival, growth, reproduction	adult survival, growth, number of pregnancies, number of births, litter size, pup growth/survival	Tarjan and Kemeny (1969)
DDT	p,p'- DDT	Sprague-Dawley rat	1.0	na	food	2 generation	reproduction	litter size and weight and uterine involution	Duby et al. (1971)
Total DDT	tech DDT	Sprague-Dawley rat	1.2	na	food	2 generation	reproduction	litter size and weight and uterine involution	Duby et al. (1971)
Total DDT	"DDT"- chem form not specified, likely DDT mixture	mouse	na	1.3	food	120 days	reproduction	litter size	Ware and Good (1967)
Total DDT	"DDT"- chem form not specified, likely DDT mixture	mouse	1.3	na	food	120 days	survival		Ware and Good (1967)
Total DDT	tech DDT	Sprague-Dawley rat	1.6	na	food	23 mos	growth, survival, reproduction	adult survival, growth, viable litter size, reproductive life-span	Ottoboni (1972)
Total DDT	tech DDT	Sprague-Dawley rat	na	2.0	food	7.5 wks	reproduction	fertility	Nickerson and Sniffen (1973)
Total DDT	"DDT"- chem form not specified, likely DDT mixture	rat	0.8	4.0	food	2 yrs	reproduction	number of young surviving to weaning (63% vs. 87% lower dose and 88% in control)	Fitzhugh (1948)
Total DDT	tech DDT	Sprague-Dawley rat	na	13.4	food	36 wks	reproduction	litter size, mating and reproductive success	Jonsson et al. (1976)
DDT	p,p'-DDT	Wistar rat	1.6	16	food	6 mos	reproduction	offspring growth	Clement and Okey (1974)

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
Total DDT	"DDT"- chem form not specified, likely DDT mixture	white lab mouse	na	37	food	F0 mating, gestation and weaning+F1 breeding	survival		Cannon and Holcomb (1968)
Total DDT	tech DDT	Oldfield mouse	2.4	na	food	15 mos	survival, reproduction	adult survival, litter size, litters per pair	Wolfe et al. (1979)
DDT	o,p'-DDT	Wistar rat	4.0	na	food	18-23 wks	reproduction	offspring survival, fertility, fecundy, growth	Wrenn et al. (1971)
Total DDT	tech DDT	Sprague-Dawley rat	6.7	na	food	36 wks	reproduction	litter size, mating and reproductive success	Jonsson et al. (1976)
Total DDT	tech DDT	mouse	9.2	46	food	6 generation	survival, reproduction	lifespan, pup survival	Turusov et al. (1973)
Total DDT	tech DDT	Sprague-Dawley rat	13	na	food	37 wks	survival		Jonsson et al. (1976)
Total DDT	tech DDT (DDD, DDE, DDT)	Sprague-Dawley rat	16	na	food	3 generation	survival, growth, reproduction	adult survival, growth, fertility, viability, stillbirths, litter size, abnormalities, pup survival	Ottoboni (1969)
DDT	p,p'-DDT	mouse	18	na	food	2 yrs	survival		Thorpe and Walker (1973)
Total DDT	pp'-DDT, '-DDD, -DDE	Wistar rat	21	na	food	6 wks	growth, survival	survival, body weight (males only)	Banerjee et al. (1996)
Dieldrin	dieldrin	Carworth rat	0.038[a]	0.19	food	3 generations	reproduction	offspring survival	Treon and Cleveland (1955)
Dieldrin	dieldrin	CFE rat	0.50	na	food	2 yrs	survival, growth		Walker et al. (1969)
Dieldrin	tech dieldrin	CFS Swiss mouse	0.92	na	food	120 d	survival		Good and Ware (1969)
Dieldrin	tech dieldrin	CFS Swiss mouse	na	0.92	food	120 d	reproduction	litter size	Good and Ware (1969)
Dieldrin	dieldrin	mouse	na	0.92	food	4 wks prior to mating through gestation	reproduction	18% pup survival	Virgo and Bellward (1977)
Dieldrin	dieldrin	Carworth rat	2.2	na	food	6 mos	growth		Treon and Cleveland (1955)
Dieldrin	dieldrin	Osborne-Mendel rat	0.8	4.1	food	2 yrs	survival		Fitzhugh et al. (1964)
Dieldrin		Osborne-Mendel rat	0.8	4.1	diet	2 years	survival	increased survival	Reuber (1980)
Dieldrin	dieldrin	Carworth rat	6.6	na	food	6 mos	survival		Treon and Cleveland (1955)

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
Dieldrin	dieldrin	Osborne-Mendel rat	12	na	food	2 yrs	growth	body weight	Fitzhugh et al. (1964)
Dieldrin	dieldrin	Carworth rat	na	26	food	6 mos	survival		Treon and Cleveland (1955)
Dieldrin	dieldrin	Carworth rat	1.9	na	food	2 yrs	survival		Treon and Cleveland (1955)
Endosulfan	technical endosulfan	mouse	0.84	2.5	food	24 mos	survival, growth	male survival (males more sensitive than females), body weight	Hack et al. (1995)
Endosulfan	technical endosulfan	rat	0.65	3.3	food	104 wks	growth		Hack et al. (1995)
Endosulfan	technical endosulfan	rat	3.5	na	food	104 wks	survival		Hack et al. (1995)
Endrin	endrin	Swiss mouse	na	0.92	food	120 days	survival, reproduction	adult survival, litter size, number of young/day	Good and Ware (1969)
Endrin	endrin	deer mouse	1.7	na	food	68 days over 7 mos	reproduction	litter survival, 3 weaned, breeding pair availability	Morris (1968)
Endrin	endrin	deer mouse	0.97	1.7	food	68 days over 7 mos	survival		Morris (1968)
Endrin	endrin	Carworth rat	0.40	2.0	food	106 wks	survival, growth	female survival; male body weight	Treon et al. (1955)
gamma-Hexachloro-cyclohexane	gamma HCH	NZ white rabbit	0.80	na	gavage	12-15wks; 3/wk	reproduction	fertilization rate, implant, embryo loss	Seiler et al. (1994)
gamma-Hexachloro-cyclohexane	gamma HCH	rat	1.00	na	gavage	6 days lactation	survival, growth	maternal survival; maternal growth	Dalsenter et al. (1997a)
gamma-Hexachloro-cyclohexane	gamma HCH	CD strain rat	6.1	na	food	3 generations	survival, growth, reproduction	adult survival, body weight, pregnancy rate, gestation period	Palmer et al. (1978)
gamma-Hexachloro-cyclohexane	gamma HCH	beagle dog	9.3	na	food	32 wks	survival, growth		Rivett et al. (1978)
gamma-Hexachloro-cyclohexane	gamma HCH	Wistar rat	10	na	gavage	7 days 1st or 2nd wk postnatal	growth	body weight until age 29 days	Rivera et al. (1990)
gamma-Hexachloro-cyclohexane	gamma HCH	Wistar rat	30	na	gavage	singe dose at day 15 post-conception	reproduction	litter size, offspring survival, offspring body weight, fertility	Dalsenter et al. (1997b)
gamma-Hexachloro-cyclohexane	gamma HCH	Wistar rat	64	na	food	2 wks	growth		Srinivasan et al. (1991)
Heptachlor	tech heptachlor	mink	1.0	na	food	181 days	survival, growth, reproduction	adult survival, female body weight, kit body weight	Crum et al. (1993)
Heptachlor	tech heptachlor	mink	na	1.8	food	181 days	survival, growth, reproduction	adult survival, female body weight, kit body weight	Crum et al. (1993)

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
Heptachlor	tech heptachlor	mink	3.1	na	food	28 days	growth	body weight	Aulerich et al. (1990)
Heptachlor	tech heptachlor	mink	na	5.7	food	28 days	growth	body weight	Aulerich et al. (1990)
Heptachlor	tech heptachlor	mink	5.7	na	food	28 days	survival		Aulerich et al. (1990)
Heptachlor	tech heptachlor	mink	na	6.2	food	28 days	survival		Aulerich et al. (1990)
Hexachlorobenzene	HCB	mink	0.026[a]	0.13	food	331 days	reproduction	birth weight	Bleavins et al. (1984)
Hexachlorobenzene	HCB	ferret	0.026[a]	0.13	food	332 days	reproduction	birth weight	Bleavins et al. (1984)
Hexachlorobenzene	HCB	Sprague-Dawley rat	na	0.80	food	4 generation	reproduction	weanling weight	Grant et al. (1977)
Hexachlorobenzene	HCB	Wistar rat	1.28	na	food	>120 d	reproduction, behavior	number of pups and weights	Lilienthal et al. (1996)
Hexachlorobenzene	HCB	rat	3.2	na	food	2 generation	survival, growth		Arnold et al. (1985)
Hexachlorobenzene	HCB	rat	0.64	3.2	food	2 generation	reproduction	decreased viability	Arnold et al. (1985)
Hexachlorobenzene	HCB	Sprague-Dawley rat	na	4.8	food	~200 d days	reproduction	pup survival	Kitchin et al. (1982)
Hexachlorobenzene	HCB	Sprague-Dawley rat	11	na	food	> 100 days	survival, growth		Kitchin et al. (1982)
Hexachlorobenzene	HCB	ferret	3.2	16	food	332 days	survival		Bleavins et al. (1984)
Hexachlorobenzene	HCB	mink	3.3	16	food	331 days	survival		Bleavins et al. (1984)
Hexachlorobenzene	HCB	Sprague-Dawley rat	13	26	food	4 generation	survival		Grant et al. (1977)
Methoxychlor	methoxychlor	Sprague-Dawley rat	na	56	food	gestation day 0 to postnatal day 22 (adult); postnatal day 28 to 100 (young)	growth, reproduction	body weight; offspring growth rate, litter size, sex development of offspring	You et al. (2002)
Methoxychlor	methoxychlor	Sprague-Dawley rat	17	86	food	gestation day 15, postnatal day 10	reproduction, growth	maternal body weight, offspring body weight, delayed onset of puberty	Masutomi et al. (2003)

Analyte	Chemical Form	Test Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Exposure Route	Exposure Duration	Endpoint	Effect Endpoint	Source
Methoxychlor	tech methoxychlor	Sprague-Dawley rat	na	80	food	parents treated, weanlings treated 8 wks thru mating	reproduction	% mated, % littered	Harris et al. (1974)
Methoxychlor	tech methoxychlor	Sprague-Dawley rat	na	168	food	pre-mating thru weaning	reproduction	% mated, litter production and size	Harris et al. (1974)

[a] NOAEL estimated using an uncertainty factor of 5 (chronic LOAEL to chronic NOAEL)

[b] NOAEL estimated using an uncertainty factor of 10 (acute/subchronic LOAEL to chronic NOAEL)

LOAEL – lowest-observed-effect concentration

NOAEL – no-observed-effect concentration

PAH – polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

REFERENCES

Adachi A, Asai K, Koyama Y, Matsumoto Y, Okano T. 2000. Subacute vanadium toxicity in rats. J Health Sci 46(6):503-508.

Agarwal DK, Maronpot RR, Lamb IV JC, Kluwe WM. 1985. Adverse effects of butyl benzyl phthalate on the reproductive and hematopoietic systems of male rats. Toxicology 35:189-206.

Ambrose AM, Christensen HE, Robbins DJ, Rather LJ. 1953. Toxicological and pharmacological studies on chlordane. Arch Indus Hyg Occup Med 7:197-210.

Ambrose AM, Larson PS, Borzelleca JF, Hennigar Jr GR. 1976. Long term toxicologic assessment of nickel in rats and dogs. J Food Sci Technol 13(4):181-187.

Anderson RA, Bryden NA, Polansky MM. 1997. Lack of toxicity of chromium chloride and chromium picolinate in rats. J Am Coll Nutr 16(3):273-279.

Arnold DL, Moodie CA, Charbonneau SM, Grice HC, McGuire PF, Bryce FR, Collins BT, Zawidzka ZZ, Krewski DR, Nera EA, Munro IC. 1985. Long-term toxicity of hexachlorobenzene in the rat and the effect of dietary vitamin A. Food Chem Toxicol 23(9):779-793.

Aulerich RJ, Ringer RK. 1977. Current status of PCB toxicity to mink, and effect on their reproduction. Arch Environ Contam Toxicol 6:279-292.

Aulerich RJ, Ringer RK, Iwamoto S. 1974. Effects of dietary mercury on mink. Arch Environ Contam Toxicol 2(1):43-51.

Aulerich RJ, Ringer RK, Bleavins MR, Napolitano A. 1982. Effects of supplemental dietary copper on growth, reproductive performance and kit survival of standard dark mink and the acute toxicity of copper to mink. J Anim Sci 55(2):337-343.

Aulerich RJ, Bursian SJ, Breslin WJ, Olson BA, Ringer RK. 1985. Toxicological manifestations of 2,4,5,2',4',5'-, 2,3,6,2',3',6'-, and 3,4,5,3',4',5'- hexachlorobiphenyl and Aroclor 1254 in mink. J Toxicol Environ Health 5:63-79.

Aulerich RJ, Ringer RK, Safronoff J. 1986. Assessment of primary vs secondary toxicity of Aroclor 1254 to mink. Arch Environ Contam Toxicol 15:393-399.

Aulerich RJ, Bursian SJ, Napolitano A. 1990. Subacute toxicity of dietary heptachlor to mink (*Mustela vison*). Arch Environ Contam Toxicol 19:913-916.

Azar A, Trochimowicz HJ, Maxfield ME. 1973. Review of lead studies in animals carried out at Haskell Laboratory-two-year feeding study and response to hemorrhage study. In: Barth D, ed, Environmental Health Aspects of Lead, International Symposium, Amsterdam. Commission of European Communities, pp 199-210.

Banerjee BD, Ray A, Pasha ST. 1996. A comparative evaluation of immunotoxicity of DDT and its metabolites in rats. Ind J Exper Biol 34:517-522.

Behne D, Kyriakopoulos A, Gessner H, Walzog B, Meinhold H. 1992. Type I iodothyronine deiodinase activity after high selenium intake, and relations between selenium and iodine metabolism in rats. J Nutr 122(7):1542-1546.

Bleavins MR, Aulerich RJ, Ringer RK. 1980. Polychlorinated biphenyls (Aroclors 1016 and 1242): Effects on survival and reproduction in mink and ferrets. Arch Environ Contam Toxicol 9:627-635.

Bleavins MR, Aulerich RJ, Ringer RK. 1984. Effects of chronic dietary hexachlorobenzene exposure on the reproductive performance and survivability of mink and European ferrets. Arch Environ Contam Toxicol 13:357-365.

Bondy GS, Newsome WH, Armstrong CL, Suzuki CAM, Doucet J, Fernie S, Hierlihy SL, Feeley MM, Barker MG. 2000. Trans-nonachlor and cis-nonachlor toxicity in Sprague-Dawley rats: comparison with technical chlordane. Toxicol Sci 58:386-398.

Brunström B, Lund BE, Bergman A, Asplund L, Athanassiadis I, Athanasiadou M, Jensen S, Örberg J. 2001. Reproductive toxicity in mink (*Mustela vison*) chronically exposed to environmentally relevant polychlorinated biphenyl concentrations. Environ Toxicol Chem 20(10):2318-2327.

Byron WR, Bierbower GW, Brouwer JB, Hansen WH. 1967. Pathologic changes in rats and dogs from two-year feeding of sodium arsenite or sodium arsenate. Toxicol Appl Pharmacol 10:132-147.

Cannon MS, Holcomb LC. 1968. The effect of DDT on reproduction in mice. Ohio J Sci 68(1):19-24.

Carpenter CP, Weil CS, Smyth Jr HF. 1953. Chronic oral toxicity of di(2-ethylhexyl) phthalate for rats, guinea pigs, and dogs. Arch Indus Hyg Occup Med 8:219-227.

Cassidy RA, Vorhees CV, Minnema DJ, Hastings L. 1994. The effects of chlordane exposure during pre- and postnatal periods at environmentally relevant levels on sex steroid-mediated behaviors and functions in the rat. Toxicol Appl Pharmacol 126:326-337.

Chetty KY, Rau S, Drummond L, Desaiah D. 1979. Cobalt induced changes in immune response and adenosine triphosphatase activities in rats. J Environ Sci Health B14(5):525-544.

Clement JG, Okey AB. 1974. Reproduction in female rats born to DDT-treated parents. Bull Environ Contam Toxicol 12:373-377.

Crum JA, Bursian SJ, Aulerich RJ, Polin D, Braselton WE. 1993. The reproductive effects of dietary heptachlor in mink (*Mustela vison*). Arch Environ Contam Toxicol 24:156-164.

Dalsenter PR, Faqi AS, Webb J, Merker HJ, Chahoud I. 1997a. Reproductive toxicity and toxicokinetics of lindane in the male offspring of rats exposed during lactation. Hum Exp Toxicol 16:146-153.

Dalsenter PR, Faqi AS, Chahoud I. 1997b. Serum testosterone and sexual behavior in rats after prenatal exposure to lindane. Bull Environ Contam Toxicol 59:360-366.

Dodds-Smith ME, Johnson MS, Thompson DJ. 1992. Trace metal accumulation by the shrew *Sorex araneus*. I. Total body burden, growth, and survival. Ecotoxicol Environ Saf 24:102-117.

Duby RT, Travis HF, Terrill CE. 1971. Uterotropic activity of DDT in rats and mink and its influence on reproduction in the rat. Toxicol Appl Pharmacol 18:348-355.

Elfant M, Keen CL. 1987. Sodium vanadate toxicity in adult and developing rats. Biol Trace Elem Res 14:193-207.

Ema M, Itami K, Kawasaki H. 1992a. Effect of period of exposure on the developmental toxicity of butyl benzyl phthalate in rats. J Appl Toxicol 12(1):57-61.

Ema M, Itami T, Kawasaki H. 1992b. Embryolethality and teratogenicity of butyl benzyl phthalate in rats. J Appl Toxicol 12(3):179-183.

Ema M, Kurosaka R, Amano H, Ogawa Y. 1994. Embryolethality of butyl benzyl phthalate during early pregnancy in rats. Reprod Toxicol 8(3):231-236.

Ema M, Harazono A, Hirose A, Kamata E. 2003. Protective effects of progesterone on implantation failure induced by dibutyltin dichloride in rats. Toxicol Let 143:233-238.

EPA. 2006. Integrated Risk Information System (IRIS) online database [online]. US Environmental Protection Agency, Washington, DC. [Cited 1/2006]. Available from: http://www.epa.gov/iris/.

Fitzhugh OG. 1948. Use of DDT insecticides on food products. Ind and Eng Chemistry 40(4):704-705.

Fitzhugh OG, Nelson AA, Quaife ML. 1964. Chronic oral toxicity of aldrin and dieldrin in rats and dogs. Food Chem Toxicol 2:551-562.

Formigli L, Scelsi R, Poggi P, Gregotti C, Di Nucci A, Sabbioni E, Gottardi L, Manzo L. 1986. Thallium-induced testicular toxicity in the rat. Environ Res 40:531-539.

Good EE, Ware GW. 1969. Effects of insecticides on reproduction in the laboratory mouse. IV. Endrin and dieldrin. Toxicol Appl Pharmacol 14:201-203.

Grant DL, Phillips WEJ, Hatina GV. 1977. Effect of hexachlorobenzene on reproduction in the rat. Arch Environ Contam Toxicol 5(2):207-216.

Hack R, Ebert E, Leist KH. 1995. Chronic toxicity and carcinogenicity studies with the insecticide endosulfan in rats and mice. Food Chem Toxicol 33(11):941-950.

Halverson AW, Palmer IS, Guss PL. 1966. Toxicity of selenium to post-weanling rats. Toxicol Appl Pharmacol 9:477-484.

Harazono A, Ema M. 2003. Suppression of decidual cell resonse induced by dibutyltin dichloride in pseudopregnant rats: as a cause of early embryonic loss. Reprod Toxicol 17:393-399.

Harris SJ, Cecil HC, Bitman J. 1974. Effect of several dietary levels of technical methoxychlor on reproduction in rats. J Agr Food Chem 22(6):969-973.

Hasten DL, Hegsted M, Keenan MJ, Morris GS. 1997. Dosage effects of chromium picolinate on growth and body composition in the rat. Nutr Res 17(7):1175-1186.

Heindel JJ, Gulati DK, Mounce RC, Russell SR, Lamb IV JC. 1989. Reproductive toxicity of three phthalic acid esters in a continuous breeding protocol. Fund Appl Toxicol 12:508-518.

Hext PM, Pinto PJ, Rimmel BA. 1999. Subchronic feeding study of antimony trioxide in rats. J Appl Toxicol 19:205-209.

Hornshaw TC, Safronoff J, Ringer RK, Aulerich RJ. 1986. LC50 test results in polychlorinated biphenyl-fed mink: age, season, and diet comparisons. Arch Environ Contam Toxicol 15:717-723.

Ingle L. 1952. Chronic oral toxicity of chlordane to rats. Arch Indus Hyg Occup Med 6:357-367.

Ivankovic S, Preussman R. 1975. Absence of toxic and carcinogenic effects after administration of high doses of chromic oxide pigment in subacute and long-term feeding experiments in rats. Food Cosmet Toxicol 13:347-351.

Jensen S, Kihlstrom JE, Olsson M, Lundberg C, Orberg J. 1977. Effects of PCB and DDT on mink (*Mustela vison*) during the reproductive season. Ambio 6(4):239.

Jia X, Lia N, Chen J. 2005. A subchronic toxicity study of elemental nano-Se in Sprague-Dawley rats. Life Sci 76:1989-2003.

Jonsson JT, Jr, Keil JE, Gaddy RG, Loadholt CB, Hennigar GR, Walker Jr EM. 1976. Prolonged ingestion of commercial DDT and PCB; Effects on progesterone levels and reproduction in the mature female rat. Arch Environ Contam Toxicol 3:479-490.

Julius AD, Davies MH, Birt DF. 1983. Toxic effects of dietary selenium in the Syrian hamster. Ann Nutr Metab 27:296-305.

Khasawinah AM, Grutsch JF. 1989. Chlordane: 24-month tumorigenicity and chronic toxicity test in mice. Regul Toxicol Pharmacol 10:244-254.

Kihlstrom JE, Olsson M, Jensen S, Johansson A, Ahlbom J, Bergman A. 1992. Effects of PCB and different fractions of PCB on the reproduction of the mink (*Mustela vison*). Ambio 2(8):563-569.

Kitchin KT, Linder RE, Scotti TM, Walsh D, Curley AO, Svendsgaard D. 1982. Offspring survival and maternal lung pathology in female rats fed hexachlorobenzene. Toxicology 23:33-39.

Lake BG, Cunningham ME, Price RJ. 1997. Comparison of the hepatic and renal effects of 1,4-dichlorobenzene in the rat and mouse. Fund Appl Toxicol 39:67-75.

Lamb JC, Chapin RE, Teague J, Lawton AD, Reel JR. 1987. Reproductive effects of four phthalic acid esters in the mouse. Toxicol Appl Pharmacol 88:255-269.

Leighton FA. 1989. Acute oral toxicity of dibenzothiophene for male CD-1 mice: LD50, lesions, and the effect of preinduction of mixed-function oxidases. Fund Appl Toxicol 12:787-792.

Lilienthal H, Benthe C, Heinzow B, Winneke G. 1996. Impairment of schedule-controlled behavior by pre- and postnatal exposure to hexachlorobenzene in rats. Arch Toxicol 70:174-181.

Loeser E, Lorke D. 1977a. Semichronic oral toxicity of cadmium. 1. Studies on rats. Toxicology 7:215-224.

Loeser E, Lorke D. 1977b. Semichronic oral toxicity of cadmium. 2. Studies on dogs. Toxicology 7:225-232.

Machemer L, Lorke D. 1981. Embryotoxic effect of cadmium on rats upon oral administration. Toxicol Appl Pharmacol 58:438-443.

MacKenzie KM, Angevine DM. 1981. Infertility in mice exposed *in utero* to benzo(a)pyrene. Biol Reprod 24:183-191.

Makita Y. 2005. Effects of perinatal combined exposure to 1,4-dichlorobenzene and 1,1-dichloro-2,2-bis (p-chlorophenyl) ethylene on rat male offspring. Basic Clin Pharmacol Toxicol 96:361-365.

Makita Y, Tanaka A, Omura M, Ogata R. 2003. Effects of simultaneous administration of tributyltin (TBT) and p, p′-DDE on female offspring of Wistar rats. J Toxicol Env Health Part A, 66:2337-2347.

Makita Y, Omura M, Ogata R. 2004. Effects of perinatal simultaneous administration of tributyltin (TBT) and p, p′-DDE (1,1-dichloro-2,2-bis(*p*-chlorophenyl) ethylene) on male offspring of Wistar rats. J Toxicol Env Health Part A, 67:385-395.

Masutomi N, Shibutani M, Takagi H, Uneyama C, Takahashi N, Hirose M. 2003. Impact of dietary exposure to methoxychlor, genestein, or diisononyl phthalate during the perinatal period on the development of the rat endocrine/reproductive systems in later life. Toxicology 192:149-170.

Mohiuddin SM, Taskar PK, Rheault M, Roy P-E, Chenard J, Morin Y. 1970. Experimental cobalt cardiomyopathy. Amer Heart J 80(4):532-543.

Morris RD. 1968. Effects of endrin feeding on survival and reproduction in the deer mouse, *Peromyscus maniculatus*. Can J Zool 46:951-958.

Murata Y, Denda A, Maruyama H, Konishi Y. 1993. Chronic toxicity and carcinogenicity studies of 1-methylnaphthalene in B6C3F$_1$ mice. Fund Appl Toxicol 21:44-51.

Murata Y, Denda A, Maruyama H, Nakae D, Tsutsumi M, Tsujiuchi T, Konishi Y. 1997. Chronic toxicity and carcinogenicity studies of 2-methylnaphthalene in B6C3F$_1$ mice. Fund Appl Toxicol 36:90-93.

Nation JR, Hare MF, Baker DM, Clark DE, Bourgeois AE. 1985. Dietary administration of nickel: effects on behavior and metallothionein levels. Physiol Behav 34:349-353.

Neal J, Rigdon RH. 1967. Gastric tumors in mice fed benzo(a)pyrene: a quantitative study. Tex Rep Biol Med 25(4):553-557.

Nickerson SC, Sniffen CJ. 1973. Effects of DDT on growth and reproduction in the rat. Res in Life Sci 20:1-6.

NTP. 1993. National Toxicology Program technical report on toxicity studies of cupric sulfate (CAS no. 7758-99-8) administered in drinking water and feed to F344/N rats and B6C3F$_1$ mice. NIH publ. 93-3352. National Toxicology Program Toxicity Report Series no. 29, National Institutes of Health, Research Triangle Park, NC.

Ogata R, Omura M, Shimasaki Y, Kubo K, Oshima Y, Aou S, Inoue N. 2001. Two-generation reproductive toxicity study of tributyltin chloride in female rats. J Toxicol Env Health Part A, 63:127-144.

Omura M, Ogata R, Kubo K, Shimasaki Y, Aou S, Oshima Y, Tanaka A, Hirata M, Makita Y, Inoue N. 2001. Two-generation reproductive toxicity study ot tributyltin chloride in male rats. Toxicol Sci 64:224-232.

Ottoboni A. 1969. Effect of DDT on reproduction in the rat. Toxicol Appl Pharmacol 14:74-81.

Ottoboni A. 1972. Effect of DDT on the reproductive life-span in the female rat. Toxicol Appl Pharmacol 22:497-502.

Palmer AK, Cozens DD, Spicer EJF, Worden AN. 1978. Effects of lindane upon reproductive function in a 3-generation study in rats. Toxicology 10:45-54.

Plasterer MR, Bradshaw WS, Booth GM, Carter MW. 1985. Developmental toxicity of nine selected compounds following prenatal exposure in the mouse: naphthalene, *p*-nitrophenol, sodium selenite, dimethyl phthalate, ethylenethiourea, and four glycol ether derivatives. J Toxicol Environ Health 15:25-38.

Pond WG, Walker EF, Jr. 1975. Effect of dietary Ca and Cd level of pregnant rats on reproduction and on dam and progeny tissue mineral concentrations. Proc Soc Exper Biol Med 148:665-668.

Poon R, Lecavalier P, Mueller R, Valli VE, Procter BG, Chu I. 1997. Subchronic oral toxicity of di-n-octyl phthalate and di(2-ethylhexyl) phthalate in the rat. Food Chem Toxicol 35:225-239.

Reuber MD. 1980. Significance of acute and chronic renal disease in Osborne-Mendel rats ingesting dieldrin or aldrin. Clin Toxicol 17(2):159-170.

Ringer RK. 1983. Toxicology of PCBs in mink and ferrets. In: D'Itri FM, Kamrin MA, eds, PCBs: human and environmental hazards. Butterworth Publishers, Boston, pp 227-240.

Rivera S, Sanfeliu C, Rodriguez-Farre E. 1990. Behavioral changes induced in developing rats by an early postnatal exposure to lindane. Neurotoxicol Teratol 12:591-595.

Rivett KF, Chesterman H, Kellett DN, Newman AJ, Worden AN. 1978. Effects of feeding lindane to dogs for periods of up to 2 years. Toxicology 9:273-289.

Schlicker SA, Cox DH. 1968. Maternal dietary zinc, and development and zinc, iron, and copper content of the rat fetus. J Nutr 95:287-294.

Schroeder HA, Balassa JJ. 1967. Arsenic, germanium, tin and vanadium in mice: effects on growth, survival and tissue levels. J Nutr 92:245-252.

Schroeder HA, Mitchener M. 1971. Toxic effects of trace elements on the reproduction of mice and rats. Arch Environ Health 23:102-106.

Seiler P, Fischer B, Lindenau A, Beier HM. 1994. Effects of persistent chlorinated hydrocarbons on fertility and embryonic development in the rabbit. Hum Reprod 9(10):1920-1926.

Shiota K, Chou MJ, Nishimura H. 1980. Embryotoxic effects of di-2-ethylhexyl phthalate (DEHP) and di-n-butyl phthalate (DBP) in mice. Environ Res 22:245-253.

Shopp GM, White Jr KL, Holsapple MP, Barnes DW, Duke SD, Anderson AC, Condie Jr LW, Hayes JR, Borzelleca JF. 1984. Naphthalene toxicity in CD-1 mice: general toxicology and immunotoxicology. Fund Appl Toxicol 4:407-419.

Smith CC. 1953. Toxicity of butyl stearate, dibutyl sebacate, dibutyl phthalate, and methoxyethyl oleate. Arch Indus Hyg Occup Med 7(4):310-318.

Srinivasan K, Mahadevappa KL, Radhakrishnamurty R. 1991. Toxicity of beta-and gamma-hexachlorocyclohexane in rats of different ages. Bull Environ Contam Toxicol 47:623-627.

Straube EF, Schuster NH, Sinclair AJ. 1980. Zinc toxicity in the ferret. J Comp Path 90:355-361.

Sutton WR, Nelson VE. 1937. Studies on zinc. Proc Soc Exper Biol Med 36:211-213.

Tarjan R, Kemeny T. 1969. Multigeneration studies on DDT in mice. Food Chem Toxicol 7:215-222.

Thorpe E, Walker AIT. 1973. The toxicology of dieldrin (HEOD). II. Comparative long-term oral toxicity studies in mice with dieldrin, DDT, phenobarbitone, beta-BHC and gamma-BHC. Food Chem Toxicol 11:433-442.

Treon JF, Cleveland FP. 1955. Toxicity of certain chlorinated hydrocarbon insecticides for laboratory animals, with special reference to aldrin and dieldrin. Agric Food Chem 3:402-408.

Treon JF, Cleveland FP, Cappel J. 1955. Toxicity of endrin for laboratory animals. Agric Food Chem 3:842-848.

Turusov VS, Day NE, Tomatis L, Gati E, Charles RT. 1973. Tumors in CF-1 mice exposed for six consecutive generations to DDT. J Nat Cancer Inst 51(983-998).

Tyl RW, Price CJ, Marr MC, Kimmel CA. 1988. Developmental toxicity evaluation of dietary di(2-ethylhexyl) phthalate in Fischer 344 rats and CD-1 mice. Fund Appl Toxicol 10:395-412.

Tyl RW, Myers CB, Marr MC, Fail PA, Seely JC, Brine DR, Barter RA, Butala JH. 2004. Reproductive toxicity evaluationof dietary butyl benzyl phthalate (BBP) in rats. Reprod Toxicol 18:241-264.

Van Velsen FL, Danse LHJC, van Leeuwen FXR, Dormans JAMA, van Logten MJ. 1986. The subchronic oral toxicity of the beta-isomer of hexachlorocyclohexane in rats. Fund Appl Toxicol 6:697-712.

Verschuuren HG, Kroes R, Den Tonkelaar EM, Berkvens JM, Helleman PW, Rauws AG, Schuller PL, Van Esch GJ. 1976. Toxicity of methylmercury chloride in rats. II. Reproduction study. Toxicology 6:97-106.

Virgo BB, Bellward GD. 1977. Effects of dietary dieldrin on offspring viability, maternal behaviour, and milk production in the mouse. Chem Pathol Pharmacol 17(3):399-409.

Walker AIT, Stevenson DE, Robinson J, Thorpe E, Roberts M. 1969. The toxicology and pharmacodynamics of dieldrin (HEOD): two-year oral exposures of rats and dogs. Toxicol Appl Pharmacol 15:345-373.

Ware GW, Good EE. 1967. Effects of insecticides on reproduction in the laboratory mouse. II. Mirex, telodrin, and DDT. Toxicol Appl Pharmacol 10:54-61.

Weber CW, Reid BL. 1969. Nickel toxicity in young growing mice. J Anim Sci 20:620-623.

Wellman PJ, Watkins PA, Nation JR, Clark DE. 1984. Conditioned taste aversion in the adult rat induced by dietary ingestion of cadmium or cobalt. Neurotoxicology 5(2):81-90.

Wester PW, Krajnc EI, van Leeuwen FX, Loeber JG, van der Heijden CA, Vaessen HAMG, Helleman PW. 1990. Chronic toxicity and carcinogenicity of bis(tri-n-butyltin) oxide (TBTO) in the rat. Fd Chem Toxicol 28(3):179-196.

Wine RN, Li LH, Barnes LH, Gulati DK, Chapin RE. 1997. Reproductive toxicity of di-n-butylphthalate in a continuous breeding protocol in Sprague-Dawley rats. Environ Health Perspect 105:101-107.

Wobeser G, Nielsen NO, Schiefer B. 1976. Mercury and mink II. Experimental methyl mercury intoxication. Can J Comp Med 40:34-45.

Wolfe JL, Esher RJ, Robinson KM, Yarbrough JD. 1979. Lethal and reproductive effects of dietary mirex and DDT on old-field mice, *Peromyscus polionotus*. Bull Environ Contam Toxicol 21:397-402.

Wren CD, Hunter DB, Leatherland JF, Stokes PM. 1987. The effects of polychlorinated biphenyls and methylmercury, singly and in combination on mink. II: Reproduction and kit development. Arch Environ Contam Toxicol 16:449-454.

Wrenn TR, Weyant JR, Fries GF, Bitman J. 1971. Effect of several dietary levels of o,p'-DDT on reproduction and lactation in the rat. Bull Environ Contam Toxicol 6:471-480.

You L, Casanova M, Bartolucci EJ, Fryczynski MW, Dorman DC, Everitt JI, Gaido KW, Ross SM, Heck HD. 2002. Combined effects of dietary phytoestrogen and synthetic endocrine-active compound on reproductive development in Sprague-Dawley rats: genistein and methoxychlor. Toxicol Sci 66:91-104.

ATTACHMENT 11
EXPOSURE CONCENTRATION CALCULATION METHODS

Attachment 11 Exposure Concentration Calculation Methods

1.0 INTRODUCTION 1
2.0 PROUCL CALCULATIONS 2
3.0 ROS METHODS 2
FIGURE 1. NORMAL QUANTILE PLOT FOR TBT CONCENTRATIONS IN CHINOOK SALMON WHOLE-BODY SAMPLES (WET WEIGHT) 3
4.0 REGRESSION-BASED CALCULATIONS 4
FIGURE 2. LINEAR REGRESSION RELATIONSHIP FOR TOTAL PCBS CONCENTRATIONS IN BENTHIC INVERTEBRATE TISSUES 4
FIGURE 3. REGRESSION RELATIONSHIP FOR ARSENIC CONCENTRATIONS IN BENTHIC INVERTEBRATE TISSUES 5
FIGURE 4. REGRESSION RELATIONSHIP FOR TBT CONCENTRATIONS IN BENTHIC INVERTEBRATE TISSUES 6
5.0 CALCULATION OF SWAC SEDIMENT CONCENTRATIONS 7
6.0 CALCULATION OF SWAC UCL 8
7.0 REFERENCES 9
TABLE 1. SUMMARY OF METHODS USED TO CALCULATE EXPOSURE CONCENTRATIONS 10
TABLE 2. EXPOSURE CONCENTRATIONS IN FISH, CRAB, CLAM, AND MUSSEL TISSUE ON A DRY-WEIGHT BASIS 13
TABLE 3. EXPOSURE CONCENTRATIONS IN SHINER SURFPERCH AND CRAB TISSUE FOR FOUR PACIFIC STAGHORN SCULPIN MODELING AREAS ON A DRY-WEIGHT BASIS 17
TABLE 4. EXPOSURE CONCENTRATIONS IN FISH AND CRAB TISSUE ON A WET WEIGHT BASIS FOR THE ENTIRE LDW 19
TABLE 5. EXPOSURE CONCENTRATIONS IN PACIFIC STAGHORN SCULPIN TISSUE ON A WET WEIGHT BASIS FOR FOUR PACIFIC STAGHORN SCULPIN MODELING AREAS 21
TABLE 6. EXPOSURE CONCENTRATIONS IN BENTHIC INVERTEBRATES CALCULATED FOR SIX EXPOSURE SCENARIOS FOR SPOTTED SANDPIPER ON A DRY WEIGHT BASIS 22
TABLE 7. EXPOSURE CONCENTRATIONS IN BENTHIC INVERTEBRATES CALCULATED FOR INTERTIDAL AREAS ON A DRY-WEIGHT BASIS 27
TABLE 8. EXPOSURE CONCENTRATIONS IN BENTHIC INVERTEBRATE TISSUE CALCULATED FOR FOUR PACIFIC STAGHORN SCULPIN MODELING AREAS ON A DRY-WEIGHT BASIS 28
TABLE 9. EXPOSURE CONCENTRATIONS IN BENTHIC INVERTEBRATES CALCULATED FOR THE ENTIRE LDW ON A DRY-WEIGHT BASIS 29
TABLE 10. EXPOSURE CONCENTRATIONS IN SEDIMENT CALCULATED FOR SIX EXPOSURE SCENARIOS FOR SANDPIPER ON A DRY-WEIGHT BASIS 30
TABLE 11. EXPOSURE CONCENTRATIONS IN SEDIMENT CALCULATED FOR INTERTIDAL AREAS ON A DRY-WEIGHT BASIS 34
TABLE 12. EXPOSURE CONCENTRATIONS IN SEDIMENT CALCULATED FOR FOUR PACIFIC STAGHORN SCULPIN MODELING AREAS ON A DRY-WEIGHT BASIS 35

TABLE 13. EXPOSURE CONCENTRATIONS FOR SEDIMENT CALCULATED FOR THE ENTIRE LDW ON A DRY-WEIGHT BASIS *36*
TABLE 14. EXPOSURE CONCENTRATIONS IN WATER CALCULATED FOR THREE EXPOSURE AREAS FOR SPOTTED SANDPIPER *37*
TABLE 15. EXPOSURE CONCENTRATIONS IN WATER CALCULATED FOR THE ENTIRE LDW *39*

1.0 INTRODUCTION

Exposure concentrations were calculated throughout the ecological risk assessment (ERA) for all receptors and exposure pathways. The purpose of this attachment is to provide a summary of the methods used for the calculation of each exposure concentration. Upper confidence limits (UCLs) were calculated using three methods:

The method used most frequently was the calculation of UCLs using ProUCL software. ProUCL tests for normality, lognormality, and a gamma distribution of the dataset and computes a 95% UCL of the unknown population mean (EPA 2004) (Section 2.0). The tissue sample dataset for the ERA consists of composite tissue samples. The analysis of composite samples results in an increase in the number of individuals represented by the sample results. However, compositing samples results in the loss of information about the variance within a population of individuals and may therefore affect exposure estimates. If the variance of data based on individual samples is greater than the variance of composite samples, the difference between the 95% UCL and the sample mean would be greater for a dataset consisting of individual samples. This distinction is relevant to the risk assessment because the reasonable maximum exposure scenarios were derived based on the 95% UCL for each medium, the 95% UCL may be lower based on a dataset of composite samples compared to a dataset of individual samples because of the reduced variance around the mean.

However, there is no reason to assume that the mean of the composites would be equal to the mean of the individual samples, nor that the variance would be greater. The sample mean and variance of analogous data sets based on individual samples is simply unknown. If the mean of the individuals were sufficiently lower than the mean of the composites, it is possible that the 95% UCL on the mean of the individual samples would be less than the 95% UCL on the mean of the composite samples, even if the individual variance was greater.

If the non-detect frequency was less than or equal to 20%, ProUCL was used with half the reporting limit used as the value for non-detected results. The ProUCL User Guide (Singh et al, 2004) recommends careful attention to data sets with < 85% detection frequency. Therefore, different non-detect methods were used for datasets with greater than 20% and less than 85% frequency of non-detects. Robust regression on order statistic (ROS) methods were used to fill-in a set of concentrations for non-detects that would result in a better estimate of central tendency than substitution methods. The non-detected concentrations were assumed to follow a log-normal distribution with parameters estimated from the detected concentrations. The full dataset, including the filled-in concentrations, was then used to calculate the 95%UCL using ProUCL (Section 3.0).

If the non-detect frequency was greater than 80% then, no non-detect method should be used (Helsel 2005). In the instances where this was the case, the maximum detected concentration was used as the exposure point concentration.

Because of the limited dataset of benthic invertebrate tissue samples, the use of regression relationships between the invertebrate tissue concentrations and co-located sediment concentrations was explored to estimate the UCLs of the mean site-wide benthic invertebrate tissue concentrations as well as the UCLs of the mean for benthic invertebrate tissue concentrations within specific exposure areas (i.e., sandpiper exposure areas and intertidal exposure areas for fish and wildlife) (Section 4.0).

Spatially weighted average concentrations (SWACs) are available for sediment total PCB concentrations in the LDW (Section 5.0). For total PCBs, the area SWAC concentrations were used with the regression relationships to estimate the benthic invertebrate tissue concentrations. SWAC UCL values were calculated for the LDW as a whole as well as the intertidal areas to provide exposure point concentrations for the dietary consumption of sediment by wildlife receptors (Section 6.0).

A summary of the methods used for each receptor and chemical of potential concern (COPC) is presented in Table 1. Detailed descriptions of the specific statistic calculated for each exposure concentration as well as summary statistics on the dataset used in the calculation are presented in Tables 2–12.

2.0 PROUCL CALCULATIONS

UCL calculations and assessment of data distributions were performed using ProUCL software (EPA 2004). ProUCL software was developed by EPA to compute an appropriate 95% UCL of an unknown population mean. ProUCL tests for normality, lognormality, and a gamma distribution of the dataset, selects a conservative distribution, and computes a UCL of the unknown population mean.

For each chemical of potential concern (COPC) that was detected in more than 85% of the dataset being considered, the UCL recommended by ProUCL was used as the exposure concentration for the risk calculations. Summary statistics, distribution type, and UCL on the mean for chemical concentrations in tissue and sediment are presented in Tables 2–12.

ProUCL is not recommended for use in determining UCLs for datasets with a percentage of non-detects greater than 15% when a simple substitution method for non-detects has been employed (EPA 2004). In the baseline dataset, each non-detect value was replaced by one-half the reporting limit (RL) as a proxy value. When datasets contain a large number of proxy values, these values have the potential to skew the overall distribution of the data.

3.0 ROS METHODS

For COPCs that had detection frequencies less than 85% and greater than 20%, robust ROS methods were used to estimate a distribution of values for the non-detected results. These calculated values are referred to as "imputed data" and comprise the censored portion of the dataset. A robust ROS method that required an assumed

distribution for the censored portions of the datasets was selected based on Helsel (2005). Lognormal distributions were assumed for the censored data. This is an appropriate default distribution for environmental concentrations because it will not generate negative concentration values and is robust to differing amounts of skewness. A linear regression was computed from the normal probability plot of the natural logarithms of the detected concentrations versus their normal scores. Values for individual censored or nondetected observations were predicted from the linear regression. An example of the results of the imputation process for TBT concentrations in juvenile chinook is displayed graphically in Figure 1. In Figure 1, the natural-log transformed detected TBT concentrations are plotted versus their normal scores, and the least-squares linear regression line fit to these data points is displayed. The non detected concentrations are displayed as inverted triangles at their respective RLs, and as "x" marks at their imputed concentrations. Note that the regression lines displayed were the original regressions on detected concentrations only ("o" symbols). The final imputed values were converted back to original units to avoid transformation bias, and combined with the original detected concentrations in the calculation of the UCL using ProUCL.

The ROS method produces datasets containing both detected and imputed data. These datasets are then used to calculate UCLs, employing ProUCL as before.



Figure 1. Normal quantile plot for TBT concentrations in chinook salmon whole-body samples (wet weight)

4.0 REGRESSION-BASED CALCULATIONS

Benthic invertebrate tissue samples were collected from 10 intertidal locations and 10 subtidal locations, with co-located sediment collected from each location (Windward 2005). The sample locations were selected to represent areas that covered the range of arsenic, lead, and polychlorinated biphenyl (PCB) concentrations measured throughout the Lower Duwamish Waterway (LDW) so that a relationship between PCB concentrations in sediment and benthic invertebrate tissue could be established (Windward 2004). The locations were not intended to be used to represent the average concentrations of chemicals for the entire site.

The relationships between COPC concentrations in benthic invertebrate tissue and co-located sediment for each COPC were examined to determine whether a significant linear regression existed. The term "significant" regression as used here means that there is sufficient statistical evidence to conclude that the slope of the true regression line is not zero. The significance of the linear regression was determined by calculating the probability (p-value) of obtaining an F-ratio greater than or equal to the observed F ratio if the true slope of the regression line was zero. The regression was deemed significant if the p-value was less than 0.05. Influential points with Cook's Distance greater than 1 were removed for this determination. Both linear and exponential regressions were evaluated, and the exponential model was used instead of the linear model if it provided a superior fit, as determined by the R^2 value. A linear model was selected for the PCBs (Figure 2), and exponential models were selected for arsenic and TBT (Figures 3 and 4).



Figure 2. Linear regression relationship for total PCBs concentrations in benthic invertebrate tissues



Figure 3. Regression relationship for arsenic concentrations in benthic invertebrate tissues



Figure 4. Regression relationship for TBT concentrations in benthic invertebrate tissues

The concentrations in sediment were often highly skewed, with the highest values identified as overly influential points using Cook's distance. The sediment spatially weighted average concentrations (SWACs) and mean concentrations were generally between the majority of the co-located sediment concentrations and the highly influential points. In these cases, the influential points were retained in the final regression models, presented in Table 3, in order to avoid extrapolation issues.

There is uncertainty in the selected exponential regressions for arsenic and TBT because of three data points that exerted undue influence on the regression relationship. The points were influential because the distribution of sediment concentrations was skewed, with few high concentrations. The removal of these points would result in a different regression relationship. However, the resulting relationship would not be appropriate for predicting tissue concentrations because extrapolation beyond the included sediment chemistry concentrations would be necessary. Also, because the existing data give some evidence that the relationship is exponential, excluding the high points in the distribution would likely result in a highly biased tissue prediction.

The regressions were used to estimate LDW-wide exposure concentrations to assess risk to benthic invertebrates from TBT exposure based on a tissue-residue approach as well as dietary exposure to arsenic for fish receptors. The PCB concentrations were used to assess risk to sandpiper in the various sandpiper exposure areas throughout

the LDW. For TBT and arsenic, site-wide arithmetic mean sediment concentrations were calculated and the TBT and arsenic regression models were used to predict the corresponding 95th UCL on the expected TBT and arsenic tissue concentrations. The 95th UCL on the TBT and arsenic tissue concentrations were used as the exposure point concentrations in the risk assessment.

5.0 CALCULATION OF SWAC SEDIMENT CONCENTRATIONS

An arithmetic mean of the sediment concentrations throughout a specified area is biased to overestimate exposure when more highly contaminated areas have been sampled more intensively than less-contaminated areas. In the LDW, this bias is most pronounced for total PCBs, which are the focus of several ongoing early action investigations in the LDW. Consequently, SWAC were calculated to estimate benthic tissue concentrations from the sediment-tissue regression model, and SWAC UCLs for total PCBs were used as exposure concentrations for the sediment component of the wildlife diets. The methods used to generate the UCL and SWAC are briefly described below. A more thorough description of the methods is provided in *Technical Memorandum: GIS Interpolation of Total PCBs in LDW Surface Sediment* (Windward 2006).

The SWAC was calculated from an inverse distance weighting (IDW) interpolation using the ESRI® ArcGIS extension Geostatistical Analyst. IDW is a technique in which interpolated estimates are made at the center of 10 x 10 ft grid cells covering the study area. Sample concentrations within a specified search radius of each grid cell are weighted by the inverse of their distance from the cell, with the effect of giving more influence to data points nearby than to those farther away. A variable weighting factor can be used to give more or less importance to distant points, relative to nearby points. The weighted mean is applied to each grid cell to create the IDW interpolation. The IDW technique creates a continuous surface of grid cells in which each cell is represented by a single estimated concentration.

The best fit of the search radius to the data can be evaluated by comparing root mean square errors (RMSEs) and mean errors of trial interpolations. These statistics are generated by removing one sample location from the dataset and using the specified search parameters and weighting factor to estimate the concentration at that location. The differences between the known concentrations and the interpolated concentrations are then used to calculate RMSEs and mean errors across the LDW.

Many interpolations were created systematically using different sets of input parameters, and the interpolation that optimized the error statistics was selected. The search shape that optimized these statistics for PCB interpolations was an ellipse angled in the direction of the river's flow. Because the LDW has three major angles, the LDW was divided into three reaches, and interpolation parameters were optimized for each reach separately. Once the interpolations were finalized for each of the three reaches, they were combined using tools available through the ESRI® ArcGIS

extension Spatial Analyst. Finally, the SWAC was calculated as the mean of all grid cell values in the LDW.

6.0 CALCULATION OF SWAC UCL

The IDW method does not include the necessary information that would permit the calculation of the UCL. Therefore, Microsoft® Visual Basic for Applications (VBA) was used to build a customizable program that generated the IDW grid as well as calculated various descriptive statistics for the IDW grid values. The interpolation parameters determined through systematic testing used in the VBA program to isolate the samples used to interpolate each cell value. Sample concentrations used to estimate the value of each grid cell were assumed to have a normal distribution because the sample size was often too low for normality testing (most cells were estimated using 4 to 16 samples). The program identified the samples within the search radius as defined by the interpolation parameters, retrieved the distance to each sample, and calculated weights based on the specified power value (p).

The UCLs and lower confidence limits (LCLs) for each grid cell are then calculated using the sample weights determined above. First, the weighted sample variance is calculated (Bing-Canar 2006):

$$\text{weighted sample variance} = \frac{\sum_{i=1}^{n} \text{adjusted sample weight}_i \left(\text{sample concentration}_i - \text{weighted mean}\right)^2}{\sum_{i=1}^{n} wt^2}$$

Equation 1

Where n is the number of sampling stations within the search area. The weighted standard deviation is calculated by taking the square root of the weighted sample variance.

The UCL per cell was then calculated as follows:

$$\text{UCL} = \text{weighted mean} + \left(t_{n-1,0.95}\right) \times \left(\frac{\text{weighted standard deviation}}{\sqrt{n}}\right)$$

Equation 2

Where:

$t_{n\text{-}1,0.95}$ = the 95th percentile of the Student's t-distribution with n-1 degrees of freedom

Each cell UCL is then applied to the cells in a new output grid. The result is a surface that represents the UCL across all cells. The cell values in the UCL surface were averaged to calculate the UCL for the LDW.

The LDW-wide total PCB interpolation was clipped to create UCL-SWACs for the intertidal area (for dietary sediment exposure for great blue heron and osprey) as well

as for the spotted sandpiper exposure areas (for dietary sediment exposure for spotted sandpiper).

7.0 REFERENCES

Bing-Canar J. 2006. Personal communication (telephone conversation with Cindy Jackson, Windward Environmental, regarding interpolation of LDW sediment chemistry and UCL calculations). US Environmental Protection Agency Region 10, Seattle, WA. February 8, 2006.

EPA. 2004. ProUCL Version 3.0 user guide. Technical Support Center for Monitoring and Site Characterization, US Environmental Protection Agency.

Helsel DR. 2005. Nondetects and data analysis: Statistics for censored environmental data. John Wiley & Sons, Hoboken, NJ.

Windward. 2004. Lower Duwamish Waterway remedial investigation. Quality assurance project plan: Benthic invertebrate sampling of the Lower Duwamish Waterway. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005. Lower Duwamish Waterway remedial investigation. Data report: Chemical analyses of benthic invertebrate and clam tissue samples and co-located sediment samples. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward. 2006. Lower Duwamish Waterway remedial investigation. Technical memorandum: GIS interpolation of total PCBs in LDW surface sediment. Windward Environmental LLC, Seattle, WA.

Table 1. Summary of methods used to calculate exposure concentrations

Receptor of Concern/Approach	Chemical of Potential Concern	Wet- or Dry-Weight Basis[a]	Media Type	Statistical Method	Table Presenting results
Benthic invertebrates/critical tissue residue	TBT	dry	benthic invertebrate tissue	regression	Table 9
Crab/critical tissue residue	zinc, methoxychlor, total DDTs, total PCBs	wet	whole-body crab tissue	UCL	Table 4
Juvenile chinook salmon/critical tissue residue	endrin	wet	juvenile chinook salmon	UCL	Table 4
English sole/critical tissue residue	alpha-endosulfan, beta-endosulfan, benzoic acid, BEHP, dimethyl-phthalate, di-n-butyl-phthalate, endrin, total PCBs, PCB TEQ	wet	whole-body English sole	UCL, maximum detected concentration, or maximum RL	Table 4
Pacific staghorn sculpin/critical tissue residue	alpha-endosulfan, beta-endosulfan, benzoic acid, BEHP, dimethyl-phthalate, di-n-butyl-phthalate, endrin, TBT, total PCBs	wet	whole-body Pacific staghorn sculpin	UCL, maximum detected concentration, or maximum RL	Tables 4, 5
Shiner surfperch/ critical tissue residue	total PCBs, TBT	wet	whole-body shiner surfperch	UCL	Table 4
Juvenile chinook salmon/dietary	arsenic, cadmium, chromium, copper, vanadium, total alkylated +non-alkylated PAH (benthic invertebrates only)	dry	benthic invertebrates	95% UCL of regression for arsenic, UCL for other COPCs	Table 7, 9
		dry	juvenile chinook salmon stomach contents	sample concentration	Table A.4-5 in Section A.4.1.2
Juvenile chinook salmon/sediment	total PAHs	dry	sediment	UCL	Table 13
English sole/dietary	arsenic, cadmium, chromium, copper, vanadium	dry	benthic invertebrates	95% UCL of regression for arsenic, UCL for other COPCs	Table 9
		dry	sediment	UCL	Table 13
English sole/sediment	total PAHs	dry	sediment	UCL	Table 13

Receptor of Concern/Approach	Chemical of Potential Concern	Wet- or Dry-Weight Basis[a]	Media Type	Statistical Method	Table Presenting results
Pacific staghorn sculpin/dietary	arsenic, cadmium, chromium, copper, vanadium	dry	benthic invertebrates	95% UCL of regression for arsenic, UCL for other COPCs	Tables 8, 9
		dry	shiner surfperch, crab	UCL	Tables 2, 3
		dry	sediment	UCL	Tables 12, 13
Pacific staghorn sculpin/sediment	total PAHs	dry	sediment	UCL	Table 13
Spotted sandpiper/dietary	arsenic, chromium, cadmium, cobalt, copper, lead, mercury, nickel, selenium, vanadium, zinc, BEHP, total PCBs, PCB TEQs, total DDTs	dry	benthic invertebrates	95% UCL of regression for arsenic and PCBs; UCL, maximum detected concentration, or maximum RL for other COPCs	Table 6
			sediment	95% UCL of SWAC for PCBs, UCL for other COPCs	Table 10
		na	water	UCL or maximum detected concentration	Table 14
Great blue heron/dietary	chromium, lead, mercury, total PCBs, PCB TEQs	dry	shiner surfperch, English sole, juvenile chinook salmon, Pacific staghorn sculpin, whole-body crab	UCL	Table 2
		dry	sediment	95% UCL of SWAC for PCBs; UCL for other COPCs	Table 11
		na	water	UCL	Table 15
Osprey/dietary	chromium, lead, mercury, total PCBs, PCB TEQs	dry	shiner surfperch, English sole, juvenile chinook salmon	UCL	Table 2
		dry	sediment	95% UCL of SWAC for PCBs; UCL for other COPCs	Table 11
		na	water	UCL or maximum detected concentration	Table 15

Receptor of Concern/Approach	Chemical of Potential Concern	Wet- or Dry-Weight Basis[a]	Media Type	Statistical Method	Table Presenting results
River otter/dietary	arsenic, cobalt, mercury, selenium, total PCBs, PCB TEQs	dry	shiner surfperch, English sole, juvenile chinook salmon, Pacific staghorn sculpin, whole-body crab, clams, mussels	UCL	Table 2
		dry	sediment	95% UCL of SWAC for PCBs; UCL for other COPCs	Table 13
		na	water	UCL or maximum detected concentration	Table 15
Harbor seal/dietary	mercury, total PCBs, PCB TEQs	dry	shiner surfperch, English sole, juvenile chinook salmon, Pacific staghorn sculpin	UCL	Table 2
		dry	sediment	95% UCL of SWAC for PCBs; UCL for other COPCs	Table 13
		na	water	UCL or maximum detected concentration	Table 15

[a] Tissue concentrations in Tables 2 through 9 are presented on a wet or dry weight basis depending on how those data were used in risk calculations.

BEHP – bis(2-ethylhexyl)phthalate
COPC – chemical of potential concern
na – not applicable
PCB – polychlorinated biphenyl
SWAC – spatially weighted average concentration
TBT – tributyltin
TEQ – toxic equivalency quotient
UCL – upper confidence limit

Table 2. Exposure concentrations in fish, crab, clam, and mussel tissue on a dry-weight basis

Receptor and COPC	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	UCL Type	Concentration Used in Risk Estimation (mg/kg dw)
			Min	Max	Mean			
Shiner surfperch								
Arsenic	27	100%	2.97	5.4	4.0	gamma (0.05)	approximate gamma	4.2
Cadmium	27	100%	0.038	0.0972	0.061	normal (0.05)	Student's-t UCL	0.066
Chromium	27	96%	0.26	1.8	0.70	gamma distribution (0.05)	approximate gamma UCL	0.82
Cobalt	24	100%	0.110	0.222	0.17	normal (0.05)	Student's-t UCL	0.18
Copper	27	100%	2.260	8.902	6.4	normal (0.05)	Student's-t UCL	7.0
Total DDTs	24	100%	0.14	3.8	0.75	lognormal (0.05)	95% H-UCL	1.0
Lead	27	100%	0.176	1.061	0.47	lognormal (0.05)	Student's-t UCL	0.53
Mercury	27	100%	0.071	0.34	0.13	nonparametric (0.05)	Student's-t UCL	0.15
Total PCBs	49	100%	1.3	69	6.9	nonparametric (0.05)	95% Chebyshev UCL	14
PCB TEQ (mammal)	9	100%	3.04×10^{-5}	2.74×10^{-4}	9.3×10^{-5}	gamma distribution (0.05)	approximate gamma UCL	1.6×10^{-4}
PCB TEQ (bird)	9	100%	1.67×10^{-4}	1.48×10^{-3}	4.5×10^{-4}	gamma distribution (0.05)	approximate gamma UCL	7.2×10^{-4}
Selenium	24	100%	0.448	0.866	0.72	gamma distribution (0.05)	approximate gamma UCL	0.77
Silver	27	89%	0.009	0.0437	0.019	gamma distribution (0.05)	approximate gamma UCL	0.021
Vanadium	24	92%	0.49	4.62	1.6	lognormal (0.05)	95% H-UCL	2.0
English sole								
Arsenic	24	100%	3.51	19.67	12	nonparametric (0.05)	Student's-t UCL	13
Chromium	24	88%	0.25	16.3	1.5	nonparametric (0.05)	95% Chebyshev UCL	4.4
Cobalt	24	100%	0.102	1	0.24	nonparametric (0.05)	95% Chebyshev UCL	0.41
Total DDTs	24	100%	0.22	1.1	0.68	data are normal (0.05)	Student's-t UCL	0.76
Lead	24	100%	0.15	3.64	1.4	normal (0.05)	Student's-t UCL	1.7
Mercury	24	100%	0.02	0.12	0.058	gamma distribution (0.05)	approximate gamma UCL	0.070
PCBs	45	100%	2.0	18	8.8	gamma distribution (0.05)	approximate gamma UCL	10
PCB TEQ (mammal)	8	100%	1.86×10^{-5}	1.02×10^{-4}	6.4×10^{-5}	data are normal (0.05)	Student's-t UCL	8.2×10^{-5}
PCB TEQ (bird)	8	100%	8.38×10^{-5}	6.03×10^{-4}	3.2×10^{-4}	normal (0.05)	Student's-t UCL	4.2×10^{-4}

Table 2, cont. Exposure concentrations in fish, crab, clam, and mussel tissue on a dry-weight basis

Receptor and COPC	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	UCL Type	Concentration Used in Risk Estimation (mg/kg dw)
			Min	Max	Mean			
Selenium	24	100%	0.4	1.24	0.76	gamma distribution (0.05)	approximate gamma UCL	0.82
Juvenile chinook salmon								
Arsenic	nd	nd	nd	nd	nd	nd	nd	nd
Chromium	nd	nd	nd	nd	nd	nd	nd	nd
Cobalt	nd	nd	nd	nd	nd	nd	nd	nd
Total DDTs	18	100%	0.0067	0.44	0.13	gamma distribution (0.05)	approximate gamma UCL	0.21
Lead	nd	nd	nd	nd	nd	nd	nd	nd
Mercury	6	100%	0.1	0.15	0.12	normal (0.05)	Student's-t UCL	0.14
PCBs	24	100%	0.031	6	0.71	nonparametric (0.05)	99% Chebyshev UCL	3.4
PCB TEQ (mammal)	nd	nd	nd	nd	nd	nd	nd	nd
PCB TEQ (bird)	nd	nd	nd	nd	nd	nd	nd	nd
Selenium	nd	nd	nd	nd	nd	nd	nd	nd
Pacific staghorn sculpin								
Arsenic	24	100%	1.78	6.59	3.5	normal (0.05)	Student's-t UCL	4.0
Chromium	24	25%	0.24	0.5	0.30	nonparametric (0.05)	Student's-t UCL	0.40[a]
Cobalt	24	100%	0.0854	0.193	0.12	gamma distribution (0.05)	approximate gamma UCL	0.13
Total DDTs	24	100%	0.15	1.1	0.40	lognormal (0.05)	95% H-UCL	0.50
Lead	24	100%	0.059	0.525	0.25	normal (0.05)	Student's-t UCL	0.30
Mercury	24	100%	0.088	0.18	0.14	normal (0.05)	Student's-t UCL	0.15
PCBs	28	100%	2.1	13	4.3	nonparametric (0.05)	Student's-t UCL	5.0
PCB TEQ (mammal)	8	100%	1.84×10^{-5}	4.88×10^{-5}	3.1×10^{-5}	normal (0.05)	Student's-t UCL	3.8×10^{-5}
PCB TEQ (bird)	8	100%	3.90×10^{-5}	1.59×10^{-4}	9.4×10^{-5}	normal (0.05)	Student's-t UCL	1.2×10^{-4}
Selenium	24	100%	0.66	1.1	0.83	normal (0.05)	Student's-t UCL	0.87
Clam								
Arsenic	14	100%	9.487	43.16	21	normal (0.05)	Student's-t UCL	25
Chromium	14	100%	2.5	8.15	4.5	normal (0.05)	Student's-t UCL	5.2

Table 2, cont. Exposure concentrations in fish, crab, clam, and mussel tissue on a dry-weight basis

Receptor and COPC	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	UCL Type	Concentration Used in Risk Estimation (mg/kg dw)
			Min	Max	Mean			
Cobalt	14	100%	1.122	5.19	2.3	nonparametric (0.05)	Student's-t UCL	2.9
Total DDTs	14	100%	0.028	0.2	0.077	gamma distribution (0.05)	approximate gamma UCL	0.10
Lead	14	100%	2.45	39.32	13	gamma distribution (0.05)	approximate gamma UCL	19
Mercury	14	100%	0.06	0.15	0.11	normal (0.05)	Student's-t UCL	0.12
PCBs	14	100%	0.18	3.6	0.97	nonparametric (0.05)	99% Chebyshev UCL	4.0
PCB TEQ (mammal)	8	100%	3.13×10^{-6}	3.49×10^{-5}	9.6×10^{-6}	nonparametric (0.05)	95% Chebyshev UCL	2.7×10^{-5}
PCB TEQ (bird)	8	100%	3.15×10^{-5}	1.90×10^{-4}	6.2×10^{-5}	nonparametric (0.05)	95% Chebyshev UCL	1.5×10^{-4}
Selenium	14	100%	1.41	2.39	1.9	normal (0.05)	Student's-t UCL	2.0
Crab whole body (estimated)								
Arsenic	21	100%	13.95 M	55 M	21	nonparametric (0.05)	Student's-t UCL	25
Cadmium	21	100%	0.19 M	1.778 M	0.95	normal (0.05)	Student's-t UCL	1.1
Chromium	21	29%	0.1 M	0.64 M	0.25	nonparametric (0.05)	95% Chebyshev UCL	0.30[a]
Copper	21	100%	47.1 M	121 M	80	gamma distribution (0.05)	approximate gamma UCL	90
Cobalt	19	100%	0.146 JM	0.6855 JM	0.31	gamma distribution (0.05)	approximate gamma UCL	0.39
Total DDTs	19	100%	0..298 M	0.92 M	0.52	nonparametric (0.05)	Student's-t UCL	0.61
Lead	21	100%	0.1 M	1.1 JM	0.38	lognormal (0.05)	95% H-UCL	0.53
Mercury	21	100%	0.17 M	0.49 M	0.26	gamma distribution (0.05)	approximate gamma UCL	0.29
PCBs	25	100%	1.4 M	9.2 M	5.1	gamma distribution (0.05)	approximate gamma UCL	6.0
PCB TEQ (mammal)	6	100%	3.18×10^{-5} M	5.93×10^{-5} M	4.2×10^{-5}	normal (0.05)	Student's-t UCL	5.0×10^{-5}
PCB TEQ (bird)	6	100%	2.28×10^{-4} M	3.71×10^{-4} M	3.2×10^{-4}	normal (0.05)	Student's-t UCL	3.6×10^{-4}
Selenium	19	100%	0.81 M	1.45 M	1.1	normal (0.05)	Student's-t UCL	1.2
Vanadium	19	62%	0.25 M	1.0 M	0.73	nonparametric (0.05)	95% Chebyshev UCL	1.1
Zinc	21	100%	130 M	216 M	180	nonparametric (0.05)	Student's-t UCL	190
Mussels								
Arsenic	22	100%	2.3	7.4	5.4	normal (0.05)	Student's-t UCL	5.9
Chromium	22	95%	0.17	2.3	1.1	normal (0.05)	Student's-t UCL	1.2

Table 2, cont. Exposure concentrations in fish, crab, clam, and mussel tissue on a dry-weight basis

Receptor and COPC	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	UCL Type	Concentration Used in Risk Estimation (mg/kg dw)
			Min	Max	Mean			
Cobalt	11	100%	0.2	0.47	0.37	normal (0.05)	Student's-t UCL	0.41
Total DDTs	11	0%	0.0087	0.0087	0.0087	all non-detects	maximum RL	0.0087
Lead	22	100%	0.87	4.8	2.8	gamma distribution (0.05)	approximate gamma UCL	3.3
Mercury	21	100%	0.059	0.15	0.087	gamma distribution (0.05)	approximate gamma UCL	0.094
PCBs	22	82%	0.078	0.4	0.23	normal (0.05)	Student's-t UCL	0.27[a]
PCB TEQ (mammal)	nd	nd	nd	nd	nd	nd	nd	nd
PCB TEQ (bird)	nd	nd	nd	nd	nd	nd	nd	nd
Selenium	nd	nd	nd	nd	nd	nd	nd	nd

[a] Calculated with ROS method because detection frequency was <85%.

[b] Detection frequency was below 20%. If detected, the maximum detected concentration was used. If not detected the maximum RL was used.

COPC – chemical of potential concern

dw – dry weight

JM – estimated calculated value

M – calculated value

N – number of samples

nd – no data

PCB – polychlorinated biphenyl

RL – reporting limit

ROS – regression on order statistic

TEQ – toxic equivalency quotient

UCL – upper confidence limit

Table 3. Exposure concentrations in shiner surfperch and crab tissue for four Pacific staghorn sculpin modeling areas on a dry-weight basis

Receptor and COPC	Area	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	UCL Type	Concentration Used in Risk Estimation (mg/kg dw)
				Minimum	Maximum	Mean			
Crab									
Arsenic	M1	8	100%	16.24 M	55 M	28	normal (0.05)	Student's-t	36
	M2	6	100%	14.68 M	17.05 M	16	normal (0.05)	Student's-t	17
	M3	6	100%	13.95 M	18.75 M	15	non-parametric (0.05)	Student's-t	17
	M4	1	100%	26.98 M	26.98 M	27	n=1	reported conc.	27
Cadmium	M1	8	100%	0.19 M	1.778 M	1.2	non-parametric (0.05)	95% Chebyshev (Mean, Sd) UCL	2.2
	M2	6	100%	0.7916 M	0.9669 M	0.89	normal (0.05)	Student's-t	0.95
	M3	6	100%	0.4871 M	0.8336 M	0.61	normal (0.05)	Student's-t	0.71
	M4	1	100%	1.197 M	1.197 M	1.2	n=1	reported conc.	1.2
Copper	M1	8	100%	93.8 M	121 M	110	normal (0.05)	Student's-t	110
	M2	6	100%	65 M	79.5 M	70	normal (0.05)	Student's-t	75
	M3	6	100%	47.1 M	65.1 M	57	normal (0.05)	Student's-t	63
	M4	1	100%	52 M	52 M	52	n=1	reported conc.	52
Vanadium	M1	6	50%	0.3 M	1 JM	0.65	gamma	approximate Gamma UCL	1.1
	M2	6	50%	0.25 M	1 M	0.63	non-parametric (0.05)	95% Chebyshev (Mean, Sd) UCL	1.3
	M3	6	100%	1 JM	1M	1.0	data all equal	not calc'd	1.0
	M4	1	0%	0.5 M	0.5 M	0.5	n=1	maximum RL	0.5
Shiner surfperch									
Arsenic	M1	9	100%	3.2	5.4	4.2	normal (0.05)	Student's-t UCL	4.7
	M2	6	100%	3.84	5.227	4.5	normal (0.05)	Student's-t UCL	5.0
	M3	6	100%	3.26	3.82	3.6	normal (0.05)	Student's-t UCL	3.7
	M4	6	100%	2.97	4.886	3.5	normal (0.05)	Student's-t UCL	4.1

Receptor and COPC	Area	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	UCL Type	Concentration Used in Risk Estimation (mg/kg dw)
				Minimum	Maximum	Mean			
Cadmium	M1	9	100%	0.046	0.0972	0.070	normal (0.05)	Student's-t UCL	0.079
	M2	6	100%	0.0527	0.075	0.064	normal (0.05)	Student's-t UCL	0.071
	M3	6	100%	0.0435	0.0749	0.059	normal (0.05)	Student's-t UCL	0.068
	M4	6	100%	0.038	0.0675	0.048	normal (0.05)	Student's-t UCL	0.056
Copper	M1	9	100%	3.7	8.4	6.6	normal (0.05)	Student's-t UCL	7.6
	M2	6	100%	5.354	8.672	6.5	normal (0.05)	Student's-t UCL	7.5
	M3	6	100%	6.067	8.902	7.6	normal (0.05)	Student's-t UCL	8.3
	M4	6	100%	2.26	6.667	4.9	normal (0.05)	Student's-t UCL	6.4
Vanadium	M1	6	66%	0.49	3.6	1.2	lognormal (0.05)	H-UCL	3.6
	M2	6	100%	1.1	4.62	2.5	normal (0.05)	Student's-t UCL	3.7
	M3	6	100%	1.3	2.1	1.7	normal (0.05)	Student's-t UCL	2.0
	M4	6	100%	0.87 J	1.2	1.1	normal (0.05)	Student's-t UCL	1.2

COPC – chemical of potential concern

dw – dry weight

JM – estimated calculated value

M – calculated value

N – number of samples

UCL – upper confidence limit

Table 4. Exposure concentrations in fish and crab tissue on a wet weight basis for the entire LDW

Receptor and COPC	N	Detection Frequency	Concentration (µg/kg ww)			Distribution Type	UCL Type	Concentration Used in Risk Estimation (µg/kg ww)
			Minimum	Maximum	Mean			
Juvenile chinook salmon								
Endrin	18	49%	0.09	6.5	0.76	nonparametric (0.05)	99% Chebyshev UCL	4.3[a]
Crab whole body								
Zinc	21	100%	24.6	37.3	31	normal (0.05)	Student's-t UCL	32
Methoxychlor	19	16%	0.4	90.2	6.9	nonparametric (0.05)	95% Chebyshev UCL	54
Total DDTs	19	100%	110	150	90	nonparametric (0.05)	Student's-t UCL	48
Total PCBs	25	100%	250	1,900	890	lognormal (0.05)	95% H-UCL	1,100
English sole[a]								
alpha-Endosulfan	24[b]	54%	2.1	6.6	3.7	gamma distribution (0.05)	approximate gamma UCL	4.2[a]
beta-Endosulfan	24[b]	17%	3.6	18	5.4	low detection frequency	maximum detected conc.	18[c]
Benzoic acid	24[b]	71%	1,900	6,500	4,400	nonparametric (0.05)	Student's-t UCL	5,100[a]
BEHP	24[b]	0%	66	3,600	1,300	all non-detects	maximum RL	3,600[c]
Dimethyl phthalate	24[b]	0%	290	580	440	all non-detects	maximum RL	580[c]
Di-n-butyl phthalate	24[b]	0%	290	1200	770	all non-detects	maximum RL	1,200[c]
Endrin	24[b]	17%	0.85	14	4.4	nonparametric (0.05)	95% Chebyshev UCL	14[c]
Total PCBs	45[b]	100%	450	4,700	2,200	gamma distribution (0.05)	approximate gamma UCL	2,600
PCB TEQ (fish)	7	100%	0.00103	0.00218	0.00154	normal (0.05)	Student's-t UCL	0.00188
Pacific staghorn sculpin								
alpha-endosulfan	24	46%	0.50	5.0	1.2	gamma distribution (0.05)	approximate gamma UCL	1.4[a]
beta-endosulfan	24	21%	0.55	6.5	2.6	nonparametric (0.05)	95% Chebyshev UCL	4.1[a]
Benzoic acid	24	96%	400	6,800	4,600	nonparametric (0.05)	95% Chebyshev UCL	7,000
BEHP	24	0%	490	5,000	3,500	all non-detects	maximum RL	5,000 [c]
Dimethyl phthalate	24	0%	40	400	140	all non-detects	maximum RL	400[c]
Di-n-butyl phthalate	24	4%	100	1,300	490	low detection frequency	maximum detected conc.	1,300[c]

Receptor and COPC	N	Detection Frequency	Concentration (µg/kg ww)			Distribution Type	UCL Type	Concentration Used in Risk Estimation (µg/kg ww)
			Minimum	Maximum	Mean			
Endrin	24	4%	0.5	36	2.3	low detection frequency	maximum detected conc.	36[c]
TBT	24	100%	23	80	32	nonparametric (0.05)	Student's-t UCL	36
Total PCBs	28	100%	430	2,800	900	nonparametric (0.05)	Student's-t UCL	1,100
Shiner surfperch								
Total PCBs	49	100%	350	18,400	1,800	nonparametric (0.05)	95% Chebyshev UCL	3,500
TBT	27	100%	33	180	58	nonparametric (0.05)	Student's-t UCL	69

[a] Calculated with ROS method because detection frequency was <85%.

[b] Includes 3 starry flounder whole-body/composite samples; starry flounder were used as a surrogate for English sole because insufficient numbers of English sole were collected in Area T4 during Phase 2 sampling (see Map A.2-3).

[c] Detection frequency was below 20% or in less than 4 samples. If detected, the maximum detected concentration was used. If not detected the maximum RL was used.

COPC – chemical of potential concern

N – number of samples

PCB – polychlorinated biphenyl

RL – reporting limit

ROS – regression on order statistic

TBT – tributyltin

UCL – upper confidence limit

ww – wet weight

Table 5. Exposure concentrations in Pacific staghorn sculpin tissue on a wet weight basis for four Pacific staghorn sculpin modeling areas

COPC	Area	N	Detection Frequency	Concentration (µg/kg ww)			Distribution Type	UCL Type	Concentration Used in Risk Estimation (µg/kg ww)
				Minimum	Maximum	Mean			
TBT	M1	6	100%	27	40	33	normal (0.05)	student's-t UCL	37
	M2	6	100%	26	39	32	normal (0.05)	student's-t UCL	36
	M3	6	100%	25	29	27	normal (0.05)	student's-t UCL	28
	M4	6	100%	23	80	35	non-parametric (0.05)	student's-t UCL	53
PCBs	M1	7	100%	580	860	720	normal (0.05)	student's-t UCL	800
	M2	7	100%	620	1,260	750	non-parametric (0.05)	student's-t UCL	920
	M3	7	100%	590	2,800	1,400	normal (0.05)	student's-t UCL	2,000
	M4	7	100%	430	1,330	730	normal (0.05)	student's-t UCL	940

COPC – chemical of potential concern

N – number of samples

PCB – polychlorinated biphenyl

RL – reporting limit

ROS – regression on order statistic

TBT – tributyltin

UCL – upper confidence limit

ww – wet weight

Table 6. Exposure concentrations in benthic invertebrates calculated for six exposure scenarios for spotted sandpiper on a dry weight basis

COPC	Exposure Scenario[a]	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
				Minimum	Maximum	Mean			
Arsenic	Area 1 – high	8	100%	5.3	26.09	11	nonlinear regression	95% UCL of regression	23
	Area 1 – high/poor	8	100%	5.3	26.09	11	nonlinear regression	95% UCL of regression	21
	Area 2 – high	3	100%	10.89	26.09	16	nonlinear regression	95% UCL of regression	25
	Area 2 – high/poor	4	100%	10.89	26.09	16	nonlinear regression	95% UCL of regression	21
	Area 3 – high	6	100%	7.88	21.01	13	nonlinear regression	95% UCL of regression	18
	Area 3 – high/poor	6	100%	7.88	21.01	13	nonlinear regression	95% UCL of regression	26
Bis(2-ethylhexyl) phthalate	Area 1 – high Area 1 – high/poor	8	25%	0.065	15	7.4	not enough samples to calculate UCL	maximum detected conc.	2.9[a]
	Area 2 – high	3	33%	8	14	12	not enough samples to calculate UCL	maximum detected conc.	14[a]
	Area 2 – high/poor	4	25%	8	19	14	not enough samples to calculate UCL	maximum detected conc.	14[a]
	Area 3 – high Area 3 – high/poor	6	0%	23	61	34	all non-detects	maximum RL	61[a]
Cadmium	Area 1 – high Area 1 – high/poor	8	100%	0.094	0.83	0.38	normal (0.05)	Student's-t UCL	0.56
	Area 2 – high	3	100%	0.193	0.401	0.32	not enough samples to calculate UCL	maximum detected conc.	0.40
	Area 2 – high/poor	4	100%	0.193	0.46	0.35	normal (0.05)	Student's-t UCL	0.49
	Area 3 – high Area 3 – high/poor	6	100%	0.261	0.831	0.46	normal (0.05)	Student's-t UCL	0.64

Table 6, cont. Exposure concentrations in benthic invertebrates calculated for six exposure scenarios for spotted sandpiper on a dry weight basis

COPC	Exposure Scenario[a]	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
				Minimum	Maximum	Mean			
Chromium	Area 1 – high Area 1 – high/poor	8	100%	1.0	3.1	2.0	normal (0.05)	Student's-t UCL	3.0
	Area 2 – high	3	100%	1.9	18.4	7.6	not enough samples to calculate UCL	maximum detected conc.	18
	Area 2 – high/poor	4	100%	1.9	58.2	20	normal (0.05)	Student's-t UCL	51
	Area 3 – high Area 3 – high/poor	6	100%	2.3	6.2	3.8	normal (0.05)	Student's-t UCL	5.0
Cobalt	Area 1 – high Area 1 – high/poor	4	100%	0.62	1.552	1.3	normal (0.05)	Student's-t UCL	1.8
	Area 2 – high	3	100%	1.459	2.02	1.8	not enough samples to calculate UCL	maximum detected conc.	2.0
	Area 2 – high/poor	4	100%	1.459	2.149	1.9	normal (0.05)	Student's-t UCL	2.2
	Area 3 – high Area 3 – high/poor	6	100%	1.059	2.408	1.8	normal (0.05)	Student's-t UCL	2.2
Copper	Area 1 – high Area 1 – high/poor	8	100%	31.24	170	86	normal (0.05)	Student's-t UCL	120
	Area 2 – high	3	100%	24.52	143	89	not enough samples to calculate UCL	maximum detected conc.	140
	Area 2 – high/poor	4	100%	24.52	143	80	normal (0.05)	Student's-t UCL	140
	Area 3 – high Area 3 – high/poor	6	100%	34.12	84.58	56	normal (0.05)	Student's-t UCL	70
DDTs	Area 1 – high Area 1 – high/poor	4	100%	0.077	0.31	0.17	normal (0.05)	Student's-t UCL	0.29
	Area 2 – high	3	100%	0.077	1.1	0.43	not enough samples to calculate UCL	maximum detected conc.	1.1
	Area 2 – high/poor	4	100%	0.077	1.1	0.40	normal (0.05)	Student's T-UCL	0.96
	Area 3 – high Area 3 – high/poor	6	100%	0.023	1.9	0.48	gamma distribution	gamma UCL	1.8

Table 6, cont. Exposure concentrations in benthic invertebrates calculated for six exposure scenarios for spotted sandpiper on a dry weight basis

COPC	Exposure Scenario[a]	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
				Minimum	Maximum	Mean			
Lead	Area 1 – high Area 1 – high/poor	8	100%	1.836	41	13	lognormal (0.05)	95% Chebyshev UCL	31
	Area 2 – high	3	100%	5.527	14.87	10	not enough samples to calculate UCL	maximum detected conc.	15
	Area 2 – high/poor	4	100%	5.527	217.9	62	gamma distribution (0.05)	approximate gamma UCL	660
	Area 3 – high Area 3 – high/poor	6	100%	1.888	8.048	3.7	normal (0.05)	Student's-t UCL	5.6
Mercury	Area 1 – high Area 1 – high/poor	8	88%	0.02	0.19	0.073	gamma distribution (0.05)	approximate gamma UCL	0.10
	Area 2 – high	3	100%	0.03	0.066	0.054	not enough samples to calculate UCL	maximum detected conc.	0.066
	Area 2 – high/poor	4	100%	0.03	0.066	0.054	normal (0.05)	Student's-t UCL	0.070
	Area 3 – high Area 3 – high/poor	6	100%	0.03	0.44	0.17	gamma distribution (0.05)	approximate gamma UCL	0.50
Nickel	Area 1 – high Area 1 – high/poor	8	100%	1.1	4.3	2.8	normal (0.05)	Student's-t UCL	3.5
	Area 2 – high	3	100%	1.96	5.84	3.3	not enough samples to calculate UCL	maximum detected conc.	5.8
	Area 2 – high/poor	4	100%	1.96	5.84	3.1	normal (0.05)	Student's-t UCL	5.3
	Area 3 – high Area 3 – high/poor	6	100%	2.33	9.907	5.0	gamma distribution (0.05)	approximate gamma UCL	7.4
Total PCBs	Area 1 – high	8	88%	0.59	2.3	1.2	linear regression	95% UCL of regression	1.5
	Area 1 – high/poor	8	88%	0.59	2.3	1.2	linear regression	95% UCL of regression	1.5
	Area 2 – high	3	67%	1.1	9.1	3.9	linear regression	95% UCL of regression	5.9
	Area 2 – high/poor	4	75%	1.1	9.1	3.3	linear regression	95% UCL of regression	3.8
	Area 3 – high	6	100%	0.61	21	4.5	linear regression	95% UCL of regression	2.7
	Area 3 – high/poor	6	100%	0.61	21	4.5	linear regression	95% UCL of regression	3.4

Table 6, cont. Exposure concentrations in benthic invertebrates calculated for six exposure scenarios for spotted sandpiper on a dry weight basis

COPC	Exposure Scenario[a]	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
				Minimum	Maximum	Mean			
PCB TEQ (bird)	Area 1 – high	1	100%	1.62×10^{-4}	1.62×10^{-4}	1.6×10^{-4}	not enough samples to calculate UCL	maximum detected conc.	1.6×10^{-4}
	Area 2 – high	1	100%	5.63×10^{-4}	5.63×10^{-4}	5.6×10^{-4}	not enough samples to calculate UCL	maximum detected conc.	5.6×10^{-4}
	Area 2 – high/poor	1	100%	5.63×10^{-4}	5.63×10^{-4}	5.6×10^{-4}	not enough samples to calculate UCL	maximum detected conc.	5.6×10^{-4}
	Area 3 – high Area 3 – high/poor	3	100%	2.21×10^{-5}	3.90×10^{-4}	1.7×10^{-4}	not enough samples to calculate UCL	maximum detected conc.	3.9×10^{-4}
Selenium	Area 1 – high Area 1 – high/poor	4	100%	1.0	1.84	1.5	normal (0.05)	Student's-t UCL	1.9
	Area 2 – high	3	100%	1.13	1.6	1.4	not enough samples to calculate UCL	maximum detected conc.	1.6
	Area 2 – high/poor	4	100%	1.13	1.79	1.5	normal (0.05)	Student's-t UCL	1.8
	Area 3 – high Area 3 – high/poor	6	100%	0.80	1.62	1.2	normal (0.05)	Student's-t UCL	1.4
Vanadium	Area 1 – high Area 1 – high/poor	4	100%	3.1	5.9	4.9	normal (0.05)	Student's-t UCL	6.4
	Area 2 – high	3	100%	5.1	10.2	7.5	not enough samples to calculate UCL	maximum detected conc.	10
	Area 2 – high/poor	4	100%	5.1	10.2	7.2	normal (0.05)	Student's-t UCL	9.8
	Area 3 – high Area 3 – high/poor	6	100%	4.6	9.8	7.9	normal (0.05)	Student's-t UCL	9.6
Zinc	Area 1 – high Area 1 – high/poor	8	100%	44	295	130	normal (0.05)	Student's-t UCL	190
	Area 2 – high	3	100%	208	295	260	not enough samples to calculate UCL	maximum detected conc.	300
	Area 2 – high/poor	4	100%	208	384	290	normal (0.05)	Student's-t UCL	380
	Area 3 – high Area 3 – high/poor	6	100%	102	346	190	normal (0.05)	Student's-t UCL	270

Table 6, cont. Exposure concentrations in benthic invertebrates calculated for six exposure scenarios for spotted sandpiper on a dry weight basis

[a] Six exposure scenarios were evaluated; in each of three exposure areas, foraging in high-quality habitat only and foraging in both high- and poor-quality habitat were evaluated. These exposure scenarios are described in detail in Section A.5.1.3.1.

[b] Detection frequency was below 20% or in less than 4 samples. If detected, the maximum detected concentration was used. If not detected the maximum RL was used.

COPC – chemical of potential concern

dw – dry weight

N – number of samples

PCB – polychlorinated biphenyl

TEQ – toxic equivalent

UCL – upper confidence limit

Table 7. Exposure concentrations in benthic invertebrates calculated for intertidal areas on a dry-weight basis

COPC	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
			Minimum	Maximum	Mean			
Arsenic	17	100%	5.3	26.09	12	nonlinear regression	95% UCL of regression	22
						gamma distribution (0.05)	approximate gamma UCL	14
Cadmium	17	100%	0.094	0.831	0.40	normal (0.05)	Student's-t UCL	0.50
Chromium	17	100%	1.0	58.2	7.0	nonparametric (0.05)	95% Chebyshev UCL	20
Copper	17	100%	24.52	170	73	gamma distribution (0.05)	approximate gamma UCL	93
Vanadium	13	100%	3.1	10.2	7.0	normal (0.05)	Student's-t UCL	8.1

COPC – chemical of potential concern

dw – dry weight

N – number of samples

UCL – upper confidence limit

Table 8. Exposure concentrations in benthic invertebrate tissue calculated for four Pacific staghorn sculpin modeling areas on a dry-weight basis

COPC	Area	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
				Minimum	Maximum	Mean			
Arsenic	M1	10	100%	5.3	116	19	nonlinear regression	95% UCL of regression	23
	M2	6	100%	9.655	31.56	17	nonlinear regression	95% UCL of regression	25
	M3	4	100%	11.9	22.41	16	nonlinear regression	95% UCL of regression	23
	M4	4	100%	7.88	21.01	12	nonlinear regression	95% UCL of regression	18
Cadmium	M1	10	100%	0.094	1.3 J	0.57	normal (0.05)	Student's-t UCL	0.81
	M2	6	100%	0.193 J	0.57 J	0.38	normal (0.05)	Student's-t UCL	0.48
	M3	4	100%	0.309	0.831	0.51	normal (0.05)	Student's-t UCL	0.77
	M4	4	100%	0.261	0.560	0.38	normal (0.05)	Student's-t UCL	0.54
Copper	M1	10	100%	31.24 J	170	90	normal (0.05)	Student's-t UCL	110
	M2	6	100%	24.52 J	151	90	normal (0.05)	Student's-t UCL	130
	M3	4	100%	37.23 J	66.11 J	51	normal (0.05)	Student's-t UCL	65
	M4	4	100%	34.12 J	84.58 J	55	normal (0.05)	Student's-t UCL	81
Vanadium	M1	6	100%	3.10	16.5	8.3	normal (0.05)	Student's-t UCL	12
	M2	6	100%	5.10	21.3	12	normal (0.05)	Student's-t UCL	18
	M3	4	100%	6.30	22.2	12	normal (0.05)	Student's-t UCL	20
	M4	4	100%	4.60	9.80	7.4	normal (0.05)	Student's-t UCL	10

COPC – chemical of potential concern

dw – dry weight

J – estimated concentration

N – number of samples

UCL – upper confidence limit

Table 9. Exposure concentrations in benthic invertebrates calculated for the entire LDW on a dry-weight basis

COPC	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
			Minimum	Maximum	Mean			
Arsenic	24	100%	5.3	116	17	nonlinear regression	95th UCL of regression	24
						non-parametric (0.05)	95% Chebyshev (Mean, SD) UCL	37
Cadmium	24	100%	0.094	1.3 J	0.48	gamma distribution (0.05)	approximate gamma UCL	0.60
Chromium	24	100%	1.0	58.2	8.2	lognormal (0.05)	95% H-UCL	12
Copper	24	100%	24.52 J	170	78	normal (0.05)	Student's-t UCL	92
Lead	24	100%	1.836	217.9	19	data are lognormal (0.05)	95% Chebyshev (MVUE) UCL	31
total alkylated + non-alkylated PAHs	24	88%	0.18	36.1 J	6.8	gamma distribution (0.05)	approximate gamma UCL	11
Total PCBs	24	96%	0.59	21	2.5	linear regression	95% UCL of regression	2.3
Selenium	20	100%	0.80	3.63	1.8	normal (0.05)	Student's-t UCL	2.1
Silver	24	100%	0.156	2.496	0.48	gamma distribution (0.05)	approximate gamma UCL	0.62
Vanadium	20	100%	3.1	22.2	10	gamma distribution (0.05)	approximate gamma UCL	12
TBT	20	95%	0.03	0.55	0.24	nonlinear regression	95% UCL of regression	0.38

[a] Total PAHs included: indeno (1,2,3-cd)perylene, benz(a)anthracene, benzo(a)pyrene, benzo(k)fluoranthene, fluoranthene, benzo(g,h,i)perylene, benzo(b)fluoranthene, chrysene, dibenzo(a,h)anthracene, pyrene.

COPC – chemical of potential concern

dw – dry weight

N – number of samples

UCL – upper confidence limit

Table 10. Exposure concentrations in sediment calculated for six exposure scenarios for sandpiper on a dry-weight basis

COPC	Exposure Scenario[a]	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
				Minimum	Maximum	Mean			
Arsenic	Area 1 – high	39	95%	3.13	161	17	nonparametric (0.05)	95% Chebyshev UCL	36
	Area 1 – high/poor	55	95%	1.2	161	15	lognormal	95% H-UCL	18
	Area 2 – high	25	92%	4.0	161	22	nonparametric (0.05)	95% Chebyshev UCL	49
	Area 2 – high/poor	49	96%	1.2	161	15	lognormal	95% H-UCL	18
	Area 3 – high	87	92%	2.5	79.4	10	nonparametric (0.05)	95% Chebyshev UCL	20
	Area 3 – high/poor	150	93%	2.5	1,100	20	nonparametric (0.05)	95% Chebyshev UCL	60
Bis(2-ethylhexyl) phthalate	Area 1 – high	39	85%	0.0054	0.65	0.089	gamma distribution (0.05)	approximate gamma UCL	0.12
	Area 1 – high/poor	55	87%	0.0054	14	0.63	nonparametric (0.05)	97.5% Chebyshev UCL	2.5
	Area 2 – high	25	96%	0.02	5.1	0.94	lognormal (0.05)	95% Chebyshev UCL	2.7
	Area 2 – high/poor	49	94%	0.01	5.1	0.54	nonparametric (0.05)	99% Chebyshev UCL	2.1
	Area 3 – high	92	78%	0.014	8.6	0.40	lognormal (0.05)	95% H-UCL	0.41[b]
	Area 3 – high/poor	145	85%	0.014	8.6	0.40	lognormal (0.05)	95% H-UCL	0.41
Cadmium	Area 1 – high	39	77%	0.050	1.0	0.36	gamma distribution (0.05)	approximate gamma UCL	0.45[b]
	Area 1 – high/poor	55	78%	0.03	2	0.51	gamma distribution (0.05)	approximate gamma UCL	0.65[b]
	Area 2 – high	25	96%	0.070	2.7	0.84	gamma distribution (0.05)	approximate gamma UCL	1.0
	Area 2 – high/poor	49	78%	0.030 J	2.7	0.52	data are lognormal (0.05)	95% H-UCL	0.93[b]
	Area 3 – high	80	61%	0.068	5.2	0.52	data are lognormal (0.05)	95% H-UCL	0.62[b]
	Area 3 – high/poor	143	69%	0.068	92	2.5	nonparametric (0.05)	97.5% Chebyshev UCL	7.8[b]
Chromium	Area 1 – high	39	100%	9.84	60.5	28	data are normal (0.05)	Student's-t UCL	32
	Area 1 – high/poor	55	100%	4.8	60.5	27	data are normal (0.05)	Student's-t UCL	30
	Area 2 – high	25	100%	9.0	122 J	35	gamma distribution (0.05)	approximate gamma UCL	43
	Area 2 – high/poor	49	100%	4.8	122 J	28	gamma distribution (0.05)	approximate gamma UCL	32
	Area 3 – high	87	100%	9.19	76 J	29	nonparametric (0.05)	Student's-t UCL	31
	Area 3 – high/poor	150	100%	9.19	1,100	67	nonparametric (0.05)	95% Chebyshev UCL	120

Table 10, cont. Exposure concentrations in sediment calculated for six exposure scenarios for sandpiper on a dry-weight basis

COPC	Exposure Scenario[a]	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
				Minimum	Maximum	Mean			
Cobalt	Area 1 – high	25	100%	2.82	18.7	7.4	gamma distribution (0.05)	approximate gamma UCL	9.0
	Area 1 – high/poor	28	100%	2.82	18.7	7.1	gamma distribution (0.05)	approximate gamma UCL	8.0
	Area 2 – high	22	100%	3.0	18.7	8.2	data are normal (0.05)	Student's-t UCL	9.5
	Area 2 – high/poor	38	100%	3.0	18.7	7.3	gamma distribution (0.05)	approximate gamma UCL	8.0
	Area 3 – high	70	100%	3.48	12	7.7	gamma distribution (0.05)	approximate gamma UCL	8.0
	Area 3 – high/poor	106	100%	3.48	37	9.2	nonparametric (0.05)	Student's-t UCL	10
Copper	Area 1 – high	39	100%	7.90	365	74	gamma distribution (0.05)	approximate gamma UCL	94
	Area 1 – high/poor	55	100%	7.90	365	65	gamma distribution (0.05)	approximate gamma UCL	80
	Area 2 – high	25	100%	16	365	87	gamma distribution (0.05)	approximate gamma UCL	120
	Area 2 – high/poor	49	100%	11.5	365	64	lognormal (0.05)	95% H-UCL	79
	Area 3 – high	87	100%	17.2	290	45	nonparametric (0.05)	95% Chebyshev UCL	63
	Area 3 – high/poor	150	100%	17.2	12,000	200	nonparametric (0.05)	97.5% Chebyshev UCL	730
Total DDTs	Area 1 – high	20	50%	7.2×10^{-4}	0.0216	0.0050	nonparametric (0.05)	99% Chebyshev UCL	0.017
	Area 1 – high/poor	24	54%	7.2×10^{-4}	0.0216	0.0055	nonparametric (0.05)	99% Chebyshev UCL	0.018
	Area 2 – high	13	69%	9.5×10^{-4}	2.9	0.28	lognormal (0.05)	99% Chebyshev (MVUE) UCL	1.0
	Area 2 – high/poor	19	63%	9.5×10^{-4}	2.9	0.19	lognormal (0.05)	99% Chebyshev (MVUE) UCL	0.52
	Area 3 – high	15	53%	9.5×10^{-4}	0.047	0.012	gamma distribution (0.05)	approximate gamma UCL	0.024
	Area 3 – high/poor	18	56%	9.5×10^{-4}	0.047	0.012	gamma distribution (0.05)	approximate gamma UCL	0.020
Lead	Area 1 – high	39	100%	7.94 J	400	70	gamma distribution (0.05)	approximate gamma UCL	90
	Area 1 – high/poor	55	100%	7.94 J	400	70	gamma distribution (0.05)	approximate gamma UCL	90
	Area 2 – high	25	100%	8.2	615	110	gamma distribution (0.05)	approximate gamma UCL	160
	Area 2 – high/poor	49	100%	8.2	615	70	lognormal (0.05)	95% H-UCL	96
	Area 3 – high	87	100%	6.3	533	50	nonparametric (0.05)	95% Chebyshev UCL	90
	Area 3 – high/poor	150	100%	6.3	23,000	300	nonparametric (0.05)	97.5% Chebyshev UCL	1,000

Table 10, cont. Exposure concentrations in sediment calculated for six exposure scenarios for sandpiper on a dry-weight basis

COPC	Exposure Scenario[a]	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
				Minimum	Maximum	Mean			
Mercury	Area 1 – high	39	87%	0.015	0.63	0.13	gamma distribution (0.05)	approximate gamma UCL	0.17
	Area 1 – high/poor	47	85%	0.01	0.63	0.13	gamma distribution (0.05)	approximate gamma UCL	0.16
	Area 2 – high	32	91%	0.025	2.46	0.30	gamma distribution (0.05)	approximate gamma UCL	0.43
	Area 2 – high/poor	49	84%	0.021	2.46	0.22	lognormal (0.05)	95% H-UCL	0.29
	Area 3 – high	87	82%	0.025	4.6 J	0.20	nonparametric (0.05)	97.5% Chebyshev UCL	0.60[b]
	Area 3 – high/poor	149	83%	0.025	4.6 J	0.20	nonparametric (0.05)	95% Chebyshev UCL	0.40[b]
Nickel	Area 1 – high	39	100%	6	37	19	data are normal (0.05)	Student's-t UCL	21
	Area 1 – high/poor	48	98%	6	37	20	nonparametric (0.05)	95% Chebyshev UCL	30
	Area 2 – high	25	100%	8.9	39	19	data are normal (0.05)	Student's-t UCL	22
	Area 2 – high/poor	42	98%	8.9	39	17	gamma distribution (0.05)	approximate gamma UCL	20
	Area 3 – high	85	100%	7.66	52	20	gamma distribution (0.05)	approximate gamma UCL	22
	Area 3 – high/poor	141	100%	7.66	910	50	nonparametric (0.05)	95% Chebyshev UCL	90
Total PCBs	Area 1 – high	56	93%	0.0026	0.81	0.15	SWAC	95% UCL	0.34
	Area 1 – high/poor	81	91%	0.0022	0.81	0.14	SWAC	95% UCL	0.33
	Area 2 – high	50	98%	0.02	25	2.8	SWAC	95% UCL	2.5
	Area 2 – high/poor	88	95%	0.0047	25	1.6	SWAC	95% UCL	1.5
	Area 3 – high	129	91%	0.0061	15	1.3	SWAC	95% UCL	0.72
	Area 3 – high/poor	228	95%	0.0061	110	3.3	SWAC	95% UCL	1.1
PCB TEQ (bird)	Area 1 – high	8	100%	6.46×10^{-7}	5.61×10^{-5}	2.3×10^{-5}	normal (0.05)	Student's-t UCL	3.7×10^{-5}
	Area 1 – high/poor	9	100%	6.46×10^{-7}	5.61×10^{-5}	2.1×10^{-5}	normal (0.05)	Student's-t UCL	3.4×10^{-5}
	Area 2 – high	7	100%	6.22×10^{-6}	2.64×10^{-3}	4.6×10^{-4}	gamma distribution (0.05)	Adjusted gamma UCL	4.3×10^{-3}
	Area 2 – high/poor	10	100%	1.61×10^{-6}	2.64×10^{-3}	3.2×10^{-4}	gamma distribution (0.05)	Adjusted gamma UCL	1.8×10^{-3}
	Area 3 – high	8	100%	2.53×10^{-6}	1.33×10^{-4}	3.6×10^{-5}	gamma distribution (0.05)	approximate gamma UCL	9.6×10^{-5}
	Area 3 – high/poor	11	100%	2.53×10^{-6}	6.21×10^{-3}	6.6×10^{-4}	lognormal (0.05)	99% Chebyshev UCL	3.0×10^{-3}

Table 10, cont. Exposure concentrations in sediment calculated for six exposure scenarios for sandpiper on a dry-weight basis

COPC	Exposure Scenario[a]	N	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
				Minimum	Maximum	Mean			
Selenium	Area 1 – high	28	43%	0.2	10	2.5	normal (0.05)	Student's-t UCL	0.59[b]
	Area 1 – high/poor	36	36%	0.2	10	2.5	gamma distribution (0.05)	approximate gamma UCL	0.56[b]
	Area 2 – high	22	55%	0.3	9	3.2	nonparametric (0.05)	99% Chebyshev UCL	7.2[b]
	Area 2 – high/poor	38	42%	0.3	10	3.3	nonparametric (0.05)	95% Chebyshev UCL	3.0[b]
	Area 3 – high	60	37%	0.5	13	5.5	nonparametric (0.05)	97.5% Chebyshev UCL	7.0[b]
	Area 3 – high/poor	96	33%	0.5	20	5.9	nonparametric (0.05)	97.5% Chebyshev UCL	6.6[b]
Vanadium	Area 1 – high	25	100%	34.8	72.6	50	lognormal (0.05)	Student's-t UCL	53
	Area 1 – high/poor	28	100%	34.8	72.6	49	nonparametric (0.05)	Student's-t UCL	52
	Area 2 – high	22	100%	15	72.6	53	normal (0.05)	Student's-t UCL	58
	Area 2 – high/poor	38	100%	15	72.6	52	normal (0.05)	Student's-t UCL	55
	Area 3 – high	70	100%	27.9	83	55	normal (0.05)	Student's-t UCL	57
	Area 3 – high/poor	106	100%	27.9	87	57	normal (0.05)	Student's-t UCL	59
Zinc	Area 1 – high	39	100%	31	607	150	gamma distribution (0.05)	approximate gamma UCL	180
	Area 1 – high/poor	55	98%	19.2	607	140	gamma distribution (0.05)	approximate gamma UCL	170
	Area 2 – high	25	100%	28	607	190	gamma distribution (0.05)	approximate gamma UCL	240
	Area 2 – high/poor	49	98%	19.2	607	140	lognormal (0.05)	95% H-UCL	180
	Area 3 – high	87	100%	35.6	343	110	nonparametric (0.05)	95% Chebyshev UCL	150
	Area 3 – high/poor	150	100%	35.6	6,400	320	nonparametric (0.05)	97.5% Chebyshev UCL	710

[a] Six exposure scenarios were evaluated; in each of three exposure areas, foraging in high-quality habitat only and foraging in both high- and poor-quality habitat were evaluated. These exposure scenarios are described in detail in Section A.5.1.3.1.

[b] Calculated with ROS method because detection frequency was < 85%

[c] Detection frequency was below 20% or in less than 4 samples. If detected, the maximum detected concentration was used. If not detected the maximum RL was used.

COPC – chemical of potential concern

dw – dry weight

N – number of samples

na – not available

PCB – polychlorinated biphenyl

ROS – regression on order statistic

SWAC – spatially weighted average concentration

UCL – upper confidence limit

Table 11. Exposure concentrations in sediment calculated for intertidal areas on a dry-weight basis

COPC	N[a]	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
			Minimum	Maximum	Mean			
Arsenic	357	86%	1.2	1,100 J	16	nonparametric (0.05)	95% Chebyshev UCL	30
Cadmium	348	63%	0.0131	120	1.7	nonparametric (0.05)	97.5% Chebyshev UCL	4.7[b]
Chromium	357	100%	4.8	1,100	50	nonparametric (0.05)	95% Chebyshev UCL	70
Cobalt	232	100%	2.82	140	9.0	nonparametric (0.05)	Student's-t UCL	10
Copper	357	100%	5	12,000	150	nonparametric (0.05)	95% Chebyshev UCL	400
Total DDTs	75	45%	5.7×10^{-5}	2.9	0.049	nonparametric (0.05)	97.5% Chebyshev UCL	0.29[b]
Lead	357	100%	2	23,000	190	nonparametric (0.05)	97.5% Chebyshev UCL	600
Mercury	356	78%	0.009	5.6	0.17	nonparametric (0.05)	97.5% Chebyshev UCL	0.40[b]
Total PCBs	552	92%	0.0022 J	220	2.3	SWAC	95 %UCL	0.98
PCB TEQ (mammal)	34	100%	9.08×10^{-8}	1.38×10^{-3}	6.1×10^{-5}	lognormal (0.05)	97.5% Chebyshev UCL	1.5×10^{-4}
PCB TEQ (bird)	34	100%	6.5×10^{-7}	0.0062	3.5×10^{-4}	lognormal (0.05)	97.5% Chebyshev UCL	9.2×10^{-4}
Selenium	249	29%	0.0958	20	2.5	nonparametric (0.05)	97.5% Chebyshev UCL	3.9[b]
Silver	345	50%	0.020	270	1.6	nonparametric (0.05)	97.5% Chebyshev UCL	6.8[b]
Total PAHs	345	97%	0.0095	128	4.0	nonparametric (0.05)	97.5% Chebyshev UCL	7.0
Vanadium	232	100%	15	150	55	gamma distribution (0.05)	approximate gamma UCL	57

[a] Number of intertidal sediment samples in the baseline surface sediment dataset as of June 26, 2006, prior to an update to the baseline dataset.

[b] Calculated with ROS method because detection frequency was <85%.

COPC – chemical of potential concern

dw – dry weight

N – number of samples

na – not available

PCB – polychlorinated biphenyl

ROS – regression on order statistic

SWAC – spatially weighted average concentration

UCL – upper confidence limit

Table 12. Exposure concentrations in sediment calculated for four Pacific staghorn sculpin modeling areas on a dry-weight basis

COPC	Modeling Area	N[a]	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
				Minimum	Maximum	Mean			
Arsenic	M1	222	98%	1.55	725	19	non-parametric (0.05)	95% Chebyshev (mean, sd) UCL	34
	M2	152	98%	1.2	807	22	non-parametric (0.05)	95% Chebyshev (mean, sd) UCL	46
	M3	239	81%	2.5	1,100	18	non-parametric (0.05)	95% Chebyshev (mean, sd) UCL	40
	M4	186	95%	1.5	51	11	gamma	approximate gamma UCL	11
Cadmium	M1	217	88%	0.050	11.7	0.78	non-parametric (0.05)	95% Chebyshev (mean, sd) UCL	1.1
	M2	152	84%	0.030	3	0.44	non-parametric (0.05)	95% Chebyshev (mean, sd) UCL	0.55
	M3	239	75%	0.060	120	2.3	non-parametric (0.05)	97.5% Chebyshev (Mean, Sd) UCL	6.8
	M4	177	37%	0.030	1	0.26	non-parametric (0.05)	95% Chebyshev (mean, sd) UCL	0.33
Copper	M1	222	100%	7.9	495	84	non-parametric (0.05)	95% Chebyshev (mean, sd) UCL	97
	M2	152	100%	10	1,420	100	non-parametric (0.05)	95% Chebyshev (mean, sd) UCL	170
	M3	239	100%	14	12,000	200	non-parametric (0.05)	95% Chebyshev (mean, sd) UCL	510
	M4	186	100%	5	89.9	38	gamma	Approximate Gamma UCL	41
Vanadium	M1	128	100%	27.7	100	61	normal (0.05)	Student's-t UCL	63
	M2	132	100%	15	86	59	normal (0.05)	Student's-t UCL	60
	M3	182	100%	30	150	58	gamma	approximate gamma UCL	60
	M4	100	100%	27.9	89.6	58	normal (0.05)	Student's-t UCL	60

[a] Number of intertidal sediment samples in the baseline surface sediment dataset as of June 26, 2006, prior to an update to the baseline dataset.

[b] Calculated with ROS method because detection frequency was <85%.

COPC – chemical of potential concern
dw – dry weight
N – number of samples
na – not available
PCB – polychlorinated biphenyl
ROS – regression on order statistic
SWAC – spatially weighted average concentration
UCL – upper confidence limit

Table 13. Exposure concentrations for sediment calculated for the entire LDW on a dry-weight basis

COPC	N[a]	Detection Frequency	Concentration (mg/kg dw)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/kg dw)
			Minimum	Maximum	Mean			
Arsenic	814	93%	1.2	1,100	20	nonparametric (0.05)	95% Chebyshev UCL	30
Cadmium	797	71%	0.02	120	1.0	nonparametric (0.05)	97.5% Chebyshev UCL	2.4[b]
Chromium	811	100%	4.8	1,100 J	40	nonparametric (0.05)	95% Chebyshev UCL	50
Cobalt	556	100%	2.82	140	9.5	nonparametric (0.05)	Student's-t UCL	10
Copper	814	100%	5	12,000 J	110	nonparametric (0.05)	95% Chebyshev UCL	200
Total DDTs	197	40%	1.3×10^{-5}	2.9	0.025	nonparametric (0.05)	97.5% Chebyshev UCL	0.12[b]
Lead	814	100%	2	23,000	110	nonparametric (0.05)	97.5% Chebyshev UCL	300
Mercury	831	86%	0.01	4.6 J	0.21	nonparametric (0.05)	95% Chebyshev UCL	0.30
Total PCBs	1288	93%	2.8×10^{-4}	220	1.0	SWAC	95% UCL	0.72
PCB TEQ (mammal)	48	100%	9.08×10^{-8}	1.38×10^{-3}	4.6×10^{-5}	lognormal (0.05)	95% Chebyshev UCL	7.2×10^{-5}
PCB TEQ (bird)	48	100%	6.46×10^{-7}	0.00621	2.9×10^{-4}	lognormal (0.05)	95% Chebyshev UCL	5.4×10^{-4}
Selenium	629	44%	0.1	28	4.3	nonparametric (0.05)	97.5% Chebyshev UCL	5.0[b]
Silver	782	62%	0.02	270	1.0	nonparametric (0.05)	97.5% Chebyshev UCL	3.4[b]
Total PAHs	790	97%	0.0095	128	4.5	nonparametric (0.05)	97.5% Chebyshev UCL	6.0
Vanadium	556	100%	15	150	59	nonparametric (0.05)	Student's-t UCL	60

[a] Number of sediment samples in the baseline surface sediment dataset as of June 26, 2006, prior to an update to two locations at the Norfolk site.

[b] Calculated with ROS method because detection frequency was <85%.

COPC – chemical of potential concern
dw – dry weight
J – estimated value
N – number of samples
na – not available
PAH – polycyclic aromatic hydrocarbon
PCB – polychlorinated biphenyl
RL – reporting limit
ROS – regression on order statistic
SWAC – spatially weighted average concentration
TEQ – toxic equivalent
UCL – upper confidence limit

Table 14. Exposure concentrations in water calculated for three exposure areas for spotted sandpiper

COPC[a]	Exposure Area[b]	N	Detection Frequency	Concentration (mg/L)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/L)
				Minimum	Maximum	Mean			
Arsenic (total)	1	167	100%	2.92×10^{-4}	0.00153	9.3×10^{-4}	nonparametric (0.05)	Student's-t UCL	9.7×10^{-4}
	2	167	100%	2.82×10^{-4}	0.00157	8.8×10^{-4}	nonparametric (0.05)	Student's-t UCL	9.3×10^{-4}
	3	56	100%	1.83×10^{-4}	8.68×10^{-4}	4.9×10^{-4}	normal (0.05)	Student's-t UCL	5.3×10^{-4}
BEHP	1	39	18%	7.0×10^{-5}	0.00351	4.3×10^{-4}	low detection frequency	maximum detected conc.	0.0035[c]
	2	42	19%	7.0×10^{-5}	0.0238	8.6×10^{-4}	low detection frequency	maximum detected conc.	0.024[c]
	3	13	31%	1.4×10^{-4}	2.5×10^{-4}	1.7×10^{-4}	normal (0.05)	Student's-t UCL	1.8×10^{-4} [b]
Cadmium (total)	1	173	100%	8.8×10^{-6}	7.8×10^{-5}	4.7×10^{-5}	nonparametric (0.05)	95% Chebyshev UCL	5.4×10^{-5}
	2	173	99%	3.35×10^{-6}	3.91×10^{-4}	4.3×10^{-5}	nonparametric (0.05)	95% Chebyshev UCL	5.5×10^{-5}
	3	56	88%	3.4×10^{-6}	3.2×10^{-5}	1.5×10^{-5}	normal (0.05)	Student's-t UCL	1.6×10^{-5}
Chromium (total)	1	165	100%	2.4×10^{-4}	0.00174	5.7×10^{-4}	lognormal (0.05)	Student's-t UCL	6.1×10^{-4}
	2	157	100%	2.77×10^{-4}	0.00232	6.8×10^{-4}	nonparametric (0.05)	Student's-t UCL	7.3×10^{-4}
	3	58	100%	2.56×10^{-4}	0.00237	9.8×10^{-4}	gamma (0.05)	approximate gamma UCL	0.0011
Cobalt (total)	1	155	100%	3.34×10^{-5}	5.75×10^{-4}	1.8×10^{-4}	nonparametric (0.05)	95% Chebyshev UCL	2.2×10^{-4}
	2	155	100%	3.11×10^{-5}	7.72×10^{-4}	2.3×10^{-4}	gamma (0.05)	approximate gamma UCL	2.5×10^{-4}
	3	49	100%	1.23×10^{-4}	0.00133	4.0×10^{-4}	gamma (0.05)	approximate gamma UCL	4.6×10^{-4}
Copper (total)	1	166	100%	5.36×10^{-4}	0.00583	0.0014	nonparametric (0.05)	Student's-t UCL	0.0015
	2	167	99%	3.215×10^{-4}	0.00403	0.0015	gamma (0.05)	approximate gamma UCL	0.0016
	3	58	100%	7.28×10^{-4}	0.00424	0.0021	normal (0.05)	Student's-t UCL	0.0023
DDTs	1	nd	nd	nd	nd	nd	nd	nd	nd
	2	nd	nd	nd	nd	nd	nd	nd	nd
	3	nd	nd	nd	nd	nd	nd	nd	nd
Lead (total)	1	171	100%	4.59×10^{-5}	0.00145	3.5×10^{-4}	lognormal (0.05)	95% H-UCL	3.8×10^{-4}
	2	167	100%	5.7×10^{-5}	0.00157	4.0×10^{-4}	nonparametric (0.05)	95% Chebyshev UCL	4.9×10^{-4}
	3	58	100%	1.43×10^{-4}	0.00281	7.8×10^{-4}	gamma (0.05)	approximate gamma UCL	8.9×10^{-4}

COPC[a]	Exposure Area[b]	N	Detection Frequency	Concentration (mg/L)			Distribution Type	Statistic Used	Concentration Used in Risk Estimation (mg/L)
				Minimum	Maximum	Mean			
Mercury (total)	1	15	60%	5.6×10^{-7}	3.4×10^{-6}	1.5×10^{-6}	gamma (0.05)	approximate gamma UCL	2.0×10^{-6} [b]
	2	6	0%	1.0×10^{-4}	2.0×10^{-4}	1.0×10^{-4}	all non-detects	maximum RL	2.0×10^{-4} [c]
	3	8	75%	1.04×10^{-6}	6.9×10^{-6}	3.0×10^{-6}	normal (0.05)	Student's-t UCL	4.3×10^{-6} [b]
Nickel (total)	1	147	99%	1.74×10^{-4}	0.00211	6.7×10^{-4}	nonparametric (0.05)	Student's-t UCL	7.1×10^{-4}
	2	143	96%	1.455×10^{-4}	0.00288	7.6×10^{-4}	nonparametric (0.05)	Student's-t UCL	8.2×10^{-4}
	3	55	100%	4.34×10^{-4}	0.00291	0.0011	lognormal (0.05)	95% H-UCL	0.0013
Selenium (total)	1	151	2%	6.5×10^{-5}	2.7×10^{-4}	7.7×10^{-5}	low detection frequency	maximum detected conc.	2.7×10^{-4} [c]
	2	155	0%	6.5×10^{-5}	8.0×10^{-5}	7.4×10^{-5}	all non-detects	maximum RL	1.6×10^{-4} [c]
	3	54	0%	7.0×10^{-5}	8.0×10^{-5}	7.5×10^{-5}	all non-detects	maximum RL	1.6×10^{-4} [c]
PCBs	1	7	100%	2.0×10^{-7}	2.0×10^{-6}	1.1×10^{-6}	normal (0.05)	Student's-t UCL	1.7×10^{-6}
	2	8	100%	1.1×10^{-7}	3.1×10^{-6}	1.3×10^{-6}	normal (0.05)	Student's-t UCL	1.9×10^{-6}
	3	8	100%	1.1×10^{-7}	3.1×10^{-6}	1.3×10^{-6}	normal (0.05)	Student's-t UCL	1.9×10^{-6}
PCB TEQ (bird)	1	nd	nd	nd	nd	nd	nd	nd	nd
	2	nd	nd	nd	nd	nd	nd	nd	nd
	3	nd	nd	nd	nd	nd	nd	nd	nd
Vanadium (Total)	1	133	100%	2.67×10^{-4}	0.00296	0.0014	normal (0.05)	Student's-t UCL	0.0014
	2	125	100%	2.2×10^{-4}	0.00399	0.0014	gamma (0.05)	approximate gamma UCL	0.0015
	3	46	100%	3.15×10^{-4}	0.00357	0.0015	lognormal (0.05)	95% H-UCL	0.0019
Zinc (total)	1	173	100%	7.0×10^{-4}	0.00834	0.0028	lognormal (0.05)	Student's-t UCL	0.0030
	2	173	100%	0.00108	0.00662	0.0028	nonparametric (0.05)	Student's-t UCL	0.0030
	3	56	100%	9.79×10^{-4}	0.00904	0.0042	normal (0.05)	Student's-t UCL	0.0046

[a] Metals were analyzed as total concentrations in water.

[b] The thee exposure areas are described in Section A.5.1.3.1 and shown on Figure 5-1.

[c] Detection frequency was below 20% or in less than 4 samples. If detected, the maximum detected concentration was used. If not detected the maximum RL was used.

[d] Calculated with ROS method because detection frequency was <85%.

COPC – chemical of potential concern
N – number of samples
nd – no data
PCB – polychlorinated biphenyl
ROS – regression on order statistic
TEQ – toxic equivalent
UCL – upper confidence limit

Table 15. Exposure concentrations in water calculated for the entire LDW

COPC[a]	N	Detection Frequency	Concentration (mg/L)			Distribution Type	Statistic Used	Concentration used in Risk Estimation (mg/L)
			Minimum	Maximum	Mean			
Arsenic (total)	390	100%	1.83×10^{-4}	0.00157	8.5×10^{-4}	nonparametric (0.05)	Student's-t UCL	8.8×10^{-4}
Bis(2-ethylhexyl) phthalate	94	20%	9.9×10^{-7}	0.0238	4.6×10^{-4}	nonparametric (0.05)	97.5% Chebyshev UCL	0.0021[b]
Cadmium (total)	402	98%	3.35×10^{-6}	3.91×10^{-4}	4.1×10^{-5}	nonparametric (0.05)	95% Chebyshev UCL	4.7×10^{-5}
Total chromium	380	100%	2.4×10^{-4}	0.00237	6.8×10^{-4}	nonparametric (0.05)	Student's-t UCL	7.1×10^{-4}
Cobalt (total)	359	100%	3.11×10^{-5}	0.00133	2.3×10^{-4}	nonparametric (0.05)	95% Chebyshev UCL	2.7×10^{-4}
Copper (total)	391	100%	3.215×10^{-4}	0.00583	0.0016	lognormal (0.05)	Student's-t UCL	0.0016
DDTs	nd	nd	nd	nd	nd	nd	nd	nd
Lead (total)	396	100%	4.59×10^{-5}	0.00281	4.3×10^{-4}	nonparametric (0.05)	95% Chebyshev UCL	5.0×10^{-4}
Mercury (total)	29	52%	5.0×10^{-7}	6.9×10^{-6}	2.1×10^{-6}	gamma distribution (0.05)	approximate gamma UCL	2.7×10^{-6} [b]
Nickel (total)	345	98%	1.455×10^{-4}	0.00291	7.8×10^{-4}	nonparametric (0.05)	Student's-t UCL	8.2×10^{-4}
Total PCBs	15	100%	1.1×10^{-7}	3.1×10^{-6}	1.2×10^{-6}	normal (0.05)	Student's-t UCL	1.6×10^{-6}
PCB TEQ (bird)	nd	nd	nd	nd	nd	nd	nd	nd
PCB TEQ (mammal)	nd	nd	nd	nd	nd	nd	nd	nd
Selenium (total)	360	1%	6.5×10^{-5}	2.7×10^{-4}	7.6×10^{-5}	low detection frequency	maximum detected conc.	2.7×10^{-4} [c]
Vanadium (total)	304	100%	2.2×10^{-4}	0.00399	0.0014	nonparametric (0.05)	Student's-t UCL	0.0014
Zinc (total)	402	100%	7.0×10^{-4}	0.00904	0.0030	nonparametric (0.05)	Student's-t UCL	0.0032

[a] Metals were analyzed as total concentrations in water

[b] Calculated with ROS method because detection frequency was <85%

[c] Detection frequency was below 20% or in less than 4 samples. If detected, the maximum detected concentration was used. If not detected the maximum RL was used.

COPC – chemical of potential concern

N – number of samples

nd – no data

PCB – polychlorinated biphenyl

ROS – regression on order statistic

TEQ – toxic equivalent

ATTACHMENT 12
WILDLIFE EXPOSURE
DOSE CALCULATIONS

Attachment 12 Wildlife Exposure Dose Calculations

Table 1.	*COPC concentrations in great blue heron food*	*1*
Table 2.	*COPC concentrations in osprey food*	*2*
Table 3.	*COPC concentrations in river otter food*	*3*
Table 4.	*COPC concentrations in harbor seal food*	*4*
Table 5.	*Calculated COPC exposure doses for spotted sandpiper*	*5*
Table 6.	*Calculated COPC exposure doses for great blue heron*	*9*
Table 7.	*Calculated COPC exposure doses for osprey*	*10*
Table 8.	*Calculated COPC exposure doses for river otter*	*11*
Table 9.	*Calculated COPC exposure doses for harbor seal*	*11*

Acronyms

Acronym	Definition
COPC	chemical of potential concern
dw	dry weight
nd	no data
PCB	polychlorinated biphenyl
TEQ	toxic equivalent
TRV	toxicity reference value

Table 1. COPC concentrations in great blue heron food

Chemical	Concentration in Prey (mg/kg dw)					Fraction in Diet[a]					Concentration in Food (mg/kg)
	Shiner Surfperch	English Sole	Chinook Salmon	Sculpin	Crab	Shiner Surfperch	English Sole	Chinook Salmon	Sculpin	Crab	
Chromium	0.82	4.4	nd	0.40	0.30	0.317	0.317	0	0.317	0.05	1.8
Lead	0.53	1.7	nd	0.30	0.53	0.317	0.317	0	0.317	0.05	0.83
Mercury	0.15	0.070	0.14	0.15	0.29	0.238	0.238	0.238	0.238	0.05	0.14
Total PCBs	14	10	3.4	5.0	6.0	0.238	0.238	0.238	0.238	0.05	8.0
PCB TEQs	7.2×10^{-4}	4.2×10^{-4}	nd	1.2×10^{-4}	3.6×10^{-4}	0.317	0.317	0	0.317	0.05	4.2×10^{-4}
Total DDTs	1.0	0.76	0.21	0.50	0.61	0.238	0.238	0.238	0.238	0.05	0.57

[a] For COPCs with no data for salmon, the fraction of shiner surfperch, English sole, and sculpin in the diet was assumed to be equal.

dw – dry weight

nd – no data

PCB – polychlorinated biphenyl

TEQ – toxic equivalency quotient

Table 2. COPC concentrations in osprey food

Chemical	Concentration in Prey (mg/kg dw)			Fraction in Diet[a]			Concentration in Food (mg/kg)
	Shiner Surfperch	English Sole	Chinook Salmon	Shiner Surfperch	English Sole	Chinook Salmon	
Chromium	0.82	4.4	nd	0.62	0.38	0	2.2
Lead	0.53	1.7	nd	0.62	0.38	0	0.97
Mercury	0.15	0.070	0.14	0.34	0.21	0.45	0.13
Total PCBs	14	10	3.4	0.34	0.21	0.45	8.4
PCB TEQs	7.2×10^{-4}	4.2×10^{-4}	nd	0.62	0.38	0	6.1×10^{-4}
Total DDTs	1.0	0.76	0.21	0.34	0.21	0.45	0.59

[a] For COPCs with no data for salmon, the fraction of salmon in the diet was reassigned to shiner surfperch and English sole in proportion to the fraction of those species in the diet (i.e., 62% of the salmon fraction was added to perch, and 38% was added to English sole).

dw – dry weight

nd – no data

PCB – polychlorinated biphenyl

TEQ – toxic equivalency quotient

Table 3. COPC concentrations in river otter food

Chemical	Concentration in Prey (mg/kg dw)							Fraction in Diet[a, b]							Concentration in Food (mg/kg)
	Shiner Surf-perch	English Sole	Chinook Salmon	Sculpin	Crab	Clam	Mussel	Shiner Surf-perch	English Sole	Chinook Salmon	Sculpin	Crab	Clam	Mussel	
Arsenic	4.2	13	nd	4.0	25	25	5.9	0.293	0.293	0	0.293	0.10	0.01	0.01	9.0
Cobalt	0.18	0.412	nd	0.13	0.39	2.9	0.41	0.293	0.293	0	0.293	0.10	0.01	0.01	0.28
Mercury	0.15	0.070	0.14	0.15	0.29	0.12	0.094	0.22	0.22	0.22	0.22	0.10	0.01	0.01	0.14
Selenium	0.77	0.82	nd	0.87	1.2	2.0	nd	0.293	0.293	0	0.293	0.10	0.02	0	0.88
Total PCBs	14	10	3.4	5.0	6.0	4.0	0.27	0.22	0.22	0.22	0.22	0.10	0.01	0.01	7.8
PCB TEQs	1.6×10^{-4}	8.2×10^{-5}	nd	3.8×10^{-5}	5.0×10^{-5}	2.7×10^{-5}	nd	0.293	0.293	0	0.293	0.10	0.02	0	8.8×10^{-5}

[a] For COPCs with no data for salmon, the fraction of shiner surfperch, English sole, and sculpin in the diet was assumed to be equal.

[b] For COPCs with no mussel data, the fraction of diet normally assigned to mussel was assigned to crab.

dw – dry weight

nd – no data

PCB – polychlorinated biphenyl

TEQ – toxic equivalency quotient

Table 4. COPC concentrations in harbor seal food

Chemical	Concentration in Prey (mg/kg dw)				Fraction in Diet[a]				Concentration in Food (mg/kg)
	Shiner Surfperch	English Sole	Chinook Salmon	Sculpin	Shiner Surfperch	English Sole	Chinook Salmon	Sculpin	
Mercury	0.15	0.070	0.14	0.15	0.25	0.25	0.25	0.25	0.13
Total PCBs	14	10	3.4	5.0	0.25	0.25	0.25	0.25	8.1
PCB TEQs	1.6×10^{-4}	8.2×10^{-5}	nd	3.8×10^{-5}	0.333	0.333	0	0.333	9.3×10^{-5}

[a] For COPCs with no data for salmon, the fraction of shiner surfperch, English sole, and sculpin in the diet was assumed to be equal.

dw – dry weight

nd – no data

PCB – polychlorinated biphenyl

TEQ – toxic equivalency quotient

Table 5. Calculated COPC exposure doses for spotted sandpiper

Chemical and Exposure Scenario[a]	Concentration in Media			Consumption Rate[c]			Site Use Factor	Body Weight (kg)	Exposure Dose (mg/kg bw/day)
	Food (mg/kg dw)[b]	Water (mg/L)	Sediment (mg/kg dw)	Food (kg/day)	Water (L/day)	Sediment (kg/day)			
Arsenic									
Area 1/high	23	9.7×10^{-4}	36	0.0074	0.0076	0.0013	1	0.047	4.6
Area 1/high and poor	21	9.7×10^{-4}	18	0.0074	0.0076	0.0013	1	0.047	3.8
Area 2/high	25	9.3×10^{-4}	49	0.0074	0.0076	0.0013	1	0.047	5.3
Area 2/high and poor	21	9.3×10^{-4}	18	0.0074	0.0076	0.0013	1	0.047	3.8
Area 3/high	18	5.3×10^{-4}	20	0.0074	0.0076	0.0013	1	0.047	3.4
Area 3/high and poor	26	5.3×10^{-4}	60	0.0074	0.0076	0.0013	1	0.047	5.8
Cadmium eggshell thinning TRV									
Area 1/high	0.56	5.4×10^{-5}	0.45	0.0067	0.0071	0.0012	1	0.043	0.10
Area 1/high and poor	0.56	5.4×10^{-5}	0.65	0.0067	0.0071	0.0012	1	0.043	0.11
Area 2/high	0.40	5.5×10^{-5}	1.0	0.0067	0.0071	0.0012	1	0.043	0.090
Area 2/high and poor	0.49	5.5×10^{-5}	0.93	0.0067	0.0071	0.0012	1	0.043	0.10
Area 3/high	0.64	1.6×10^{-5}	0.62	0.0067	0.0071	0.0012	1	0.043	0.12
Area 3/high and poor	0.64	1.6×10^{-5}	7.8	0.0067	0.0071	0.0012	1	0.043	0.32
Cadmium growth TRV									
Area 1/high	0.56	5.4×10^{-5}	0.45	0.0067	0.0071	0.0012	1	0.043	0.10
Area 1/high and poor	0.56	5.4×10^{-5}	0.65	0.0067	0.0071	0.0012	1	0.043	0.11
Area 2/high	0.40	5.5×10^{-5}	1.0	0.0067	0.0071	0.0012	1	0.043	0.090
Area 2/high and poor	0.49	5.5×10^{-5}	0.93	0.0067	0.0071	0.0012	1	0.043	0.10
Area 3/high	0.64	1.6×10^{-5}	0.62	0.0067	0.0071	0.0012	1	0.043	0.12
Area 3/high and poor	0.64	1.6×10^{-5}	7.8	0.0067	0.0071	0.0012	1	0.043	0.32
Chromium									
Area 1/high	3.0	6.1×10^{-4}	32	0.0067	0.0071	0.0012	1	0.043	1.4
Area 1/high and poor	3.0	6.1×10^{-4}	30	0.0067	0.0071	0.0012	1	0.043	1.3
Area 2/high	18	7.3×10^{-4}	43	0.0067	0.0071	0.0012	1	0.043	4.0
Area 2/high and poor	51	7.3×10^{-4}	32	0.0067	0.0071	0.0012	1	0.043	8.8

Table 5, cont. Calculated COPC exposure doses for spotted sandpiper

Chemical and Exposure Scenario[a]	Concentration in Media			Consumption Rate[c]			Site Use Factor	Body Weight (kg)	Exposure Dose (mg/kg bw/day)
	Food (mg/kg dw)[b]	Water (mg/L)	Sediment (mg/kg dw)	Food (kg/day)	Water (L/day)	Sediment (kg/day)			
Area 3/high	5.0	1.1×10^{-3}	31	0.0067	0.0071	0.0012	1	0.043	1.6
Area 3/high and poor	5.0	1.1×10^{-3}	120	0.0067	0.0071	0.0012	1	0.043	4.1
Cobalt									
Area 1/high	1.8	2.2×10^{-4}	9.0	0.0067	0.0071	0.0012	1	0.043	0.53
Area 1/high and poor	1.8	2.2×10^{-4}	8.0	0.0067	0.0071	0.0012	1	0.043	0.50
Area 2/high	2.0	2.5×10^{-4}	9.5	0.0067	0.0071	0.0012	1	0.043	0.58
Area 2/high and poor	2.2	2.5×10^{-4}	8.0	0.0067	0.0071	0.0012	1	0.043	0.57
Area 3/high	2.2	4.6×10^{-4}	8.0	0.0067	0.0071	0.0012	1	0.043	0.57
Area 3/high and poor	2.2	4.6×10^{-4}	10	0.0067	0.0071	0.0012	1	0.043	0.62
Copper									
Area 1/high	120	0.0015	94	0.0067	0.0071	0.0012	1	0.043	21
Area 1/high and poor	120	0.0015	80	0.0067	0.0071	0.0012	1	0.043	21
Area 2/high	140	0.0016	120	0.0067	0.0071	0.0012	1	0.043	25
Area 2/high and poor	140	0.0016	79	0.0067	0.0071	0.0012	1	0.043	24
Area 3/high	70	0.0023	63	0.0067	0.0071	0.0012	1	0.043	13
Area 3/high and poor	70	0.0023	730	0.0067	0.0071	0.0012	1	0.043	31
Lead									
Area 1/high	31	3.8×10^{-4}	90	0.0074	0.0076	0.0013	1	0.047	7.4
Area 1/high and poor	31	3.8×10^{-4}	90	0.0074	0.0076	0.0013	1	0.047	7.4
Area 2/high	15	4.9×10^{-4}	160	0.0074	0.0076	0.0013	1	0.047	6.8
Area 2/high and poor	660	4.9×10^{-4}	96	0.0074	0.0076	0.0013	1	0.047	110
Area 3/high	5.6	8.9×10^{-4}	90	0.0074	0.0076	0.0013	1	0.047	3.4
Area 3/high and poor	5.6	8.9×10^{-4}	1000	0.0074	0.0076	0.0013	1	0.047	29
Mercury									
Area 1/high	0.10	2.0×10^{-6}	0.17	0.0074	0.0076	0.0013	1	0.047	0.020
Area 1/high and poor	0.10	2.0×10^{-6}	0.16	0.0074	0.0076	0.0013	1	0.047	0.020
Area 2/high	0.066	2.0×10^{-4}	0.43	0.0074	0.0076	0.0013	1	0.047	0.022
Area 2/high and poor	0.070	2.0×10^{-4}	0.29	0.0074	0.0076	0.0013	1	0.047	0.019
Area 3/high	0.50	4.3×10^{-6}	0.60	0.0074	0.0076	0.0013	1	0.047	0.095

Table 5, cont. Calculated COPC exposure doses for spotted sandpiper

Chemical and Exposure Scenario[a]	Concentration in Media			Consumption Rate[c]			Site Use Factor	Body Weight (kg)	Exposure Dose (mg/kg bw/day)
	Food (mg/kg dw)[b]	Water (mg/L)	Sediment (mg/kg dw)	Food (kg/day)	Water (L/day)	Sediment (kg/day)			
Area 3/high and poor	0.50	4.3×10^{-6}	0.40	0.0074	0.0076	0.0013	1	0.047	0.090
Nickel									
Area 1/high	3.5	7.1×10^{-4}	21	0.0067	0.0071	0.0012	1	0.043	1.1
Area 1/high and poor	3.5	7.1×10^{-4}	30	0.0067	0.0071	0.0012	1	0.043	1.4
Area 2/high	5.8	8.2×10^{-4}	22	0.0067	0.0071	0.0012	1	0.043	1.5
Area 2/high and poor	5.3	8.2×10^{-4}	20	0.0067	0.0071	0.0012	1	0.043	1.4
Area 3/high	7.4	1.3×10^{-3}	22	0.0067	0.0071	0.0012	1	0.043	1.8
Area 3/high and poor	7.4	1.3×10^{-3}	90	0.0067	0.0071	0.0012	1	0.043	3.7
Selenium									
Area 1/high	1.9	2.7×10^{-4}	0.59	0.0074	0.0076	0.0013	1	0.047	0.32
Area 1/high and poor	1.9	2.7×10^{-4}	0.56	0.0074	0.0076	0.0013	1	0.047	0.31
Area 2/high	1.6	1.6×10^{-4}	7.2	0.0074	0.0076	0.0013	1	0.047	0.45
Area 2/high and poor	1.8	1.6×10^{-4}	3.0	0.0074	0.0076	0.0013	1	0.047	0.37
Area 3/high	1.4	1.6×10^{-4}	7.0	0.0074	0.0076	0.0013	1	0.047	0.41
Area 3/high and poor	1.4	1.6×10^{-4}	6.6	0.0074	0.0076	0.0013	1	0.047	0.40
Zinc									
Area 1/high	190	0.0030	180	0.0067	0.0071	0.0012	1	0.043	35
Area 1/high and poor	190	0.0030	170	0.0067	0.0071	0.0012	1	0.043	34
Area 2/high	300	0.0030	240	0.0067	0.0071	0.0012	1	0.043	53
Area 2/high and poor	380	0.0030	180	0.0067	0.0071	0.0012	1	0.043	64
Area 3/high	270	0.0046	150	0.0067	0.0071	0.0012	1	0.043	46
Area 3/high and poor	270	0.0046	710	0.0067	0.0071	0.0012	1	0.043	62
Vanadium									
Area 1/high	6.4	0.0014	53	0.0067	0.0071	0.0012	1	0.043	2.5
Area 1/high and poor	6.4	0.0014	52	0.0067	0.0071	0.0012	1	0.043	2.4
Area 2/high	10	0.0015	58	0.0067	0.0071	0.0012	1	0.043	3.2
Area 2/high and poor	9.8	0.0015	55	0.0067	0.0071	0.0012	1	0.043	3.1
Area 3/high	9.6	0.0019	57	0.0067	0.0071	0.0012	1	0.043	3.1
Area 3/high and poor	9.6	0.0019	59	0.0067	0.0071	0.0012	1	0.043	3.1

Table 5, cont. Calculated COPC exposure doses for spotted sandpiper

Chemical and Exposure Scenario[a]	Concentration in Media			Consumption Rate[c]			Site Use Factor	Body Weight (kg)	Exposure Dose (mg/kg bw/day)
	Food (mg/kg dw)[b]	Water (mg/L)	Sediment (mg/kg dw)	Food (kg/day)	Water (L/day)	Sediment (kg/day)			
BEHP									
Area 1/high	2.9	0.0035	0.12	0.0074	0.0076	0.0013	1	0.047	0.46
Area 1/high and poor	2.9	0.0035	2.5	0.0074	0.0076	0.0013	1	0.047	0.53
Area 2/high	14	0.024	2.7	0.0074	0.0076	0.0013	1	0.047	2.3
Area 2/high and poor	14	0.024	2.1	0.0074	0.0076	0.0013	1	0.047	2.3
Area 3/high	61	1.8×10^{-4}	0.41	0.0074	0.0076	0.0013	1	0.047	9.6
Area 3/high and poor	61	1.8×10^{-4}	0.41	0.0074	0.0076	0.0013	1	0.047	9.6
Total PCBs									
Area 1/high	1.5	1.7×10^{-6}	0.34	0.0074	0.0076	0.0013	1	0.047	0.25
Area 1/high and poor	1.5	1.7×10^{-6}	0.33	0.0074	0.0076	0.0013	1	0.047	0.25
Area 2/high	5.9	1.9×10^{-6}	2.5	0.0074	0.0076	0.0013	1	0.047	1.0
Area 2/high and poor	3.8	1.9×10^{-6}	1.5	0.0074	0.0076	0.0013	1	0.047	0.64
Area 3/high	2.7	1.9×10^{-6}	0.72	0.0074	0.0076	0.0013	1	0.047	0.45
Area 3/high and poor	3.4	1.9×10^{-6}	1.1	0.0074	0.0076	0.0013	1	0.047	0.57
PCB TEQs									
Area 1/high	1.6×10^{-4}	0	3.7×10^{-5}	0.0074	0.0076	0.0013	1	0.047	2.6×10^{-5}
Area 1/high and poor	1.6×10^{-4}	0	3.4×10^{-5}	0.0074	0.0076	0.0013	1	0.047	2.6×10^{-5}
Area 2/high	5.6×10^{-4}	0	4.3×10^{-3}	0.0074	0.0076	0.0013	1	0.047	2.1×10^{-4}
Area 2/high and poor	5.6×10^{-4}	0	1.8×10^{-3}	0.0074	0.0076	0.0013	1	0.047	1.4×10^{-4}
Area 3/high	3.9×10^{-4}	0	9.6×10^{-5}	0.0074	0.0076	0.0013	1	0.047	6.4×10^{-5}
Area 3/high and poor	3.9×10^{-4}	0	3.0×10^{-3}	0.0074	0.0076	0.0013	1	0.047	1.4×10^{-4}

[a] Six exposure scenarios were evaluated; in each of three exposure areas, foraging in high-quality habitat only and foraging in both high- and poor-quality habitat were evaluated. These exposure scenarios are described in detail in Section A.5.1.3.1.

[b] Sandpiper food consists entirely of benthic invertebrates.

[c] Consumption rates are for females if the TRV endpoint is reproduction and for average of males and females if the TRV endpoint is for growth or survival.

bw – body weight

dw – dry weight

PCB – polychlorinated biphenyl

TEQ – toxic equivalency quotient

Table 6. Calculated COPC exposure doses for great blue heron

Chemical and Exposure Scenario	Concentration in Media			Consumption Rate[a]			Site Use Factor	Body Weight (kg)	Exposure Dose (mg/kg bw/day)
	Food (mg/kg dw)	Water (mg/L)	Sediment (mg/kg dw)	Food (kg/day)	Water (L/day)	Sediment (kg/day)			
Chromium	1.8	7.1×10^{-4}	70	0.10	0.11	0.0020	0.5	2.4	0.067
Lead	0.83	5.0×10^{-4}	600	0.093	0.10	0.0019	0.5	2.2	0.27
Mercury	0.14	2.7×10^{-6}	0.40	0.093	0.10	0.0019	0.5	2.2	0.0031
Total PCBs	8.0	1.6×10^{-6}	0.72	0.093	0.10	0.0019	0.5	2.2	0.17
PCB TEQs	4.2×10^{-4}	0	9.2×10^{-4}	0.093	0.10	0.0019	0.5	2.2	9.3×10^{-6}
Total DDTs	0.57	0	0.29	0.093	0.10	0.0019	0.5	2.2	0.012

[a] Consumption rates are for females if the TRV endpoint is reproduction and for an average of males and females if the TRV endpoint is for growth or survival.

bw – body weight

dw – dry weight

PCB – polychlorinated biphenyl

TEQ – toxic equivalency quotient

Table 7. Calculated COPC exposure doses for osprey

Chemical and Exposure Scenario	Concentration in Media			Consumption Rate[a]			Site Use Factor	Body Weight (kg)	Exposure Dose (mg/kg bw/day)
	Food (mg/kg dw)	Water (mg/L)	Sediment (mg/kg dw)	Food (kg/day)	Water (L/day)	Sediment (kg/day)			
Chromium	2.0	7.1×10^{-4}	70	0.083	0.083	8.3×10^{-4}	0.75	1.7	0.11
Lead	0.97	5.0×10^{-4}	600	0.091	0.087	9.1×10^{-4}	0.75	1.8	0.26
Mercury	0.13	2.7×10^{-6}	0.40	0.091	0.087	9.1×10^{-4}	0.75	1.8	0.0051
Total PCBs	8.4	1.6×10^{-6}	0.72	0.091	0.087	9.1×10^{-4}	0.75	1.8	0.32
PCB TEQs	6.1×10^{-4}	0	9.2×10^{-4}	0.091	0.087	9.1×10^{-4}	0.75	1.8	2.3×10^{-5}
Total DDTs	0.59	0	0.29	0.091	0.087	9.1×10^{-4}	0.75	1.8	0.022

[a] Consumption rates are for females if the TRV endpoint is reproduction and for an average of males and females if the TRV endpoint is for growth or survival

bw – body weight

dw – dry weight

PCB – polychlorinated biphenyl

TEQ – toxic equivalency quotient

Table 8. Calculated COPC exposure doses for river otter

Chemical and Exposure Scenario	Concentration in Media			Consumption Rate[a]			Site Use Factor	Body Weight (kg)	Exposure Dose (mg/kg bw/day)
	Food (mg/kg dw)	Water (mg/L)	Sediment (mg/kg dw)	Food (kg/day)	Water (L/day)	Sediment (kg/day)			
Arsenic	9.0	8.8×10^{-4}	30	0.26	0.64	0.0052	1	7.9	0.32
Cobalt	0.28	2.7×10^{-4}	10	0.28	0.68	0.0056	1	8.6	0.016
Mercury	0.14	2.7×10^{-6}	0.30	0.28	0.68	0.0056	1	8.6	0.0048
Selenium	0.88	2.7×10^{-4}	5.0	0.28	0.68	0.0056	1	8.6	0.032
Total PCBs	7.8	1.6×10^{-6}	0.98	0.26	0.64	0.0052	1	7.9	0.26
PCB TEQs	8.8×10^{-5}	0	7.2×10^{-5}	0.26	0.64	0.0052	1	7.9	2.9×10^{-6}

[a] Consumption rates are for females if the TRV endpoint is reproduction and for an average of males and females if the TRV endpoint is for growth or survival.

bw – body weight

dw – dry weight

PCB – polychlorinated biphenyl

TEQ – toxic equivalency quotient

Table 9. Calculated COPC exposure doses for harbor seal

Chemical and Exposure Scenario	Concentration in Media			Consumption Rate[a]			Site Use Factor	Body Weight (kg)	Exposure Dose (mg/kg bw/day)
	Food (mg/kg dw)	Water (mg/L)	Sediment (mg/kg dw)	Food (kg/day)	Water (L/day)	Sediment (kg/day)			
Mercury	0.13	2.7×10^{-6}	0.30	0.60	5.1	0.012	0.33	81	3.3×10^{-4}
Total PCBs	8.1	1.6×10^{-6}	0.98	0.58	4.9	0.012	0.33	77	0.020
PCB TEQs	9.3×10^{-5}	0	7.2×10^{-5}	0.58	4.9	0.012	0.33	77	2.3×10^{-7}

[a] Consumption rates are for females if the TRV endpoint is reproduction and for an average of males and females if the TRV endpoint is for growth or survival.

bw – body weight

dw – dry weight

PCB – polychlorinated biphenyl

TEQ – toxic equivalency quotient