Field DM-4

Years 2 thru 5

Alternate Crop Options

Year 2

Dairy Annual Nutrient Manager

Date	12/3/2009	NM STATE UNIVERSITY Dairy Extension Program	Victor E. Cabrera Dairy Specialist dairy.nmsu.edu: Tools vcabrera@nmsu.edu	PLANNED		[redacted]	
Dairy	Dominguez Dairy				N Loss	0%	
Crop_Year	2010-2011 (alt)				Goal/Real	Nutrient Needed	
Field_ID	DM-4						
Area (ac)	48						
	Month - Month	Crop		Unit	Yield	N	P
1st Crop	Nov'10-May'11	Triticale (Wheat for green chop 70%)		t/ac	20	13455	1786
2nd Crop	Jun'11-Oct'11	Cotton		lb/ac	1250	1824	228
						0	0
Total		Nutrient Needed				15,279	2,014
						Soil Analyses	
Texture by Feel		Clay				N	P
Nutrient Available in Soil				lb/ac		63	226.2
		Nutrient Still Needed				12,255	-8,844
						Effluent Analyses	
						N	P
Effluent Manure Application		NM Dairy Ponds Net from J.S. 590 L&MLA		ac-in	0.712	854	388
				ac-in		0	0
				ac-in		0	0
		Nutrient Still Needed				11,401	-9,232
						Manure Analyses	
						N	P
Dry Manure Application				t/ac		0	0
				t/ac		0	0
				t/ac		0	0
		Nutrient Still Needed				11,401	-9,232
						Fertilizer Content	
						N	P
Chemical Fertilizers Applied		22-0-0		lb/ac	950	11400	0
				lb/ac		0	0
				lb/ac		0	0
		Annual Nutrient Balance				**1**	**-9,232**

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	**Acres:**	48	**Date:**	11/2/2009	**Field ID:**	DM-4

Application information (*enter the units that will be or has been applied to the field*):					
Crop Rotation: Triticale Silage				**Needed for field (acin):**	13.68
Liquid Applied:	0.285	Acin/ac		**(gal):**	371,412
Solids Applied:		ton/ac		**Needed for field (tons):**	
Liquid Loads Applied:		1000gal/ac		**Loads needed for field:**	

Nutrient Content of Organic Material

Solid-Lab Report	**% Moisture**	**TKN (%) (dry)**	**NH_4-N (ppm) (dry)**	**P_2O_5 (%) (dry)**	**K_2O (%) (dry)**
Fill in Lab data:					

Solid Book Values (select even if test values are used)	**% Moisture**		**TKN (lbs/wet ton)**		**NH_4-N (lbs/ton)**		**P_2O_5 (lbs/wet ton)**		**K_2O (lbs/wet ton)**	
	Book	**Test**	**Book**	**Test**	**Book**	**Test**	**Book**	**Test**	**Book**	**Test**
Beef (DM)		0		0		0.0		0		0

Liquid-Lab Report	**NH_3-N (mg/L)**	**TKN (mg/L)**	**NO_3-N (mg/L)**	**Tot-PO_4 (mg/L)**	**K (mg/L)**
Fill in Lab data:		205	0.17		

Liquid	**% Moisture**		**TKN (lbs/acin)**		**NH_4-N (lbs/acin)**		**P_2O_5 (lbs/acin)**		**K_2O (lbs/acin)**	
	Book	**Test**	**Book**	**Test**	**Book**	**Test**	**Book**	**Test**	**Book**	**Test**
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		**NH_4-N (lbs/1000gal)**		**P_2O_5 (lbs/1000gal)**		**K_2O (lbs/1000gal)**	
			Book	**Test**	**Book**	**Test**	**Book**	**Test**	**Book**	**Test**
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	**Type of Climate**	**Percent Remaining**	**NH_4-N Remaining**
ıdcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	20.8 (lbs/acin) NH4-N
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	**Percent Nutrient Available the 1st Year**			
	Organic N	**P**	**K**	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	**Solid Source**
Lagoon or diluted Pond	40 %	75 %	80 %	**Liquid Source**

Solid	**Organic N (lbs/ton)**	**P_2O_5 (lbs/ton)**	**K_2O (lbs/ton)**
	0	0	0
Liquid	**Organic N (lbs/acin)**	**P_2O_5 (lbs/acin)**	**K_2O (lbs/acin)**
	8	26	205
	Organic N (lbs/100gal)	**P_2O_5 (lbs/1000gal)**	**K_2O (lbs/1000gal)**
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	**Soil Drainage Class** (See Survey Information)	**Percent Remaining (%)**
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	**lbs/1000gal**	**lbs/ac in**	**lbs/ton**
N	0.0	25	0
P_2O_5	0.0	26	0
K_2O	0.0	205	0

Total Nutrients Applied (net to the field)	**All Forms N (lbs/ac)**	**P_2O_5 (lbs/ac)**	**K_2O (lbs/ac)**
	7.3	7.5	58.3

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	Acres:	48	Date:	11/2/2009	Field ID:	DM-4

Application Information (*enter the units that will be or has been applied to the field*):	Crop Rotation: Cotton			Needed for field (acin):	20.496
	Liquid Applied:	0.427	Acin/ac	(gal):	556,466
	Solids Applied:		ton/ac	Needed for field (tons):	
	Liquid Loads Applied:		1000gal/ac	Loads needed for field:	

Nutrient Content of Organic Material

Solid-Lab Report	% Moisture	TKN (%) (dry)	NH_4-N (ppm) (dry)	P_2O_5 (%) (dry)	K_2O (%) (dry)
Fill in Lab data:					

Solid Book Values (select even if test values are used)	% Moisture		TKN (lbs/wet ton)		NH_4-N (lbs/ton)		P_2O_5 (lbs/wet ton)		K_2O (lbs/wet ton)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
Beef (DM)		0		0		0.0		0		0

Liquid-Lab Report	NH_3-N (mg/L)	TKN (mg/L)	NO_3-N (mg/L)	Tot-PO_4 (mg/L)	K (mg/L)
Fill in Lab data:		205	0.17		

Liquid	% Moisture		TKN (lbs/acin)		NH_4-N (lbs/acin)		P_2O_5 (lbs/acin)		K_2O (lbs/acin)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		NH_4-N (lbs/1000gal)		P_2O_5 (lbs/1000gal)		K_2O (lbs/1000gal)	
			Book	Test	Book	Test	Book	Test	Book	Test
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	Type of Climate	Percent Remaining	NH_4-N Remaining
dcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	20.8 (lbs/acin) NH4-N
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	Percent Nutrient Available the 1st Year			
	Organic N	P	K	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	Solid Source
Lagoon or diluted Pond	40 %	75 %	80 %	Liquid Source

Solid	Organic N (lbs/ton)	P_2O_5 (lbs/ton)	K_2O (lbs/ton)
	0	0	0
Liquid	Organic N (lbs/acin)	P_2O_5 (lbs/acin)	K_2O (lbs/acin)
	8	26	205
	Organic N (lbs/100gal)	P_2O_5 (lbs/1000gal)	K_2O (lbs/1000gal)
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	Soil Drainage Class (See Survey Information)	Percent Remaining (%)
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	lbs/1000gal	lbs/ac in	lbs/ton
N	0.0	25	0
P_2O_5	0.0	26	0
K_2O	0.0	205	0

Total Nutrients Applied (net to the field)	All Forms N (lbs/ac)	P_2O_5 (lbs/ac)	K_2O (lbs/ac)
	10.9	11.2	87.4

USDA Natural Resources Conservation Service

PLANTS

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

- Return to the crop list and make a new selection of crop(s)

used for Triticale

Wheat, for green chop

Forage

Triticum sp.

Plant part harvested: Aboveground biomass

Crop yield unit: ton

Nutrients in harvested part (lb/ton) at 70% moisture percentage.

Nitrogen	Phosphorus	Potassium
14.0160	1.8600	

Nutrients removed in harvested part (lb/acre) at 20 ton yield level.

Nitrogen	Phosphorus	Potassium
280.3200	37.2000	

Nutrients removed in harvested part (lb/ton) at 20 ton yield level and 48 acres.

Nitrogen	Phosphorus	Potassium
13455.3600	1785.6000	

Element-Fertilizer Equivalents

Average NPK Percentages

«Contents»

- Intellectual Property Statement / How to Cite the PLANTS Database
- Disclaimers
- USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: Mon 4:19 PM - 11/30/2009

USDA Natural Resources Conservation Service

PLANTS

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

Return to the crop list and make a new selection of crop(s)

Cotton, for seed with lint or seed cotton
Fiber and miscellaneous
Gossypium sp.
Plant part harvested: Seed and lint
Crop yield unit: lb of seed and lint

Nutrients in harvested part (lb/lb of seed and lint) at 7.80% moisture percentage.

Nitrogen	Phosphorus	Potassium
.0304	.0038	.0045

Nutrients removed in harvested part (lb/acre) at 1250 lb of seed and lint yield level.

Nitrogen	Phosphorus	Potassium
37.9864	4.7253	5.6473

Nutrients removed in harvested part (lb/lb of seed and lint) at 1250 lb of seed and lint yield level and 48 acres.

Nitrogen	Phosphorus	Potassium
1823.3472	226.8120	271.0680

Element-Fertilizer Equivalents

Average NPK Percentages

«Contents»

Intellectual Property Statement / How to Cite the PLANTS Database

Disclaimers

USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: Mon 4:20 PM - 11/30/2009

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-4	Date:	Nov'10 (alt)
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Triticale
Soil Permeability (in/hr):	0.3	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the approprate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <8 ppm	Low 8-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
					X	8
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
		X				1
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
				X		4
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >500-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		X				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 80% Dry Matter as Supplemental Feed	Pasture 80 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>65-100 ft wide	20-65 feet wide	< 20 feet wide	No Buffer	
			X			3

P Hazard Class: Medium **Total Index Points:** 20.5

Phosphorus Application Classification: N Based

Notes:

This evaluation has a Medium P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 11/10 to 5/11) 0.35 t/ac + (RUSLE 2) 0.20 t/ac = 0.55 t/ac

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-4	Date:	Jun'11 (alt)
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Cotton
Soil Permeablity (in/hr):	0.3	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the approprate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <8 ppm	Low 8-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
					X	8
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
		X				1
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
				X		4
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >500-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		x				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 80% Dry Matter as Supplemental Feed	Pasture 80 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>65-100 ft wide	20-65 feet wide	< 20 feet wide	No Buffer	
			x			3

P Hazard Class: Medium **Total Index Points:** 20.5

Phosphorus Application Classification: N Based

Notes:

This evaluation has a Medium P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 6/11 to 10/11) 0.45 t/ac + (RUSLE 2) 0.52 t/ac = 0.97 t/ac

NRCS Natural Resources Conservation Service

Irrigation Water Management

Conservation Practice Job Sheet 449

Natural Resources Conservation Service (NRCS) Jan, 2006

Client: Dominguez Dairy
Planner: Chet Wyant
Current Land Use: Triticale Silage / Cotton
Date: 11/20/2009

Tract: 1579
Field(s) No.: DM-4
Total Acres: 48
Date to apply: Nov'10 alt

See the Conservation Plan map for the location of the field(s) to applying IWM.

Purposes (check all that apply)

- ☐ Manage soil moisture to improve crops
- ☑ Optimize use of water
- ☑ Minimize Irrigation Erosion
- ☐ Decrease non-point source pollution
- ☐ Manage salt in the root zone
- ☐ Manage air, soil, or plant microclimate

Conditions where practice applies

This practice is applicable to all irrigated lands. An irrigation system adapted for the site conditions is available and capable of applying water to meet the intended purpose(s).

Soil (Series, Texture, and Map Unit) *Select the soil to manage for:*

Critical soil to manage: **Glendale L; 23-36** Intake Family (in/hr): **0.30**

Soil Interpretations for Irrigaiton

Crop Name	Rooting Depth	Moisture Replacement Depth (ft)	Water Holding Capacity (in)	Mgt Allowed Depletion (MAD) (%)	Net Water to Replace (in)	Time Needed to Infiltrate (hrs)
Wheat, winter, silage; Las Cruces	Medium	3.0	6.0	50%	3.0	6.3
Cotton; Las Cruces	Medium	3.0	6.0	50%	3.0	6.3

Crop Consumptive Use (CU) Information *(inches/month needed)*

Crop: Wheat, winter, silage; Las Cruces — **Total Irrigation Needed:** 10.2 ac in/ac

Month	Est. Frequency (days between irr.)	In/Mo	Month	Est. Frequency (days between irr.)	In/Mo
Jan	>1 Mo.	1.2	Jul		0.0
Feb	>1 Mo.	2.1	Aug		0.0
Mar	27	3.3	Sep	>1 Mo.	0.3
Apr		0.0	Oct	>1 Mo.	1.3
May		0.0	Nov	>1 Mo.	1.2
Jun		0.0	Dec	>1 Mo.	0.9

Crop: Cotton; Las Cruces — **Total Irrigation Needed:** 30.2 ac in/ac

Month	Est. Frequency (days between irr.)	In/Mo	Month	Est. Frequency (days between irr.)	In/Mo
Jan		0.0	Jul	12	7.5
Feb		0.0	Aug	12	7.3
Mar		0.0	Sep	17	5.4
Apr	>1 Mo.	0.3	Oct	>1 Mo.	2.8
May	>1 Mo.	2.0	Nov	>1 Mo.	0.6
Jun	21	4.4	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** (days between irr.)	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Farm Irrigation Rating System (FIRS, irrigation system efficiency)

Type of System: **Flood, controlled** System Efficiency (%): **80%**

System Capacity (Ditch/Pump/Well): **6000** GPM

Crop Grown	Days of Operation for CU by crop (days)	Est. System Application Rate (in/day)	Needed Application Rate (in/day)	System Evaluation
Wheat, winter, silage; Las Cruces	2	5.33	0.11	System meets CU for crop
Cotton; Las Cruces	6	5.33	0.25	System meets CU for crop

Operation and Maintenance Requirements

1. Irrigate when the soils reaches the MAD level, determined by soil moisture monitoring. Use one of the following methods to monitor soil moisture; the feel method, tensiometers, electrical resistance blocks, or moisture probes. Drip or Center Pivots can be irrigated on an as needed basis to meet daily CU.

2. Test irrigation water for Nitrate and Salts (Total Desolved Solids/Electrical Conductivity)

3. Do not exceed the net water to replace listed above when irrigating, unless salts are being managed.

4. Do not apply water at rates that cause runoff or erosion.

5. Monitor the soil to maintain: pH, permeability, salinity, and structure.

6. Application of pond effluent shall not exceed the crop needs (water and nutrient), and will not exceed the water holding capacity listed above.

7. Consider using crops such as sorghum, cotton, or winter wheat when water supplies are short.

8. Avoid traffic on wet soils to minimize soil compaction.

Additional Requirements

Job Approval and Completion

Client: Isaac Domingez **Date:** 11-23-2009

Conservationist: _____ **Date:** 11-23-09

Completed by: _____ **Date:** _____

NRCS - WEQ CALCULATIONS

Version 9.03 11-19-2007

Producer: Dominguez Dairy | Planner: Chet Wyant | Location: Mesquite, NM | Tract: 1579 | Field: DM-4

Crop Rotation: Triticale Silage / Cotton | Climate Data Station: NM, LAS CRUCES | Site "C" Value: 150

Tillage Direction (NS/EW): NS | Length/width ratio: 2 | Field Direction (NS/EW): NS | Field Width (Ft.): 1233

Irrigation (Y or N): Y | Soil "I": 86 | Wind Erodibility Group: 3 (1-7) | TWF: 1 (see instr.)

Sum Period Erosion (t/ac): 0.8 | No. Yrs in Rotation: 1.0 | Av. Annual Wind Erosion: 0.8 (t/ac/yr)

Calculations and Output

Mgt Periods		Irr.	Soil	Ridge Roughness				Random Roughness	Unsheltered Distance				SGe	Erosion			
Dates		No. of	"I"	Dev.	Ht.	Sp.	"Krd"	"Krr"	Dev.	Prep.	WED	"L"	"V"	"E"	EWE	"IF"	Loss
Begin	End	(#)	(t/ac)	(deg)	(in.)	(in.)	(factor)	(factor)	(deg)	(factor)	(factor)	(ft)	(lbs/ac)	(t/ac)	(%)	(%)	(t/ac)
11/1	11/02/10	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	0	20.5	0.2	1.00	0.03
11/2	12/02/10	0	56	0.0	2	10	0.44	0.99	0.0	4.0	1.010	1245	3544	0.0	5.0	1.00	0.00
12/2	01/01/11	1	56	0.0	2	10	0.44	0.99	0.0	4.0	1.010	1245	3898	0.0	7.4	0.97	0.00
1/1	01/31/11	0	56	0.0	2	10	0.44	0.99	0.0	4.0	1.010	1245	6010	0.0	4.0	1.00	0.00
1/31	03/02/11	0	56	0.0	2	10	0.44	0.99	0.0	4.0	1.010	1245	6999	0.0	24.7	1.00	0.00
3/2	03/17/11	2	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	6999	0.0	12.9	0.87	0.00
3/17	04/01/11	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	3544	0.0	12.9	1.00	0.00
4/1	04/16/11	1	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	3544	0.0	8.8	0.93	0.00
4/16	05/01/11	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	3544	0.0	8.8	1.00	0.00
5/1	05/16/11	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	4282	0.0	5.3	1.00	0.00
5/16	05/23/11	1	56	0.0	4	13	0.24	0.52	0.0	4.0	1.010	1245	89	4.2	2.5	0.86	0.09
5/23	05/29/11	1	56	0.0	4	30	0.37	0.50	0.0	4.0	1.010	1245	27	7.2	2.1	0.83	0.13
5/29	06/13/11	0	56	0.0	8	30	0.21	0.80	0.0	4.0	1.010	1245	23	6.3	1.7	1.00	0.11
6/13	06/28/11	1	56	22.5	2	10	0.62	0.86	22.5	1.2	1.370	1689	16	35.2	0.8	0.93	0.26
6/28	07/13/11	1	56	22.5	2	10	0.62	0.86	22.5	1.2	1.370	1689	21	35.1	0.3	0.93	0.10
7/13	07/28/11	1	56	67.5	2	10	0.65	0.86	67.5	1.1	1.475	1819	203	33.8	0.2	0.93	0.06
7/28	08/12/11	1	56	67.5	2	10	0.65	0.86	67.5	1.1	1.475	1819	668	19.5	0.1	0.93	0.02
8/12	08/27/11	1	56	90.0	2	10	0.81	0.86	90.0	2.8	2.00	2466	1367	5.4	0.1	0.93	0.00
8/27	09/11/11	1	56	90.0	2	10	0.81	0.86	90.0	2.8	2.00	2466	2319	0.0	0.3	0.93	0.00
9/11	09/20/11	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.01	1245	943	4.8	0.3	1.00	0.01
9/20	09/21/11	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.01	1245	1178	2.2	0.0	1.00	0.00
9/21	10/31/11	0	56	0.0	2	10	0.44	0.99	0.0	4.0	1.01	1245	5304	0.0	1.6	1.00	0.00
		0			2	10											
		0															
		0															
		0															

USDA NRCS Natural Resources Conservation Service

RUSLE2 Worksheet Erosion Calculation Record

Info: (Alt) Triticale Silage Nov'10 to May'11, Cotton Jun'11 to Oct'11

Tract #: 1579
Owner name: DOMINGUEZ DAIRY
Field name: DM-4

Location: New Mexico\DonaAna County\NM_Dona Ana R 9
Soil: Ge GLENDALE LOAM\GLENDALE loam 85%
Slope length (horiz): 100.0 ft
Avg. slope steepness: 0.50 %
T value: 5.0 t/ac/yr

Alternatives:

Management	*Contouring*	*Strips / barriers*	*Diversion/terrace, sediment basin*	*Cons. plan. soil loss, t/ac/yr*	*Description*
a.Single Year/Single Crop Templates\Triticale, silage, plow, irr CT	b. absolute row grade 0.5 percent	(none)	(none)	0.20	
a.Single Year/Single Crop Templates\Cotton, CT 2 bale	b. absolute row grade 0.5 percent	(none)	(none)	0.52	

Year 3

Dairy Annual Nutrient Manager

Date	12/3/2009	NM STATE UNIVERSITY Dairy Extension Program	Victor E. Cabrera Dairy Specialist dairy.nmsu.edu: Tools vcabrera@nmsu.edu	PLANNED		[redacted]	
Dairy	Dominguez Dairy				N Loss	0%	
Crop_Year	2011-2012 (alt)				Goal/Real	Nutrient Needed	
Field_ID	DM-4						
Area (ac)	48						
	Month - Month	Crop		Unit	Yield	N	P
1st Crop	Nov'11-May'12	Barley-6 row, for green chop (boot)		t/ac	20	8614	1446
2nd Crop	Jun'11-Oct'12	Corn-Field for Silage (dough 68%)		t/ac	22	8351	1250
						0	0
Total		Nutrient Needed				16,965	2,697
						Soil Analyses	
Texture by Feel		Clay				N	P
Nutrient Available in Soil				lb/ac		63	226.2
		Nutrient Still Needed				13,941	-8,161
						Effluent Analyses	
						N	P
Effluent Manure Application		NM Dairy Ponds Net from J.S. 590 L&MLA		ac-in	0.712	854	388
				ac-in		0	0
				ac-in		0	0
		Nutrient Still Needed				13,086	-8,549
						Manure Analyses	
						N	P
Dry Manure Application				t/ac		0	0
				t/ac		0	0
				t/ac		0	0
		Nutrient Still Needed				13,086	-8,549
						Fertilizer Content	
						N	P
Chemical Fertilizers Applied		22-0-0		lb/ac	1138	13656	0
				lb/ac		0	0
				lb/ac		0	0
		Annual Nutrient Balance				**-570**	**-8,549**

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	Acres:	48	Date:	11/2/2009	Field ID:	DM-4

Application Information (*enter the units that will be or has been applied to the field*):					
Crop Rotation:	Barley Silage			Needed for field (acin):	13.68
Liquid Applied:	0.285	AcIn/ac		(gal):	371,412
Solids Applied:		ton/ac		Needed for field (tons):	
Liquid Loads Applied:		1000gal/ac		Loads needed for field:	

Nutrient Content of Organic Material

Solid-Lab Report	% Moisture	TKN (%) (dry)	NH_4-N (ppm) (dry)	P_2O_5 (%) (dry)	K_2O (%) (dry)
Fill in Lab data:					

Solid Book Values (select even if test values are used)	% Moisture		TKN (lbs/wet ton)		NH_4-N (lbs/ton)		P_2O_5 (lbs/wet ton)		K_2O (lbs/wet ton)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
Beef (DM)		0		0		0.0		0		0

Liquid-Lab Report	NH_3-N (mg/L)	TKN (mg/L)	NO_3-N (mg/L)	Tot-PO_4 (mg/L)	K (mg/L)
Fill in Lab data:		205	0.17		

Liquid	% Moisture		TKN (lbs/acin)		NH_4-N (lbs/acin)		P_2O_5 (lbs/acin)		K_2O (lbs/acin)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		NH_4-N (lbs/1000gal)		P_2O_5 (lbs/1000gal)		K_2O (lbs/1000gal)	
			Book	Test	Book	Test	Book	Test	Book	Test
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	Type of Climate	Percent Remaining	NH_4-N Remaining
dcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	20.8 (lbs/acin) NH4-N
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	Percent Nutrient Available the 1st Year			
	Organic N	P	K	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	Solid Source
Lagoon or diluted Pond	40 %	75 %	80 %	Liquid Source

Solid	Organic N (lbs/ton)	P_2O_5 (lbs/ton)	K_2O (lbs/ton)
	0	0	0
Liquid	**Organic N (lbs/acin)**	**P_2O_5 (lbs/acin)**	**K_2O (lbs/acin)**
	8	26	205
	Organic N (lbs/100gal)	**P_2O_5 (lbs/1000gal)**	**K_2O (lbs/1000gal)**
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	Soil Drainage Class (See Survey Information)	Percent Remaining (%)
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	lbs/1000gal	lbs/ac in	lbs/ton
N	0.0	25	0
P_2O_5	0.0	26	0
K_2O	0.0	205	0

Total Nutrients Applied (net to the field)	All Forms N (lbs/ac)	P_2O_5 (lbs/ac)	K_2O (lbs/ac)
	7.3	7.5	58.3

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	Acres:	48	Date:	11/2/2009	Field ID:	DM-4

Application Information (*enter the units that will be or has been applied to the field*):					
Crop Rotation: Corn Silage				Needed for field (acin):	20.496
Liquid Applied:	0.427	Acin/ac		(gal):	556,466
Solids Applied:		ton/ac		Needed for field (tons):	
Liquid Loads Applied:		1000gal/ac		Loads needed for field:	

Nutrient Content of Organic Material

Solid-Lab Report	% Moisture	TKN (%) (dry)	NH_4-N (ppm) (dry)	P_2O_5 (%) (dry)	K_2O (%) (dry)
Fill in Lab data:					

Solid Book Values (select even if test values are used)	% Moisture		TKN (lbs/wet ton)		NH_4-N (lbs/ton)		P_2O_5 (lbs/wet ton)		K_2O (lbs/wet ton)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
Beef (DM)		0		0		0.0		0		0

Liquid-Lab Report	NH_3-N (mg/L)	TKN (mg/L)	NO_3-N (mg/L)	Tot-PO_4 (mg/L)	K (mg/L)
Fill in Lab data:		205	0.17		

Liquid	% Moisture		TKN (lbs/acin)		NH_4-N (lbs/acin)		P_2O_5 (lbs/acin)		K_2O (lbs/acin)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		NH_4-N (lbs/1000gal)		P_2O_5 (lbs/1000gal)		K_2O (lbs/1000gal)	
			Book	Test	Book	Test	Book	Test	Book	Test
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	Type of Climate	Percent Remaining	NH_4-N Remaining
dcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	20.8 (lbs/acin) NH4-N
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	Percent Nutrient Available the 1st Year			
	Organic N	P	K	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	Solid Source
Lagoon or diluted Pond	40 %	75 %	80 %	Liquid Source

Solid	Organic N (lbs/ton)	P_2O_5 (lbs/ton)	K_2O (lbs/ton)
	0	0	0
Liquid	**Organic N (lbs/acin)**	**P_2O_5 (lbs/acin)**	**K_2O (lbs/acin)**
	8	26	205
	Organic N (lbs/100gal)	**P_2O_5 (lbs/1000gal)**	**K_2O (lbs/1000gal)**
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	Soil Drainage Class (See Survey Information)	Percent Remaining (%)
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	lbs/1000gal	lbs/ac in	lbs/ton
N	0.0	25	0
P_2O_5	0.0	26	0
K_2O	0.0	205	0

Total Nutrients Applied (net to the field)	All Forms N (lbs/ac)	P_2O_5 (lbs/ac)	K_2O (lbs/ac)
	10.9	11.2	87.4

USDA Natural Resources Conservation Service

PLANTS

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

Return to the crop list and make a new selection of crop(s)

Barley-6 row, for green chop (boot)

Forage

Hordeum vulgare

Plant part harvested: Aboveground biomass

Crop yield unit: ton

Nutrients in harvested part (lb/ton) at 69% moisture percentage.

Nitrogen	Phosphorus	Potassium
13.6688	2.2951	9.7854

Nutrients removed in harvested part (lb/acre) at 20 ton yield level.

Nitrogen	Phosphorus	Potassium
273.3764	45.9023	195.7077

Nutrients removed in harvested part (lb/ton) at 20 ton yield level and 48 acres.

Nitrogen	Phosphorus	Potassium
13122.0688	2203.3124	9393.9719

Element-Fertilizer Equivalents

Average NPK Percentages

«Contents»

Intellectual Property Statement / How to Cite the PLANTS Database

Disclaimers

USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: Mon 4:27 PM - 11/30/2009

USDA Natural Resources Conservation Service

PLANTS

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

Return to the crop list and make a new selection of crop(s)

Corn-Field, for silage (dough stage)

Forage

Zea mays ssp. mays

Plant part harvested: Aboveground biomass

Crop yield unit: ton

Nutrients in harvested part (lb/ton) at 68% moisture percentage.

Nitrogen	Phosphorus	Potassium
7.9078	1.1840	6.6560

Nutrients removed in harvested part (lb/acre) at 22 ton yield level.

Nitrogen	Phosphorus	Potassium
173.9725	26.0480	146.4320

Nutrients removed in harvested part (lb/ton) at 22 ton yield level and 48 acres.

Nitrogen	Phosphorus	Potassium
8350.6790	1250.3040	7028.7360

Element-Fertilizer Equivalents

Average NPK Percentages

«Contents»

Intellectual Property Statement / How to Cite the PLANTS Database

Disclaimers

USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: Mon 4:25 PM - 11/30/2009

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-4	Date:	Nov'11 (alt)
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Barley Silage
Soil Permeablity (in/hr):	0.3	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the approprate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <8 ppm	Low 8-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
					X	8
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
		X				1
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
				X		4
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >500-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		x				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 60% Dry Matter as Supplemental Feed	Pasture 60 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>65-100 ft wide	20-65 feet wide	< 20 feet wide	No Buffer	
			x			3

P Hazard Class:	Medium	Total Index Points:	20.5
Phosphorus Application Classification:		N Based	

Notes:

This evaluation has a Medium P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 11/11 to 5/12) 0.25 t/ac + (RUSLE 2) 0.11 t/ac = 0.36 t/ac

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-4	Date:	Jun'12 (alt)
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Corn Silage
Soil Permeability (in/hr):	0.3	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the approprate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <8 ppm	Low 8-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
					X	8
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
		X				1
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
				X		4
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >500-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		x				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 60% Dry Matter as Supplemental Feed	Pasture 60 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>65-100 ft wide	20-65 feet wide	< 20 feet wide	No Buffer	
			x			3

P Hazard Class: Medium **Total Index Points:** 20.5

Phosphorus Application Classification: N Based

Notes:

This evaluation has a Medium P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 6/12 to 10/12) 0.17 t/ac + (RUSLE 2) 0.10 t/ac = 0.27 t/ac

NRCS Natural Resources Conservation Service

Irrigation Water Management

Conservation Practice Job Sheet 449

Natural Resources Conservation Service (NRCS) Jan, 2006

Client: Dominguez Dairy
Tract: 1579
Planner: Chet Wyant
Field(s) No.: DM-4
Current Land Use: Barley Silage / Corn Silage
Total Acres: 48
Date: 11/20/2009
Date to apply: Nov'11 alt

See the Conservation Plan map for the location of the field(s) to applying IWM.

Purposes (check all that apply)

- ☐ Manage soil moisture to improve crops
- ☑ Optimize use of water
- ☑ Minimize Irrigation Erosion
- ☐ Decrease non-point source pollution
- ☐ Manage salt in the root zone
- ☐ Manage air, soil, or plant microclimate

Conditions where practice applies

This practice is applicable to all irrigated lands. An irrigation system adapted for the site conditions is available and capable of applying water to meet the intended purpose(s).

Soil (Series, Texture, and Map Unit) *Select the soil to manage for:*

Critical soil to manage: **Glendale L; 23-36** Intake Family (in/hr): **0.30**

Soil Interpretations for Irrigaiton

Crop Name	Rooting Depth	Moisture Replacement Depth (ft)	Water Holding Capacity (in)	Mgt Allowed Depletion (MAD) (%)	Net Water to Replace (in)	Time Needed to Infiltrate (hrs)
Wheat, winter, silage; Las Cruces	Medium	3.0	6.0	50%	3.0	6.3
Corn, silage; Las Cruces	Medium	3.0	6.0	50%	3.0	6.3

Crop Consumptive Use (CU) Information *(inches/month needed)*

Crop: Wheat, winter, silage; Las Cruces — **Total Irrigation Needed:** 10.2 ac in/ac

Month	Est. Frequency (days between irr.)	In/Mo	Month	Est. Frequency (days between irr.)	In/Mo
Jan	>1 Mo.	1.2	Jul		0.0
Feb	>1 Mo.	2.1	Aug		0.0
Mar	27	3.3	Sep	>1 Mo.	0.3
Apr		0.0	Oct	>1 Mo.	1.3
May		0.0	Nov	>1 Mo.	1.2
Jun		0.0	Dec	>1 Mo.	0.9

Crop: Corn, silage; Las Cruces — **Total Irrigation Needed:** 29.3 ac in/ac

Month	Est. Frequency (days between irr.)	In/Mo	Month	Est. Frequency (days between irr.)	In/Mo
Jan		0.0	Jul	10	9.4
Feb		0.0	Aug	14	6.6
Mar		0.0	Sep		0.0
Apr	>1 Mo.	1.2	Oct		0.0
May	22	4.2	Nov		0.0
Jun	11	7.9	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** (days between irr.)	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Farm Irrigation Rating System (FIRS, irrigation system efficiency)

Type of System: **Flood, controlled** System Efficiency (%): **80%**

System Capacity (Ditch/Pump/Well): **6000** GPM

Crop Grown	Days of Operation for CU by crop (days)	Est. System Applicaiton Rate (in/day)	Needed Application Rate (in/day)	System Evaluation
Wheat, winter, silage; Las Cruces	2	5.33	0.11	System meets CU for crop
Corn, silage; Las Cruces	6	5.33	0.31	System meets CU for crop

Operation and Maintenance Requirements

1. Irrigate when the soils reaches the MAD level, determined by soil moisture monitoring. Use one of the following methods to monitor soil moisture; the feel method, tensiometers, electrical resistance blocks, or moisture probes. Drip or Center Pivots can be irrigated on an as needed basis to meet daily CU.

2. Test irrigation water for Nitrate and Salts (Total Desolved Solids/Electrical Conductivity)

3. Do not exceed the net water to replace listed above when irrigating, unless salts are being managed.

4. Do not apply water at rates that cause runoff or erosion.

5. Monitor the soil to maintain: pH, permeability, salinity, and structure.

6. Application of pond effluent shall not exceed the crop needs (water and nutrient), and will not exceed the water holding capacity listed above.

7. Consider using crops such as sorghum, cotton, or winter wheat when water supplies are short.

8. Avoid traffic on wet soils to minimize soil compaction.

Additional Requirements

Job Approval and Completion

Client: Joe and Dominique **Date:** 11-23-2009

Conservationist: _____ **Date:** 11-23-09

Completed by: _____ **Date:** _____

NRCS - WEQ CALCULATIONS

Version 9.03 11-19-2007

Producer: Dominguez Dairy | Planner: Chet Wyant | Location: Anthony, NM | Tract: 1579 | Field: DM-4

Crop Rotation: Barley Silage/Corn Silage | Climate Data Station: NM, LAS CRUCES | Site "C" Value: 150

Tillage Direction (NS/EW): NS | Length/width ratio: 2 | Field Direction (NS/EW): NS | Field Width (Ft.): 1233

Irrigation (Y or N): Y | Soil "I": 86 | Wind Erodibility Group: 3 (1-7) | TWF: 1 (see instr.)

Sum Period Erosion (t/ac): 0.4 | No. Yrs in Rotation: 1.0 | Av. Annual Wind Erosion: 0.4 (t/ac/yr)

Calculations and Output

Mgt Periods		Irr.	Soil	Ridge Roughness				Random Roughness	Unsheltered Distance				SGe	Erosion			
Dates		No. of	"I"	Dev.	Ht.	Sp.	"Krd"	"Krr"	Dev.	Prep.	WED	"L"	"V"	"E"	EWE	"F"	Loss
Begin	End	(#)	(t/ac)	(deg)	(in.)	(in.)	(factor)	(factor)	(deg)	(factor)	(factor)	(ft)	(lbs/ac)	(t/ac)	(%)	(%)	(t/ac)
11/1	11/02/11	0	56	0.0	2	18	0.58	0.86	0.0	4.0	1.010	1245	0	29.9	0.2	1.00	0.05
11/2	11/17/11	1	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	3774	0.0	2.5	0.93	0.00
11/17	12/02/11	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	3774	0.0	2.5	1.00	0.00
12/2	12/17/11	1	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	3774	0.0	3.7	0.93	0.00
12/17	01/01/12	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	3774	0.0	3.7	1.00	0.00
1/1	01/16/12	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	3774	0.0	2.0	1.00	0.00
1/16	05/15/12	3	56	0.0	2	10	0.44	0.99	0.0	4.0	1.010	1245	3774	0.0	75.1	0.98	0.00
5/15	05/22/12	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	4282	0.0	2.5	1.00	0.00
5/22	05/29/12	0	56	0.0	4	13	0.24	0.52	0.0	4.0	1.010	1245	789	1.2	2.5	1.00	0.03
5/29	05/29/12	0	56	0.0	4	30	0.37	0.50	0.0	4.0	1.010	1245	285	5.6	0.0	1.00	0.00
5/29	06/29/12	0	56	0.0	8	30	0.21	0.99	0.0	4.0	1.010	1245	251	6.6	2.6	1.00	0.17
6/29	07/14/12	1	56	22.5	2	18	0.71	0.86	22.5	1.2	1.370	1689	237	36.2	0.3	0.93	0.09
7/14	07/29/12	2	56	67.5	2	18	0.74	0.86	67.5	1.1	1.475	1819	242	38.2	0.2	0.87	0.06
7/29	08/13/12	1	56	67.5	2	18	0.74	0.86	67.5	1.1	1.475	1819	763	20.0	0.1	0.93	0.02
8/13	08/28/12	1	56	90.0	2	18	0.86	0.86	90.0	2.8	2.000	2466	1671	2.4	0.1	0.93	0.00
8/28	09/12/12	1	56	90.0	2	18	0.86	0.86	90.0	2.8	2.000	2466	5473	0.0	0.4	0.93	0.00
9/12	10/01/12	0	56	0.0	2	18	0.58	0.99	0.0	4.0	1.010	1245	6999	0.0	0.6	1.00	0.00
10/1	11/01/12	0	56	0.0	2	18	0.59	0.99	0.0	3.3	1.01	1245	5317	0.0	1.3	1.00	0.00
		0			2	18											
		0															
		0															
		0															
		0															
		0															
		0															
		0															

USDA NRCS

RUSLE2 Worksheet Erosion Calculation Record

Info: (Alt) Barley Silage Nov'11 to May'12, Corn Silage Jun'12 to Oct'12

Tract #: 1579
Owner name: DOMINGUEZ DAIRY
Field name: DM-4

Location: New Mexico\DonaAna County\NM_Dona Ana R 9
Soil: Ge GLENDALE LOAM\GLENDALE loam 85%
Slope length (horiz): 100.0 ft
Avg. slope steepness: 0.50 %
T value: 5.0 t/ac/yr

Alternatives:

Management	*Contouring*	*Strips / barriers*	*Diversion/terrace, sediment basin*	*Cons. plan. soil loss, t/ac/yr*	*Description*
a.Single Year/Single Crop Templates\WWheat, silage, plow, irr CT	b. absolute row grade 0.5 percent	(none)	(none)	0.11	
a.Single Year/Single Crop Templates\Corn, silage, plow	b. absolute row grade 0.5 percent	(none)	(none)	0.10	

Used winter Dry Barley

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	**Acres:**	48	**Date:** 11/2/2009	**Field ID:** DM-4

Application information (*enter the units that will be or has been applied to the field*):					
Crop Rotation: Winter Wheat Silage				**Needed for field (acin):**	13.68
Liquid Applied:	0.285	AcIn/ac		**(gal):**	371,412
Solids Applied:		ton/ac		**Needed for field (tons):**	
Liquid Loads Applied:		1000gal/ac		**Loads needed for field:**	

Nutrient Content of Organic Material

Solid-Lab Report	**% Moisture**	**TKN (%) (dry)**	**NH_4-N (ppm) (dry)**	**P_2O_5 (%) (dry)**	**K_2O (%) (dry)**
Fill in Lab data:					

Solid Book Values (select even if test values are used)	**% Moisture**		**TKN (lbs/wet ton)**		**NH_4-N (lbs/ton)**		**P_2O_5 (lbs/wet ton)**		**K_2O (lbs/wet ton)**	
	Book	**Test**	**Book**	**Test**	**Book**	**Test**	**Book**	**Test**	**Book**	**Test**
Beef (DM)		0		0		0.0		0		0

Liquid-Lab Report	**NH_3-N (mg/L)**	**TKN (mg/L)**	**NO_3-N (mg/L)**	**Tot-PO_4 (mg/L)**	**K (mg/L)**
Fill in Lab data:		205	0.17		

Liquid	**% Moisture**		**TKN (lbs/acin)**		**NH_4-N (lbs/acin)**		**P_2O_5 (lbs/acin)**		**K_2O (lbs/acin)**	
	Book	**Test**	**Book**	**Test**	**Book**	**Test**	**Book**	**Test**	**Book**	**Test**
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		**NH_4-N (lbs/1000gal)**		**P_2O_5 (lbs/1000gal)**		**K_2O (lbs/1000gal)**	
			Book	**Test**	**Book**	**Test**	**Book**	**Test**	**Book**	**Test**
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	**Type of Climate**	**Percent Remaining**	**NH_4-N Remaining**
dcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	20.8 (lbs/acin) NH4-N
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	**Percent Nutrient Available the 1st Year**			
	Organic N	**P**	**K**	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	**Solid Source**
Lagoon or diluted Pond	40 %	75 %	80 %	**Liquid Source**

Solid	**Organic N (lbs/ton)**	**P_2O_5 (lbs/ton)**	**K_2O (lbs/ton)**
	0	0	0
Liquid	**Organic N (lbs/acin)**	**P_2O_5 (lbs/acin)**	**K_2O (lbs/acin)**
	8	26	205
	Organic N (lbs/100gal)	**P_2O_5 (lbs/1000gal)**	**K_2O (lbs/1000gal)**
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	**Soil Drainage Class** (See Survey Information)	**Percent Remaining (%)**
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	**lbs/1000gal**	**lbs/ac in**	**lbs/ton**
N	0.0	25	0
P_2O_5	0.0	26	0
K_2O	0.0	205	0

Total Nutrients Applied (net to the field)	**All Forms N (lbs/ac)**	**P_2O_5 (lbs/ac)**	**K_2O (lbs/ac)**
	7.3	7.5	58.3

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	Acres:	48	Date: 11/2/2009	Field ID: DM-4

Application information (*enter the units that will be or has been applied to the field*):					
Crop Rotation:	Sorghum Silage			Needed for field (acin):	20.496
Liquid Applied:	0.427	Acin/ac		(gal):	556,466
Solids Applied:		ton/ac		Needed for field (tons):	
Liquid Loads Applied:		1000gal/ac		Loads needed for field:	

Nutrient Content of Organic Material

Solid-Lab Report	% Moisture	TKN (%) (dry)	NH_4-N (ppm) (dry)	P_2O_5 (%) (dry)	K_2O (%) (dry)
Fill in Lab data:					

Solid Book Values (select even if test values are used)	% Moisture		TKN (lbs/wet ton)		NH_4-N (lbs/ton)		P_2O_5 (lbs/wet ton)		K_2O (lbs/wet ton)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
Beef (DM)		0		0		0.0		0		0

Liquid-Lab Report	NH_3-N (mg/L)	TKN (mg/L)	NO_3-N (mg/L)	Tot-PO_4 (mg/L)	K (mg/L)
Fill in Lab data:		205	0.17		

Liquid	% Moisture		TKN (lbs/acin)		NH_4-N (lbs/acin)		P_2O_5 (lbs/acin)		K_2O (lbs/acin)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		NH_4-N (lbs/1000gal)		P_2O_5 (lbs/1000gal)		K_2O (lbs/1000gal)	
			Book	Test	Book	Test	Book	Test	Book	Test
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	Type of Climate	Percent Remaining	NH_4-N Remaining
dcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	**NH_4-N Remaining**
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	20.8 (lbs/acin) NH4-N
			0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	Percent Nutrient Available the 1st Year			
	Organic N	P	K	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	Solid Source
Lagoon or diluted Pond	40 %	75 %	80 %	Liquid Source

Solid	Organic N (lbs/ton)	P_2O_5 (lbs/ton)	K_2O (lbs/ton)
	0	0	0
Liquid	**Organic N (lbs/acin)**	**P_2O_5 (lbs/acin)**	**K_2O (lbs/acin)**
	8	26	205
	Organic N (lbs/100gal)	**P_2O_5 (lbs/1000gal)**	**K_2O (lbs/1000gal)**
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	Soil Drainage Class (See Survey Information)	Percent Remaining (%)
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	lbs/1000gal	lbs/ac in	lbs/ton
N	0.0	25	0
P_2O_5	0.0	26	0
K_2O	0.0	205	0

Total Nutrients Applied (net to the field)	All Forms N (lbs/ac)	P_2O_5 (lbs/ac)	K_2O (lbs/ac)
	10.9	11.2	87.4

USDA Natural Resources Conservation Service

PLANTS

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

Return to the crop list and make a new selection of crop(s)

Wheat, for green chop
Forage
Triticum sp.
Plant part harvested: Aboveground biomass
Crop yield unit: ton

Nutrients in harvested part (lb/ton) at 70% moisture percentage.

Nitrogen	Phosphorus	Potassium
14.0160	1.8600	

Nutrients removed in harvested part (lb/acre) at 20 ton yield level.

Nitrogen	Phosphorus	Potassium
280.3200	37.2000	

Nutrients removed in harvested part (lb/ton) at 20 ton yield level and 48 acres.

Nitrogen	Phosphorus	Potassium
13455.3600	1785.6000	

Element-Fertilizer Equivalents

Average NPK Percentages

«Contents»

Intellectual Property Statement / How to Cite the PLANTS Database

Disclaimers

USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: Mon 4:19 PM - 11/30/2009

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

Return to the crop list and make a new selection of crop(s)

Sorghum/Sudangrass, for silage
Forage
Sorghum bicolor ssp. drummondii
Plant part harvested: Aboveground biomass
Crop yield unit: ton

Nutrients in harvested part (lb/ton) at 70% moisture percentage.

Nitrogen	Phosphorus	Potassium
9.8280	1.2600	15.6600

Nutrients removed in harvested part (lb/acre) at 21 ton yield level.

Nitrogen	Phosphorus	Potassium
206.3880	26.4600	328.8600

Nutrients removed in harvested part (lb/ton) at 21 ton yield level and 48 acres.

Nitrogen	Phosphorus	Potassium
9906.6240	1270.0800	15785.2800

Element-Fertilizer Equivalents

Average NPK Percentages

«Contents»

Intellectual Property Statement / How to Cite the PLANTS Database

Disclaimers

USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: Mon 4:21 PM - 11/30/2009

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-4	Date:	Nov'12 (alt)
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Wheat Silage
Soil Permeablity (in/hr):	0.3	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the approprate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <6 ppm	Low 6-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
					X	8
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
		X				1
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
				X		4
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >500-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		x				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 80% Dry Matter as Supplemental Feed	Pasture 80 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>85-100 ft wide	20-85 feet wide	< 20 feet wide	No Buffer	
			x			3

P Hazard Class:	Medium	Total Index Points:	20.5
Phosphorus Application Classification:	N Based		

Notes:

This evaluation has a Medium P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 11/12 to 5/13) 0.27 t/ac + (RUSLE 2) 0.11 t/ac = 0.38 t/ac

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-4	Date:	Jun'13 (alt)
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Sudangrass
Soil Permeablity (in/hr):	0.3	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the approprate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <8 ppm	Low 8-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
					X	8
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
		X				1
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
				X		4
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >500-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		x				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 80% Dry Matter as Supplemental Feed	Pasture 80 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>65-100 ft wide	20-65 feet wide	< 20 feet wide	No Buffer	
			x			3

P Hazard Class:	Medium	Total Index Points:	20.5
Phosphorus Application Classification:	N Based		

Notes:

This evaluation has a Medium P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 6/13 to 10/13) 0.19 t/ac + (RUSLE 2) 0.00078 t/ac = 0.19 t/ac

NRCS Natural Resources Conservation Service

Irrigation Water Management

Conservation Practice Job Sheet 449

Natural Resources Conservation Service (NRCS) Jan, 2006

Client: Dominguez Dairy **Tract:** 1579
Planner: Chet Wyant **Field(s) No.:** DM-4
Current Land Use: Winter Wheat Silage / Sorghum Silage **Total Acres:** 48
Date: 11/20/2009 **Date to apply:** Nov'12 alt

See the Conservation Plan map for the location of the field(s) to applying IWM.

Purposes (check all that apply)

- ☐ Manage soil moisture to improve crops
- ☑ Optimize use of water
- ☑ Minimize Irrigation Erosion
- ☐ Decrease non-point source pollution
- ☐ Manage salt in the root zone
- ☐ Manage air, soil, or plant microclimate

Conditions where practice applies

This practice is applicable to all irrigated lands. An irrigation system adapted for the site conditions is available and capable of applying water to meet the intended purpose(s).

Soil (Series, Texture, and Map Unit) *Select the soil to manage for:*

Critical soil to manage: **Glendale L; 23-36** Intake Family (in/hr): **0.30**

Soil Interpretations for Irrigaiton

Crop Name	Rooting Depth	Moisture Replacement Depth (ft)	Water Holding Capacity (in)	Mgt Allowed Depletion (MAD) (%)	Net Water to Replace (in)	Time Needed to Infiltrate (hrs)
Wheat, winter, silage; Las Cruces	Medium	3.0	6.0	50%	3.0	6.3
Sorghum, silage; Deming	Medium	3.0	6.0	65%	3.9	9.7

Crop Consumptive Use (CU) Information *(inches/month needed)*

Crop: Wheat, winter, silage; Las Cruces **Total Irrigation Needed:** 10.2 ac in/ac

Month	Est. Frequency *(days between irr.)*	In/Mo	Month	Est. Frequency *(days between irr.)*	In/Mo
Jan	>1 Mo.	1.2	Jul		0.0
Feb	>1 Mo.	2.1	Aug		0.0
Mar	27	3.3	Sep	>1 Mo.	0.3
Apr		0.0	Oct	>1 Mo.	1.3
May		0.0	Nov	>1 Mo.	1.2
Jun		0.0	Dec	>1 Mo.	0.9

Crop: Sorghum, silage; Deming **Total Irrigation Needed:** 23.5 ac in/ac

Month	Est. Frequency *(days between irr.)*	In/Mo	Month	Est. Frequency *(days between irr.)*	In/Mo
Jan		0.0	Jul	14	8.5
Feb		0.0	Aug	21	5.5
Mar		0.0	Sep	>1 Mo.	1.0
Apr		0.0	Oct		0.0
May	>1 Mo.	1.6	Nov		0.0
Jun	17	7.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** (days between irr.)	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Farm Irrigation Rating System (FIRS, irrigation system efficiency)

Type of System:	**Flood, controlled**	System Efficiency (%):	**80%**
System Capacity (Ditch/Pump/Well):	**6000**	GPM	

Crop Grown	Days of Operation for CU by crop (days)	Est. System Applicaiton Rate (in/day)	Needed Application Rate (in/day)	System Evaluation
Wheat, winter, silage; Las Cruces	2	5.33	0.11	System meets CU for crop
Sorghum, silage; Deming	4	5.33	0.28	System meets CU for crop

Operation and Maintenance Requirements

1. Irrigate when the soils reaches the MAD level, determined by soil moisture monitoring. Use one of the following methods to monitor soil moisture; the feel method, tensiometers, electrical resistance blocks, or moisture probes. Drip or Center Pivots can be irrigated on an as needed basis to meet daily CU.

2. Test irrigation water for Nitrate and Salts (Total Desolved Solids/Electrical Conductivity)

3. Do not exceed the net water to replace listed above when irrigating, unless salts are being managed.

4. Do not apply water at rates that cause runoff or erosion.

5. Monitor the soil to maintain: pH, permeability, salinity, and structure.

6. Application of pond effluent shall not exceed the crop needs (water and nutrient), and will not exceed the water holding capacity listed above.

7. Consider using crops such as sorghum, cotton, or winter wheat when water supplies are short.

8. Avoid traffic on wet soils to minimize soil compaction.

Additional Requirements

Job Approval and Completion

Client: Joan Dominguez **Date:** 11-23-2009

Conservationist: [illegible] **Date:** 11-23-09

Completed by: [illegible] **Date:**

NRCS - WEQ CALCULATIONS

Version 9.03 11-19-2007

Producer: Dominguez Dairy | Planner: Chet Wyant | Location: Anthony, NM | Tract: 1579 | Field: DM-4

Crop Rotation: Wheat Silage/Sudangrass Hay | Climate Data Station: NM, LAS CRUCES | Site "C" Value: 150

Tillage Direction (NS/EW): NS | Length/width ratio: 2 | Field Direction (NS/EW): NS | Field Width (Ft.): 1233

Irrigation (Y or N): Y | Soil "I": 86 | Wind Erodibility Group: 3 (1-7) | TWF: 1 (see instr.)

Sum Period Erosion (t/ac): 0.5 | No. Yrs in Rotation: 1.0 | Av. Annual Wind Erosion: 0.5 (t/ac/yr)

Calculations and Output

Mgt Periods		Irr.	Soil	Ridge Roughness				Random Roughness	Unsheltered Distance				SGe	Erosion			
Dates		No. of	"I"	Dev.	Ht.	Sp.	"Krd"	"Krr"	Dev.	Prep.	WED	"L"	"V"	"E"	EWE	"IF"	Loss
Begin	End	(#)	(t/ac)	(deg)	(in.)	(in.)	(factor)	(factor)	(deg)	(factor)	(factor)	(ft)	(lbs/ac)	(t/ac)	(%)	(%)	(t/ac)
11/1	11/02/12	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	0	20.5	0.2	1.00	0.03
11/2	11/17/12	1	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	6999	0.0	2.5	0.93	0.00
11/17	12/02/12	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	6999	0.0	2.5	1.00	0.00
12/2	12/17/12	1	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	6999	0.0	3.7	0.93	0.00
12/17	01/01/13	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	6999	0.0	3.7	1.00	0.00
1/1	01/16/13	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	6999	0.0	2.0	1.00	0.00
1/16	05/15/13	3	56	0.0	2	10	0.44	0.99	0.0	4.0	1.010	1245	6999	0.0	75.1	0.97	0.00
5/15	05/22/13	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	4282	0.0	2.5	1.00	0.00
5/22	05/29/13	0	56	0.0	4	13	0.24	0.52	0.0	4.0	1.010	1245	789	1.2	2.5	1.00	0.03
5/29	06/29/13	0	56	0.0	4	30	0.37	0.65	0.0	4.0	1.010	1245	285	8.2	2.6	1.00	0.21
6/29	07/14/13	1	56	22.5	2	10	0.62	0.86	22.5	1.2	1.370	1689	202	31.1	0.3	0.93	0.08
7/14	07/29/13	2	56	67.5	2	10	0.65	0.86	67.5	1.1	1.475	1819	227	33.2	0.2	0.87	0.06
7/29	08/13/13	1	56	67.5	2	10	0.65	0.86	67.5	1.1	1.475	1819	313	30.7	0.1	0.93	0.03
8/13	08/28/13	1	56	90.0	2	10	0.81	0.86	90.0	2.8	2.000	2466	706	26.9	0.1	0.93	0.02
8/28	09/12/13	1	56	90.0	2	10	0.81	0.86	90.0	2.8	2.000	2466	1869	1.2	0.4	0.93	0.00
9/12	10/01/13	1	56	0.0	2	10	0.44	0.99	0.0	4.0	1.010	1245	2297	0.0	0.6	0.95	0.00
10/1	11/01/13	1	56	0.0	2	10	0.46	0.99	0.0	3.3	1.010	1245	6999	0.0	1.3	0.97	0.00
		0			2	10											
		0															
		0															
		0															
		0															
		0															
		0															
		0															
		0															

USDA NRCS Natural Resources Conservation Service

RUSLE2 Worksheet Erosion Calculation Record

Info: (Alt) Winter Wheat Silage Nov'12 to May'13, Sudangrass Silage Jun'13 to Oct'13

Tract #: 1579
Owner name: DOMINGUEZ DAIRY
Field name: DM-4

Location: New Mexico\DonaAna County\NM_Dona Ana R 9
Soil: Ge GLENDALE LOAM\GLENDALE loam 85%
Slope length (horiz): 100.0 ft
Avg. slope steepness: 0.50 %
T value: 5.0 t/ac/yr

Alternatives:

Management	*Contouring*	*Strips / barriers*	*Diversion/terrace, sediment basin*	*Cons. plan. soil loss, t/ac/yr*	*Description*
a.Single Year/Single Crop Templates\WWheat, silage, plow, irr CT	b. absolute row grade 0.5 percent	(none)	(none)	0.11	
a.Single Year/Single Crop Templates\Haygrazer, sudangrass, disk	b. absolute row grade 0.5 percent	(none)	(none)	0.00078	

Year 5

Dairy Annual Nutrient Manager

Date	12/3/2009	NM STATE UNIVERSITY Dairy Extension Program	Victor E. Cabrera Dairy Specialist dairy.nmsu.edu: Tools vcabrera@nmsu.edu	PLANNED		[redacted]	
Dairy	Dominguez Dairy				N Loss	0%	
Crop_Year	2013-2014				Goal/Real	Nutrient Needed	
Field_ID	DM-4						
Area (ac)	48			Unit	Yield	N	P
	Month - Month	Crop					
1st Crop	Nov'13-May'14	Oats for green chop (boot to early bloom 69%)		t/ac	20	21608	4034
2nd Crop	Jun'14-Oct'14	Alfalfa, for hay		t/ac	1	2419	227
						0	0
Total		Nutrient Needed				24,027	4,261
						Soil Analyses	
Texture by Feel		Clay				N	P
Nutrient Available in Soil				lb/ac		63	226.2
		Nutrient Still Needed				21,003	-6,597
						Effluent Analyses	
						N	P
Effluent Manure Application		NM Dairy Ponds Net from J.S. 590 L&MLA		ac-in	0.712	854	388
				ac-in		0	0
				ac-in		0	0
		Nutrient Still Needed				20,149	-6,985
						Manure Analyses	
						N	P
Dry Manure Application		NM Dairy Cattle Net from J.S. 590 L&MLA		t/ac	29.9	20093	4392
				t/ac		0	0
				t/ac		0	0
		Nutrient Still Needed				56	-11,377
						Fertilizer Content	
						N	P
Chemical Fertilizers Applied				lb/ac		0	0
				lb/ac		0	0
				lb/ac		0	0
		Annual Nutrient Balance				**56**	**-11,377**

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	Acres:	48	Date:	11/2/2009	Field ID:	DM-4

Application Information (*enter the units that will be or has been applied to the field*):					
Crop Rotation:	Oat Silage			Needed for field (acin):	13.68
Liquid Applied:	0.285	Acin/ac		(gal):	371,412
Solids Applied:	29.9	ton/ac		Needed for field (tons):	1435.2
Liquid Loads Applied:		1000gal/ac		Loads needed for field:	

Nutrient Content of Organic Material

Solid-Lab Report	% Moisture	TKN (%) (dry)	NH_4-N (ppm) (dry)	P_2O_5 (%) (dry)	K_2O (%) (dry)
Fill in Lab data:		2.3	89	0.5	

Solid Book Values (select even if test values are used)	% Moisture		TKN (lbs/wet ton)		NH_4-N (lbs/ton)		P_2O_5 (lbs/wet ton)		K_2O (lbs/wet ton)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
Beef (DM)	0	0	0	46	0.0	0.2	0	9	60	0

Liquid-Lab Report	NH_3-N (mg/L)	TKN (mg/L)	NO_3-N (mg/L)	Tot-PO_4 (mg/L)	K (mg/L)
Fill in Lab data:		205	0.17		

Liquid	% Moisture		TKN (lbs/acin)		NH_4-N (lbs/acin)		P_2O_5 (lbs/acin)		K_2O (lbs/acin)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		NH_4-N (lbs/1000gal)		P_2O_5 (lbs/1000gal)		K_2O (lbs/1000gal)	
			Book	Test	Book	Test	Book	Test	Book	Test
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	Type of Climate	Percent Remaining	NH_4-N Remaining
dcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	**NH_4-N Remaining**
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	20.8 (lbs/acin) NH4-N 0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	Percent Nutrient Available the 1st Year			
	Organic N	P	K	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	Solid Source
Lagoon or diluted Pond	40 %	75 %	80 %	Liquid Source

Solid	Organic N (lbs/ton)	P_2O_5 (lbs/ton)	K_2O (lbs/ton)
	16	7	48
Liquid	**Organic N (lbs/acin)**	**P_2O_5 (lbs/acin)**	**K_2O (lbs/acin)**
	8	26	205
	Organic N (lbs/100gal)	**P_2O_5 (lbs/1000gal)**	**K_2O (lbs/1000gal)**
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	Soil Drainage Class (See Survey Information)	Percent Remaining (%)
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	lbs/1000gal	lbs/ac in	lbs/ton
N	0.0	25	14
P_2O_5	0.0	26	7
K_2O	0.0	205	48

Total Nutrients Applied (net to the field)	All Forms N (lbs/ac)	P_2O_5 (lbs/ac)	K_2O (lbs/ac)
	433.9	213.8	1500.6

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	Acres:	48	Date:	11/2/2009	Field ID:	DM-4

Application information (*enter the units that will be or has been applied to the field*):					
Crop Rotation:	Alfalfa			Needed for field (acin):	20.496
Liquid Applied:	0.427	Acin/ac		(gal):	556,466
Solids Applied:		ton/ac		Needed for field (tons):	
Liquid Loads Applied:		1000gal/ac		Loads needed for field:	

Nutrient Content of Organic Material

Solid-Lab Report	% Moisture	TKN (%) (dry)	NH_4-N (ppm) (dry)	P_2O_5 (%) (dry)	K_2O (%) (dry)
Fill in Lab data:					

Solid Book Values (select even if test values are used)	% Moisture		TKN (lbs/wet ton)		NH_4-N (lbs/ton)		P_2O_5 (lbs/wet ton)		K_2O (lbs/wet ton)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
Beef (DM)		0		0		0.0		0		0

Liquid-Lab Report	NH_3-N (mg/L)	TKN (mg/L)	NO_3-N (mg/L)	Tot-PO_4 (mg/L)	K (mg/L)
Fill in Lab data:		205	0.17		

Liquid	% Moisture		TKN (lbs/acin)		NH_4-N (lbs/acin)		P_2O_5 (lbs/acin)		K_2O (lbs/acin)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		NH_4-N (lbs/1000gal)		P_2O_5 (lbs/1000gal)		K_2O (lbs/1000gal)	
			Book	Test	Book	Test	Book	Test	Book	Test
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	Type of Climate	Percent Remaining	NH_4-N Remaining
dcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	**NH_4-N Remaining**
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	20.8 (lbs/acin) NH4-N 0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	Percent Nutrient Available the 1st Year			
	Organic N	P	K	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	Solid Source
Lagoon or diluted Pond	40 %	75 %	80 %	Liquid Source

Solid	Organic N (lbs/ton)	P_2O_5 (lbs/ton)	K_2O (lbs/ton)
	0	0	0
Liquid	**Organic N (lbs/acin)**	**P_2O_5 (lbs/acin)**	**K_2O (lbs/acin)**
	8	26	205
	Organic N (lbs/100gal)	**P_2O_5 (lbs/1000gal)**	**K_2O (lbs/1000gal)**
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	Soil Drainage Class (See Survey Information)	Percent Remaining (%)
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	lbs/1000gal	lbs/ac in	lbs/ton
N	0.0	25	0
P_2O_5	0.0	26	0
K_2O	0.0	205	0

Total Nutrients Applied (net to the field)	All Forms N (lbs/ac)	P_2O_5 (lbs/ac)	K_2O (lbs/ac)
	10.9	11.2	87.4

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

Return to the crop list and make a new selection of crop(s)

Oat, for green chop (boot to early bloom)
Forage
Avena sativa
Plant part harvested: Aboveground biomass
Crop yield unit: ton

Nutrients in harvested part (lb/ton) at 69% moisture percentage.

Nitrogen	Phosphorus	Potassium
12.2508	2.1046	11.6554

Nutrients removed in harvested part (lb/acre) at 20 ton yield level.

Nitrogen	Phosphorus	Potassium
245.0165	42.0916	233.1076

Nutrients removed in harvested part (lb/ton) at 20 ton yield level and 48 acres.

Nitrogen	Phosphorus	Potassium
11760.7940	2020.3982	11189.1633

Element-Fertilizer Equivalents

Average NPK Percentages

«Contents»

Intellectual Property Statement / How to Cite the PLANTS Database

Disclaimers

USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: Mon 4:23 PM - 11/30/2009

USDA Natural Resources Conservation Service

PLANTS

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

Return to the crop list and make a new selection of crop(s)

Alfalfa, for hay
Forage
Medicago sativa
Plant part harvested: Aboveground biomass
Crop yield unit: ton

Nutrients in harvested part (lb/ton) at 9.65% moisture percentage.

Nitrogen	Phosphorus	Potassium
50.3935	4.7217	38.2939

Nutrients removed in harvested part (lb/acre) at 1 ton yield level.

Nitrogen	Phosphorus	Potassium
50.3935	4.7217	38.2939

Nutrients removed in harvested part (lb/ton) at 1 ton yield level and 48 acres.

Nitrogen	Phosphorus	Potassium
2418.8897	226.6397	1838.1084

Element-Fertilizer Equivalents

Average NPK Percentages

«Contents»

Intellectual Property Statement / How to Cite the PLANTS Database

Disclaimers

USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: Mon 4:35 PM - 11/30/2009

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-4	Date:	Nov'13 alt
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Oat Silage
Soil Permeablity (in/hr):	0.3	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the approprate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <8 ppm	Low 8-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
					X	8
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
					X	8
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
			X			2
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >500-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		x				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 80% Dry Matter as Supplemental Feed	Pasture 80 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>65-100 ft wide	20-65 feet wide	< 20 feet wide	No Buffer	
			x			3

P Hazard Class: Medium **Total Index Points:** 25.5

Phosphorus Application Classification: N Based

Notes:

This evaluation has a Medium P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 11/13 to 5/14) 0.29 t/ac + (RUSLE 2) 0.16 t/ac = 0.45 t/ac

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-2	Date:	Jun'14 alt
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Alfalfa
Soil Permeablity (in/hr):	0.3	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the appropriate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <8 ppm	Low 8-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
					X	8
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
		X				1
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
				X		4
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >600-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		x				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 80% Dry Matter as Supplemental Feed	Pasture 80 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>65-100 ft wide	20-65 feet wide	< 20 feet wide	No Buffer	
			x			3

P Hazard Class: Medium **Total Index Points:** 20.5

Phosphorus Application Classification: N Based

Notes:

This evaluation has a Medium P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 6/11 to 10/11) 0.40 t/ac + (RUSLE 2) 0.042 t/ac = 0.44 t/ac

NRCS Natural Resources Conservation Service

Irrigation Water Management

Conservation Practice Job Sheet 449

Natural Resources Conservation Service (NRCS) Jan, 2006

Client: Dominguez Dairy
Planner: Chet Wyant
Current Land Use: Oat Silage / Alfalfa
Date: 11/20/2009
Tract: 1579
Field(s) No.: DM-4
Total Acres: 48
Date to apply: Nov'13 alt

See the Conservation Plan map for the location of the field(s) to applying IWM.

Purposes (check all that apply)

- ☐ Manage soil moisture to improve crops
- ☑ Optimize use of water
- ☑ Minimize Irrigation Erosion
- ☐ Decrease non-point source pollution
- ☐ Manage salt in the root zone
- ☐ Manage air, soil, or plant microclimate

Conditions where practice applies

This practice is applicable to all irrigated lands. An irrigation system adapted for the site conditions is available and capable of applying water to meet the intended purpose(s).

Soil (Series, Texture, and Map Unit) *Select the soil to manage for:*

Critical soil to manage: **Glendale L; 23-36** Intake Family (in/hr): **0.30**

Soil Interpretations for Irrigaiton

Crop Name	Rooting Depth	Moisture Replacement Depth (ft)	Water Holding Capacity (in)	Mgt Allowed Depletion (MAD) (%)	Net Water to Replace (in)	Time Needed to Infiltrate (hrs)
Oat, hay; Deming	Medium	3.0	6.0	50%	3.0	6.3
Alfalfa, hay, southern; Las Cruces	Deep	4.0	8.0	50%	4.0	10.1

Crop Consumptive Use (CU) Information *(inches/month needed)*

Crop: Oat, hay; Deming — **Total Irrigation Needed:** 9.4 ac in/ac

Month	Est. Frequency (days between irr.)	In/Mo	Month	Est. Frequency (days between irr.)	In/Mo
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar	>1 Mo.	2.3	Sep		0.0
Apr	18	5.0	Oct		0.0
May	>1 Mo.	2.1	Nov		0.0
Jun		0.0	Dec		0.0

Crop: Alfalfa, hay, southern; Las Cruces — **Total Irrigation Needed:** 47.7 ac in/ac

Month	Est. Frequency (days between irr.)	In/Mo	Month	Est. Frequency (days between irr.)	In/Mo
Jan		0.0	Jul	12	9.7
Feb		0.0	Aug	16	7.7
Mar	>1 Mo.	1.6	Sep	21	5.7
Apr	29	4.2	Oct	>1 Mo.	2.7
May	17	6.9	Nov		0.0
Jun	13	9.3	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** (days between irr.)	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Farm Irrigation Rating System (FIRS, irrigation system efficiency)

Type of System:	Flood, controlled		System Efficiency (%):	80%
System Capacity (Ditch/Pump/Well):		6000	GPM	
Crop Grown	**Days of Operation for CU by crop (days)**	**Est. System Applicalton Rate (in/day)**	**Needed Application Rate (in/day)**	**System Evaluation**
Oat, hay; Deming	2	5.33	0.17	System meets CU for crop
Alfalfa, hay, southern; Las Cruces	9	5.33	0.32	System meets CU for crop

Operation and Maintenance Requirements

1. Irrigate when the soils reaches the MAD level, determined by soil moisture monitoring. Use one of the following methods to monitor soil moisture; the feel method, tensiometers, electrical resistance blocks, or moisture probes. Drip or Center Pivots can be irrigated on an as needed basis to meet daily CU.

2. Test irrigation water for Nitrate and Salts (Total Desolved Solids/Electrical Conductivity)

3. Do not exceed the net water to replace listed above when irrigating, unless salts are being managed.

4. Do not apply water at rates that cause runoff or erosion.

5. Monitor the soil to maintain: pH, permeability, salinity, and structure.

6. Application of pond effluent shall not exceed the crop needs (water and nutrient), and will not exceed the water holding capacity listed above.

7. Consider using crops such as sorghum, cotton, or winter wheat when water supplies are short.

8. Avoid traffic on wet soils to minimize soil compaction.

Additional Requirements

Job Approval and Completion

Client: Isaac Domínguez **Date:** 11-23-2009

Conservationist: _____ **Date:** 11-23-09

Completed by: _____ **Date:** _____

NRCS - WEQ CALCULATIONS

Version 9 03 11-19-2007

Producer: Dominguez Dairy
Planner: Chet Wyant
Location: Mesquite, NM
Tract: 1579
Field: DM-4
Crop Rotation: Oat Silage / Alfalfa
Climate Data Station: NM, LAS CRUCES
Site "C" Value: 150
Tillage Direction (NS/EW): NS
Length/width ratio: 2
Field Direction (NS/EW): NS
Field Width (Ft.): 1233
Irrigation (Y or N): Y
Soil "I": 86
Wind Erodibility Group: 3 (1-7)
TWF: 1 (see instr.)
Sum Period Erosion (t/ac): 0.7
No. Yrs in Rotation: 1.0
Av. Annual Wind Erosion: 0.7 (t/ac/yr)

Calculations and Output

Mgt. Periods Dates		Irr.	Soil	Ridge Roughness				Random Roughness	Unsheltered Distance				SCe	Erosion			
Begin	End	No. of (#)	"I" (t/ac)	Dev. (deg)	Ht. (in.)	Sp. (in.)	"Krd" (factor)	"Krr" (factor)	Dev. (deg)	Prep. (factor)	WED (factor)	"L" (ft)	"V" (lbs/ac)	"E" (t/ac)	EWE (%)	"F" (%)	Loss (t/ac)
11/1	11/02/13	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	0	20.5	0.2	1.00	0.03
11/2	12/02/13	0	56	0.0	2	10	0.44	0.99	0.0	4.0	1.010	1245	1989	0.0	5.0	1.00	0.00
12/2	01/01/14	1	56	0.0	2	10	0.44	0.99	0.0	4.0	1.010	1245	1989	0.0	7.4	0.97	0.00
1/1	01/31/14	0	56	0.0	2	10	0.44	0.99	0.0	4.0	1.010	1245	1989	0.0	4.0	1.00	0.00
1/31	03/02/14	1	56	0.0	2	10	0.44	0.99	0.0	4.0	1.010	1245	1989	0.0	24.7	0.97	0.00
3/2	03/17/14	1	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	1989	0.0	12.9	0.93	0.00
3/17	04/01/14	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	1989	0.0	12.9	1.00	0.00
4/1	04/16/14	1	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	1989	0.0	8.8	0.93	0.00
4/16	05/01/14	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	1989	0.0	8.8	1.00	0.00
5/1	05/16/14	0	56	0.0	2	10	0.44	0.86	0.0	4.0	1.010	1245	3587	0.0	5.3	1.00	0.00
5/16	05/23/14	0	56	0.0	4	13	0.24	0.52	0.0	4.0	1.010	1245	518	2.2	2.5	1.00	0.06
5/23	05/29/14	1	56	0.0	4	30	0.37	0.50	0.0	4.0	1.010	1245	187	6.3	2.1	0.83	0.11
5/29	06/13/14	1	56	0.0	8	30	0.21	0.80	0.0	4.0	1.010	1245	165	5.6	1.7	0.93	0.09
6/13	06/28/14	1	56	22.5	2	10	0.62	0.86	22.5	1.2	1.370	1689	117	33.0	0.8	0.93	0.25
6/28	07/13/14	1	56	22.5	2	10	0.62	0.86	22.5	1.2	1.370	1689	163	32.0	0.3	0.93	0.09
7/13	07/28/14	1	56	67.5	2	10	0.65	0.86	67.5	1.1	1.475	1819	220	33.4	0.2	0.93	0.06
7/28	08/12/14	1	56	67.5	2	10	0.65	0.86	67.5	1.1	1.475	1819	1792	0.5	0.1	0.93	0.00
8/12	08/27/14	1	56	90.0	2	10	0.81	0.86	90.0	2.8	2.00	2466	5501	0.0	0.1	0.93	0.00
8/27	09/11/14	1	56	90.0	2	10	0.81	0.86	90.0	2.8	2.00	2466	5501	0.0	0.3	0.93	0.00
9/11	09/29/14	0	56	0.0	2	10	0.44	0.99	0.0	4.0	1.01	1245	2799	0.0	0.5	1.00	0.00
9/29	10/04/14	1	56	0.0	2	10	0.44	0.86	0.0	4.0	1.01	1245	2799	0.0	0.2	0.80	0.00
10/4	10/31/14	1	56	0.0	2	10	0.46	0.99	0.0	3.3	1.01	1245	2799	0.0	1.1	0.96	0.00
		0			2	10											
		0															
		0															
		0															

USDA NRCS

RUSLE2 Worksheet Erosion Calculation Record

Info: (alt) Oat Silage Nov'13 to May'14, Alfalfa Jun'14 to Oct'14

Tract #: 1579
Owner name: DOMINGUEZ DAIRY
Field name: DM-4

Location: New Mexico\DonaAna County\NM_Dona Ana R 9
Soil: Ge GLENDALE LOAM\GLENDALE loam 85%
Slope length (horiz): 100.0 ft
Avg. slope steepness: 0.50 %
T value: 5.0 t/ac/yr

Alternatives:

Management	*Contouring*	*Strips / barriers*	*Diversion/terrace, sediment basin*	*Cons. plan. soil loss, t/ac/yr*	*Description*
a.Single Year/Single Crop Templates\Forage\Oats, hay and grain, conv till, irr, CMZ29	b. absolute row grade 0.5 percent	(none)	(none)	0.16	
b.Mullti-year Rotation Templates\Alfalfa-Small Grain, 5yr, 5cuts CMZ 23	b. absolute row grade 0.5 percent	(none)	(none)	0.042	

Field DM-5

Years 1 thru 5

NMSU Annual Nutrient Manager Calculation
590 Nutrient Mgt. Jobsheet for Organic & Manure Land Application
USDA-NRCS PLANTS Crop Nutrient Tool
NRCS Phosphorus Index Worksheet for New Mexico
NRCS Irrigation Water Mgt. Conservation Practice Jobsheet 449
NRCS - WEQ Calculations
NRCS RUSLE2 Worksheet Erosion Calculation Record

Year 1

Dairy Annual Nutrient Manager

Date	11/30/2009	NM STATE UNIVERSITY Dairy Extension Program	Victor E. Cabrera Dairy Specialist dairy.nmsu.edu: Tools vcabrera@nmsu.edu	PLANNED		
Dairy	Dominguez Dairy					
Crop_Year	2009-2010					
Field_ID	DM-5			N Loss	0%	
Area (ac)	52			Goal/Real	Nutrient Needed	
	Month - Month	Crop	Unit	Yield	N	P
1st Crop	Nov'09-Oct'10	Alfalfa, for hay	t/ac	12	31446	2946
2nd Crop					0	0
					0	0
Total		Nutrient Needed			31,446	2,946
					Soil Analyses	
Texture by Feel		Sandy Clay			N	P
Nutrient Available in Soil			lb/ac		142	97.38
		Nutrient Still Needed			24,062	-2,117
					Effluent Analyses	
					N	P
Effluent Manure Application		NM Dairy Ponds Net from J.S. 590 L&MLA	ac-in	0.712	926	420
			ac-in		0	0
			ac-in		0	0
		Nutrient Still Needed			23,136	-2,538
					Manure Analyses	
					N	P
Dry Manure Application			t/ac		0	0
			t/ac		0	0
			t/ac		0	0
		Nutrient Still Needed			23,136	-2,538
					Fertilizer Content	
					N	P
Chemical Fertilizers Applied			lb/ac		0	0
			lb/ac		0	0
			lb/ac		0	0
		Annual Nutrient Balance			23,136	-2,538

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	Acres:	52	Date:	11/2/2009	Field ID:	DM-5

Application information	Crop Rotation: Alfalfa			Needed for field (acin):	37.024
***nter the units that will be or has been applied to the field*):**	Liquid Applied:	0.712	Acln/ac	(gal):	1,005,202
	Solids Applied:		ton/ac	Needed for field (tons):	
	Liquid Loads Applied:		1000gal/ac	Loads needed for field:	

Nutrient Content of Organic Material

Solid-Lab Report	% Moisture	TKN (%) (dry)	NH_4-N (ppm) (dry)	P_2O_5 (%) (dry)	K_2O (%) (dry)
Fill in Lab data:					

Solid Book Values (select even if test values are used)	% Moisture		TKN (lbs/wet ton)		NH_4-N (lbs/ton)		P_2O_5 (lbs/wet ton)		K_2O (lbs/wet ton)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
Beef (DM)		0		0		0.0		0		0

Liquid-Lab Report	NH_3-N (mg/L)	TKN (mg/L)	NO_3-N (mg/L)	Tot-PO_4 (mg/L)	K (mg/L)
Fill in Lab data:		205	0.17		

Liquid	% Moisture		TKN (lbs/acin)		NH_4-N (lbs/acin)		P_2O_5 (lbs/acin)		K_2O (lbs/acin)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		NH_4-N (lbs/1000gal)		P_2O_5 (lbs/1000gal)		K_2O (lbs/1000gal)	
			Book	Test	Book	Test	Book	Test	Book	Test
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	Type of Climate	Percent Remaining	NH_4-N Remaining
Broadcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	20.8 (lbs/acin) NH4-N
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	Percent Nutrient Available the 1st Year			
	Organic N	P	K	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	Solid Source
Lagoon or diluted Pond	40 %	75 %	80 %	Liquid Source

Solid	Organic N (lbs/ton)	P_2O_5 (lbs/ton)	K_2O (lbs/ton)
	0	0	0
Liquid	**Organic N (lbs/acin)**	**P_2O_5 (lbs/acin)**	**K_2O (lbs/acin)**
	8	26	205
	Organic N (lbs/100gal)	**P_2O_5 (lbs/1000gal)**	**K_2O (lbs/1000gal)**
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	Soil Drainage Class (See Survey Information)	Percent Remaining (%)
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	lbs/1000gal	lbs/ac in	lbs/ton
N	0.0	25	0
P_2O_5	0.0	26	0
K_2O	0.0	205	0

Total Nutrients Applied (net to the field)	All Forms N (lbs/ac)	P_2O_5 (lbs/ac)	K_2O (lbs/ac)
	18.1	18.7	145.7

USDA Natural Resources Conservation Service

PLANTS

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

Return to the crop list and make a new selection of crop(s)

Alfalfa, for hay
Forage
Medicago sativa
Plant part harvested: Aboveground biomass
Crop yield unit: ton

Nutrients in harvested part (lb/ton) at 9.65% moisture percentage.

Nitrogen	Phosphorus	Potassium
50.3935	4.7217	38.2939

Nutrients removed in harvested part (lb/acre) at 12 ton yield level.

Nitrogen	Phosphorus	Potassium
604.7224	56.6599	459.5271

Nutrients removed in harvested part (lb/ton) at 12 ton yield level and 52 acres.

Nitrogen	Phosphorus	Potassium
31445.5665	2946.3165	23895.4086

Element-Fertilizer Equivalents

Average NPK Percentages

«Contents»

Intellectual Property Statement / How to Cite the PLANTS Database

Disclaimers

USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: Mon 4:38 PM - 11/30/2009

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-5	Date:	Nov'09
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Alfalfa
Soil Permeablity (in/hr):	0.5	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the approprate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <8 ppm	Low 8-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
				X		4
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
		X				1
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
				X		4
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >500-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		x				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 80% Dry Matter as Supplemental Feed	Pasture 80 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>65-100 ft wide	20-65 feet wide	< 20 feet wide	No Buffer	
			x			3

P Hazard Class:	Low	Total Index Points:	16.5
Phosphorus Application Classification:	N Based		

Notes:

This evaluation has a Low P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 11'09 to 10/10) 0.08 t/ac + (RUSLE 2) 0.042 t/ac = 0.12 t/ac

NRCS Natural Resources Conservation Service

Irrigation Water Management

Conservation Practice Job Sheet 449

Natural Resources Conservation Service (NRCS) Jan, 2006

Client: Dominguez Dairy **Tract:** 1579
Planner: Chet Wyant **Field(s) No.:** DM-5
Current Land Use: Alfalfa **Total Acres:** 52
Date: 11/20/2009 **Date to apply:** Nov'09

See the Conservation Plan map for the location of the field(s) to applying IWM.

Purposes (check all that apply)

- ☐ Manage soil moisture to improve crops
- ☑ Optimize use of water
- ☑ Minimize Irrigation Erosion
- ☐ Decrease non-point source pollution
- ☐ Manage salt in the root zone
- ☐ Manage air, soil, or plant microclimate

Conditions where practice applies

This practice is applicable to all irrigated lands. An irrigation system adapted for the site conditions is available and capable of applying water to meet the intended purpose(s).

Soil (Series, Texture, and Map Unit) *Select the soil to manage for:*

Critical soil to manage: **Harkey L; 33-41** Intake Family (in/hr): **0.50**

Soil Interpretations for Irrigaiton

Crop Name	Rooting Depth	Moisture Replacement Depth (ft)	Water Holding Capacity (in)	Mgt Allowed Depletion (MAD) (%)	Net Water to Replace (in)	Time Needed to Infiltrate (hrs)
Alfalfa, hay, southern; Las Cruces	Deep	4.0	8.0	50%	4.0	5.4

Crop Consumptive Use (CU) Information *(inches/month needed)*

Crop: Alfalfa, hay, southern; Las Cruces **Total Irrigation Needed:** 47.7 ac in/ac

Month	Est. Frequency *(days between irr.)*	In/Mo	Month	Est. Frequency *(days between irr.)*	In/Mo
Jan		0.0	Jul	12	9.7
Feb		0.0	Aug	16	7.7
Mar	>1 Mo.	1.6	Sep	21	5.7
Apr	29	4.2	Oct	>1 Mo.	2.7
May	17	6.9	Nov		0.0
Jun	13	9.3	Dec		0.0

Crop: **Total Irrigation Needed:** ac in/ac

Month	Est. Frequency *(days between irr.)*	In/Mo	Month	Est. Frequency *(days between irr.)*	In/Mo
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** (days between irr.)	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Farm Irrigation Rating System (FIRS, irrigation system efficiency)

Type of System:	**Flood, controlled**	System Efficiency (%):	**80%**
System Capacity (Ditch/Pump/Well):	**6000**	GPM	

Crop Grown	Days of Operation for CU by crop (days)	Est. System Applicaiton Rate (in/day)	Needed Application Rate (in/day)	System Evaluation
Alfalfa, hay, southern; Las Cruces	10	4.92	0.32	System meets CU for crop

Operation and Maintenance Requirements

1. Irrigate when the soils reaches the MAD level, determined by soil moisture monitoring. Use one of the following methods to monitor soil moisture; the feel method, tensiometers, electrical resistance blocks, or moisture probes. Drip or Center Pivots can be irrigated on an as needed basis to meet daily CU.

2. Test irrigation water for Nitrate and Salts (Total Desolved Solids/Electrical Conductivity)

3. Do not exceed the net water to replace listed above when irrigating, unless salts are being managed.

4. Do not apply water at rates that cause runoff or erosion.

5. Monitor the soil to maintain: pH, permeability, salinity, and structure.

6. Application of pond effluent shall not exceed the crop needs (water and nutrient), and will not exceed the water holding capacity listed above.

7. Consider using crops such as sorghum, cotton, or winter wheat when water supplies are short.

8. Avoid traffic on wet soils to minimize soil compaction.

Additional Requirements

Job Approval and Completion

Client: Isaac Domingues **Date:** 11-23-2009

Conservationist: [signature] **Date:** 11-23-09

Completed by: [signature] **Date:**

NRCS - WEQ CALCULATIONS

Version 9 03 11-19-2007

Producer: Dominguez Dairy #1 | Planner: Chet Wyant | Location: Mesquite, NM | Tract: 1579 | Field: DM-5

Crop Rotation: Alfalfa | Climate Data Station: NM, LAS CRUCES | Site "C" Value: 150

Tillage Direction (NS/EW): NS | Length/width ratio: 1 | Field Direction (NS/EW): NS | Field Width (Ft.): 1412

Irrigation (Y or N): Y | Soil "I": 86 | Wind Erodibility Group: 4L (1-7) | TWF: 2 (see instr.)

Sum Period Erosion (t/ac): 0.0 | No. Yrs in Rotation: 1.0 | Av. Annual Wind Erosion: (t/ac/yr)

Calculations and Output

Mgt Periods Dates		Irr.	Soil	Ridge Roughness				Random Roughness	Unsheltered Distance				SCe	Erosion			
Begin	End	No. of (#)	"I" (t/ac)	Dev. (deg)	Ht. (in.)	Sp. (in.)	"Krd" (factor)	"Krr" (factor)	Dev. (deg)	Prep. (factor)	WED (factor)	"L" (ft)	"V" (lbs/ac)	"E" (t/ac)	EWE (%)	"IF" (%)	Loss (t/ac)
11/1	01/02/10	0	56	0.0	0	0	1.00	0.99	0.0	4.0	1.000	1412	0	62.7	12.6	1.00	0.08
1/2	01/17/10	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			2.0	1.00	
1/17	02/01/10	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			2.0	1.00	
2/1	02/16/10	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.7	0.87	
2/16	03/03/10	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.7	1.00	
3/3	03/18/10	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.9	0.87	
3/18	04/02/10	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.6	1.00	
4/2	04/17/10	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			8.8	0.87	
4/17	05/02/10	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			8.6	0.87	
5/2	05/17/10	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			5.3	1.00	
5/17	06/06/10	2	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			5.6	0.80	
6/6	06/21/10	0	56	22.5	0	0	1.00	0.94	22.5	1.2	1.030	1454			0.8	1.00	
6/21	07/11/10	2	56	22.5	0	0	1.00	0.94	22.5	1.2	1.030	1454			0.7	0.80	
7/11	07/26/10	0	56	67.5	0	0	1.00	0.94	67.5	1.1	1.030	1454			0.2	1.00	
7/26	08/15/10	2	56	67.5	0	0	1.00	0.94	67.5	1.1	1.030	1454			0.2	0.80	
8/15	08/30/10	0	56	90.0	0	0	1.00	0.94	90.0	2.8	1.010	1426			0.1	1.00	
8/30	09/19/10	2	56	90.0	0	0	1.00	0.94	90.0	2.8	1.010	1426			0.6	0.80	
9/19	11/01/10	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.00	1412			1.7	0.95	
		0															
		0															
		0															
		0															
		0															
		0															
		0															
		0															

NRCS

RUSLE2 Worksheet Erosion Calculation Record

Info: Alfalfa Nov'09 to Oct'10

Tract #: 1579
Owner name: DOMINGUEZ DAIRY
Field name: DM-5

Location: New Mexico\DonaAna County\NM_Dona Ana R 9
Soil: Hg HARKEY LOAM\HARKEY loam 85%
Slope length (horiz): 100.0 ft
Avg. slope steepness: 0.50 %
T value: 5.0 t/ac/yr

Alternatives:

Management	*Contouring*	*Strips / barriers*	*Diversion/terrace, sediment basin*	*Cons. plan. soil loss, t/ac/yr*	*Description*
b.Mullti-year Rotation Templates\Alfalfa-Small Grain, 5yr, 5cuts CMZ 23	b. absolute row grade 0.5 percent	(none)	(none)	0.042	

Year 2

Dairy Annual Nutrient Manager

Date	11/30/2009	NM STATE UNIVERSITY	Dairy Extension Program	Victor E. Cabrera Dairy Specialist dairy.nmsu.edu: Tools vcabrera@nmsu.edu	PLANNED		[redacted]
Dairy	Dominguez Dairy						
Crop_Year	2010-2011						
Field_ID	DM-5				N Loss	0%	
Area (ac)	52				Goal/Real	Nutrient Needed	
	Month - Month	Crop		Unit	Yield	N	P
1st Crop	Nov'10-Oct'11	Alfalfa, for hay		t/ac	12	31446	2946
2nd Crop						0	0
						0	0
Total		Nutrient Needed				31,446	2,946
						Soil Analyses	
Texture by Feel		Sandy Clay				N	P
Nutrient Available in Soil				lb/ac		142	97.38
		Nutrient Still Needed				24,062	-2,117
						Effluent Analyses	
						N	P
Effluent Manure Application		NM Dairy Ponds Net from J.S. 590 L&MLA		ac-in	0.712	926	420
				ac-in		0	0
				ac-in		0	0
		Nutrient Still Needed				23,136	-2,538
						Manure Analyses	
						N	P
Dry Manure Application				t/ac		0	0
				t/ac		0	0
				t/ac		0	0
		Nutrient Still Needed				23,136	-2,538
						Fertilizer Content	
						N	P
Chemical Fertilizers Applied				lb/ac		0	0
				lb/ac		0	0
				lb/ac		0	0
		Annual Nutrient Balance				**23,136**	**-2,538**

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	Acres:	52	Date:	11/2/2009	Field ID:	DM-5

Application Information *(Enter the units that will be or has been applied to the field)*:					
Crop Rotation:	Alfalfa			Needed for field (acin):	37.024
Liquid Applied:	0.712	Acin/ac		(gal):	1,005,202
Solids Applied:		ton/ac		Needed for field (tons):	
Liquid Loads Applied:		1000gal/ac		Loads needed for field:	

Nutrient Content of Organic Material

Solid-Lab Report	% Moisture	TKN (%) (dry)	NH_4-N (ppm) (dry)	P_2O_5 (%) (dry)	K_2O (%) (dry)
Fill in Lab data:					

Solid Book Values (select even if test values are used)	% Moisture		TKN (lbs/wet ton)		NH_4-N (lbs/ton)		P_2O_5 (lbs/wet ton)		K_2O (lbs/wet ton)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
Beef (DM)		0		0		0.0		0		0

Liquid-Lab Report	NH_3-N (mg/L)	TKN (mg/L)	NO_3-N (mg/L)	Tot-PO_4 (mg/L)	K (mg/L)
Fill in Lab data:		205	0.17		

Liquid	% Moisture		TKN (lbs/acin)		NH_4-N (lbs/acin)		P_2O_5 (lbs/acin)		K_2O (lbs/acin)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		NH_4-N (lbs/1000gal)		P_2O_5 (lbs/1000gal)		K_2O (lbs/1000gal)	
			Book	Test	Book	Test	Book	Test	Book	Test
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	Type of Climate	Percent Remaining	NH_4-N Remaining
Broadcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	**NH_4-N Remaining**
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	20.8 (lbs/acin) NH4-N 0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	Percent Nutrient Available the 1st Year			
	Organic N	P	K	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	Solid Source
Lagoon or diluted Pond	40 %	75 %	80 %	Liquid Source

Solid	Organic N (lbs/ton)	P_2O_5 (lbs/ton)	K_2O (lbs/ton)
	0	0	0
Liquid	**Organic N (lbs/acin)**	**P_2O_5 (lbs/acin)**	**K_2O (lbs/acin)**
	8	26	205
	Organic N (lbs/100gal)	**P_2O_5 (lbs/1000gal)**	**K_2O (lbs/1000gal)**
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	Soil Drainage Class (See Survey Information)	Percent Remaining (%)
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	lbs/1000gal	lbs/ac in	lbs/ton
N	0.0	25	0
P_2O_5	0.0	26	0
K_2O	0.0	205	0

Total Nutrients Applied (net to the field)	All Forms N (lbs/ac)	P_2O_5 (lbs/ac)	K_2O (lbs/ac)
	18.1	18.7	145.7

USDA Natural Resources Conservation Service
PLANTS

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

Return to the crop list and make a new selection of crop(s)

Alfalfa, for hay
Forage
Medicago sativa
Plant part harvested: Aboveground biomass
Crop yield unit: ton

Nutrients in harvested part (lb/ton) at 9.65% moisture percentage.

Nitrogen	Phosphorus	Potassium
50.3935	4.7217	38.2939

Nutrients removed in harvested part (lb/acre) at 12 ton yield level.

Nitrogen	Phosphorus	Potassium
604.7224	56.6599	459.5271

Nutrients removed in harvested part (lb/ton) at 12 ton yield level and 52 acres.

Nitrogen	Phosphorus	Potassium
31445.5665	2946.3165	23895.4086

Element-Fertilizer Equivalents | Average NPK Percentages

«Contents»

Intellectual Property Statement / How to Cite the PLANTS Database

Disclaimers

USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: **Mon 4:38 PM - 11/30/2009**

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-5	Date:	Nov'10
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Alfalfa
Soil Permeablity (in/hr):	0.5	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the approprate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <8 ppm	Low 8-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
				X		4
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
		X				1
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
				X		4
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >500-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		X				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 80% Dry Matter as Supplemental Feed	Pasture 80 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>65-100 ft wide	20-65 feet wide	< 20 feet wide	No Buffer	
			X			3

P Hazard Class:	Low	Total Index Points:	16.5
Phosphorus Application Classification:	N Based		

Notes:

This evaluation has a Low P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 11'10 to 10/11) 0.08 t/ac + (RUSLE 2) 0.042 t/ac = 0.12 t/ac

NRCS Natural Resources Conservation Service

Irrigation Water Management

Conservation Practice Job Sheet 449

Natural Resources Conservation Service (NRCS) Jan, 2006

Client: Dominguez Dairy **Tract:** 1579
Planner: Chet Wyant **Field(s) No.:** DM-5
Current Land Use: Alfalfa **Total Acres:** 52
Date: 11/20/2009 **Date to apply:** Nov'10

See the Conservation Plan map for the location of the field(s) to applying IWM.

Purposes (check all that apply)

- ☐ Manage soil moisture to improve crops
- ☑ Optimize use of water
- ☑ Minimize Irrigation Erosion
- ☐ Decrease non-point source pollution
- ☐ Manage salt in the root zone
- ☐ Manage air, soil, or plant microclimate

Conditions where practice applies

This practice is applicable to all irrigated lands. An irrigation system adapted for the site conditions is available and capable of applying water to meet the intended purpose(s).

Soil (Series, Texture, and Map Unit) *Select the soil to manage for:*

Critical soil to manage: **Harkey L; 33-41** Intake Family (in/hr): **0.50**

Soil Interpretations for Irrigaiton

Crop Name	Rooting Depth	Moisture Replacement Depth (ft)	Water Holding Capacity (in)	Mgt Allowed Depletion (MAD) (%)	Net Water to Replace (in)	Time Needed to Infiltrate (hrs)
Alfalfa, hay, southern; Las Cruces	Deep	4.0	8.0	50%	4.0	5.4

Crop Consumptive Use (CU) Information *(inches/month needed)*

Crop:	Alfalfa, hay, southern; Las Cruces		Total Irrigation Needed:	47.7 ac in/ac	
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul	12	9.7
Feb		0.0	Aug	16	7.7
Mar	>1 Mo.	1.6	Sep	21	5.7
Apr	29	4.2	Oct	>1 Mo.	2.7
May	17	6.9	Nov		0.0
Jun	13	9.3	Dec		0.0

Crop:			Total Irrigation Needed:	ac in/ac	
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency (days between irr.)**	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Farm Irrigation Rating System (FIRS, irrigation system efficiency)

Type of System:	**Flood, controlled**		System Efficiency (%):	**80%**
System Capacity (Ditch/Pump/Well):		**6000**	GPM	
Crop Grown	**Days of Operation for CU by crop (days)**	**Est. System Application Rate (in/day)**	**Needed Application Rate (in/day)**	**System Evaluation**
Alfalfa, hay, southern; Las Cruces	10	4.92	0.32	System meets CU for crop

Operation and Maintenance Requirements

1. Irrigate when the soils reaches the MAD level, determined by soil moisture monitoring. Use one of the following methods to monitor soil moisture; the feel method, tensiometers, electrical resistance blocks, or moisture probes. Drip or Center Pivots can be irrigated on an as needed basis to meet daily CU.

2. Test irrigation water for Nitrate and Salts (Total Desolved Solids/Electrical Conductivity)

3. Do not exceed the net water to replace listed above when irrigating, unless salts are being managed.

4. Do not apply water at rates that cause runoff or erosion.

5. Monitor the soil to maintain: pH, permeability, salinity, and structure.

6. Application of pond effluent shall not exceed the crop needs (water and nutrient), and will not exceed the water holding capacity listed above.

7. Consider using crops such as sorghum, cotton, or winter wheat when water supplies are short.

8. Avoid traffic on wet soils to minimize soil compaction.

Additional Requirements

Job Approval and Completion

Client: Joe Domingez **Date:** 11-23-2009

Conservationist: [signature] **Date:** 11-23-09

Completed by: [signature] **Date:**

NRCS - WEQ CALCULATIONS

Version 9.03 11-19-2007

Producer:	Dominguez Dairy #1	Planner:	Chet Wyant	Location:	Mesquite, NM	Tract:	1579	Field:	DM-5		
Crop Rotation:	Alfalfa	Climate Data Station:	NM, LAS CRUCES	Site "C" Value:	150						
Tillage Direction (NS/EW):	NS	Length/width ratio:	1	Field Direction (NS/EW):	NS	Field Width (Ft.):	1412				
Irrigation (Y or N):	Y	Soil "I":	86	Wind Erodibility Group:	4L	(1-7)	TWF:	2	(see instr.)		
Sum Period Erosion (t/ac):	0.0	No. Yrs in Rotation:	1.0	Av. Annual Wind Erosion:		(t/ac/yr)					

Calculations and Output

Mgt Periods		Irr.	Soil	Ridge Roughness				Random Roughness	Unsheltered Distance				SGe	Erosion			
Dates		No. of	"I"	Dev.	Ht.	Sp.	"Krd"	"Krr"	Dev.	Prep.	WED	"L"	"V"	"E"	EWE	"IF"	Loss
Begin	End	(#)	(t/ac)	(deg)	(in.)	(in.)	(factor)	(factor)	(deg)	(factor)	(factor)	(ft)	(lbs/ac)	(t/ac)	(%)	(%)	(t/ac)
11/1	01/02/11	0	56	0.0	0	0	1.00	0.99	0.0	4.0	1.000	1412	0	62.7	12.6	1.00	0.08
1/2	01/17/11	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			2.0	1.00	
1/17	02/01/11	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			2.0	1.00	
2/1	02/16/11	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.7	0.87	
2/16	03/03/11	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.7	1.00	
3/3	03/18/11	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.9	0.87	
3/18	04/02/11	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.6	1.00	
4/2	04/17/11	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			8.8	0.87	
4/17	05/02/11	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			8.6	0.87	
5/2	05/17/11	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			5.3	1.00	
5/17	06/06/11	2	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			5.6	0.80	
6/6	06/21/11	0	56	22.5	0	0	1.00	0.94	22.5	1.2	1.030	1454			0.8	1.00	
6/21	07/11/11	2	56	22.5	0	0	1.00	0.94	22.5	1.2	1.030	1454			0.7	0.80	
7/11	07/26/11	0	56	67.5	0	0	1.00	0.94	67.5	1.1	1.030	1454			0.2	1.00	
7/26	08/15/11	2	56	67.5	0	0	1.00	0.94	67.5	1.1	1.030	1454			0.2	0.80	
8/15	08/30/11	0	56	90.0	0	0	1.00	0.94	90.0	2.8	1.010	1426			0.1	1.00	
8/30	09/19/11	2	56	90.0	0	0	1.00	0.94	90.0	2.8	1.010	1426			0.6	0.80	
9/19	11/01/11	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.00	1412			1.7	0.95	
		0															
		0															
		0															
		0															
		0															
		0															
		0															
		0															

RUSLE2 Worksheet Erosion Calculation Record

Info: Alfalfa Nov'10 to Oct'11

Tract #: 1579
Owner name: DOMINGUEZ DAIRY
Field name: DM-5

Location: New Mexico\DonaAna County\NM_Dona Ana R 9
Soil: Hg HARKEY LOAM\HARKEY loam 85%
Slope length (horiz): 100.0 ft
Avg. slope steepness: 0.50 %
T value: 5.0 t/ac/yr

Alternatives:

Management	*Contouring*	*Strips / barriers*	*Diversion/terrace, sediment basin*	*Cons. plan. soil loss, t/ac/yr*	*Description*
b.Mullti-year Rotation Templates\Alfalfa-Small Grain, 5yr, 5cuts CMZ 23	b. absolute row grade 0.5 percent	(none)	(none)	0.042	

Year 3

Dairy Annual Nutrient Manager

Date	11/30/2009	NM STATE UNIVERSITY	Dairy Extension Program	Victor E. Cabrera Dairy Specialist dairy.nmsu.edu: Tools vcabrera@nmsu.edu	PLANNED	[redacted]	
Dairy	Dominguez Dairy						
Crop_Year	2011-2012						
Field_ID	DM-5				N Loss	0%	
Area (ac)	52				Goal/Real	Nutrient Needed	
	Month - Month	Crop		Unit	Yield	N	P
1st Crop	Nov'11-Oct'12	Alfalfa, for hay		t/ac	12	31446	2946
2nd Crop						0	0
						0	0
Total		Nutrient Needed				31,446	2,946
						Soil Analyses	
Texture by Feel		Sandy Clay				N	P
Nutrient Available in Soil				lb/ac		142	97.38
		Nutrient Still Needed				24,062	-2,117
						Effluent Analyses	
						N	P
Effluent Manure Application		NM Dairy Ponds Net from J.S. 590 L&MLA		ac-in	0.712	926	420
				ac-in		0	0
				ac-in		0	0
		Nutrient Still Needed				23,136	-2,538
						Manure Analyses	
						N	P
Dry Manure Application				t/ac		0	0
				t/ac		0	0
				t/ac		0	0
		Nutrient Still Needed				23,136	-2,538
						Fertilizer Content	
						N	P
Chemical Fertilizers Applied				lb/ac		0	0
				lb/ac		0	0
				lb/ac		0	0
		Annual Nutrient Balance				23,136	-2,538

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	Acres:	52	Date:	11/2/2009	Field ID:	DM-5

Application information (Enter the units that will be or has been applied to the field):					
Crop Rotation:	Alfalfa			Needed for field (acin):	37.024
Liquid Applied:	0.712	Acin/ac		(gal):	1,005,202
Solids Applied:		ton/ac		Needed for field (tons):	
Liquid Loads Applied:		1000gal/ac		Loads needed for field:	

Nutrient Content of Organic Material

Solid-Lab Report	% Moisture	TKN (%) (dry)	NH_4-N (ppm) (dry)	P_2O_5 (%) (dry)	K_2O (%) (dry)
Fill in Lab data:					

Solid Book Values (select even if test values are used)	% Moisture		TKN (lbs/wet ton)		NH_4-N (lbs/ton)		P_2O_5 (lbs/wet ton)		K_2O (lbs/wet ton)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
Beef (DM)		0		0		0.0		0		0

Liquid-Lab Report	NH_3-N (mg/L)	TKN (mg/L)	NO_3-N (mg/L)	Tot-PO_4 (mg/L)	K (mg/L)
Fill in Lab data:		205	0.17		

Liquid	% Moisture		TKN (lbs/acin)		NH_4-N (lbs/acin)		P_2O_5 (lbs/acin)		K_2O (lbs/acin)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		NH_4-N (lbs/1000gal)		P_2O_5 (lbs/1000gal)		K_2O (lbs/1000gal)	
			Book	Test	Book	Test	Book	Test	Book	Test
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	Type of Climate	Percent Remaining	NH_4-N Remaining
Broadcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	20.8 (lbs/acin) NH4-N 0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	Percent Nutrient Available the 1st Year			
	Organic N	P	K	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	Solid Source
Lagoon or diluted Pond	40 %	75 %	80 %	Liquid Source

Solid	Organic N (lbs/ton)	P_2O_5 (lbs/ton)	K_2O (lbs/ton)
	0	0	0
Liquid	**Organic N (lbs/acin)**	**P_2O_5 (lbs/acin)**	**K_2O (lbs/acin)**
	8	26	205
	Organic N (lbs/100gal)	**P_2O_5 (lbs/1000gal)**	**K_2O (lbs/1000gal)**
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	Soil Drainage Class (See Survey Information)	Percent Remaining (%)
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	lbs/1000gal	lbs/ac in	lbs/ton
N	0.0	25	0
P_2O_5	0.0	26	0
K_2O	0.0	205	0

Total Nutrients Applied (net to the field)	All Forms N (lbs/ac)	P_2O_5 (lbs/ac)	K_2O (lbs/ac)
	18.1	18.7	145.7

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

- Return to the crop list and make a new selection of crop(s)

Alfalfa, for hay
Forage
Medicago sativa
Plant part harvested: Aboveground biomass
Crop yield unit: ton

Nutrients in harvested part (lb/ton) at 9.65% moisture percentage.

Nitrogen	Phosphorus	Potassium
50.3935	4.7217	38.2939

Nutrients removed in harvested part (lb/acre) at 12 ton yield level.

Nitrogen	Phosphorus	Potassium
604.7224	56.6599	459.5271

Nutrients removed in harvested part (lb/ton) at 12 ton yield level and 52 acres.

Nitrogen	Phosphorus	Potassium
31445.5665	2946.3165	23895.4086

Element-Fertilizer Equivalents | Average NPK Percentages

«Contents»

- Intellectual Property Statement / How to Cite the PLANTS Database
- Disclaimers
- USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: Mon 4:38 PM - 11/30/2009

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-5	Date:	Nov'11
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Alfalfa
Soil Permeablity (in/hr):	0.5	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the approprate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <8 ppm	Low 8-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
				X		4
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
		X				1
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
				X		4
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >500-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		x				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 80% Dry Matter as Supplemental Feed	Pasture 80 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>65-100 ft wide	20-65 feet wide	< 20 feet wide	No Buffer	
			x			3

P Hazard Class:	Low	Total Index Points:	16.5
Phosphorus Application Classification:	N Based		

Notes:

This evaluation has a Low P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 11'11 to 10/12) 0.08 t/ac + (RUSLE 2) 0.042 t/ac = 0.12 t/ac

NRCS Natural Resources Conservation Service

Irrigation Water Management

Conservation Practice Job Sheet 449

Natural Resources Conservation Service (NRCS) Jan, 2006

Client: Dominguez Dairy **Tract:** 1579
Planner: Chet Wyant **Field(s) No.:** DM-5
Current Land Use: Alfalfa **Total Acres:** 52
Date: 11/20/2009 **Date to apply:** Nov'11

See the Conservation Plan map for the location of the field(s) to applying IWM.

Purposes (check all that apply)

- [ ] Manage soil moisture to improve crops
- [x] Optimize use of water
- [x] Minimize Irrigation Erosion
- [ ] Decrease non-point source pollution
- [ ] Manage salt in the root zone
- [ ] Manage air, soil, or plant microclimate

Conditions where practice applies

This practice is applicable to all irrigated lands. An irrigation system adapted for the site conditions is available and capable of applying water to meet the intended purpose(s).

Soil (Series, Texture, and Map Unit) *Select the soil to manage for:*

Critical soil to manage:	Harkey L; 33-41	Intake Family (in/hr):	0.50

Soil Interpretations for Irrigaiton

Crop Name	Rooting Depth	Moisture Replacement Depth (ft)	Water Holding Capacity (in)	Mgt Allowed Depletion (MAD) (%)	Net Water to Replace (in)	Time Needed to Infiltrate (hrs)
Alfalfa, hay, southern; Las Cruces	Deep	4.0	8.0	50%	4.0	5.4

Crop Consumptive Use (CU) Information *(inches/month needed)*

Crop:	Alfalfa, hay, southern; Las Cruces		Total Irrigation Needed:	47.7 ac in/ac	
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul	12	9.7
Feb		0.0	Aug	16	7.7
Mar	>1 Mo.	1.6	Sep	21	5.7
Apr	29	4.2	Oct	>1 Mo.	2.7
May	17	6.9	Nov		0.0
Jun	13	9.3	Dec		0.0

Crop:			Total Irrigation Needed:	ac in/ac	
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** (days between irr.)	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Farm Irrigation Rating System (FIRS, irrigation system efficiency)

Type of System: **Flood, controlled** System Efficiency (%): **80%**

System Capacity (Ditch/Pump/Well): **6000** GPM

Crop Grown	Days of Operation for CU by crop (days)	Est. System Applicaiton Rate (in/day)	Needed Application Rate (in/day)	System Evaluation
Alfalfa, hay, southern; Las Cruces	10	4.92	0.32	System meets CU for crop

Operation and Maintenance Requirements

1. Irrigate when the soils reaches the MAD level, determined by soil moisture monitoring. Use one of the following methods to monitor soil moisture; the feel method, tensiometers, electrical resistance blocks, or moisture probes. Drip or Center Pivots can be irrigated on an as needed basis to meet daily CU.

2. Test irrigation water for Nitrate and Salts (Total Desolved Solids/Electrical Conductivity)

3. Do not exceed the net water to replace listed above when irrigating, unless salts are being managed.

4. Do not apply water at rates that cause runoff or erosion.

5. Monitor the soil to maintain: pH, permeability, salinity, and structure.

6. Application of pond effluent shall not exceed the crop needs (water and nutrient), and will not exceed the water holding capacity listed above.

7. Consider using crops such as sorghum, cotton, or winter wheat when water supplies are short.

8. Avoid traffic on wet soils to minimize soil compaction.

Additional Requirements

Job Approval and Completion

Client: Israel Dominguez **Date:** 11-23-2009

Conservationist: [illegible] **Date:** 11-23-09

Completed by: [illegible] **Date:**

NRCS - WEQ CALCULATIONS

Version 9.03 11-19-2007

Producer:	Dominguez Dairy #1	Planner:	Chet Wyant	Location:	Mesquite, NM	Tract:	1579	Field:	DM-5		
Crop Rotation:	Alfalfa	Climate Data Station:	NM, LAS CRUCES	Site "C" Value:	150						
Tillage Direction (NS/EW):	NS	Length/width ratio:	1	Field Direction (NS/EW):	NS	Field Width (Ft.):	1412				
Irrigation (Y or N):	Y	Soil "I":	86	Wind Erodibility Group:	4L	(1-7)	TWF:	2	(see instr.)		
Sum Period Erosion (t/ac):	0.0	No. Yrs in Rotation:	1.0	Av. Annual Wind Erosion:		(t/ac/yr)					

Calculations and Output

Mgt Periods		Irr.	Soil	Ridge Roughness				Random Roughness	Unsheltered Distance				SGe	Erosion			
Dates		No. of	"I"	Dev.	Ht.	Sp.	"Krd"	"Krr"	Dev.	Prep.	WED	"L"	"V"	"E"	EWE	"IF"	Loss
Begin	End	(#)	(t/ac)	(deg)	(in.)	(in.)	(factor)	(factor)	(deg)	(factor)	(factor)	(ft)	(lbs/ac)	(t/ac)	(%)	(%)	(t/ac)
11/1	01/02/12	0	56	0.0	0	0	1.00	0.99	0.0	4.0	1.000	1412	0	62.7	12.6	1.00	0.08
1/2	01/17/12	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			2.0	1.00	
1/17	02/01/12	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			2.0	1.00	
2/1	02/16/12	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.7	0.87	
2/16	03/02/12	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			11.9	1.00	
3/2	03/17/12	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.9	0.87	
3/17	04/01/12	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.9	1.00	
4/1	04/16/12	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			8.8	0.87	
4/16	05/01/12	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			8.8	0.87	
5/1	05/16/12	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			5.3	1.00	
5/16	06/05/12	2	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			5.9	0.80	
6/5	06/20/12	0	56	22.5	0	0	1.00	0.94	22.5	1.2	1.030	1454			0.8	1.00	
6/20	07/10/12	2	56	22.5	0	0	1.00	0.94	22.5	1.2	1.030	1454			0.7	0.80	
7/10	07/25/12	0	56	67.5	0	0	1.00	0.94	67.5	1.1	1.030	1454			0.2	1.00	
7/25	08/14/12	2	56	67.5	0	0	1.00	0.94	67.5	1.1	1.030	1454			0.2	0.80	
8/14	08/29/12	0	56	90.0	0	0	1.00	0.94	90.0	2.8	1.010	1426			0.1	1.00	
8/29	09/18/12	2	56	90.0	0	0	1.00	0.94	90.0	2.8	1.010	1426			0.5	0.80	
9/18	11/01/12	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.00	1412			1.7	0.95	
		0															
		0															
		0															
		0															
		0															
		0															
		0															
		0															

NRCS

RUSLE2 Worksheet Erosion Calculation Record

Info: Alfalfa Nov'11 to Oct'12

Tract #: 1579
Owner name: DOMINGUEZ DAIRY
Field name: DM-5

Location: New Mexico\DonaAna County\NM_Dona Ana R 9
Soil: Hg HARKEY LOAM\HARKEY loam 85%
Slope length (horiz): 100.0 ft
Avg. slope steepness: 0.50 %
T value: 5.0 t/ac/yr

Alternatives:

Management	*Contouring*	*Strips / barriers*	*Diversion/terrace, sediment basin*	*Cons. plan. soil loss, t/ac/yr*	*Description*
b.Mullti-year Rotation Templates\Alfalfa-Small Grain, 5yr, 5cuts CMZ 23	b. absolute row grade 0.5 percent	(none)	(none)	0.042	

Year 4

Dairy Annual Nutrient Manager

Date	11/30/2009	NM STATE UNIVERSITY Dairy Extension Program	Victor E. Cabrera Dairy Specialist dairy.nmsu.edu: Tools vcabrera@nmsu.edu	PLANNED		[redacted]	
Dairy	Dominguez Dairy						
Crop_Year	2012-2013				N Loss	0%	
Field_ID	DM-5				Goal/Real	Nutrient Needed	
Area (ac)	52						
	Month - Month	Crop		Unit	Yield	N	P
1st Crop	Nov'12-Oct'13	Alfalfa, for hay		t/ac	12	31446	2946
2nd Crop						0	0
						0	0
Total		Nutrient Needed				31,446	2,946
						Soil Analyses	
Texture by Feel		Sandy Clay				N	P
Nutrient Available in Soil				lb/ac		142	97.38
		Nutrient Still Needed				24,062	-2,117
						Effluent Analyses	
						N	P
Effluent Manure Application		NM Dairy Ponds Net from J.S. 590 L&MLA		ac-in	0.712	926	420
				ac-in		0	0
				ac-in		0	0
		Nutrient Still Needed				23,136	-2,538
						Manure Analyses	
						N	P
Dry Manure Application				t/ac		0	0
				t/ac		0	0
				t/ac		0	0
		Nutrient Still Needed				23,136	-2,538
						Fertilizer Content	
						N	P
Chemical Fertilizers Applied				lb/ac		0	0
				lb/ac		0	0
				lb/ac		0	0
		Annual Nutrient Balance				**23,136**	**-2,538**

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	Acres:	52	Date:	11/2/2009	Field ID:	DM-5

Application information					
nter the units that will be or has been applied to the field):	Crop Rotation: Alfalfa			Needed for field (acin):	37.024
	Liquid Applied:	0.712	Acin/ac	(gal):	1,005,202
	Solids Applied:		ton/ac	Needed for field (tons):	
	Liquid Loads Applied:		1000gal/ac	Loads needed for field:	

Nutrient Content of Organic Material

Solid-Lab Report	% Moisture	TKN (%) (dry)	NH_4-N (ppm) (dry)	P_2O_5 (%) (dry)	K_2O (%) (dry)
Fill in Lab data:					

Solid Book Values (select even if test values are used)	% Moisture		TKN (lbs/wet ton)		NH_4-N (lbs/ton)		P_2O_5 (lbs/wet ton)		K_2O (lbs/wet ton)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
Beef (DM)		0		0		0.0		0		0

Liquid-Lab Report	NH_3-N (mg/L)	TKN (mg/L)	NO_3-N (mg/L)	Tot-PO_4 (mg/L)	K (mg/L)
Fill in Lab data:		205	0.17		

Liquid	% Moisture		TKN (lbs/acin)		NH_4-N (lbs/acin)		P_2O_5 (lbs/acin)		K_2O (lbs/acin)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		NH_4-N (lbs/1000gal)		P_2O_5 (lbs/1000gal)		K_2O (lbs/1000gal)	
			Book	Test	Book	Test	Book	Test	Book	Test
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	Type of Climate	Percent Remaining	NH_4-N Remaining
Broadcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	**NH_4-N Remaining**
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	20.8 (lbs/acin) NH4-N 0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	Percent Nutrient Available the 1st Year			
	Organic N	P	K	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	Solid Source
Lagoon or diluted Pond	40 %	75 %	80 %	Liquid Source

Solid	Organic N (lbs/ton)	P_2O_5 (lbs/ton)	K_2O (lbs/ton)
	0	0	0
Liquid	**Organic N (lbs/acin)**	**P_2O_5 (lbs/acin)**	**K_2O (lbs/acin)**
	8	26	205
	Organic N (lbs/100gal)	**P_2O_5 (lbs/1000gal)**	**K_2O (lbs/1000gal)**
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	Soil Drainage Class (See Survey Information)	Percent Remaining (%)
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	lbs/1000gal	lbs/ac in	lbs/ton
N	0.0	25	0
P_2O_5	0.0	26	0
K_2O	0.0	205	0

Total Nutrients Applied (net to the field)	All Forms N (lbs/ac)	P_2O_5 (lbs/ac)	K_2O (lbs/ac)
	18.1	18.7	145.7

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

Return to the crop list and make a new selection of crop(s)

Alfalfa, for hay
Forage
Medicago sativa
Plant part harvested: Aboveground biomass
Crop yield unit: ton

Nutrients in harvested part (lb/ton) at 9.65% moisture percentage.			**Nutrients removed in harvested part (lb/acre) at** 12 ton yield level.		
Nitrogen	**Phosphorus**	**Potassium**	**Nitrogen**	**Phosphorus**	**Potassium**
50.3935	4.7217	38.2939	604.7224	56.6599	459.5271

Nutrients removed in harvested part (lb/ton) at 12 ton yield level and 52 acres.

Nitrogen	**Phosphorus**	**Potassium**
31445.5665	2946.3165	23895.4086

Element-Fertilizer Equivalents

Average NPK Percentages

«Contents»

Intellectual Property Statement / How to Cite the PLANTS Database

Disclaimers

USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: Mon 4:38 PM - 11/30/2009

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-5	Date:	Nov'12
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Alfalfa
Soil Permeablity (in/hr):	0.5	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the approprate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <8 ppm	Low 8-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
				X		4
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
		X				1
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
				X		4
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >500-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		x				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 80% Dry Matter as Supplemental Feed	Pasture 80 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>65-100 ft wide	20-65 feet wide	< 20 feet wide	No Buffer	
			x			3

P Hazard Class:	Low	Total Index Points:	16.5
Phosphorus Application Classification:	N Based		

Notes:

This evaluation has a Low P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 11'12 to 10/13) 0.08 t/ac + (RUSLE 2) 0.042 t/ac = 0.12 t/ac

NRCS Natural Resources Conservation Service

Irrigation Water Management

Conservation Practice Job Sheet *449*

Natural Resources Conservation Service (NRCS) Jan, 2006

Client: Dominguez Dairy **Tract:** 1579
Planner: Chet Wyant **Field(s) No.:** DM-5
Current Land Use: Alfalfa **Total Acres:** 52
Date: 11/20/2009 **Date to apply:** Nov'12

See the Conservation Plan map for the location of the field(s) to applying IWM.

Purposes (check all that apply)

- ☐ Manage soil moisture to improve crops
- ☑ Optimize use of water
- ☑ Minimize Irrigation Erosion
- ☐ Decrease non-point source pollution
- ☐ Manage salt in the root zone
- ☐ Manage air, soil, or plant microclimate

Conditions where practice applies

This practice is applicable to all irrigated lands. An irrigation system adapted for the site conditions is available and capable of applying water to meet the intended purpose(s).

Soil (Series, Texture, and Map Unit) *Select the soil to manage for:*

Critical soil to manage: **Harkey L; 33-41** Intake Family (in/hr): **0.50**

Soil Interpretations for Irrigaiton

Crop Name	Rooting Depth	Moisture Replacement Depth (ft)	Water Holding Capacity (in)	Mgt Allowed Depletion (MAD) (%)	Net Water to Replace (in)	Time Needed to Infiltrate (hrs)
Alfalfa, hay, southern; Las Cruces	Deep	4.0	8.0	50%	4.0	5.4

Crop Consumptive Use (CU) Information *(inches/month needed)*

Crop: Alfalfa, hay, southern; Las Cruces **Total Irrigation Needed:** 47.7 ac in/ac

Month	Est. Frequency *(days between irr.)*	In/Mo	Month	Est. Frequency *(days between irr.)*	In/Mo
Jan		0.0	Jul	12	9.7
Feb		0.0	Aug	16	7.7
Mar	>1 Mo.	1.6	Sep	21	5.7
Apr	29	4.2	Oct	>1 Mo.	2.7
May	17	6.9	Nov		0.0
Jun	13	9.3	Dec		0.0

Crop: **Total Irrigation Needed:** ac in/ac

Month	Est. Frequency *(days between irr.)*	In/Mo	Month	Est. Frequency *(days between irr.)*	In/Mo
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** (days between irr.)	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Farm Irrigation Rating System (FIRS, irrigation system efficiency)

Type of System:	**Flood, controlled**		System Efficiency (%):	**80%**
System Capacity (Ditch/Pump/Well):		**6000**	GPM	
Crop Grown	**Days of Operation for CU by crop (days)**	**Est. System Applicalton Rate (in/day)**	**Needed Application Rate (in/day)**	**System Evaluation**
Alfalfa, hay, southern; Las Cruces	10	4.92	0.32	System meets CU for crop

Operation and Maintenance Requirements

1. Irrigate when the soils reaches the MAD level, determined by soil moisture monitoring. Use one of the following methods to monitor soil moisture; the feel method, tensiometers, electrical resistance blocks, or moisture probes. Drip or Center Pivots can be irrigated on an as needed basis to meet daily CU.

2. Test irrigation water for Nitrate and Salts (Total Desolved Solids/Electrical Conductivity)

3. Do not exceed the net water to replace listed above when irrigating, unless salts are being managed.

4. Do not apply water at rates that cause runoff or erosion.

5. Monitor the soil to maintain: pH, permeability, salinity, and structure.

6. Application of pond effluent shall not exceed the crop needs (water and nutrient), and will not exceed the water holding capacity listed above.

7. Consider using crops such as sorghum, cotton, or winter wheat when water supplies are short.

8. Avoid traffic on wet soils to minimize soil compaction.

Additional Requirements

Job Approval and Completion

Client: [illegible] **Date:** 11-23-2009

Conservationist: [illegible] **Date:** 11-23-09

Completed by: [illegible] **Date:**

NRCS - WEQ CALCULATIONS

Version 9.03 11-19-2007

Producer: Dominguez Dairy #1 | Planner: Chet Wyant | Location: Mesquite, NM | Tract: 1579 | Field: DM-5

Crop Rotation: Alfalfa | Climate Data Station: NM, LAS CRUCES | Site "C" Value: 150

Tillage Direction (NS/EW): NS | Length/width ratio: 1 | Field Direction (NS/EW): NS | Field Width (Ft.): 1412

Irrigation (Y or N): Y | Soil "I": 86 | Wind Erodibility Group: 4L (1-7) | TWF: 2 (see instr.)

Sum Period Erosion (t/ac): 0.0 | No. Yrs in Rotation: 1.0 | Av. Annual Wind Erosion: (t/ac/yr)

Calculations and Output

Mgt. Periods Dates		Irr.	Soil	Ridge Roughness				Random Roughness	Unsheltered Distance				SGe	Erosion			
Begin	End	No. of (#)	"I" (t/ac)	Dev. (deg)	Ht. (in.)	Sp. (in.)	"Krd" (factor)	"Krr" (factor)	Dev. (deg)	Prep. (factor)	WED (factor)	"L" (ft)	"V" (lbs/ac)	"E" (t/ac)	EWE (%)	"IF" (%)	Loss (t/ac)
11/1	01/02/13	0	56	0.0	0	0	1.00	0.99	0.0	4.0	1.000	1412	0	62.7	12.6	1.00	0.08
1/2	01/17/13	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			2.0	1.00	
1/17	02/01/13	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			2.0	1.00	
2/1	02/16/13	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.7	0.87	
2/16	03/03/13	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.7	1.00	
3/3	03/18/13	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.9	0.87	
3/18	04/02/13	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			12.6	1.00	
4/2	04/17/13	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			8.8	0.87	
4/17	05/02/13	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			8.6	0.87	
5/2	05/17/13	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			5.3	1.00	
5/17	06/06/13	2	56	0.0	0	0	1.00	0.94	0.0	4.0	1.000	1412			5.6	0.80	
6/6	06/21/13	0	56	22.5	0	0	1.00	0.94	22.5	1.2	1.030	1454			0.8	1.00	
6/21	07/11/13	2	56	22.5	0	0	1.00	0.94	22.5	1.2	1.030	1454			0.7	0.80	
7/11	07/26/13	0	56	67.5	0	0	1.00	0.94	67.5	1.1	1.030	1454			0.2	1.00	
7/26	08/15/13	2	56	67.5	0	0	1.00	0.94	67.5	1.1	1.030	1454			0.2	0.80	
8/15	08/30/13	0	56	90.0	0	0	1.00	0.94	90.0	2.8	1.010	1426			0.1	1.00	
8/30	09/19/13	2	56	90.0	0	0	1.00	0.94	90.0	2.8	1.010	1426			0.6	0.80	
9/19	11/01/13	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.00	1412			1.7	0.95	
		0															
		0															
		0															
		0															
		0															
		0															
		0															
		0															

NRCS

RUSLE2 Worksheet Erosion Calculation Record

Info: Alfalfa Nov'12 to Oct'13

Tract #: 1579
Owner name: DOMINGUEZ DAIRY
Field name: DM-5

Location: New Mexico\DonaAna County\NM_Dona Ana R 9
Soil: Hg HARKEY LOAM\HARKEY loam 85%
Slope length (horiz): 100.0 ft
Avg. slope steepness: 0.50 %
T value: 5.0 t/ac/yr

Alternatives:

Management	*Contouring*	*Strips / barriers*	*Diversion/terrace, sediment basin*	*Cons. plan. soil loss, t/ac/yr*	*Description*
b.Mullti-year Rotation Templates\Alfalfa-Small Grain, 5yr, 5cuts CMZ 23	b. absolute row grade 0.5 percent	(none)	(none)	0.042	

Year 5

Dairy Annual Nutrient Manager

Date	11/30/2009	NM STATE UNIVERSITY Dairy Extension Program	Victor E. Cabrera Dairy Specialist dairy.nmsu.edu: Tools vcabrera@nmsu.edu	PLANNED		[redacted]	
Dairy	Dominguez Dairy						
Crop_Year	2013-2014				N Loss	0%	
Field_ID	DM-5				Goal/Real	Nutrient Needed	
Area (ac)	52						
	Month - Month	Crop		Unit	Yield	N	P
1st Crop	Nov'13-Oct'14	Alfalfa, for hay		t/ac	12	31446	2946
2nd Crop						0	0
						0	0
Total		Nutrient Needed				31,446	2,946
						Soil Analyses	
Texture by Feel		Sandy Clay				N	P
Nutrient Available in Soil				lb/ac		142	97.38
		Nutrient Still Needed				24,062	-2,117
						Effluent Analyses	
						N	P
Effluent Manure Application		NM Dairy Ponds Net from J.S. 590 L&MLA		ac-in	0.712	926	420
				ac-in		0	0
				ac-in		0	0
		Nutrient Still Needed				23,136	-2,538
						Manure Analyses	
						N	P
Dry Manure Application				t/ac		0	0
				t/ac		0	0
				t/ac		0	0
		Nutrient Still Needed				23,136	-2,538
						Fertilizer Content	
						N	P
Chemical Fertilizers Applied				lb/ac		0	0
				lb/ac		0	0
				lb/ac		0	0
		Annual Nutrient Balance				**23,136**	**-2,538**

590 Nutrient Mgt. Jobsheet for Organic and Manure Land Application

Client Name:	Dominguez Dairy	Acres:	52	Date:	11/2/2009	Field ID:	DM-5

Application information (enter the units that will be or has been applied to the field):					
Crop Rotation: Alfalfa				Needed for field (acin):	37.024
Liquid Applied:	0.712	Acin/ac		(gal):	1,005,202
Solids Applied:		ton/ac		Needed for field (tons):	
Liquid Loads Applied:		1000gal/ac		Loads needed for field:	

Nutrient Content of Organic Material

Solid-Lab Report	% Moisture	TKN (%) (dry)	NH_4-N (ppm) (dry)	P_2O_5 (%) (dry)	K_2O (%) (dry)
Fill in Lab data:					

Solid Book Values (select even if test values are used)	% Moisture		TKN (lbs/wet ton)		NH_4-N (lbs/ton)		P_2O_5 (lbs/wet ton)		K_2O (lbs/wet ton)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
Beef (DM)		0		0		0.0		0		0

Liquid-Lab Report	NH_3-N (mg/L)	TKN (mg/L)	NO_3-N (mg/L)	Tot-PO_4 (mg/L)	K (mg/L)
Fill in Lab data:		205	0.17		

Liquid	% Moisture		TKN (lbs/acin)		NH_4-N (lbs/acin)		P_2O_5 (lbs/acin)		K_2O (lbs/acin)	
	Book	Test	Book	Test	Book	Test	Book	Test	Book	Test
NM Dairy Ponds (99-99.4% liq.)	99		0	46	0	26	35	0	256	0
			TKN (lbs/1000gal)		NH_4-N (lbs/1000gal)		P_2O_5 (lbs/1000gal)		K_2O (lbs/1000gal)	
			Book	Test	Book	Test	Book	Test	Book	Test
			0.0			0.0		0.0		0.0

N Volatilization

Solid (type of application)	Type of Climate	Percent Remaining	NH_4-N Remaining
Broadcast-incorporated in 4 days	Warm Dry	60 %	0 (lbs/ton) NH4-N
Liquid (type of application)	**Type of Climate**	**Percent Remaining**	
Surface Irr w/o incorp & w/crop canopy	Warm Dry	80 %	20.8 (lbs/acin) NH4-N 0.0 (lbs/1000gal) NH4-N

Mineralization of N, P, & K

Manure Source	Percent Nutrient Available the 1st Year			
	Organic N	P	K	
Beef & Dairy Solid w/o bedding	35 %	75 %	80 %	Solid Source
Lagoon or diluted Pond	40 %	75 %	80 %	Liquid Source

Solid	Organic N (lbs/ton)	P_2O_5 (lbs/ton)	K_2O (lbs/ton)
	0	0	0
Liquid	**Organic N (lbs/acin)**	**P_2O_5 (lbs/acin)**	**K_2O (lbs/acin)**
	8	26	205
	Organic N (lbs/100gal)	**P_2O_5 (lbs/1000gal)**	**K_2O (lbs/1000gal)**
	0.00	0.0	0.0

Denitrification of N

Organic Matter Content (%)	Soil Drainage Class (See Survey Information)	Percent Remaining (%)
<2	Well Drained	88

Summary of Nutrients

Net by Form as applied	lbs/1000gal	lbs/ac in	lbs/ton
N	0.0	25	0
P_2O_5	0.0	26	0
K_2O	0.0	205	0

Total Nutrients Applied (net to the field)	All Forms N (lbs/ac)	P_2O_5 (lbs/ac)	K_2O (lbs/ac)
	18.1	18.7	145.7

USDA Natural Resources Conservation Service
PLANTS

Nutrient Report Summary for the Crop(s) Selected

The table below summarizes the nutrients removed for the crops selected in the previous page. Crop nutrient information for individual crops follow the summary table.

Return to the crop list and make a new selection of crop(s)

Alfalfa, for hay
Forage
Medicago sativa
Plant part harvested: Aboveground biomass
Crop yield unit: ton

Nutrients in harvested part (lb/ton) at 9.65% moisture percentage.			**Nutrients removed in harvested part (lb/acre) at** 12 ton yield level.		
Nitrogen	**Phosphorus**	**Potassium**	**Nitrogen**	**Phosphorus**	**Potassium**
50.3935	4.7217	38.2939	604.7224	56.6599	459.5271

Nutrients removed in harvested part (lb/ton) at 12 ton yield level and 52 acres.

Nitrogen	**Phosphorus**	**Potassium**
31445.5665	2946.3165	23895.4086

Element-Fertilizer Equivalents | Average NPK Percentages

«Contents»

Intellectual Property Statement / How to Cite the PLANTS Database

Disclaimers

USDA non-discrimination policy

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Time Generated: Mon 4:38 PM - 11/30/2009

PHOSPHORUS INDEX WORKSHEET for New Mexico

Client Name:	Dominguez Dairy	Field(s):	DM-5	Date:	Nov'13
Planner:	Chet Wyant	Location:	DonaAna	Crop:	Alfalfa
Soil Permeablity (in/hr):	0.5	Slope (%):	0.5	Planned/Exist.:	planned

Site Characteristic	Place an X in the approprate box for each of the Site Characteristic listed below.					Sub Total
Soil Test P Level	Very Low <8 ppm	Low 8-15 ppm	Moderate >15-23 ppm	High >23-30 ppm	Very High >30 ppm	
				X		4
Phosphorus (P_2O_5) Application Rate	None Applied	<30 lbs/ac P_2O_5	30-90 lbs/ac P_2O_5	>90-150 lbs/ac P_2O_5	>150 lbs/ac P_2O_5	
		X				1
Organic Phosphorus Source Application Method	None Applied	Injected Deeper than 2 inches	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
				X		4
Phosphorus Fertilizer Application Method	None Applied	Placed with Planter Deeper than 2 in.	Incorporated Immediately before Planting	Incorp. >3 Mo. Before Planting or Surface Applied <3 Mo. before Planting	Surface Applied >3 Months Before Planting	
	X					0
Proximity of Nearest Field Edge to Named Stream or Lake	Very Low >1000 feet	Low >500-1000 feet	Medium >200-500 feet	High 30-200 feet	Very High <30 feet	
	X					0
Soil Erosion (wind & water)	Very Low <1 t/ac	Low 1-3 t/ac	Medium >3-5 t/ac	High >5-15 t/ac	Very High >15 t/ac	
	X					0
Runoff Class (Runoff Class Table 2)	Very Low or Negligible	Low	Medium	High	Very High	
		x				1.5
Irrigation Erosion (See QS note)	Not Irrigated or No Furrow Irrigation	Tailwater Recovery or QS<6 for very erodible soils or QS<10 for resistant soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QS>6 for very erodible soils	
			X			3
Grazing Management	Not Grazed	Graze Crop Residues	Pasture <30% Dry Matter as Supplemental Feed	Pasture 30 to 80% Dry Matter as Supplemental Feed	Pasture 80 to 100% Dry Matter as Supplemental Feed	
	X					0
Vegetative Buffer	> 100 ft wide	>65-100 ft wide	20-65 feet wide	< 20 feet wide	No Buffer	
			x			3

P Hazard Class:	Low	Total Index Points:	16.5
Phosphorus Application Classification:		N Based	

Notes:

This evaluation has a Low P hazard class and the nutrient application can be based on N.

Comments: Soil Erosion = (WEQ 11'13 to 10/14) 0.08 t/ac + (RUSLE 2) 0.042 t/ac = 0.12 t/ac

NRCS Natural Resources Conservation Service

Irrigation Water Management

Conservation Practice Job Sheet 449

Natural Resources Conservation Service (NRCS) Jan, 2006

Client: Dominguez Dairy **Tract:** 1579
Planner: Chet Wyant **Field(s) No.:** DM-5
Current Land Use: Alfalfa **Total Acres:** 52
Date: 11/20/2009 **Date to apply:** Nov'13

See the Conservation Plan map for the location of the field(s) to applying IWM.

Purposes (check all that apply)

- ☐ Manage soil moisture to improve crops
- ☑ Optimize use of water
- ☑ Minimize Irrigation Erosion
- ☐ Decrease non-point source pollution
- ☐ Manage salt in the root zone
- ☐ Manage air, soil, or plant microclimate

Conditions where practice applies

This practice is applicable to all irrigated lands. An irrigation system adapted for the site conditions is available and capable of applying water to meet the intended purpose(s).

Soil (Series, Texture, and Map Unit) *Select the soil to manage for:*

Critical soil to manage: **Harkey L; 33-41** Intake Family (in/hr): **0.50**

Soil Interpretations for Irrigaiton

Crop Name	Rooting Depth	Moisture Replacement Depth (ft)	Water Holding Capacity (in)	Mgt Allowed Depletion (MAD) (%)	Net Water to Replace (in)	Time Needed to Infiltrate (hrs)
Alfalfa, hay, southern; Las Cruces	Deep	4.0	8.0	50%	4.0	5.4

Crop Consumptive Use (CU) Information *(inches/month needed)*

Crop:	Alfalfa, hay, southern; Las Cruces		Total Irrigation Needed:	47.7 ac in/ac	
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul	12	9.7
Feb		0.0	Aug	16	7.7
Mar	>1 Mo.	1.6	Sep	21	5.7
Apr	29	4.2	Oct	>1 Mo.	2.7
May	17	6.9	Nov		0.0
Jun	13	9.3	Dec		0.0

Crop:			Total Irrigation Needed:	ac in/ac	
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** (days between irr.)	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Crop:			Total Irrigation Needed:		ac in/ac
Month	**Est. Frequency** *(days between irr.)*	**In/Mo**	**Month**	**Est. Frequency** *(days between irr.)*	**In/Mo**
Jan		0.0	Jul		0.0
Feb		0.0	Aug		0.0
Mar		0.0	Sep		0.0
Apr		0.0	Oct		0.0
May		0.0	Nov		0.0
Jun		0.0	Dec		0.0

Farm Irrigation Rating System (FIRS, irrigation system efficiency)

Type of System:	**Flood, controlled**		System Efficiency (%):	**80%**
System Capacity (Ditch/Pump/Well):		**6000**	GPM	
Crop Grown	**Days of Operation for CU by crop (days)**	**Est. System Applicaiton Rate (in/day)**	**Needed Application Rate (in/day)**	**System Evaluation**
Alfalfa, hay, southern; Las Cruces	10	4.92	0.32	System meets CU for crop

Operation and Maintenance Requirements

1. Irrigate when the soils reaches the MAD level, determined by soil moisture monitoring. Use one of the following methods to monitor soil moisture; the feel method, tensiometers, electrical resistance blocks, or moisture probes. Drip or Center Pivots can be irrigated on an as needed basis to meet daily CU.

2. Test irrigation water for Nitrate and Salts (Total Desolved Solids/Electrical Conductivity)

3. Do not exceed the net water to replace listed above when irrigating, unless salts are being managed.

4. Do not apply water at rates that cause runoff or erosion.

5. Monitor the soil to maintain: pH, permeability, salinity, and structure.

6. Application of pond effluent shall not exceed the crop needs (water and nutrient), and will not exceed the water holding capacity listed above.

7. Consider using crops such as sorghum, cotton, or winter wheat when water supplies are short.

8. Avoid traffic on wet soils to minimize soil compaction.

Additional Requirements

Job Approval and Completion

Client: Isaac Domingues **Date:** 11-23-2009

Conservationist: [signature] **Date:** 11-27-09

Completed by: **Date:**

NRCS - WEQ CALCULATIONS

Version 9 03 11-19-2007

Producer:	Dominguez Dairy #1	Planner:	Chet Wyant	Location:	Mesquite, NM	Tract:	1579	Field:	DM-4
Crop Rotation:	Alfalfa	Climate Data Station:	NM, LAS CRUCES			Site "C" Value:			150
Tillage Direction (NS/EW):	NS	Length/width ratio:	2	Field Direction (NS/EW):	NS	Field Width (Ft.):			1233
Irrigation (Y or N):	Y	Soil "I":	86	Wind Erodibility Group:	3 (1-7)	TWF:	1	(see instr.)	
Sum Period Erosion (t/ac):	0.0	No. Yrs in Rotation:	1.0	Av. Annual Wind Erosion:		(t/ac/yr)			

Calculations and Output

Mgt Periods		Irr.	Soil	Ridge Roughness				Random Roughness	Unsheltered Distance				SGe	Erosion			
Dates		No. of	"I"	Dev.	Ht.	Sp.	"Krd"	"Krr"	Dev.	Prep.	WED	"L"	"V"	"E"	EWE	"IF"	Loss
Begin	End	(#)	(t/ac)	(deg)	(in.)	(in.)	(factor)	(factor)	(deg)	(factor)	(factor)	(ft)	(lbs/ac)	(t/ac)	(%)	(%)	(t/ac)
11/1	01/02/14	0	56	0.0	0	0	1.00	0.99	0.0	4.0	1.010	1245	0	62.7	12.6	1.00	0.08
1/2	01/17/14	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.010	1245			2.0	1.00	
1/17	02/01/14	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.010	1245			2.0	1.00	
2/1	02/16/14	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.010	1245			12.7	0.93	
2/16	03/03/14	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.010	1245			12.7	1.00	
3/3	03/18/14	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.010	1245			12.9	0.93	
3/18	04/02/14	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.010	1245			12.6	1.00	
4/2	04/17/14	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.010	1245			8.8	0.93	
4/17	05/02/14	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.010	1245			8.6	0.93	
5/2	05/17/14	0	56	0.0	0	0	1.00	0.94	0.0	4.0	1.010	1245			5.3	1.00	
5/17	06/06/14	2	56	0.0	0	0	1.00	0.94	0.0	4.0	1.010	1245			5.6	0.90	
6/6	06/21/14	0	56	22.5	0	0	1.00	0.94	22.5	1.2	1.370	1689			0.8	1.00	
6/21	07/11/14	2	56	22.5	0	0	1.00	0.94	22.5	1.2	1.370	1689			0.7	0.90	
7/11	07/26/14	0	56	67.5	0	0	1.00	0.94	67.5	1.1	1.475	1819			0.2	1.00	
7/26	08/15/14	2	56	67.5	0	0	1.00	0.94	67.5	1.1	1.475	1819			0.2	0.90	
8/15	08/30/14	0	56	90.0	0	0	1.00	0.94	90.0	2.8	2.000	2466			0.1	1.00	
8/30	09/19/14	2	56	90.0	0	0	1.00	0.94	90.0	2.8	2.000	2466			0.6	0.90	
9/19	11/01/14	1	56	0.0	0	0	1.00	0.94	0.0	4.0	1.01	1245			1.7	0.98	
		0															
		0															
		0															
		0															
		0															
		0															
		0															
		0															

RUSLE2 Worksheet Erosion Calculation Record

Info: Alfalfa Nov'13 to Oct'14

Tract #: 1579
Owner name: DOMINGUEZ DAIRY
Field name: DM-5

Location: New Mexico\DonaAna County\NM_Dona Ana R 9
Soil: Hg HARKEY LOAM\HARKEY loam 85%
Slope length (horiz): 100.0 ft
Avg. slope steepness: 0.50 %
T value: 5.0 t/ac/yr

Alternatives:

Management	*Contouring*	*Strips / barriers*	*Diversion/terrace, sediment basin*	*Cons. plan. soil loss, t/ac/yr*	*Description*
b.Mullti-year Rotation Templates\Alfalfa-Small Grain, 5yr, 5cuts CMZ 23	b. absolute row grade 0.5 percent	(none)	(none)	0.042	

Field DM-5

Years 2 thru 5

Alternate Crop Options