

A photograph of the St. Louis skyline featuring the Gateway Arch in the background. The arch is a large, white, catenary-shaped structure. In the foreground, there are trees and a road. The sky is clear and blue.

Measurement and Data Analysis for Source Attribution of Ambient Particulate Matter Arsenic and Other Air Toxics Metals in St. Louis

*Varun Yadav , JAY TURNER
Washington University in St. Louis*

*Jerry Downs, Terry Rowles
Missouri Department of Natural Resources*

**National Air Monitoring Conference
Nashville, TN
November 5, 2009**

Motivation

The St. Louis Community Air Project (CAP)

- identified six hazardous air pollutants of concern including arsenic, *however...*
- poor data quality for PM_{2.5} arsenic from speciation network data
- PM₁₀ air toxics metals routinely measured at only one site (1-in-6 day)




Objectives

Community Air Toxics Grant from USEPA




- **Phase I**
 - four site network of HiVol PM₁₀ samplers
 - one year at 1-in-3 days
 - hot acid extraction and analysis by ICP-MS
- **Phase II**
 - high(er) time resolution measurements
 - six one-month deployments of CES Xact 620
 - *Xact performance evaluation
(already shown this week)*

Filter Network Monitoring Sites

4 × Missouri (1-in-3), 2 × Illinois (1-in-6)




Intersite Variability



variability (proximate urban sites)
>
variability (distance suburban sites)

Baseline and Excess Contributions to PM₁₀ Arsenic




Baseline = regionally transported?


Excess = locally emitted?

Baseline PM₁₀ Arsenic


conditional probability function (CPF)
(24-hour As, hourly winds)



quantitative transport bias analysis
(QTBA)
(24-hour As, 3-day back trajectories)




Excess PM_{10} Arsenic at Urban Sites



conditional probability function plots (24-hour As, hourly winds)

Baseline and Excess Contributions to PM_{10} Selenium



Cooper Environmental Services (CES) Xact 620

- particle collection on a filter tape
- analysis by x-ray fluorescence (XRF)
- continuous data time series at user-defined intervals
- Missouri DNR instrument optimized for As, Hg, and Pb at remote areas



ELEMENTS THE XACT CAN MEASURE (IN BLUE)

1	1																	18
	H																	He
	1.0079																	4.0026
2	3	4											5	6	7	8	9	10
	Li	Be											B	C	N	O	F	Ne
	6.941	9.0122											10.811	12.011	14.007	15.999	18.998	20.18
3	11	12										13	14	15	16	17	18	
	Na	Mg										Al	Si	P	S	Cl	Ar	
	22.99	24.305										26.982	28.086	30.974	32.066	35.453	39.948	
4	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
	39.098	40.078	44.956	47.88	50.942	51.996	54.938	55.847	58.933	58.693	63.546	65.39	69.723	72.61	74.922	78.96	79.904	83.8
5	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
	85.468	87.62	88.906	91.224	92.906	95.94	(97.91)	101.07	102.91	106.42	107.87	112.41	114.82	118.71	121.76	127.6	126.9	131.29
6	55	56	57	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
	Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
	132.91	137.33	138.91	178.49	180.95	183.84	186.21	190.23	192.22	195.08	196.97	200.59	204.38	207.2	208.98	(209)	(210)	(222)
7	87	88	89	104	105	106	107	108	109	110	111							
	Fr	Ra	Ac	Rf	Ha	Sg	Ns	Hs	Mt	Unn	Unu							
	(223)	(226)	(227)	(261.1)	(262.1)	(263.1)	(262.1)	(265.1)	(266.1)	(268)	(269)							

Lanthanide Series	58	59	60	61	62	63	64	65	66	67	68	69	70	71
	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
	140.12	140.91	144.24	(144.9)	150.36	151.97	157.25	158.93	162.5	164.93	167.26	168.93	173.04	174.97
Actinide Series	90	91	92	93	94	95	96	97	98	99	100	101	102	103
	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
	232.04	231.04	238.03	(237)	(244.1)	(243.1)	(247.1)	(247.1)	(251.1)	(252.1)	(257.1)	(258.1)	(259.1)	(262.1)

-  measured by Xact in this study
-  EPA Air Toxics PM metals

Xact Performance Evaluation







HiVol PM₁₀ / quartz filter,
NATTS digestion protocol
ICP-MS: As, Pb, Se...




LowVol PM₁₀ (FRM) / Teflon filter
XRF: Ca, Fe, K, Mn, Pb, Ti...

Xact vs. LowVol PM₁₀ FRM / XRF




Selenium: Xact vs. Filter-Based Measurements

Xact vs. LowVol filter / lab XRF




Xact vs. HiVol filter / lab ICP-MS




Se: favorable comparison between Xact and PM₁₀ HiVol samples with analysis by ICP-MS

Arsenic – Methods Comparisons

Collocated HiVol Samplers Blair site, 4th Quarter 2008




Xact vs. WUSTL HiVol Dec 2008 / Jan 2009




As: favorable comparison between Xact and PM₁₀ HiVol samples with analysis by ICP-MS.

Optimizing the Sampling Time Interval

- depends on study objectives!
- trade-offs between time resolution and frequency above MDL
- Blair Street (STL)
 - 1-hour: 56% > MDL
 - 2-hour: 86% > MDL




Xact Monitoring Sites




PM₁₀ Lead by Xact (2-hour resolution)

BLAIR




ARNOLD




Dashed line is bearing of Doe Run – Herculaneum lead smelter

PM₁₀ Arsenic by Xact (2-hour resolution)

S. BROADWAY




ARNOLD




PM₁₀ Arsenic by Xact (2-hour resolution)

MARGARETTA




HALL



PM₁₀ Arsenic by Xact (2-hour resolution)


EAST ST. LOUIS




2-hour spike exceeding 2.5 $\mu\text{g}/\text{m}^3$ arsenic!

PM₁₀ Arsenic at Blair by Xact (2-hour resolution)



wind direction vs. concentration



nonparametric wind regression





PM₁₀ Arsenic at Blair for Winds from Northwest



red markers = 8 AM – 6PM weekdays

PM₁₀ Arsenic at Blair for Winds from Northwest

nonparametric wind regression



likely microscale impacts

Summary

- Observed urban-scale spatial gradients in PM_{10} arsenic not captured in the emissions inventory
 - Regional transport from the eastern U.S.
 - Local sources along industrial riverfront
 - Microscale impacts at the Blair Street NATTS site
- CES Xact 620 for automated ambient PM elemental analysis
 - Very promising, based on initial performance evaluation
 - Identify intermittent sources not detected by time-integrated measurements
 - How long to stay at a site? “information content” of plumes, separating local from regional contributions?

The Next Steps

- Collocated Xact measurements (MDNR and CES)
 - Herculaneum, August 2009... **done**
 - Measurement precision, crucial data for receptor modeling
- Receptor modeling of the data sets
- Additional Xact deployments
 - Currently near lead smelter in Herculaneum (source profile)
- Expanded performance evaluation
 - Forty additional low-volume PM₁₀ FRM samples, analysis by XRF and ICP-MS

Acknowledgements

- Missouri Department of Natural Resources
 - Jim Brunnert, Celeste Koon, Will Wetherell
- Cooper Environmental Services
 - John Cooper, Krag Petterson
- U.S. EPA / Office of Research and Development
 - Teri Conner, Gary Norris, Bob Willis
- U.S. EPA / Office of Air Quality Planning and Standards
 - Mike Jones
- U.S. EPA / Region VII
 - Gwen Yoshimura
- Washington University
 - Stephen Feinberg